

**THE GENERATING FUNCTION
FOR THE FIBONACCI SEQUENCE**

Mohammad K. Azarian

University of Evansville

Definition. Let a_0, a_1, a_2, \dots , be a sequence of real numbers.

The function

$$f(x) = a_0 + a_1x + a_2x^2 + \dots = \sum_{i=0}^{\infty} a_i x^i$$

is called the generating function for the given sequence.

Let F_n ($n \geq 1$) represent the general term of the Fibonacci sequence

$$1, 1, 2, 3, 5, 8, 13, \dots .$$

The generating function for this sequence is

$$\sum_{n=1}^{\infty} F_n x^n,$$

and it is well-known that

$$(1) \quad \frac{x}{1 - x - x^2} = \sum_{n=1}^{\infty} F_n x^n .$$

In [1, p. 1] it has been stated that (1) can be verified by long division. But, the method of long division is a long process, especially for large n . The purpose of this note is to verify (1) by the method of generating functions which is quicker, regardless of the value of n .

Now, from the fact that

$$\frac{1}{1-ax} = 1 + ax + a^2x^2 + \cdots + a^nx^n + \cdots,$$

we deduce that the coefficient of x^n in

$$\frac{x}{1-x-x^2} = \frac{1}{\sqrt{5}} \left(\frac{1}{1 - \left(\frac{1+\sqrt{5}}{2}\right)x} - \frac{1}{1 - \left(\frac{1-\sqrt{5}}{2}\right)x} \right)$$

is

$$A_n = \frac{1}{\sqrt{5}} \left(\left(\frac{1+\sqrt{5}}{2} \right)^n - \left(\frac{1-\sqrt{5}}{2} \right)^n \right), \quad n \geq 1.$$

But, if F_n represents the solution to the recurrence relation

$$F_{n+1} = F_n + F_{n-1}, \quad F_1 = F_2 = 1, \quad n \geq 1,$$

for the Fibonacci sequence, then clearly $F_n = A_n$.

Reference

1. M. Bicknell and V. E. Hoggatt, *Fibonacci's Problem Book*, The Fibonacci Association, San Jose State University, San Jose, California, 1974.