

SCHOOL OF ARTIFICIAL INTELLIGENCE

UNIVERSITY OF EDINBURGH

Memorandum: SAI-RM-4

Date:- October, 1973

Subject: Lambda-definability and logical relations

Author: G.D. Plotkin

1. Introduction

In [4] we showed that in every model, D_{∞} , of the λ -calculus as constructed in [8] the strict ordering, \prec , is first-order definable using only application. Here we look at the, perhaps more pertinent, question of definability by pure λ -terms of such lattice-theoretic entities as \perp , \top , \sqcup , \sqcap and γ , the least fixed-point operator.

The main method will be to construct certain, so-called, logical relations which are satisfied by all (constant vectors of) λ -definable elements and yet are not satisfied by the lattice-theoretic entity under discussion. The definition of logical is derived from a corresponding one of M. Gordon for the typed λ -calculus. This in turn generalised the idea of an invariant functional [2]. R. Milne [3] has independently developed analogues of the logical relations for use in equivalence proofs about programming languages.

It is not known whether logical relations also provide sufficient conditions for definability. In the second half of this memorandum we discuss this question for the typed case, obtaining necessary and sufficient conditions by using the more inclusive concept of an I-logical relation.

This memorandum is by no means self-contained. The reader should have some knowledge of both the typed and untyped λ -calculi and be fairly familiar with Scott's models of the untyped λ -calculus.

2. Pure definability in D_∞

A structure $\langle D, K, S, [\cdot] \rangle$ is called a (non-trivial) model of the λ -calculus if K and S have the usual properties and extensionality holds (and $|D| > 1$). Such structures give a denotational semantics for the λ K-calculus which we will use informally, confusing use and mention. Generally we will consider only the models D_∞ , from [8], given by a Park retraction, $\psi_0 = \lambda f: D_1.f(t)$, where t is an isolated element of D_0 . We will often use facts about such models, accompanied by a reference to the proof for the case $t = \perp$. The general proof is always similar. Also needed is the fact that if Y_λ is the paradoxical combinator, $\lambda f(\lambda x f(xx))(\lambda x.f(xx))$, then, in D_∞ , $Y_\lambda[f] = \bigcup_{n \geq 0} f^n[f[t] \ni t \rightarrow t, \perp](f \in D_\infty)$ [5].

A relation $R \subseteq D^\kappa$, (κ an ordinal) on such a structure is logical iff:

$$\forall \vec{f} \in D^\kappa. (R(\vec{f}) \equiv (\forall \vec{x} \in D^\kappa. (R(\vec{x}) \rightarrow R(\vec{f}[\vec{x}])))).$$

Here κ is any ordinal and application of vectors is defined pointwise. An element $x \in D$ satisfies R iff $R(\hat{x})$ is true, where $\hat{x} \in D^\kappa$ is the constant vector such that $(\hat{x})_\lambda = x$ ($\lambda < \kappa$).

An element $x \in D$ is λ -definable if $x=M$, for some closed λ -term M ; it is λ -definable from $X \subseteq D$ iff there is a closed term M and x_1, \dots, x_n in X such that $x=Mx_1, \dots, x_n$.

Theorem 1

1. Any closed λ -term satisfies any logical relation.
2. If x is λ -definable from $X \subseteq D$, and each element in X satisfies the logical relation R , then so does x .

Proof Clearly, if x and y satisfy a logical relation R , so does $x[y]$. So to finish the proof we need only show that K and S satisfy any such relation. Suppose R is logical. To show K satisfies R , assuming $R(\vec{x})$ we must show that $R(K[\vec{x}])$. This, in turn, follows if $R(K[\vec{x}][\vec{y}])$ when $R(\vec{y})$. But this holds as $K[\vec{x}][\vec{y}] = \vec{x}$.

In/

In the same way we see that S satisfies R if $R(\hat{S}[\bar{x}][\bar{y}][\bar{z}])$ when $R(\bar{x})$, $R(\bar{y})$, and $R(\bar{z})$. But then we have successively, by the remark made at the beginning of the proof that $R(\bar{x}[\bar{z}])$, $R(\bar{y}[\bar{z}])$ and $R(x[\bar{z}][\bar{y}][\bar{z}])$, concluding the proof.

Nothing is known about the converse of theorem 1. However it will be very useful for particular cases of undefinability. Here is a way of constructing logical relations $R \subseteq D_{\infty}^2$.

Suppose $R_0 \subseteq D_0^2$. Define $R_n \subseteq D_n^2$ by:

$$\forall f, g \in D_{n+1}. (R_{n+1}(f, g) \equiv \forall x, y \in D_n (R_n(x, y) \rightarrow R_n(fx, gy)))$$

Define $R_{\infty} \subseteq D_{\infty}^2$ by:

$$\forall d, e \in D_{\infty}. (R_{\infty}(d, e) \equiv \forall n R_n(d_n, e_n))$$

Theorem 2. Suppose that $R_0(t, t)$, that $R_0(d, e)$ implies $R_1(\phi_0 d, \phi_0 e)$, for any d, e in D_0 , and that R_0 is closed under unions of increasing sequences. Then:

1. R_{∞} is logical.
2. R_{∞} is closed under increasing sequences.
3. If R_0 is closed under \cup (\cap) so is R_{∞} ; if $R_0(\perp, \perp)$ ($R_0(T, T)$) then $R_{\infty}(\perp, \perp)$ ($R_{\infty}(T, T)$).

The construction also works for any $R_0 \subseteq D^K$ under the corresponding conditions, and the theorem analogous to theorem 2 can be proved; this extension will be assumed.

Lemma 1.1 Suppose that $R_0(t, t)$ and $R_0(d, e)$ implies $R_1(\phi_0 d, \phi_0 e)$ for any d, e in D_0 . Then,

$$\begin{aligned} &\forall n \forall f, g \in D_n (R_n(f, g) \rightarrow R_{n+1}(\phi_n(f), \phi_n(g))) \text{ and} \\ &\forall n \forall f, g \in D_{n+1} (R_{n+1}(f, g) \rightarrow R_n(\phi_n(f), \phi_n(g))) \end{aligned}$$

- 1.2 If R_0 is closed under increasing sequences so is each R_n .

Proof/

Proof 1.1 By induction on n .

For $n=0$, note that if $R_1(f,g)$ then $R_0(ft,gt)$ from the definition of R_1 and the fact that $R_0(t,t)$.

For $n+1$, suppose $R_{n+1}(f,g)$ and suppose $R_{n+1}(f',g')$. By induction hypothesis $R_n(\psi_n f', \psi_n g')$. Therefore $R_n(f(\psi_n f'), g(\psi_n g'))$, and by the induction hypothesis, $R_{n+1}(\phi_n \circ f \circ \psi_n(f'), \phi_n \circ f \circ \psi_n(g'))$, which shows that $R_{n+2}(\phi_{n+1} f, \phi_{n+1} g)$. The other half is similar.

1.2 By induction on n . For $n+1$, let $\langle f^m, g^m \rangle_{m=0}^{\infty}$ be an (infinite) increasing sequence in R_{n+1} and suppose $R_n(x,y)$. Then $\langle f^m x, g^m y \rangle_{m=0}^{\infty}$ is an increasing sequence in R_n and so $\langle (\bigcup_m f^m)x, (\bigcup_m g^m)y \rangle$ is in R_n by induction hypothesis and the complete additivity of application in its first argument. This concludes the proof.

Proof of theorem 2 1 First suppose that $R_{\infty}(f,g)$ and $R_{\infty}(x,y)$. We will show that $R_{\infty}(fx,gy)$.

Now $(fx)_n = \bigcup_{m=n}^{\infty} \psi_{mn}(f_{m+1}x_m)$ and similarly for $(gy)_n$, [7]. Since $R_m(f_{m+1}x_m, g_{m+1}y_m)$ is true for any $m \geq n$, $R_n(\psi_{mn}(f_{m+1}x_m), \psi_{mn}(g_{m+1}y_m))$ follows by $m - n$ applications of lemma 1.1, and then we see that $R_n((fx)_n, (gy)_n)$ by lemma 1.2 and the above formulae for $(fx)_n$ and $(gy)_n$.

Conversely, suppose that whenever $R_{\infty}(x,y)$ then $R_{\infty}(fx,gy)$ and yet for some n , $R_n(f_n, g_n)$ is false. By lemma 1.1 we can assume that $n > 0$, and so for some $\langle x_{n-1}, y_{n-1} \rangle \in R_{n-1}$, $R_{n-1}(f_n x_{n-1}, g_n y_{n-1})$ is false. Let $x = \bigcup_{n-1}^{\infty} x_{n-1}$ and define y similarly. By lemma 1.1, $R_{\infty}(x,y)$ is true and so therefore is $R_{\infty}(fx,gy)$ and, consequently, $R_{n-1}((fx)_{n-1}, (gy)_{n-1})$. But $(fx)_{n-1} = (f_n x_{n-1})$ and similarly for $(gy)_{n-1}$ (cf. the laws of application in [9]) and so $R_{n-1}(f_n x_{n-1}, g_n y_{n-1})$, a contradiction).

2 Suppose $\langle x^m, y^m \rangle_{m=0}^{\infty}$ is an (infinite) increasing sequence in R_{∞} .

Now/

Now, $(\bigcup_{m=0}^{\infty} x^m)_n = \bigcup_{n'=n}^{\infty} \psi_{n'n}(\bigcup_{m=0}^{\infty} (x^m)_n)$ and similarly for the y 's, (cf. [7]).

Then one sees, successively that, $R_n((x^m)_n, (y^m)_n)$ for all m and n , $R_n(\bigcup_{m=0}^{\infty} (x^m)_n, \bigcup_{m=0}^{\infty} (y^m)_n)$, for all n , by lemma 1.2, $R_n(\psi_{n'n}(\bigcup_{m=0}^{\infty} (x^m)_n), \psi_{n'n}(\bigcup_{m=0}^{\infty} (y^m)_n))$ for $n' \geq n$, by lemma 1.1 and finally $R_n(\bigcup_{m=0}^{\infty} (x^m)_n, \bigcup_{m=0}^{\infty} (y^m)_n)$ by lemma 1.2.

3 A straightforward inductive argument shows that if R_0 is closed under \bigcup so is each R_n . Then, clearly, $R_{\infty}(\lambda x \lambda y (x_n \bigcup y_n), \lambda x \lambda y (x_n \bigcup y_n))$ and so R_{∞} is closed under \bigcup as $\bigcup_{n \geq 0} (\lambda x \lambda y (x_n \bigcup y_n))$ expresses \bigcup as an increasing sequence in D_{∞} .

The argument for D_{∞} is similar; it uses the fact that if $f, g: X \rightarrow Y$ where X and Y are continuous lattices then $(f \sqcap g)_x = (fx) \sqcap (gx)$.

If $R_0(\perp, \perp)$ then $R_n(\perp, \perp)$, for any n , by lemma 1.1.

If $R_0(T_0, T_0)$ an easy inductive argument shows that $R_n(T_n, T_n)$ for all n , concluding the proof.

As an example, let $P_0 = \{t\}$. Then by the assumed extension of) theorem 2, P_{∞} is logical and so the 0th component of any closed λ -term is t . Therefore if $T \neq \perp$ neither \perp nor, since $\perp = YI$, Y is λ -definable; this is a result of Park [6].

The next example establishes all the definabilities among $\perp, T, \bigcup, \sqcap$ and Y for all possible values of t .

- Theorem 3
- 1.1 If $t = \perp$, $Y = Y_{\lambda}$ and $\perp = Y_{\lambda} I$.
 - .2 In general, $\perp = YI$ and $Y = \lambda f (Y_{\lambda} (\lambda g \lambda x f(g \perp))) I$.
 - .3 If $t = T_0$, then $T = Y_{\lambda} K$.
 - .4 If $t = T_0$ and $D_0 = \mathcal{D} = \{\perp, T\}$ then $\sqcap = Y_{\lambda} (\lambda g \lambda x \lambda y \lambda z g(xz)(yz))$.

2. The only definabilities among \perp, T, U, Π and Y are those implied by 1.

Proof 1.1 This result is known - see [9].

.2. $\perp = YI$ is obvious.

Suppose $f \in D_\infty$ and let $T = \lambda g \lambda x f(g \perp)$. As $T t = T \perp = \lambda x f \perp$, we see that $Y \lambda T = \bigcup_{n \geq 0} T^n (\lambda x f \perp)$. By induction on n , $T^n (\lambda x f \perp) = \lambda x f^{n+1}(\perp)$, giving $Y \lambda T = \lambda x (Yf)$, and the result follows.

.3. As $Kt \supseteq t$, $Y \lambda K = \bigcup_{n \geq 0} K^n t$. As $K^n t \supseteq T_n$ ($t = T_0$, here), for all n , $Y \lambda K = T$.

.4. Let $T = \lambda g \lambda x \lambda y \lambda z g(xz)(yz)$. Since, in this D_∞ , $x \supseteq t$ iff $xt \supseteq t$, one sees that $T t \supseteq t$. Now, $t = \lambda x \lambda y x_0 \Pi y_0$ is true in this lattice and then $\Pi = Y \lambda T$ follows by the usual inductive argument.

2. As \perp and Y are interdefinable, only definabilities among \perp, T, U and Π need be considered.

We must show that if $t \neq \perp$, then \perp is not λ -definable from $\{T, U, \Pi\}$; that if $t \neq T_0$, T is not λ -definable from $\{\perp, U, \Pi\}$; that U is not λ -definable from $\{\perp, T, \Pi\}$ in all cases; and that if $t \neq T_0$ or $D_0 \neq \emptyset$ then Π is not λ -definable from $\{\perp, T, U\}$.

To show that \perp is not λ -definable from $\{T, U, \Pi\}$, when $t \neq \perp$ let $R_0 = \{t, T_0\}$. The conditions of theorem 2 are easily checked and so P_∞ is logical. It also follows from theorem 2 that U, Π and T satisfy R_∞ . Clearly \perp does not. The conclusion then follows from theorem 1.2.

In the rest of the proof we shall first display an appropriate R_0 and leave the (admittedly tedious) details to the reader.

To show that if $t \neq T_0$, T is not λ -definable from $\{\perp, \Pi, U\}$ take $R_0 = \{\perp, t\}$.

To/

To show that, in all cases, \perp is not λ -definable from $\{\perp, T, \Pi\}$, take $R_0 = \{\langle T_0, T_0, T_0 \rangle, \langle \perp, T_0, \perp \rangle, \langle T_0, \perp, \perp \rangle, \langle t, t, t \rangle, \langle \perp, t, \perp \rangle, \langle t, \perp, \perp \rangle, \langle \perp, \perp, \perp \rangle\}$. Note that $R_\infty(\perp, T_0, \perp)$ and $R_\infty(T_0, \perp, \perp)$ but not $R_\infty(T_0, T_0, \perp)$.


To show that if $t \neq T_0$ then Π is not λ -definable from $\{\perp, T, \perp\}$, take $R_0 = \{\langle x, y, z \rangle \mid \{x, y\} \subseteq \{t, T_0\}, x \neq t \text{ or } y \neq t, z \in \{\perp, t, T_0\}\} \cup \{\langle \perp, \perp, \perp \rangle\}$. Note that $R_\infty(t, T_0, \perp)$ and $R_\infty(T_0, t, \perp)$ but not $R_\infty(t, t, \perp)$.

To show that if $t = T_0$ and $D_0 \neq \emptyset$ then Π is not λ -definable from $\{\perp, T, \perp\}$, choose $u \in D_0$ distinct from \perp and t and take $R_0 = \{\langle t, u, u \rangle, \langle u, t, u \rangle, \langle t, t, u \rangle, \langle t, t, t \rangle\} \cup \{\langle x, y, \perp \rangle \mid x, y \in \{\perp, u, t\}\}$. Note that $R_\infty(t, u, u)$ and $R_\infty(u, t, u)$ but not $R_\infty(u, u, u)$. This concludes the proof.

It is interesting to note that when $t = T_0$ and $D_0 = \emptyset$ then a normal term can even equal an unsolvable term, for example, $I = Y_\lambda (\lambda f \lambda x \lambda y f(xy))$ (cf. $I = J$, when $t = \perp$ [9]).

Our method of constructing logical relations is by no means all-powerful. For example, we believe that if $t \neq T_0$ or $D_0 \neq \emptyset$ then ψ_0 is not λ -definable. Clearly, for the R_∞ 's constructed so far, if $R_\infty(\bar{x})$ then $R_0(\lambda \alpha . ((\bar{x})_\alpha)_0)$ and so $R_\infty(\lambda \alpha . ((\bar{x})_\alpha)_0)$. Therefore $R_\infty(\hat{\psi}_0)$. On the other hand, suppose ψ_0 were λ -definable by a closed term M when $t = \perp$. Clearly (see [10]) M is not unsolvable, as $\psi_0 \neq \perp$. So there are closed terms $M_1 \dots M_k$ ($k \geq 0$) such that $M M_1 \dots M_k = I$, but as mentioned above the 0th component of M_1 must be \perp and so either $\perp = I$ or $\psi_0 = I$, a contradiction. Perhaps an extension of Wadsworth's methods to the other D_∞ 's would sort this out.

The last example concerns interdefinabilities among the members of $\{tt, ff, T, \perp, \Pi, \supset\}$ in T_∞ [9] which is gotten by taking $t = \perp$ and D_0 to be the truth-value lattice displayed in figure 1.

fig. 1

The conditional, \supset , is in $T_\infty^3 \rightarrow T_\infty$, and is regarded as being in T_∞ , in the usual way. It is defined by:

$$(z \supset x, y) = \begin{cases} x \sqcup y & (\text{if } z=T) \\ x & (\text{if } tt \in z \neq T) \\ y & (\text{if } ff \in z \neq T) \\ \perp & (\text{otherwise}) \end{cases}$$

It is known that T is λ -definable from $\{tt, ff, \sqcup\}$, \sqcup is λ -definable from $\{\supset, T\}$ and \sqcap can be defined from $\{tt, ff, \sqcup, \supset\}$. We will show that there are no more λ -definabilities of tt, ff, T, \sqcup or \sqcap other than those implied by the above ones; the situation for \sqcap has only been partly clarified.

First, \supset is not definable from $\{tt, ff, \sqcup, \sqcap, T\}$. Take $R_0 = \{\langle \perp, \perp \rangle, \langle tt, ff \rangle, \langle ff, ff \rangle, \langle tt, tt \rangle, \langle tt, T \rangle, \langle T, T \rangle, \langle \perp, ff \rangle, \langle tt, \perp \rangle, \langle T, ff \rangle\}$ and note that $R_0(\supset tt ff tt, \supset ff ff tt)$ is false. Here and later theorem 2 is used implicitly.

tt is not definable from $\{ff, \sqcup, \supset, T\}$; take $R_0 = \{\perp, ff, T\}$.

ff is not definable from $\{tt, \sqcup, \supset, T, \sqcap\}$; take $R_0 = \{\perp, tt, T\}$.

T is not definable from any one of $\{tt, ff, \supset, \sqcap\}$, $\{tt, \sqcup, \supset, \sqcap\}$ or $\{ff, \sqcup, \supset, \sqcap\}$; take $R_0 = \{\perp, tt, ff\}$, $\{\perp, tt\}$ or $\{\perp, ff\}$ respectively.

\sqcup is not definable from either one of $\{tt, ff, T, \sqcap\}$ or $\{tt, ff, \sqcap, \supset\}$; take/

take $R_0 = \{\langle tt, tt \rangle, \langle ff, ff \rangle, \langle \perp, \perp \rangle, \langle T, T \rangle, \langle tt, ff \rangle, \langle tt, \perp \rangle, \langle \perp, ff \rangle\}$
and note that $R_0(\bigcup tt\ tt, \bigcup tt\ ff)$ is false in the first case and
take $R_0 = \{\perp, tt, ff\}$ in the second case.

In the case of Π , we would like to show that Π is not definable from any of the sets $\{tt, ff, \bigcup, T\}$, $\{tt, ff, \supset\}$, $\{ff, \bigcup, \supset, T\}$ or $\{tt, \bigcup, \supset, T\}$. For the first of these take $R_0 = \{\langle tt, tt \rangle, \langle ff, ff \rangle, \langle T, T \rangle, \langle \perp, \perp \rangle, \langle tt, ff \rangle, \langle tt, T \rangle, \langle T, ff \rangle\}$ and note that $R_0(\Pi\ ff\ tt, \Pi\ ff\ ff)$ is false.

The trouble with the others is that if $R_\infty(\supset, \supset)$ then $R_\infty(\Pi, \Pi)$, for the R's considered here. For if $x, y \in T_0$ then $(x \supset (y \supset y, \perp), (y \supset \perp, y)) = x \Pi y$, and so one can define from \supset terms M_n ($n \geq 0$) such that $M_n xy = x \Pi y$ if x and y are in T_n . Therefore if $R_\infty(\supset, \supset)$ then $R_\infty(\lambda x \lambda y x_n \Pi y_n)$ for any n and so $R_\infty(\Pi, \Pi)$.

On the other hand, Π is, in fact, not λ -definable from \supset . For suppose $\Pi = M \supset$ for some closed λ -term M . If M is unsolvable then $\Pi = \perp$, a contradiction; therefore M has the form $\lambda x_1 \dots \lambda x_n . x_j M_1 \dots M_k$ where $n > 0$ and $1 \leq j \leq n$. One can assume that $j \leq 3$ since one can always apply the identity $\Pi = M \supset (M \supset)(M \supset)$. If $j=1$, then $\Pi x_2 \dots x_n = \supset M_1' \dots M_k'$ where $M_r' = (\lambda x_1 M_r) \supset (1 \leq r \leq k)$. Taking the $x_1 = I$ and $P_0 = \{\perp\}$ we see that $(M_1')_0 = \perp$ and so $\Pi I \dots I = \perp$, a contradiction. If $j \neq 1$ then since $\Pi xy = \Pi yx$, for any x, y in T_∞ , we have $x M_1' \dots M_k' \supset xy = M \supset yx = y M_1'' \dots M_k''$ where the M_i' and M_i'' are λ -definable from x, y, \supset and \perp . Since x, y are arbitrary members of T_∞ , this is a contradiction.

Perhaps an extension of Wadsworth's ideas to LAMBDA [9], would settle these questions.

3. λ -definability in the full type hierarchy

For the sake of clarity, we will be a little more formal than in the last section.

The set of type symbols is the least set containing ϵ and containing $(\sigma \rightarrow \tau)$ whenever it contains σ and τ ; σ and τ are metavariables, possibly suffixed, ranging over type symbols and $(\sigma_1, \dots, \sigma_n, \tau)$ abbreviates $(\sigma_1 \rightarrow (\sigma_2 \rightarrow \dots (\sigma_n \rightarrow \tau) \dots))$ ($n \geq 0$).

The language of the typed λ -calculus has denumerably many variables α_i^τ ($i \geq 0$) of each type τ . We will use α and β , with or without various decorations as metavariables over variables. The language has a set of terms which is given by:

1. α_i^τ is a term of type τ , ($i \geq 0$),
2. if M and N are terms of type $(\sigma \rightarrow \tau)$ and σ respectively then (MN) is a term of type τ ,
3. if M is a term of type τ then $(\lambda \alpha_i^\sigma.M)$ is a term of type $(\sigma \rightarrow \tau)$, ($i \geq 0$);

M and N, possibly with suffices, will be used as metavariables over terms. The reader is assumed to know what a β, η -normal form of a term is and the elementary properties of normal forms; $M \approx N$ means that M and N have identical β, η -normal forms. By the Church-Rosser theorem this is an equivalence relation. Suppose that $K_{\sigma\tau} = (\lambda \alpha_0^\sigma (\lambda \alpha_1^\tau \alpha_0^\sigma))$ and $S_{\sigma_1\sigma_2\sigma_3} = (\lambda \alpha_0^{\sigma_1, \sigma_2, \sigma_3} (\lambda \alpha_1^{\sigma_1, \sigma_2} (\lambda \alpha_2^{\sigma_1} ((\alpha_0^{\sigma_1, \sigma_2, \sigma_3})_1 \alpha_0^{\sigma_1} (\alpha_0^{\sigma_1, \sigma_2})_2 \alpha_0^{\sigma_1, \sigma_3}))))$. Then, as is well-known, the K's and S's generate all closed terms under application, to within \approx . The type subscripts in K and S will often be omitted, as will be as many other type symbols as is convenient; the resulting propositions are to be understood as being asserted for every consistent way of putting the symbols back in.

Our language also has a semantics based on the full type hierarchy $\{D_\sigma\}$ defined by:

D/

$D(\sigma \rightarrow \tau) = (D\sigma \rightarrow D\tau)$ (the set of all functions from $D\sigma$ to $D\tau$), where Γ_i is some given set.

The semantics is a function $\llbracket \cdot \rrbracket : \text{Terms} \rightarrow (\text{Env} \rightarrow \bigcup_{\sigma} D\sigma)$ where Env, the set of environments, is the set of type respecting functions from the set of variables to $\bigcup D\tau$, and is ranged over by ρ . Then, $\llbracket \cdot \rrbracket$ is the unique function of that type such that:

1. $\llbracket \alpha_i^{\tau} \rrbracket(\rho) = \rho(\alpha_i^{\tau}) \quad (i \geq 0)$
2. $\llbracket (MN) \rrbracket(\rho) = \llbracket M \rrbracket(\rho)(\llbracket N \rrbracket(\rho))$
3. $\llbracket (\lambda \alpha_i^{\tau}. M) \rrbracket(\rho)(x) = \llbracket M \rrbracket(\rho[x/\alpha_i^{\tau}]) \quad (i \geq 0, x \in D\tau)$

where $\rho[x/\alpha_i^{\tau}]$ is the environment ρ' such that

$$\rho'(\alpha_{i'}^{\tau'}) = \begin{cases} x & (\alpha_{i'}^{\tau'} = \alpha_i^{\tau}) \\ \rho(\alpha_{i'}^{\tau'}) & (\text{otherwise}). \end{cases}$$

Note that if M has type σ , $\llbracket M \rrbracket(\rho) \in D\sigma$. If M is closed then $\llbracket M \rrbracket(\rho) = \llbracket M \rrbracket(\rho')$ for any ρ and ρ' - so we often drop the reference to ρ for closed M . If $M \approx M'$ then $\llbracket M \rrbracket(\rho) = \llbracket M' \rrbracket(\rho)$ for any ρ ; we will give a converse later.

Suppose $\pi_{\iota} \in D(\iota \rightarrow \iota)$ is a permutation. Permutations π_{σ} in any $D(\sigma \rightarrow \sigma)$ can be defined by:

$$\pi_{(\sigma \rightarrow \tau)}(f) = \pi_{\tau} \circ f \circ \pi_{\sigma}^{-1} \quad (f \in D(\sigma \rightarrow \tau)).$$

If M is closed term then $\pi(\llbracket M \rrbracket) = \llbracket M \rrbracket$ (see [2]). However this does not characterise λ -definability.

For example ground equality, $=_{\iota}$, is permutation-invariant, but is certainly not λ -definable. Explicitly let 0 abbreviate (ι, ι, ι) and let tt and ff be $\lambda \alpha_0^{\iota} \lambda \alpha_1^{\iota}. \alpha_0^{\iota}$ and $\lambda \alpha_0^{\iota} \lambda \alpha_1^{\iota}. \alpha_1^{\iota}$ respectively. Then $=_{\iota}$ is defined by:

$$=_{\iota}^{xy} = \begin{cases} \text{tt} & (\text{if } x=y) \\ \text{ff} & (\text{if } x \neq y) \end{cases} \quad (x, y \in D_{\iota})$$

But/

But the only λ -definable functionals of type $(\iota, \iota, 0)$ are $\lambda\alpha^{\iota}_0 \lambda\alpha^{\iota}_1 \lambda\alpha^{\iota}_2 \lambda\alpha^{\iota}_3 \alpha^{\iota}_j$ for $0 \leq j \leq 3$ none of which are $=_{\iota}$ if $|D_{\iota}| > 1$.

M. Gordon proposed, as a possible remedy, that relations $R_{\iota} \subseteq D_{\iota}^2$ should be extended - not just permutations. Starting with such an R_{ι} , the R_{σ} 's are defined by:

$$R_{(\sigma \rightarrow \tau)}(f, g) \equiv \forall x, y \in D_{\sigma}. (R_{\sigma}(x, y) \rightarrow R_{\tau}(fx, gy)).$$

When R_{ι} is a permutation $\pi_{\iota}, R_{\sigma} = \pi_{\sigma}$ for all σ . The definition generalises, in the obvious way, if one starts with $R_{\iota} \subseteq D^{\kappa}$, for any ordinal κ . If $R_{\sigma} \subseteq D_{\sigma}^{\kappa}$ is obtained from an R_{ι} in that way it is called κ -logical; $f \in D_{\sigma}$ satisfies it iff $R_{\sigma}(\hat{f})$ holds. With the obvious definitions of λ -definability and λ -definability from a set $X \subseteq \cup D_{\sigma}$, one shows that any λ -definable functional satisfies any κ -logical relation, of the right type and that if $\{R_{\sigma}\}$ is the system of relations obtained from some R_{ι} , and each member of X satisfies the appropriate R_{σ} and f is λ -definable from X , then x satisfies the appropriate R_{σ} . The proof is like that of theorem 1.1.

One can now see why $=_{\iota}$ is not λ -definable if $|D_{\iota}| > 1$. Let $0, 1$ be distinct elements of D_{ι} . Let $R_{\iota} = \{\langle 0, 0 \rangle, \langle 0, 1 \rangle, \langle 1, 0 \rangle\}$. Then $R(tt, ff)$ is false for $R(1, 0)$ and $R(0, 1)$ but not $R(tt10, ff01)$. Therefore $R(=_{\iota}, =_{\iota})$ is false for $R(0, 0)$ and $R(0, 1)$ but not $R(=_{\iota}(0)(0), =_{\iota}(0)(1))$.

As an example of non-relative definability, consider the universal quantifier \forall_{ι} of type $((\iota \rightarrow 0) \rightarrow 0)$ defined by:

$$\forall_{\iota}(f) = \begin{cases} tt & (\text{if } fx=tt \text{ for all } x \text{ in } D_{\iota}) \\ ff & (\text{otherwise}). \end{cases}$$

Now \forall_{ι} is permutation-invariant; however if $|D_{\iota}| \geq 3$ it is not λ -definable from $=_{\iota}$. To see this let $R_{\iota} = \{\langle 0, 0 \rangle, \langle 1, 1 \rangle\}$ where $0, 1$ are distinct elements of D_{ι} . $R(=_{\iota}, =_{\iota})$ is true, but if $f, g \in D_{(\iota \rightarrow 0)}$ are such that $f(x)$ is always tt but $g(x)$ is tt iff x is 0 or 1 then

R/

$R(f,g)$ but not $R(\forall_{\mathcal{L}}(f), \forall_{\mathcal{L}}(g))$. Incidentally, if $|D| < 3$, $\forall_{\mathcal{L}}$ is λ -definable from $=_{\mathcal{L}}$.

We can only characterise definability using logical relations, for types of level ≤ 2 .

Theorem 1 Suppose τ has the form $(\tau_1, \dots, \tau_n, \mathcal{L})$ where each τ_i has the form $(\mathcal{L}, \dots, \mathcal{L}, \mathcal{L})$. Then if $|D_{\mathcal{L}}| \geq \aleph_0$ and $f \in D_{\tau}$ satisfies every 2-logical relation, it is λ -definable.

Proof We will just give two cases since this should give the idea without overmuch detail.

Suppose $\tau = (\mathcal{L}, \mathcal{L}, \mathcal{L})$. Let $x, y, 0, 1$ be elements of D , with 0 and 1 distinct and take $R_{\mathcal{L}} = \{ \langle x, 1 \rangle, \langle y, 0 \rangle \}$. Then $R(fxy, f10)$. So for every $x, y \in D_{\mathcal{L}}$ either $fxy = x$ and $f10 = 1$ or else $fxy = y$ and $f10 = 0$. Therefore either $f10 = 1$ or $f10 = 0$ and so, since $1 \neq 0$ either $f = \llbracket \lambda \alpha'_0 \lambda \alpha'_1 \alpha'_0 \rrbracket$ or $f = \llbracket \lambda \alpha'_0 \lambda \alpha'_1 \alpha'_1 \rrbracket$.

The other case we consider is $\tau = ((\mathcal{L}, \mathcal{L}), \mathcal{L}, \mathcal{L})$. Identify the integers, with a subset of $D_{\mathcal{L}}$ and let the restriction of s to the integers be the successor function. Given g in $D_{(\mathcal{L} \rightarrow \mathcal{L})}$ and $x \in D_{\mathcal{L}}$ let $R_{\mathcal{L}} = \{ \langle g^n(x), s^n(0) \rangle \mid n \geq 0 \}$. Clearly $R(x, 0)$ and $R(g, s)$. Therefore $R(f(g)(x), f(s)(0))$ and so for every $g \in D_{(\mathcal{L} \rightarrow \mathcal{L})}$ and $x \in D_{\mathcal{L}}$ there is an n such that $f(g)(x) = g^n(x)$ and $f(s)(0) = s^n(0)$. Since $s^n(0) = s^{n'}(0)$ iff $n = n'$, n must be independent of g and x and so for some n , $f = \llbracket \lambda \alpha'_0 \underbrace{(\mathcal{L} \rightarrow \mathcal{L})}_{n \text{ times}} (\alpha'_0 \dots (\alpha'_0 (\mathcal{L} \rightarrow \mathcal{L}) (\alpha'_0)) \dots) \rrbracket$.

We believe the theorem holds without the restriction on $D_{\mathcal{L}}$. The simplest type which has us baffled when $|D_{\mathcal{L}}| \geq 2$ is $((\mathcal{L}, \rightarrow \mathcal{L}) \rightarrow \mathcal{L}) \rightarrow \mathcal{L}$.

Some characterisation of definability can be obtained by strengthening the implication in the definition of $R_{\mathcal{L}}$ to an intuitionistic one, à la Kripke [1].

To this end, suppose we have a set W (of worlds) a reflexive, transitive binary relation \leq which is a subset of W^2 (alternativeness) and a relation $R_{\mathcal{L}} \subseteq D^3 \times W$ such that:

\forall

$$\forall x, y, z \in D_{\mathcal{L}} \forall w' \in W (R(x, y, z, w) \wedge (w \leq w') \rightarrow R(x, y, z, w')).$$

Define relations $R_{\sigma} \subseteq D_{\sigma}^3 \times W$ by:

$$R_{(\sigma \rightarrow \tau)}(f, g, h, w) \equiv (\forall x, y, z \in D_{\sigma} \forall w' \in W ((R(x, y, z, w') \wedge (w \leq w')) \rightarrow R(f(x), g(y), h(z), w'))), (f, g, h \in D_{(\sigma \rightarrow \tau)}, w \in W).$$

Any such R_{σ} is called 3-I-logical; $f \in D_{\sigma}$ satisfies R_{σ} iff $R_{\sigma}(f, f, f)$ is true. It is clear how the definition of κ -logical goes, for any ordinal κ . The reason for the magic number 3 is:

Theorem 2 If $|D_{\mathcal{L}}| \geq \aleph_0$, then $f \in D_{\sigma}$ satisfies every 3-I-logical relation iff it is λ -definable.

We don't know if the restriction on $D_{\mathcal{L}}$ can be dropped, or if 3 can be reduced to 2; it cannot be reduced to 1 because if $D_{\mathcal{L}}$ is the integers and $f \in D_{((\mathcal{L} \rightarrow \mathcal{L}) \rightarrow (\mathcal{L} \rightarrow \mathcal{L}))}$ is defined by:

$f(g)(x) = g^{g(x)}(x)$ ($g \in D_{(\mathcal{L} \rightarrow \mathcal{L})}$, $x \in D_{\mathcal{L}}$), then f satisfies every 1-I-logical predicate but is not λ -definable.

Neither do we know anything about characterising relative definability, even in interesting special cases, or what happens in other models of the typed λ -calculus or, of course, what happens in the case of the untyped λ -calculus. The rest of the memorandum is devoted to the proof of theorem 2.

Lemma 1 Suppose $R_{\sigma} \subseteq D_{\sigma}^3 \times W$ is 3-I-logical. Then, for the appropriate \leq : $\forall f, g, h \in D_{\sigma} \forall w, w' \in W ((R_{\sigma}(f, g, h, w) \wedge w \leq w') \rightarrow R_{\sigma}(f, g, h, w'))$.

Proof By induction on σ .

Lemma 2 If $f \in D_{\sigma}$ is λ -definable, it satisfies every 3-I-logical relation.

Proof Let $\{R_{\sigma}\}$ be the collection of 3-I-logical relations built up, as above, from some W, \leq and $R_{\mathcal{L}} \subseteq D_{\mathcal{L}}^3 \times W$. Clearly, if $f \in D_{(\sigma \rightarrow \tau)}$ satisfies $R_{\sigma \rightarrow \tau}$ and $g \in D_{\sigma}$ satisfies R_{σ} then $f(g)$ satisfies R_{τ} . So we need only/

only show, by a remark made above, that $\llbracket K \rrbracket$ and $\llbracket S \rrbracket$ satisfy the appropriate R's.

For K, suppose that $w \leq w'$, $R(f, g, h, w')$, $w' \leq w''$ and $R(x, y, z, w'')$. Then $R(\llbracket K \rrbracket(f)(x), \llbracket K \rrbracket(g)(y), \llbracket K \rrbracket(h)(z), w'')$ by the previous lemma, since $R(f, g, h, w')$ and $w' \leq w''$.

For S, suppose that $w \leq w'$, $R(f, f', f'', w')$, $w' \leq w''$, $R(g, g', g'', w'')$, $w'' \leq w'''$ and $R(x, x', x'', w''')$. As $w' \leq w''$, $R(f(x), f'(x'), f''(x''), w'')$ and as $w'' \leq w'''$, $R(g(x), g'(x'), g''(x''), w''')$. Therefore, as $w''' \leq w'''$, $R(f(x)(g(x)), f'(x)(g'(x')), f''(x'')(g''(x''')), w''')$, concluding the proof.

This establishes the "consistency" half of theorem 2. Of course, the lemma also holds if \exists is replaced by any \mathcal{K} and, further, an analogue of theorem 1.1.2 also holds in general.

To obtain the "completeness" half we use a special W , \leq and R which in turn requires us to give a "standard" environment, ρ_0 , which assigns to each α_i^γ an element of D_γ which behaves like α_i^γ , in a way to be made clear by lemmas 3 and 4.

Now we suppose that $|D_\zeta| \geq \aleph_0$ and $(())$ is a map from the set of terms of type ζ to D_ζ such that:

$$((M)) = ((N)) \text{ iff } M \approx N.$$

A vector, $\vec{\alpha}$, of variables is non-repeating if no variable occurs twice in it. If M is a term and $\vec{N} = \langle N_1, \dots, N_n \rangle$ ($n \geq 0$) is a vector of terms, $M \vec{N}$ abbreviates $(\dots((M N_1) N_2) \dots N_n)$; similarly if \vec{x} is a vector of elements of $\bigcup D_\gamma$ then $f \vec{x}$ abbreviates $f(x_1) \dots (x_n)$, where $\vec{x} = \langle x_1, \dots, x_n \rangle$ and f is a functional of the appropriate type. If $\vec{\alpha} = \langle \alpha_1, \dots, \alpha_n \rangle$ is a vector of variables, then $\rho(\vec{\alpha}) = \langle \rho(\alpha_1), \dots, \rho(\alpha_n) \rangle$ for any environment ρ .

The standard environment, ρ_0 , is defined by:

$$1. \quad \rho_0(\alpha_i^\zeta) = ((\alpha_i^\zeta)) \quad (i \geq 0)$$

2./

2. If there is a vector \vec{N} of terms of length n such that for any j ($1 \leq j \leq n$) and almost every* non-repeating $\vec{\alpha}_j$ containing no variable free in $(\vec{N})_j$, $f_j \rho_0(\vec{\alpha}_j) = ((\vec{N})_j \vec{\alpha}_j)$, then $\rho_0(\alpha_i^{(r_1, \dots, r_n)})_{(f_1, \dots, f_n)} = ((\alpha_i^{(r_1, \dots, r_n \vec{N})}))$, where \vec{N} is one such vector; if there is no such vector then

$$\rho_0(\alpha_i^{(r_1, \dots, r_n, c)})_{(f_1, \dots, f_n)} = ((\alpha_i^c)). \quad (i \geq 0; f_j \in D_{r_j} \text{ if } 1 \leq j \leq n)$$

Notice that if \vec{N}_1 and \vec{N}_2 satisfy the conditions of the second clause then they are, componentwise, \approx . So ρ_0 is well-defined.

Lemma 3 For every term M and vector $\vec{\alpha}$ of variables of the appropriate type, $[[M\vec{\alpha}]](\rho_0) = ((M\vec{\alpha}))$.

Proof First we prove the lemma for variables, by induction on types. If $M = \alpha_i^c$, the result is immediate from the definition of ρ_0 . For $\alpha^{(r_1, \dots, r_n)}$, let $\vec{\alpha} = \alpha_1 \dots \alpha_n$. For $1 \leq j \leq n$, let $\vec{\beta}_j$ be any vector of variables such that $\alpha_j \vec{\beta}_j$ has type c . By induction hypothesis, $[[\alpha_j \vec{\beta}_j]](\rho_0) = ((\alpha_j \vec{\beta}_j))$. Therefore $\vec{\alpha}$ satisfies the condition in clause 2 of the definition of ρ_0 , when $\langle f_1, \dots, f_n \rangle = \rho_0(\vec{\alpha})$. Therefore, $[[M\vec{\alpha}]](\rho_0) = \rho_0(\alpha^{(r_1, \dots, r_n)})_{\rho_0(\vec{\alpha})} = ((M\vec{\alpha}))$.

Finally the lemma is proved for terms in β, η -normal form by induction on their size. If M is a variable, we are finished. Otherwise, if $M\vec{\alpha} \approx (\dots(\alpha_{M_1}) \dots M_k)^{(k>0)}$ where α is some variable and the M_j ($1 \leq j \leq k$) all are β, η -normal forms of smaller size than M , the proof then proceeds by applying the induction hypothesis to the M_j , in a way similar to the above. The only other case is when $M \approx \lambda \beta_1 \dots \lambda \beta_k \alpha_i^c$ for some i , and this is very easy.

* Here and elsewhere assertions of the form "for almost every $\vec{\alpha} \dots$ ". should be read as "there is a finite set of variables such that for every $\vec{\alpha}$, none of whose component variables are in the set, \dots ".

From now on, we drop references to ρ_0 in $\llbracket M \rrbracket (\rho_0)$. Lemma 3 implies that $\llbracket M \rrbracket = \llbracket M' \rrbracket$ iff $M \approx M'$. If D_c were finite then there would always be closed M, M' , even of type $((c \rightarrow c) \rightarrow (c \rightarrow c))$, such that $\llbracket M \rrbracket = \llbracket M' \rrbracket$ but $M \not\approx M'$. The above methods can be used to show that given closed M, M' , there is an integer n , such that if $|D_c| \geq n$ and $\llbracket M \rrbracket = \llbracket M' \rrbracket$ then $M \approx M'$. These remarks form the converse to the consistency of \approx , mentioned above.

A term M is of order zero iff it has the form $\alpha M_1 \dots M_k$.

Lemma 4 Suppose M has order-zero and $x \rho_0(\vec{\alpha}) = y \rho_0(\vec{\alpha})$ for almost all non-repeating vectors $\vec{\alpha}$ such that $x \rho_0(\vec{\alpha}) \in D_c$. Then $\llbracket M \rrbracket (x) = \llbracket M \rrbracket (y)$.

Proof Immediate from the definition of ρ_0 .

Now W, \leq and R_c can be defined. The worlds, W are triples, $\langle \vec{\rho}, \vec{\alpha}, \vec{\beta} \rangle$, where $\vec{\rho}$ is a vector of members of $\cup D_c$, $\vec{\alpha}, \vec{\beta}$ are non-repeating and $\vec{\rho}, \vec{\alpha}, \vec{\beta}$ have the same length and corresponding members have the same type. If $w_i = \langle \vec{\rho}_i, \vec{\alpha}_i, \vec{\beta}_i \rangle$ ($i=1,2$) then $w_1 \hat{w}_2 = \langle \vec{\rho}_1 \hat{\vec{\rho}}_2, \vec{\alpha}_1 \hat{\vec{\alpha}}_2, \vec{\beta}_1 \hat{\vec{\beta}}_2 \rangle$, where the component vectors have been concatenated. Then $w_1 \leq w_2$ iff $w_2 = w_1 \hat{w}_3$, for some world, w_3 . Finally, R_c is defined by:

$R_c(x, y, z, \langle \vec{\rho}, \vec{\alpha}, \vec{\beta} \rangle)$ iff \exists a closed term M such that $x = \llbracket M \rrbracket \vec{\rho}$, $y = \llbracket M \rrbracket \rho_0(\vec{\alpha})$ and $z = \llbracket M \rrbracket \rho_0(\vec{\beta})$.

Clearly W, \leq and R_c satisfy the required conditions for obtaining a system of 3-I-logical R_σ 's.

Lemma 5 1 If $R_\sigma(f, g, h, w)$ where $w = \langle \vec{\rho}, \vec{\alpha}, \vec{\beta} \rangle$ then there is a closed term M such that whenever $\vec{\alpha}^+ \vec{\alpha}^+, \vec{\beta}^+ \vec{\beta}^+$ are non-repeating vectors

$$f = \llbracket M \rrbracket \vec{\rho}, g \rho_0(\vec{\alpha}^+) = \llbracket (M \vec{\alpha}) \vec{\alpha}^+ \rrbracket \text{ and } h \rho_0(\vec{\beta}^+) = \llbracket (M \vec{\beta}) \vec{\beta}^+ \rrbracket$$

2 Suppose that $f = \llbracket M \rrbracket \vec{\rho}, g = \llbracket M \vec{\alpha} \rrbracket$ and $h = \llbracket M \vec{\beta} \rrbracket$ where M is a closed term and g and h are denotations of order-zero terms and $w/$

$w = \langle \vec{\phi}, \vec{\alpha}, \vec{\beta} \rangle \in W$. Then $R_\sigma(f, g, h, w)$ for the appropriate σ .

Proof Both parts are proved together by induction on σ .

1 For \langle the result is immediate from the definition of R_σ .

So suppose f has type $(\sigma \rightarrow \tau)$.

Suppose $R_{(\sigma \rightarrow \tau)}(f, g, h, w)$ where $w = \langle \vec{\phi}, \vec{\alpha}, \vec{\beta} \rangle$.

Suppose next that $w^+ = w \langle x, \alpha_1, \beta_1 \rangle$, where x has type σ , is in W .

By induction hypothesis using 2 we see that, $R_\sigma(x, \rho_0(\alpha_1), \rho_0(\beta_1), w^+)$.

Therefore $R_\tau(fx, (g \rho_0(\alpha_1)), (h \rho_0(\beta_1)), w^+)$.

By induction hypothesis, using 1, there is a closed term $M_{(x, \alpha_1, \beta_1)}$ such that if $\vec{\alpha} \sim \alpha_1 \sim \vec{\alpha}^+$ and $\vec{\beta} \sim \beta_1 \sim \vec{\beta}^+$ are non-repeating vectors then $fx = \llbracket M_{(x, \alpha_1, \beta_1)} \rrbracket \vec{\phi} x, (g \rho_0(\alpha_1)) \rho_0(\vec{\alpha}^+) = \llbracket ((M_{(x, \alpha_1, \beta_1)} \vec{\alpha}) \alpha_1) \vec{\alpha}^+ \rrbracket$ and $(h \rho_0(\beta_1)) \rho_0(\vec{\beta}^+) = \llbracket ((M_{(x, \alpha_1, \beta_1)} \vec{\beta}) \beta_1) \vec{\beta}^+ \rrbracket$.

The subscripts on $M_{(x, \alpha_1, \beta_1)}$ indicate its dependence on x, α_1 and β_1 . Then, $\llbracket ((M_{(x, \alpha_1, \beta_1)} \vec{\alpha}) \alpha_1) \vec{\alpha}^+ \rrbracket = (g \rho_0(\alpha_1)) \rho_0(\vec{\alpha}^+) = \llbracket ((M_{(x', \alpha_1, \beta_1)} \vec{\alpha}) \alpha_1) \vec{\alpha}^+ \rrbracket$ for any $x, x' \in D_\sigma$ if $\vec{\alpha} \sim \alpha_1 \sim \vec{\alpha}^+$ is non-repeating. Therefore, by lemma 3, $M_{(x, \alpha_1, \beta_1)} \approx M_{(x', \alpha_1, \beta_1)}$. A similar argument shows that $M_{(x, \alpha_1, \beta_1)}$ is also independent of α_1 and β_1 , and the conclusion follows.

2 Suppose $f = \llbracket M \rrbracket \vec{\phi}, g = \llbracket M \vec{\alpha} \rrbracket$ and $h = \llbracket M \vec{\beta} \rrbracket$ where M is a closed term and g and h are denotations of order-zero terms, and $w = \langle \vec{\phi}, \vec{\alpha}, \vec{\beta} \rangle \in W$.

Let $w^+ = \langle \vec{\phi} \sim \vec{\phi}^+, \vec{\alpha} \sim \vec{\alpha}^+, \vec{\beta} \sim \vec{\beta}^+ \rangle$ be a world and suppose $R_\sigma(x, y, z, w^+)$. Then by induction hypothesis for part 1, there is a closed term M_1 such that wherever $\vec{\alpha} \sim \vec{\alpha}^+ \sim \vec{\alpha}^{++}$, and $\vec{\beta} \sim \vec{\beta}^+ \sim \vec{\beta}^{++}$ are non-repeating vectors,

$$x = \llbracket M_1 \rrbracket \vec{\phi} \sim \vec{\phi}^+, y \rho_0(\vec{\alpha}^{++}) = \llbracket M_1 \vec{\alpha} \sim \vec{\alpha}^+ \sim \vec{\alpha}^{++} \rrbracket \text{ and } z \rho_0(\vec{\beta}^{++}) = \llbracket M_1 \vec{\beta} \sim \vec{\beta}^+ \sim \vec{\beta}^{++} \rrbracket.$$

$$\begin{aligned} \text{Then we have } fx &= (\llbracket M \rrbracket \vec{\phi}) (\llbracket M_1 \rrbracket (\vec{\phi} \sim \vec{\phi}^+)) \\ &= \llbracket M_2 \rrbracket \vec{\phi} \sim \vec{\phi}^+ \text{ for a certain closed term } M_2. \end{aligned}$$

Since $y \rho_0(\vec{\alpha}^{++}) = \llbracket M_1 \vec{\alpha} \sim \vec{\alpha}^+ \rrbracket \rho_0(\vec{\alpha}^{++})$ for almost all non-repeating vectors/

vectors $\vec{\alpha}^{\rightarrow++}$ such that $y \rho_0(\vec{\alpha}^{\rightarrow++})$ is in D_c and g is the denotation of a term of order zero, lemma 4 applies and we have,

$$\begin{aligned} gy &= g(\llbracket M_1 \vec{\alpha} \sim \vec{\alpha}^{\rightarrow+} \rrbracket) \\ &= (\llbracket M \rrbracket \rho_0(\vec{\alpha})) (\llbracket M_1 \rrbracket \rho_0(\vec{\alpha} \sim \vec{\alpha}^{\rightarrow+})) \\ &= \llbracket M_2 \rrbracket \rho_0(\vec{\alpha} \sim \vec{\alpha}^{\rightarrow+}), \text{ where } M_2 \text{ is the same term as above.} \end{aligned}$$

In the same way,

$$hz = \llbracket M_2 \rrbracket \rho_0(\vec{\beta} \sim \vec{\beta}^{\rightarrow+}).$$

It is now clear that gy and hz are themselves denotations of order zero terms, and so applying the induction hypothesis for part 2, we conclude that

$$R_{\tau}(fx, gy, hz, w^+).$$

Therefore $R_{(\sigma \rightarrow \tau)}(f, g, h, w)$, as was required, concluding the proof.

Lemma 5 gives us the second part of theorem 2. For suppose f satisfies every 3-I-logical R_{σ} . Then in particular we have $R_{\sigma}(f, f, f, w_0)$ where R_{σ} is the one to which lemma 5 applies and w_0 has all its components empty. By part 1 of lemma 5, there is a closed term M such that $f = \llbracket M \rrbracket$, which is just what was wanted.

Acknowledgements

I would like to thank Mike Gordon for his help and encouragement. The work was carried out with the aid of an S.R.C. research grant.

References

- [1] Kripke, S.A. (1965). Semantical analysis of intuitionistic logic I. Formal systems and recursive functions, Proc. 8th Logic Colloquium, Oxford, 1963. Amsterdam: North-Holland.
- [2] Läuchli, H. (1970). An abstract notion of realizability for which intuitionistic predicate calculus is complete. Intuitionism and proof theory. Amsterdam: North-Holland.
- [3] Milne, R. (1973). The formal semantics of computer languages and their implementations. Ph.D. thesis. University of Cambridge.
- [4] Plotkin, G.D. (1972) A set-theoretical definition of application. Memorandum MIP-R-95, School of Artificial Intelligence, University of Edinburgh.

[5]/

- [5] Park, D. (1970). The Y-combinator in Scott's lambda-calculus models. Symposium on Theory of Programming. University of Warwick. Unpublished.
- [6] Park, D. (1973). Personal communication.
- [7] Scott, D. (1969). Lattice-theoretic models for the λ -calculus. University of Princeton. Unpublished.
- [8] Scott, D. (1970). Continuous lattices. Technical Monograph PRG-7, Oxford University Computing Laboratory Programming Research Group.
- [9] Scott, D. (1972). Data types as lattices. University of Amsterdam. Unpublished lecture notes.
- [10] Wadsworth, C.P. (1973). The relationship between lambda-expressions and their denotations in Scott's models for the lambda-calculus, in The Proceedings of the Orleans conference (eds. Calais, Derrick and Sabbagh). To appear.