

# LEONHARD EULER: HIS LIFE, THE MAN, AND HIS WORK

WALTER GAUTSCHI

## Abstract

On the occasion of the 300th anniversary of Euler's birth (on April 15), an attempt is made to bring Euler's genius to the attention of a broad segment of the educated public. The three stations of his life—Basel, St. Petersburg, and Berlin—are sketched and the principal works identified in more or less chronological order. To convey a flavor of his work and its impact on modern science, a selection of some of Euler's memorable contributions is made and discussed in more detail. Remarks on Euler's personality, intellect, and craftsmanship, will round out the presentation.

## 1 Introduction

It is a virtually impossible task to do justice, in a short span of time and space, to the great genius of Leonhard Euler. We attempt, in this lecture, to bring across to a broad segment of the educated public, at least some glimpses of Euler's incredibly diverse and profound ideas. We begin in §2 with a brief outline of his life, going through the three stations of his life: Basel, St. Petersburg (twice), and Berlin. In §3, we identify in more or less chronological order Euler's principal works and try to convey a flavor and some of the characteristic features of his work by selecting, and describing in some more detail, a few of his many outstanding contributions. We conclude in §4 with remarks on Euler's personality, character, and intellect, as gained from testimonials of his contemporaries and from his vast correspondence, and in §5 with some bibliographic information for further reading.

## 2 His Life

### 2.1 Basel 1707–1727: Auspicious beginnings

Leonhard Euler was born on April 15, 1707, the first child of Paulus Euler and Margaretha Brucker. The father at the time was a vicar at the church of St. Jakob, just outside the old city walls of Basel. Although a theologian, Paulus had interests in mathematics and took courses from the famous Jakob Bernoulli during the first two years of his study at the university. About a year-and-a-half after Leonhard's birth, the family moved to Riehen, a suburb of Basel, where Paulus Euler assumed the position of Protestant minister at the local parish. He served in that capacity faithfully and devotedly for the rest of his life.

The parish residence, as it looks today (Fig. 1) seems comfortable enough, but at the time the quarters it provided were rather cramped, especially after the family increased by another child, Anna Maria, in 1708. Two more children, Maria Magdalena and Johann Heinrich, were to follow later on.



Figure 1: The parish residence and church in Riehen

Leonhard received his first schooling at home from his mother, in classics, and from his father, in mathematics. Around the age of eight he was sent to the Latin school in Basel and given room and board at his maternal grandmother's house. To compensate for the poor quality then prevailing at the school, Paulus Euler hired a private tutor for his son, a young theologian, himself an enthusiastic lover of mathematics. In October of 1720, at the age of thirteen (not unusual at the time), Leonhard enrolled at the University of Basel, first at the Philosophical Faculty, where he took the freshman courses

on elementary mathematics given by Johann Bernoulli, the younger brother of Jakob. He graduated three years later with a master's degree and a public lecture (in Latin) comparing Descartes's system of natural philosophy with the one of Newton.

Following the wishes of his parents, he then entered the Theological Faculty, devoting, however, most of his time to mathematics and attending Johann Bernoulli's courses on higher mathematics. So successful did he pursue his mathematical studies that he soon caught the attention of his teacher, who in turn convinced Euler's father that Leonhard's vocation was mathematics, not theology. This is how Euler himself recounts this early learning experience at the university in his brief autobiography of 1767 (here freely translated from German):



Figure 2: The old university of Basel and Johann I Bernoulli

In 1720 I was admitted to the university as a public student, where I soon found the opportunity to become acquainted with the famous professor Johann Bernoulli, who made it a special pleasure for himself to help me along in the mathematical sciences. Private lessons, however, he categorically ruled out because of his busy schedule. However, he gave me a far more beneficial advice, which consisted in myself taking a look at some of the more difficult mathematical books and work through them with great diligence, and should I encounter some objections or difficulties, he offered me free access to him every Saturday afternoon, and he was gracious enough to comment on the collected difficulties, which was done with such a desired advantage that, when he resolved one of my objections, ten others at once disappeared,

which certainly is the best method of making auspicious progress in the mathematical sciences.

These personal meetings have become known, and famous, as the *privatissima*, and they continued well beyond his graduation. It was during these *privatissima* that Johann Bernoulli discovered, and more and more began to admire, the extraordinary mathematical talents of the young Euler.

Barely nineteen years old, Euler dared to compete with the greatest scientific minds of the time by responding to a prize question of the Paris Academy of Sciences with a memoir on the optimal placing of masts on a ship. He did not win first prize, but still a respectable second. A year later, when the physics chair at the University of Basel became vacant, the young Euler, dauntlessly again, but with the full support of his mentor, Johann Bernoulli, competed for the position, but failed, undoubtedly because of his youth and lack of an extensive record of publications. In a sense, this was a blessing in disguise, because in this way he was free to accept a call to the Academy of Sciences in St. Petersburg, founded a few years earlier by the czar Peter I (the Great). The groundwork for this appointment had been laid by Johann Bernoulli and two of his sons, Niklaus II and Daniel I, both of whom were already active at the Academy.

## **2.2 St. Petersburg 1727–1741: Meteoric rise to world fame and academic advancement**

Euler spent the winter of 1726 in Basel studying anatomy and physiology in preparation for his anticipated duties at the Academy. When he arrived in St. Petersburg, and started his position as an adjunct of the Academy, it was determined, however, that he should devote himself entirely to the mathematical sciences. In addition, he was to participate in examinations for the cadet corps and act as a consultant to the Russian state in a variety of scientific and technological questions.

Euler adjusted easily and quickly to the new and sometimes harsh life in the northern part of Europe. Contrary to most other foreign members of the Academy he began immediately to study the Russian language and mastered it quickly, both in writing and speaking. For six years he shared a dwelling with Daniel Bernoulli, and he was also on friendly terms with Christian Goldbach, the permanent Secretary of the Academy and best known today for his—still open—conjecture in number theory. The extensive correspondence




Figure 3: The Academy in St. Petersburg and Peter I

between Euler and Goldbach that ensued has become an important source for the history of science in the 18th century.

Euler's years at the Academy of St. Petersburg proved to be a period of extraordinary productivity and creativity. Many spectacular results achieved during this time (more on this later) brought him instant world fame and increased status and esteem within the Academy. A portrait of Euler stemming from this period is shown in Fig. 4.

In January of 1734 Euler married Katharina Gsell and they moved into a house of their own. The marriage brought forth thirteen children, of whom, however, only five reached the age of adulthood. The first-born child, Johann Albrecht, was to become a mathematician himself and later in life was to serve Euler as one of his assistants.



Figure 4: Euler ca. 1737

Euler was not spared misfortunes. In 1735, he fell seriously ill and almost lost his life. To the great relief of all, he recovered, but suffered a repeat attack three years later of (probably) the same infectious disease. This time it cost him his right eye, which is clearly visible on all portraits of Euler from this time on (for example the one in Fig. 6).

The political turmoil in Russia that followed the death of the czarina Anna Ivanovna induced Euler to seriously consider, and eventually decide, to leave St. Petersburg. This all the more as he already had an invitation from the Prussian king Frederick II to come to Berlin and help establish an Academy of Sciences there. This is how Euler put it in his autobiography:

... in 1740, when His still gloriously reigning Royal Majesty [Frederick II] came to power in Prussia, I received a most gracious call to Berlin, which, after the illustrious Empress Anne had died and it began to look rather dismal in the regency that followed, I accepted without much hesitation ...

In June of 1741, Euler together with his wife Katharina, the six-year-old Johann Albrecht, and the one-year-old toddler Karl, set out on the journey from St. Petersburg to Berlin.

### **2.3 Berlin 1741–1766: The emergence of epochal treatises**

Because of his preoccupation with the war campaign in Silesia, Frederick II took his time to set up the Academy. It was not until 1746 that the Academy finally took on shape, with Pierre-Louis Moreau de Maupertuis its president and Euler the director of the Mathematics Class. In the interim, Euler did not remain idle; he completed some twenty memoirs, five major treatises, now classics, and he composed over 200 letters!

Even though Euler was entrusted with manifold duties at the Academy—he had to oversee the Academy’s observatory and botanical gardens, deal with personnel matters, attend to financial affairs, notably the sale of almanacs which constituted the major source of income for the Academy, not to speak of a variety of technological and engineering projects—his mathematical productivity did not slow down. Nor was he overly distracted by an ugly priority dispute that erupted in the early 1750s over Euler’s principle of least action, which was also claimed by Maupertuis and which the Swiss fellow mathematician and newly elected academician Johann Samuel König



Figure 5: The Berlin Academy and Frederick II

affirmed to have already been formulated by Leibniz in a letter to the mathematician Jakob Hermann. He even accused Maupertuis of plagiarism. When challenged to produce the letter, König was unable to do so, and Euler was asked to investigate. Not sympathetic to Leibniz’s philosophy, Euler sided with Maupertuis and in turn accused König of fraud. This all came to a boil when Voltaire, aligned with König, came forth with a scathing satire ridiculing Maupertuis and not sparing Euler either. So distraught was Maupertuis



Figure 6: Euler 1753

that he left Berlin soon thereafter, and Euler had to conduct the affairs of the Academy as *de facto*, if not *de jure*, president of the Academy.

By now, Euler was sufficiently well-off that he could purchase a country estate in Charlottenburg, in the western outskirts of Berlin, which was large

enough to provide a comfortable home for his widowed mother, whom he had come to Berlin in 1750, his sister-in-law, and all the children. His first-born son Johann Albrecht was elected 1754, only twenty years old, to the Berlin Academy on the recommendation of Maupertuis. With a memoir on the perturbation of cometary orbits by planetary attraction he won in 1762 a prize of the Petersburg Academy, but had to share it with Alexis-Claude Clairaut. Euler's second son, Karl, went to study medicine in Halle, whereas the third, Christoph, became an officer in the military. His daughter Charlotte married into Dutch nobility and her older sister Helene a Russian officer later in 1777.

Euler's relation with Frederick II was not an easy one. In part, this was due to the marked difference in personality and philosophical inclination: Frederick—proud, self-assured, worldly, a smooth and witty conversationalist, sympathetic to French Enlightenment; Euler—modest, inconspicuous, down to earth, and a devout protestant. Another, perhaps more important, reason was Euler's resentment for never having been offered the presidency of the Berlin Academy. This resentment was only reinforced after Maupertuis's departure and Euler's subsequent efforts to keep the Academy afloat, when Frederick tried to interest Jean le Rond d'Alembert in the presidency. The latter indeed came to Berlin, but only to inform the king of his disinterest and to recommend Euler for the position instead. Still, Frederick not only ignored d'Alembert's advice, but ostentatiously declared himself the head of the Academy! This, together with many other royal rebuffs finally led Euler in 1766 to leave Berlin. He in fact already had a most cordial invitation from Empress Catherine II (the Great) to return to the Academy of St. Petersburg.

## **2.4 St. Petersburg 1766–1783: The glorious final stretch**

Highly respected at the Academy and adored at Catherine's court, Euler now held a position of great prestige and influence that had been denied him in Berlin for so long. He in fact was the spiritual leader of the Academy. Unfortunately, however, there were setbacks on a personal level. A cataract in his left (good) eye, which already began to bother him in Berlin, now became increasingly worse, so that in 1771 Euler decided to undergo an operation. The operation, though successful, led to the formation of an abscess, which soon destroyed Euler's vision almost entirely. Later in the same year, his wooden house burned down during the great fire of St. Petersburg, and the almost blind Euler escaped from being burnt alive only by a heroic rescue by





Figure 7: The Euler house and Catherine II

Peter Grimm, a workman from Basel. To ease the misfortune, the Empress granted funds to build a new house (the one shown in Fig. 7). Another heavy blow hit Euler in 1773 when his wife Katharina died. Euler remarried three years later so as not to be dependent on his children.

In spite of all these fateful events, Euler remained mathematically as active as ever, if not more. Indeed, about half of his scientific output was published, or originated, during this second St. Petersburg period, among which his two “bestsellers”, the *Letters to a German princess* and the *Algebra*. Naturally, he could not have done it without good secretarial and



Figure 8: Euler 1778

technical help, which he received among others from Niklaus Fuss, a compatriot from Basel and grandson-in-law of Euler, and from Euler’s own son, Johann Albrecht. The latter, by now secretary of the Academy, also acted

as the protocolist of the Academy sessions, over which Euler, as the oldest member of the Academy, had to preside.

The high esteem in which Euler was held at the Academy and at the Court is touchingly revealed by a passage in the memoirs of the Countess Dashkova, a directress of the Academy appointed by the Empress. She recounts the first time she accompanied the old Euler to one of the sessions of the Academy, probably Euler's last. Before the session started, a prominent professor and State Councilor as a matter of course claimed the chair of honor, next to the director's chair. The countess then turned to Euler and said: "Please be seated wherever you want; the seat you select will of course become the first of all."

Leonhard Euler died from a stroke on September 18, 1783 while playing with one of his grandchildren. Formulae that he had written down on two of his large slates describing the mathematics underlying the spectacular balloon flight undertaken on June 5, 1783 by the brothers Montgolfier in Paris, were found on the day of his death. Worked out and prepared for publication by Euler's son Johann Albrecht, they became the last article of Euler; it appeared in the 1784 volume of the *Mémoires*. A stream of memoirs, however, all queued up at the presses of the Academy, were still to be published for nearly fifty years after Euler's death.

### 3 His Work

In the face of the enormous volume of Euler's writings, we content ourselves with briefly identifying his principal works, and then select, and describe in some more detail, a few of Euler's prominent results in order to convey a flavor of his work and some of its characteristic features. The papers will be cited by their Eneström-Index numbers (E-numbers).

#### 3.1 The period in Basel

During the relatively short time of Euler's creative activity in Basel, he published two papers (E1, E3) in the *Acta Eruditorum* (Leipzig), one on isochronous curves, the other on so-called "reciprocal" curves, both influenced by Johann Bernoulli, and the work on the Paris Academy prize question (E4). The major work of this period is probably his *Dissertatio physica de sono* (E2), which he submitted in support of his application to the physics

chair at the University of Basel and had it printed 1727 in Basel. In it, Euler




Figure 9: Physical dissertation on sound, 1727

discusses the nature and propagation of sound, in particular the speed of sound, and also the generation of sound by musical instruments. Some of this work is preliminary and has been revisited by Euler in his *Tentamen* (cf. §3.2.1) and, thirty years later, in several memoirs (E305–E307).

## 3.2 First St. Petersburg period

In spite of the serious setbacks in health, Euler’s creative output during this period is nothing short of astonishing. Major works on mechanics, music theory, and naval architecture are interspersed with some 50 memoirs on a great variety of topics that run the gamut from the purest parts of mathematics to concrete problems in physics and astronomy. An account of the mathematical work during this period is given in [13].

### 3.2.1 Major works

The two-volume *Mechanica* (E15, E16) is the beginning of a far-reaching program, outlined by Euler in Vol. I, §98, of composing a comprehensive treatment of all aspects of mechanics, including the mechanics of rigid, flexible, and elastic bodies, fluid mechanics, and the many-body problem. The present work is restricted almost entirely to the dynamics of a point mass, to its free motion in Vol. I and constrained motion in Vol. II. In either case the motion may take place either in vacuum or in a resisting medium. The




Figure 10: Mechanics, 1736

novelty of the *Mechanica* consists in the systematic use of (the then new) differential and integral calculus, including differential equations, and in this sense it represents the first treatise on what is now called analytic (or rational) mechanics. It has won the praise of many leading scientists of the time, Johann Bernoulli among them, who said of the work that "... it does honor to Euler's genius and acumen". Also Lagrange, who in 1788 wrote his own *Mécanique analytique*, acknowledges Euler's mechanics to be "... the first great work where Analysis has been applied to the science of motion". Implementation and systematic treatment of the rest of Euler's program, never entirely completed, occupied him throughout much of his life.

It is evident from Euler's note books, that he thought a great deal about music and musical composition while still in Basel and had plans to write a book on the subject. These plans matured only later in St. Petersburg and gave rise to the *Tentamen novae theoriae musicae* (E33), usually referred to as the *Tentamen*, published in 1739 but completed already in 1731. (An English translation is available in [14, pp. 21–347]). The work opens with a discussion of the nature of sound as vibrations of air particles, including the propagation of sound, the physiology of auditory perception, and the generation of sound by string and wind instruments. The core of the work, however, deals with a theory of pleasure that music can evoke, which Euler develops by assigning to a tone interval, a chord, or a succession of such, a numerical value—the "degree"—which is to measure the agreeableness, or pleasure, of the respective musical construct: the lower the degree, the more pleasure. This is done in the context of Euler's favorite diatonic-chromatic



Figure 11: Tentamen, 1739 (1731)

temperament, but a complete mathematical theory of temperaments, both antique and contemporary ones, is also given.

In trying to make music an exact science, Euler was not alone: Descartes, Galileo, and Mersenne did the same before him, d’Alembert and many others after him (cf. Bailhache [2], Assayag [1]). In 1766 and 1774, Euler returns to music in three memoirs E314, E315, and E457.

Euler’s two-volume *Scientia navalis* (E110, E111) is a second milestone in his development of rational mechanics. In it, he sets forth the principles of hydrostatics and develops a theory of equilibrium and oscillations about the




Figure 12: Naval science, 1749 (1740–1741)

equilibrium of three-dimensional bodies submerged in water. This already contains the beginnings of the mechanics of rigid bodies, which much later is to culminate in his *Theoria motus corporum*, the third major treatise on mechanics (cf. §4.3.1). The second volume applies the theory to ships, to ship building and navigation.

### 3.2.2 Selecta Euleriana

#### Selectio 1. *The Basel problem*

This is the name that has become attached to the problem of determining the sum of the reciprocal squares,

$$(3.1) \quad 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots .$$

In modern terminology, this is called the zeta function of 2, where more generally

$$(3.2) \quad \zeta(s) = 1 + \frac{1}{2^s} + \frac{1}{3^s} + \frac{1}{4^s} + \cdots .$$

The problem has stumped the leading mathematicians of the time—Leibniz, Stirling, de Moivre, and all the Bernoullis—until Euler came along. Typically for Euler, he started using his stupendous capabilities of mental calculations and his adroitness in speeding up slowly converging series to calculate  $\zeta(2)$  to seven decimal places (E20). (Stirling, already in 1730, actually calculated the series to nine decimal places, but Euler did not yet know this.) The breakthrough came in 1735 (published as E41 in 1740) when he showed by a brilliant but daring procedure (using Newton’s identities for polynomials of infinite degree!) that

$$\zeta(2) = \frac{\pi^2}{6} .$$

Spectacular as this achievement was, Euler went on to use the same method to determine  $\zeta(s)$  for all even  $s = 2n$  up to 12. He found  $\zeta(2n)$  to be always a rational number multiplied by the  $2n$ th power of  $\pi$ . It was in connection with the Basel problem that Euler in 1732 discovered a general summation procedure, found independently by Maclaurin in 1738, and promptly used it to calculate  $\zeta(2)$  to twenty decimal places! Eventually, Euler managed to place his approach on a more solid footing, and he succeeded, in E130 (see

also E212, Part II, Ch. 5), to prove the general formula

$$(3.3) \quad \zeta(2n) = \frac{2^{2n-1}}{(2n)!} |B_{2n}| \pi^{2n}.$$

Here,  $B_{2n}$  are the Bernoulli numbers (introduced by Jakob Bernoulli in his *Ars conjectandi*), which Euler already encountered in his general summation formula.

Euler also tried odd values of  $s$ , but wrote in a letter to Johann Bernoulli that “. . . the odd powers I cannot sum, and I don’t believe that their sums depend on the quadrature of the circle [that is, on  $\pi$ ]” (Fellmann [6, p. 84, footnote 56]). The problem in this case, as a matter of fact, is still open today.

**Selectio 2.** *Prime numbers and the zeta function*

Let

$$\mathcal{P} = \{2, 3, 5, 7, 11, 13, 17, \dots\}$$

be the set of all prime numbers larger than 1, i.e., the integers  $> 1$  that have no divisors other than 1 and themselves. Euler’s fascination with prime numbers started quite early and continued throughout his life, even though the rest of the mathematical world at the time (Lagrange excluded!) was rather indifferent to problems of this kind. An example of his profound insight into the theory of numbers is the discovery in 1737 (E72) of the fabulous product formula

$$(3.4) \quad \prod_{p \in \mathcal{P}} \frac{1}{1 - 1/p^s} = \zeta(s)$$

connecting prime numbers with the zeta function (3.2). How did he do it?—Simply by starting with the zeta function and peeling away, layer after layer, all the terms whose integers in the denominators are divisible by a prime! Thus, from

$$\zeta(s) = 1 + \frac{1}{2^s} + \frac{1}{3^s} + \frac{1}{4^s} + \dots,$$

dividing by  $2^s$  and subtracting, one gets

$$\frac{2^s - 1}{2^s} \zeta(s) = 1 + \frac{1}{3^s} + \frac{1}{5^s} + \frac{1}{7^s} + \dots.$$

All the denominator integers divisible by 2 are gone. Do the same with the next prime 3, i.e., divide the last equation by  $3^s$  and subtract, and one gets

$$\frac{2^s - 1}{2^s} \cdot \frac{3^s - 1}{3^s} \zeta(s) = 1 + \frac{1}{5^s} + \frac{1}{7^s} + \frac{1}{11^s} + \dots,$$

where all integers divisible by 3 are gone. Continue in this way *ad infinitum*; everything eventually will be gone except for the first term 1,

$$\left( \prod_{p \in \mathcal{P}} \frac{p^s - 1}{p^s} \right) \zeta(s) = 1.$$

But this is the same as (3.4)!

The result provides a neat analytic proof of the fact (already known to the Greek) that the number of primes is infinite. Indeed, since  $\zeta(1)$ —the harmonic series—diverges to  $\infty$  (cf. Selectio 4), the product on the left of (3.4), if  $s = 1$ , cannot be finite.

The formula—the beginning of “analytic number theory”—in fact paved the way to important later developments in the distribution of primes.

**Selectio 3.** *The gamma function*

Following a correspondence in 1729 with Goldbach, Euler in E19 considers the problem of interpolating the sequence of factorials

$$(3.5) \quad n! = 1 \cdot 2 \cdot 3 \cdots n, \quad n = 1, 2, 3, \dots,$$

at noninteger values of the argument. Euler quickly realized that this cannot be done algebraically, but requires “transcendentals”, that is, calculus. He writes  $n!$  as an infinite product,

$$(3.6) \quad \frac{1 \cdot 2^n}{1+n} \cdot \frac{2^{1-n} \cdot 3^n}{2+n} \cdot \frac{3^{1-n} \cdot 4^n}{3+n} \cdot \frac{4^{1-n} \cdot 5^n}{4+n} \cdots,$$

which formally, by multiplying out the numerators, can be seen to be the ratio of two infinite products,  $1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdots$  and  $(n+1)(n+2)(n+3) \cdots$ , which indeed reduces to (3.5). Now for  $n = \frac{1}{2}$ , Euler manages to manipulate the infinite product (3.6) into the square root of an infinite product for  $\pi/4$  due to Wallis; therefore,  $\frac{1}{2}! = \frac{1}{2}\sqrt{\pi}$ . This is why Euler knew that some kind of integration was necessary to solve the problem.



By a stroke of insight, Euler takes the integral  $\int_0^1 x^e(1-x)^n dx$ —up to the factor  $1/n!$  the  $n$ -times iterated integral of  $x^e$ , where  $e$  is an arbitrary number (not the basis of the natural logarithms!)—and finds the formula

$$(3.7) \quad (e+n+1) \int_0^1 x^e(1-x)^n dx = \frac{n!}{(e+1)(e+2)\cdots(e+n)}.$$

He now lets  $e = f/g$  be a fraction, so that

$$\frac{f+(n+1)g}{g^{n+1}} \int_0^1 x^{f/g}(1-x)^n dx = \frac{n!}{(f+g)(f+2g)\cdots(f+ng)}.$$

If  $f = 1$ ,  $g = 0$ , then on the right we have  $n!$ ; on the left, we have to determine the limit as  $f \rightarrow 1$ ,  $g \rightarrow 0$ , which Euler takes to be the desired interpolant, since it is meaningful also for noninteger  $n$ . Skillfully, as always, Euler carries out the limit by first changing variables,  $x = t^{g/(f+g)}$ , to obtain

$$\frac{f+(n+1)g}{f+g} \int_0^1 \left( \frac{1-t^{g/(f+g)}}{g} \right)^n dt,$$

and then doing the limit as  $g \rightarrow 0$  with  $f = 1$  by the Bernoulli-L'Hospital rule. The result is  $\int_0^1 (-\ln t)^n dt$ . Here we can set  $n = x$  to be any positive number, and thus obtain  $x! = \int_0^1 (-\ln t)^x dt$ , which today is written as

$$(3.8) \quad x! = \int_0^\infty e^{-t} t^x dt = \Gamma(x+1)$$

in terms of the gamma function  $\Gamma$ . It is easily verified that

$$(3.9) \quad \Gamma(x+1) = x\Gamma(x), \quad \Gamma(1) = 1,$$

so that indeed  $\Gamma(n+1) = n!$  if  $n$  is an integer  $\geq 0$ .

Euler's unflinching intuition in producing the gamma function has been vindicated early in the 20th century when it was shown by Bohr and Mollerup that there is no other function interpolating the factorials if, in addition to satisfying the difference equation (3.9), it is also required to be logarithmically convex. The gamma function indeed has become one of the most fundamental functions in analysis.

The integral in (3.8) is often referred to as the second Eulerian integral, the first being

$$B(x, y) = \int_0^1 t^{x-1}(1-t)^{y-1} dt,$$

also called the beta function. The latter can be beautifully expressed in terms of the gamma function by

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)},$$

which is nothing but (3.7) for  $e = x - 1$ ,  $n = y - 1$ .

**Selectio 4.** *Euler's constant*

It is generally acknowledged that the two most important constants in mathematics are  $\pi = 3.1415\dots$ , the ratio of the circumference of a circle to its diameter, and  $e = 2.7182\dots$ , the basis of the natural logarithms, sometimes named after Euler. They pop up everywhere, often quite unexpectedly. The 19th-century logician Auguste de Morgan said about  $\pi$  that "...it comes on many occasions through the window and through the door, sometimes even down the chimney." The third most important constant is undoubtedly Euler's constant  $\gamma$  introduced by him 1740 in E43. Of the three together—the "holy trinity", as they are sometimes called—the last one,  $\gamma$ , is the most mysterious one, since its arithmetic nature, in contrast to  $\pi$  and  $e$ , is still shrouded in obscurity. It is not even known whether  $\gamma$  is rational, even though most likely it is not; if it were, say equal to  $p/q$  in reduced form, then according to a 1973 result of R. Brent,  $q$  would have to be larger than  $10^{10,000}$ .

Euler's constant arises in connection with the harmonic series  $\zeta(1) = 1 + \frac{1}{2} + \frac{1}{3} + \dots$  (so called because each of its terms is the harmonic mean of the two neighboring terms) and is defined as the limit

$$(3.10) \quad \gamma = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{2} + \frac{1}{3} \cdots + \frac{1}{n} - \ln n \right) = 0.57721\dots$$

It has been known as early as the 14th century that the harmonic series diverges, but a rigorous proof of it is usually attributed to Jakob Bernoulli, who also mentioned another proof by his younger brother Johann, which however is not entirely satisfactory. At any rate, Euler, in defining his constant and showing it to be finite, puts in evidence not only the divergence of the harmonic series, but also its logarithmic rate of divergence. Beyond this, using his general summation formula (mentioned in Selectio 1), he computes  $\gamma$  to 15 correct decimal places, and to equally many decimals the sum of the first million terms of the harmonic series! Since later (in 1790) Lorenzo Mascheroni also considered Euler's constant and computed it to 32 decimal

places (of which, curiously, the 19th, 20th, and 21st are incorrect), the name “Euler-Mascheroni constant” for  $\gamma$  is also found in the literature. As of today, it appears that  $\gamma$  has been computed to 108 million decimal places, compared to over  $2 \times 10^{11}$  decimals for  $\pi$  and 50.1 billion for  $e$ .

A recent encyclopedic account of Euler’s constant—its history and mathematical properties—can be found in Krämer [10].

After all these spectacular achievements, the numerous other memoirs written on many different topics, and his responsibilities at the Academy, it is unbelievable that Euler still had the time and stamina to write a 300-page volume on elementary arithmetic for use in the St. Petersburg gymnasia. How fortunate were those St. Petersburg kids for having had Euler as their teacher!

### 3.3 Berlin

Next to a multitude of memoirs and consultation on engineering and technology projects, this period saw the creation of a number of epochal scientific treatises and a highly successful and popular work on the philosophy of science.

#### 3.3.1 Major works

The brachystochrone problem—finding the path of a mass point moving under the influence of gravity from one point of a vertical plane to another in the shortest amount of time—is an early example of an optimization problem, posed by Johann Bernoulli, which seeks a function (or a curve) that renders optimal an analytic expression that depends on this function. In 1744 (E65), and later in 1766 (E296) adopting an improved approach of Lagrange, Euler vastly generalizes this problem, thereby creating an entirely new branch of mathematics, called (already by Euler) the “calculus of variations”. He derives his famous Euler equation: a necessary condition in the form of a differential equation that any solution of the problem must satisfy. Typically for Euler, he illustrates this by many—some hundred!—examples, among them the principle of least action that caused so much turmoil in the mid 1700s (cf. §2.3).

Two smaller treatises, one on planetary and cometary trajectories (E66), and another on optics (E88), appeared at about the same time (1744 resp.




Figure 13: Calculus of variations, 1744, and Artillery, 1745

1746). The latter is insofar of historical interest as it started the debate of Newton’s particle vs Euler’s own wave theory of light.

In deference to his master, king Frederick II, Euler translated an important work on ballistics by the Englishman Benjamin Robins, even though the latter had been unfairly critical of Euler’s *Mechanica* of 1736. He added, however, so many commentaries and explanatory notes (also corrections!) that the resulting book—his *Artillerie* of 1745 (E77)—is about five times the size of the original! Niklaus Fuss in his 1783 *Eulogy* of Euler (cf. *Opera omnia*, Ser. I, Vol. 1, pp. xliii–xcv) remarks: “. . . the only revenge [Euler] took against his adversary because of the old injustice consists in having made [Robin’s] work so famous as, without him, it would never have become.”

The two-volume *Introductio* of 1748 (E101, E102) together with the *Institutiones calculi differentialis* of 1755 (E212) and the four-volume *Institutiones calculi integralis* of 1763 and 1773 (E342, E366, E385, E660)—a “magnificent trilogy” (Fellmann [6, §4])—establishes analysis as we know it today. In the first volume of the *Introductio*, after a treatment of elementary functions, Euler summarizes his many discoveries in the areas of infinite series, infinite products, partition of numbers, and continued fractions. On several occasions, he uses the fundamental theorem of algebra, clearly states it, but does not prove it. He develops a clear concept of function—real as well as complex-valued—and emphasizes the fundamental role played in analysis by the number  $e$  and the exponential and logarithm functions. The second volume is devoted to analytic geometry: the theory of algebraic curves and surfaces.



ture (i.e., indefinite integration) of elementary functions and techniques for reducing the solution of linear ordinary differential equations to quadratures. In the second volume, he presents, among other things, a detailed theory of the important linear second-order differential equations, and in the third volume a treatment, to the extent known at the time (mostly through Euler’s own work), of linear partial differential equations. The fourth volume contains supplements to the preceding volumes, mostly the first. Euler’s method—a well-known approximate method for solving an arbitrary first-order differential equation—is embedded in Chapter 7 of the second part of Volume 1. Systems of differential equations, interestingly, are not considered in this work.

Euler’s program for mechanics (cf. §4.2.1) progressed steadily but slowly, as he tried to develop a theory of the motion of solids. Major technical difficulties had to be overcome, and were gradually dealt with in a series of memoirs. These, in 1765, eventually led to his great treatise *Theoria motus corporum* (E289), also called the *Second mechanics*, which crowned Euler’s work on mechanics. In addition to an improved exposition of his earlier mechanics of mass points (cf. §4.2.1), it now contains the differential equations (Euler’s equations) of motion of a rigid body subject to external forces. Here, Euler introduces the original idea of employing two coordinate systems—one fixed, the other moving, attached to the body—and deriving differential equations for the angles between the respective coordinate axes, now called the Euler angles. The intriguing motion of the spinning top is one of many examples worked out by Euler in detail.




Figure 16: *Theoria motus corporum*, 1765

Throughout his years in Berlin and beyond, Euler was deeply occupied with geometric optics. His memoirs and books on this topic, including the monumental three-volume *Dioptrics* (E367, E386, E404), written mostly




Figure 17: Optics, 1769–1771, and Letters, 1768, 1772 (1760–1762)

while still in Berlin, fill no fewer than seven volumes in his *Opera omnia*. A central theme and motivation of this work was the improvement of optical instruments like telescopes and microscopes, notably ways of eliminating chromatic and spherical aberration.

Euler’s philosophical views on science, religion, and ethics are expressed in over 200 letters written 1760–1762 (in French) to a German princess and published later in 1768 and 1772 (E343, E344, E417). While Euler’s role as a philosopher may be controversial (even his best friend Daniel Bernoulli advised him to better deal with “more sublime matters”), his *Letters* became an instant success and were translated into all major languages.

### 3.3.2 Selecta Euleriana

#### Selectio 5. *The Königsberg bridge problem*

The river Pregel, which flows through the Prussian city of Königsberg, divides the city into an island and three distinct land masses, one in the north, one in the east, and one in the south. There are altogether seven bridges, arranged as shown in green on the left of Fig. 18, connecting the three land masses with each other and with the island. The problem is: can one take a stroll from one point in the city to another by traversing each

bridge exactly once? In particular, can one return to the starting point in the same manner?

Evidently, this is a problem that cannot be dealt with by the traditional methods of analysis and algebra. It requires a new kind of analysis, an analysis that Leibniz already has termed *analysis situs*, and today is called “topology”. Euler solved the problem in 1735, published as E53 in 1741, by showing that such paths cannot exist. He does this by an ingenious process of abstraction, associating with the given land and bridge configuration (what today is called) a connected graph, i.e., a network of vertices and connecting edges, each vertex representing a piece of land and each edge a bridge connecting the respective pieces of land. In the problem at hand, there are four distinct pieces of land, hence four vertices, and they are connected with




Figure 18: The Königsberg bridge problem

edges as shown on the right of Fig. 18. It is obvious what is meant by a path from one vertex to another. A closed path is called a circuit, and paths or circuits are (today) called Eulerian, if each edge is traversed exactly once.

Euler recognized that a crucial concept here is the *degree* of a vertex, i.e., the number of edges emanating from it. If, in an arbitrary connected graph,  $n$  denotes the number of vertices of odd degree, he proves that (a) if  $n = 0$ , the graph has at least one Eulerian circuit; (b) if  $n = 2$ , it has at least one Eulerian path, but no circuit; (c) if  $n > 2$ , it has neither. (The case  $n = 1$  is impossible.) Since the Königsberg bridge graph has  $n = 4$ , we are in case (c), hence it is impossible to traverse the city in the manner required in the problem.

Here again, like in the calculus of variations, one can admire Euler’s



powerful drive and capacity of starting with a concrete example and deriving from it, by a process of sweeping generalization, a whole new theory. In the present case, it is the theory of graphs and topological networks.

**Selectio 6.** *Euler's buckling formula* (1744)

In a first supplement to his *Methodus* (cf. Fig. 13, left), Euler applies the calculus of variations to elasticity theory, specifically to the bending of a rod subject to an axial load. He derives the critical load under which the rod buckles. This load depends on the stiffness constant of the material, on the way the rod is supported at either end, and it is inversely proportional to the square of the length of the rod. A particular configuration of two rods loaded on top by a connecting bar (assumed to be of infinite stiffness) is shown in Fig. 19, during the initial phase (left), and at the time of buckling (right). Here, the top end of the rods is slidably supported, and the bottom end clamped.




Figure 19: The buckling of a rod

The critical load is the first *elastostatic* eigenvalue of the problem. Euler also calculates the *elastokinetic* eigenvalues, the eigenfrequencies of the rod's transversal oscillations, and the associated eigenfunctions, which determine the shapes of the deformed rod.

**Selectio 7.** *Euler flow*

In a series of three memoirs, E225–E227, all published in 1757, and another three papers (E258, E396, E409), Euler gave his definitive treatment of continuum and fluid mechanics, culminating a number of earlier memoirs on the subject. It contains the celebrated Euler equations, expressing

the conservation of mass, momentum, and energy. In two [three] dimensions, these constitute a system of four [five] nonlinear hyperbolic partial differential equations, which have to be solved, given appropriate initial and boundary conditions. Naturally, in Euler’s time, this was virtually impossible, except in very special cases, and indeed Euler in the introduction to E226 had to write “... if there remain any difficulties, they shall not be on the side of mechanics, but solely on the side of analysis: for this science has not yet been carried to the degree of perfection which would be necessary in order to develop analytic formulae which include the principles of the motion of fluids.” Nowadays, however, the Euler equations are widely being used in computer simulation of fluids.




Figure 20: Transonic Euler flow at Mach .85 about a cylinder

An example is the asymmetric flow of a compressible, inviscid fluid about a circular cylinder at transonic speed, calculated in 1995 by N. Botta [3]. A still picture of the two-dimensional flow (vorticity contour lines) is shown in Fig. 20.

**Selectio 8.** *Euler’s polyhedral formula* (1758)

In a three-dimensional convex polyhedron, let  $V$  denote the number of vertices,  $E$  the number of edges, and  $F$  the number of faces. Mentioned 1750 in a letter to Goldbach, and later published in E231, Euler proves for the first time the extremely simple, but stunning formula

$$V - E + F = 2.$$

Descartes, some 100 years earlier, already knew, but did not prove, the formula. It is one of the first contributions to combinatorial topology.

Here, too, as in the Königsberg bridge problem (cf. *Selectio 5*), there is a connection with graph theory, as to each polyhedron one can associate a connected graph. The construction of the graph, however, is not so obvious, and Euler was unaware of it; the first to observe this seems to have been Cauchy in 1813.

Much later, the formula has been generalized to polytopes in spaces of arbitrary dimension.

**Selectio 9.** *Euler and  $q$ -theory*

The story here begins with a letter Euler wrote in 1734 to Daniel Bernoulli, in which he considered the (somewhat bizarre) problem of interpolating the common logarithm  $\log x$  at the points  $x_r = 10^r$ ,  $r = 0, 1, 2, \dots$ . He essentially writes down Newton's interpolation series  $S(x)$  (without mentioning Newton by name) and remarks that, when  $x = 9$ , the series converges quickly, but to a wrong value,  $S(9) \neq \log 9$  (cf. [8]). Rather than losing interest in the problem, Euler must have begun pondering the question about the nature of the limit function  $S(x)$ ; what is it, if not the logarithm?

Almost twenty years later, in 1753, he returned to this problem in E190, now more generally for the logarithm to base  $a > 1$ , and studied the respective limit function  $S(x; a)$  in great detail. Intuitively, he must have perceived its importance. Today we know indeed (Koelink and Van Assche [9]) that it can be thought of as a  $q$ -analogue of the logarithm, where  $q = 1/a$ , and some of the identities derived by Euler (in part already contained in Vol. 1, Ch. 16 of his *Introductio*) are in fact special cases of the  $q$ -binomial theorem—a centerpiece of  $q$ -theory in combinatorial analysis and physics. Thus, Euler must be counted among the precursors of  $q$ -theory, which was only developed about 100 years later by Heine.

### 3.4 Second St. Petersburg period

This may well be Euler's most productive period, with over 400 published works to his credit, not only on each of the topics already mentioned, but also on geometry, probability theory and statistics, cartography, even widow's pension funds and agriculture. In this enormous body of work there figure three treatises on algebra, lunar theory, and naval science, and what appear to be fragments of major treatises on number theory (E792), natural philosophy (E842), and dioptrics (E845).

### 3.4.1 Major works

Soon into this second St. Petersburg period, another of Euler's "bestsellers" appeared: the *Vollständige Anleitung zur Algebra* (E387, E388), or *Algebra* for short. Even before publication of the German original, a translation into Russian came out, and translations into all major languages were soon to follow. (The French translation by Johann III Bernoulli includes a long




Figure 21: Algebra, 1770

supplement by Lagrange containing an exposé on the arithmetic theory of continued fractions and many addenda to the last section of the *Algebra* dealing with Diophantine equations.)

Euler wrote this 500-page volume to introduce the absolute beginner into the realm of algebra. He dictated the work to a young man—a tailor by profession—whom he brought with him from Berlin, and who (according to the preface of the work) “. . . was fairly good at computing, but beyond that did not have the slightest notion about mathematics . . . As far as his intellect is concerned, he belonged among the mediocre minds.” Nevertheless, it is said that, when the work was completed, he understood everything perfectly well and was able to solve algebraic problems posed to him with great ease.

It is indeed a delight to witness in this work Euler's magnificent didactic skill, to watch him progress in ever so small steps from the basic principles of arithmetic to algebraic (up to quartic) equations, and finally to the beautiful art of Diophantine analysis. Equally delightful is to see how the theory is illustrated by numerous well-chosen examples, many taken from everyday life.

The orbit of the moon with all its irregularities has fascinated mathematicians like Clairaut, d’Alembert and others for a long time, including Euler, who already in 1753 published his *Theoria motus lunae* (E187), the *First lunar theory*. The theory he developed, while tentative, provided astronomers with formulae needed to prepare lunar tables, which in turn served seafaring nations for over a century with accurate navigational aids. Euler’s definitive work on the subject, however, is his *Second lunar theory* (E418) of 1772, a monumental work dealing in a more effective way than before with




Figure 22: Second lunar theory, 1772 and Second theory of ships, 1773

the difficult three-body problem, i.e., with the study of the motion of three bodies—in this case the sun, the earth, and the moon, thought of as point masses—moving under the influence of mutual gravitational forces. Already Newton is reputed to have said that “. . . an exact solution of the three-body problem exceeds, if I am not mistaken, the power of any human mind.” Today it is known, indeed, that an exact solution is not possible. Euler grapples with the problem by introducing appropriate variables, again choosing two coordinate systems—one fixed, the other moving—applying processes of successive approximation, and making use, when needed, of observational data.

According to L. Courvoisier (cf. *Opera omnia*, Ser. II, Vol. 22, p. xxviii), “. . . all later progress in celestial mechanics is based, more or less, on the ideas contained in the works of Euler, [and the later works of] Laplace and Lagrange.”

The *Théorie complète de la construction et de la manœuvre des vaisseaux* (E426), also called the *Second theory of ships*, is a work that treats the topic indicated in the title for people having no or little mathematical knowledge,

in particular for the sailors themselves. Not surprisingly, given the level of presentation and the author's extraordinary didactic skill, the work proved to be very successful. The French maritime and finance minister (and famous economist), Anne Robert Jacques Turgot, proposed to king Louis XVI to require all students in marine schools (and also those in schools of artillery) to study Euler's relevant treatises. The king even paid Euler 1,000 Rubles for the privilege of having the work reprinted, and czarina Catherine II, not wanting to be outdone by the king, doubled the amount and pitched in an additional 2,000 Rubles!

### 3.4.2 *Selecta Euleriana*

#### **Selectio 10** *Euler's disk*

In a number of memoirs (E257, E292, E386, E585) from the 20-year period 1761–1781, Euler analyzes the motion of a rigid body around a moving axis, including the effects of friction. An interesting example is the “Euler disk”, a circular (homogeneous) metal disk being spun on a clean smooth surface. At first, it will rotate around its vertical axis, but owing to friction, the axis is beginning to tilt and the disk to roll on a circular path. The more the axis is tilting, the wider the circular path and the higher the pitch of the whirring sound produced by the point of contact of the disk with the surface. Thus, paradoxically, the speed of the motion seems to increase, judging from the rising pitch of the sound, although energy is being dissipated through friction. The disk, eventually, comes to an abrupt halt, flat on the surface.

The key toward explaining the motion are Euler's equations, a set of differential equations involving the Euler angles (cf. §3.3.1). The technical details of the motion, though, are still being debated today (cf. e.g. the publications listed on the web site <http://www.eulersdisk.com>).

#### **Selectio 11** *Euler's gear transmission*

In connection with the design of water turbines, Euler developed optimal profiles for teeth in cogwheels that transmit motion with a minimum of resistance and noise (E330, OII.17, pp. 196–219). These profiles involve segments of cycloidal curves as shown in Fig. 23.

The technical realization of this design took shape only later in what is called the involute gear. Euler not only is the inventor of this kind of gear, but he also anticipated the underlying geometric equations now usually called the Euler–Savary equations.




Figure 23: Euler gear, 1767

## 4 The Man

### 4.1 Personality

From various testimonials of Euler's contemporaries, and also, of course, from Euler's extensive correspondence, one has a fairly accurate picture of Euler's personality. A valuable source is the eulogy of Nicklaus Fuchs (*Opera omnia*, Ser. I, Vol. 1, pp. xliii–xcv), who during the last ten years of Euler's life has seen him regularly, almost on a daily basis, as one of his assistants. Also based on personal acquaintance is the eulogy of the marquis Nicolas de Condorcet (*Opera omnia*, Ser. III, Vol. 12, pp. 287–310), which, however, deals more with Euler's work. Euler comes across as a modest, inconspicuous, uncomplicated, yet cheerful and sociable person. He was down-to-earth and upright; "honesty and uncompromising rectitude, acknowledged Swiss national virtues, he possessed to a superior degree" writes Fuchs. Euler never disavowed, in fact was proud of, his Swiss heritage; he, so Fuchs (who also originates from Basel): "...always retained the Basel dialect with all the peculiarities of its idiom. Often he amused himself to recall for me certain provincialisms and reversals [of word order], or mix into his parlance Basel expressions whose use and meaning I had long forgotten."

Feelings of rancor, be it because of priority issues or because of unfair criticism, are totally foreign to Euler. When Maclaurin, for example, discovered the well-known summation formula which Euler obtained six years earlier, and for some time the formula was generally referred to as the "Maclaurin

summation formula”, Euler did not object, let alone complain. It may even have pleased him that others hit upon the same fortunate idea. In due time, of course, the formula became justly known as the “Euler–Maclaurin summation formula”. Another example is Maupertuis’s claim for the principle of least action (cf. §2.3), which Euler had already enunciated before, much more clearly and exhaustively; yet Euler remained supportive of Maupertuis. Euler’s noble way of reacting to Robin’s criticism of the *Mechanica* has already been mentioned in §3.2.1.

Sharing ideas with others and letting others take part in the process of discovery is another noble trait of Euler. A case in point is the way he put on hold his already extensive work on hydrodynamics, so that his friend Daniel Bernoulli, who was working on the same topic, could complete and publish his own *Hydrodynamics* first! It became a classic.

An important aspect of Euler’s personality is his religiosity: By his upbringing in the Riehen parish environment, he was a devout protestant and even served as an elder in one of the Protestant communities in Berlin. Indeed, he felt increasingly uncomfortable and frustrated in the company of so many “free-thinkers”—as he called them—, the believers in French enlightenment, that populated and began to dominate the Berlin Academy. He gave vent to his feelings in the (anonymously published) pamphlet *Rettung der göttlichen Offenbarung gegen die Einwürfe der Freygeister* (E92, *Opera omnia*, Ser. III, Vol. 12, pp. 267–286). This frustration may well have been one, and not the least, of the reasons why Euler left Berlin and returned to St. Petersburg.

## 4.2 Intellect

There are two outstanding qualities in Euler’s intellect: a phenomenal memory, coupled with an unusual power of mental calculation, and an ease in concentrating on mental work irrespective of any hustle and bustle going on around him: “A child on the knees, a cat on his back, that’s how he wrote his immortal works”, recounts Dieudonné Thiébauld, the French linguist and confidant of Frederick II. With regard to memory, the story is well known of Euler’s ability, even at an advanced age, to recite by heart all the verses of Virgil’s *Aeneid*. One of these, Euler says in a memoir, has given him the first ideas in solving a problem in mechanics. Niklaus Fuss also tells us that during a sleepless night, Euler mentally calculated the first six powers of all the numbers less than twenty (less than 100 in Concordet’s account), and



several days later was able to recall the answers without hesitation.

Equipped with such intellectual gifts, it is not surprising that Euler was extremely well read. In Fuchs's words, "he possessed to a high degree what commonly is called erudition; he had read the best writers of antique Rome; the older mathematical literature was very well known to him; he was well versed in the history of all times and all people. Even about medical and herbal remedies, and chemistry, he knew more than one could expect from a scholar who doesn't make these sciences a special subject of his study." Many visitors who came to see Euler went away "... with a mixture of astonishment and admiration. They could not understand how a man who during half a century seemed to have occupied himself solely with discoveries in the natural sciences and mathematics could retain so many facts that to him were useless and foreign to the subject of his researches."

### 4.3 Craftsmanship

Euler's writings had the marks of a superb expositor. He always strove for utmost clarity and simplicity, and he often revisited earlier work when he felt they were lacking in these qualities. At the same time, in his quest for discovery, he could be fearless, even reckless, but owing to his secure instinct, he rarely went astray when his argumentation became shaky. He had an eye for what is essential and unifying. In mechanics, as writes Gleb Konstantinovich Mikhailov [11, p. 67], "... Euler possessed a rare gift of systematizing and generalizing scientific ideas, which allowed him to present large parts of mechanics in a relatively definitive form." Euler was open and receptive to new ideas. In the words of André Weil [16, pp. 132–133], "... what at first is striking about Euler is his extraordinary quickness in catching hold of any suggestion, wherever it came from ... There is not one of these suggestions which in Euler's hands has not become the point of departure of an impressive series of researches ... Another thing, not less striking, is that Euler never abandons a research topic, once it has excited his curiosity; on the contrary, he returns to it, relentlessly, in order to deepen and broaden it on each revisit. Even if all problems related to such a topic seem to be resolved, he never ceases until the end of his life to find proofs that are 'more natural', 'simpler', 'more direct'."

## 4.4 Epilogue

In closing, let me cite the text—concise but to the point—that Otto Spiess had inscribed on a memorial plaque attached next to the house in which Euler grew up:



Figure 24: Euler memorial plaque in Riehen

LEONHARD EULER  
1707–1783  
mathematician, physicist, engineer,  
astronomer and philosopher, spent his  
youth in Riehen. He was a great scholar  
and a kind man.

## 5 Further reading

For readers interested in more details, we recommend the authoritative scientific (yet formula-free!) biography by Fellmann [7]. More recently, a number of popular and semipopular accounts on Euler and parts of his work have appeared; we mention Dunham [4], [5], Nahin [12], Varadarajan [15], and Sandifer [13].

The web site of the U.S. Euler Archive,

<http://www.math.dartmouth.edu/~euler>,

also provides detailed information of Euler's complete works.

## References

- [1] ASSAYAG, GERARD, FEICHTINGER, HANS GEORG, AND RODRIGUES, JOSE FRANCISCO (eds.) 2002. *Mathematics and music: A Diderot mathematical forum*. Springer, Berlin.
- [2] BAILHACHE, PATRICE 1995. Deux mathématiciens musiciens: Euler et d’Alembert. *Physis Riv. Internaz. Storia Sci. (N.S.)* 32(1), 1–35.
- [3] BOTTA, NIOCOLA 1995 *Numerical investigations of two-dimensional Euler flows: cylinder at transonic speed*, Ph. D. Dissertation, Swiss Federal Institute of Technology Zurich, xxii+153 pp.
- [4] DUNHAM, WILLIAM 1999. *Euler: the master of us all*. The Dolciani Mathematical Expositions, vol. 22, Mathematical Association of America, Washington, DC.
- [5] DUNHAM, WILLIAM, ED. 2007. *The genius of Euler: reflections on his life and work*. The MAA Tercentenary Euler Celebration, vol. 2, The Mathematical Association of America, Washington, DC.
- [6] FELLMANN, EMIL A. 1983 Leonhard Euler—Ein Essay über Leben und Werk, in *Leonhard Euler 1707–1783: Beiträge zu Leben und Werk*, Gedenkband des Kantons Basel-Stadt, Birkhäuser, Basel, pp. 13–98.
- [7] FELLMANN, EMIL A. 1995. *Leonhard Euler*. Rowohlt, Reinbek bei Hamburg. [English translation by Birkhäuser, Basel, 2007.]
- [8] GAUTSCHI, WALTER 2007. On Euler’s attempt to compute logarithms by interpolation: A commentary to his letter of February 16, 1734 to Daniel Bernoulli. *J. Comput. Appl. Math.*, to appear.
- [9] KOELINK, E. AND VAN ASSCHE, W. 2007 Leonhard Euler and a  $q$ -analogue of the logarithm, submitted for publication.
- [10] KRÄMER, STEFAN 2005. *Die Eulersche Konstante  $\gamma$  und verwandte Zahlen: eine mathematische und historische Betrachtung*, Diplomarbeit Universität Göttingen, Göttingen.
- [11] MIKHAILOV, GLEB KONSTANTINOVICH 1985. Euler und die Entwicklung der Mechanik. *Ceremony and scientific conference on the occasion*

*of the 200th anniversary of the death of Leonhard Euler (Berlin, 1983)*, pp. 64–82, Abh. Akad. Wiss. DDR, Abt. Math. Naturwiss. Tech., 85-1, Akademie Verlag, Berlin, 1985.

- [12] NAHIN, PAUL J. 2006. *Dr. Euler's fabulous formula. Cures many mathematical ills*. Princeton University Press, Princeton, NJ.
- [13] SANDIFER, C. EDWARD 2007. *The early mathematics of Leonhard Euler*. The MAA Tercentenary Euler Celebration, vol. 1, The Mathematical Association of America, Washington, DC.
- [14] SMITH, CHARLES SAMUEL 1960. *Leonhard Euler's Tentamen novae theoriae musicae: A translation and commentary*, Ph.D. Thesis, Indiana University. [Accessible through UMI Dissertation Services, Ann Arbor, MI.]
- [15] VARADARAJAN, V. S. 2006. *Euler through time: A new look at old themes*. Amer. Math. Soc., Providence, RI.
- [16] WEIL, ANDRÉ 1983. L'œuvre arithmétique d'Euler, in: *Leonhard Euler 1707–1783: Beiträge zu Leben und Werk*, Gedenkband des Kantons Basel-Stadt, Birkhäuser, Basel, pp. 111–133.