

1

Rapid #: -2663496



Ariel

IP: 129.2.17.58



Status	Rapid Code	Branch Name	Start Date
Pending	IND	Math	7/13/2009 8:30:46 AM

CALL #: QA 1 .Ac82
LOCATION: IND :: Math :: MATH PER
 TYPE: Article CC:CCL
 JOURNAL TITLE: Acta mathematica Academiae Scientiarum Hungaricae
 USER JOURNAL TITLE: Acta Math. Sci. Hung.
 IND CATALOG TITLE: Acta mathematica Academiae Scientiarum Hungaricae.
 ARTICLE TITLE: On difference sets of sequences of integers I
 ARTICLE AUTHOR: Sarkozy
 VOLUME: 31
 ISSUE:
 MONTH:
 YEAR: 1977
 PAGES: 125-
 ISSN: 0001-5954
 OCLC #:
 CROSS REFERENCE ID: 414855
 VERIFIED:

BORROWER: UMC :: Main Library
PATRON: Gasarch, William

PATRON ID: -
 PATRON ADDRESS: -
 PATRON PHONE: -
 PATRON FAX: -
 PATRON E-MAIL: -
 PATRON DEPT: -
 PATRON STATUS: -
 PATRON NOTES: -



This material may be protected by copyright law (Title 17 U.S. Code)
 System Date/Time: 7/13/2009 8:46:04 AM MST

ON DIFFERENCE SETS OF SEQUENCES OF INTEGERS. I

By

A. SÁRKÖZY (Budapest)

1. A set of integers $u_1 < u_2 < \dots$ will be called an \mathcal{A} -set if its difference set does not contain the square of a positive integer; in other words, if $u_x - u_y = z^2$ (where x, y, z are integers) implies that $x=y, z=0$. Let $A(x)$ denote the greatest number of integers that can be selected from $1, 2, \dots, x$ to form an \mathcal{A} -set and let us write

$$a(x) = \frac{A(x)}{x}.$$

L. Lovász conjectured that

$$(1) \quad a(x) = o(1)$$

(oral communication). The aim of this paper is to prove the following sharper form of (1):

THEOREM.

$$(2) \quad a(x) = O\left(\frac{(\log \log x)^{2/3}}{(\log x)^{1/3}}\right).$$

(We remark that (1) has been proved independently also by H. Fürstenberg; his proof is unpublished yet.)

To prove this theorem, we are going to use that version of the Hardy-Littlewood method which has been elaborated by K. F. ROTH in [2] and [3].

Throughout this paper, we use the following notations:

We denote the distance of the real number x from the nearest integer by $\|x\|$, i.e. $\|x\| = \min\{x - [x], [x] + 1 - x\}$. We write $e(\alpha) = e^{2\pi i \alpha}$ where α is real. $L_0, L_1, \dots, X_0, X_1, \dots$ denote absolute constants. If a, b are real numbers and $b > 0$, then we define the symbol $\min\left\{a, \frac{b}{0}\right\}$ by

$$(3) \quad \min\left\{a, \frac{b}{0}\right\} = a.$$

Finally, if $|g(x_1, x_2, \dots, x_n)| \cong f(x_1, x_2, \dots, x_n)$ then we write

$$g(x_1, x_2, \dots, x_n) = \theta(f(x_1, x_2, \dots, x_n)).$$

2. Following Roth's method, we are going to deduce a functional inequality for the function $a(x)$.

Let N be a large integer and let us write $M = \lfloor \sqrt{N} \rfloor$. Let

$$T(\alpha) = \sum_{z=1}^{\lfloor \sqrt{N} \rfloor} e(z^2 \alpha) = \sum_{z=1}^M e(z^2 \alpha).$$

Let $u_1, u_2, \dots, u_{A(N)}$ be a maximal \mathcal{A} -set selected from $1, 2, \dots, N$ and let

$$F(\alpha) = \sum_{x=1}^{A(N)} e(u_x \alpha).$$

We are going to investigate the integral

$$(4) \quad E = \int_0^1 |F(\alpha)|^2 T(\alpha) d\alpha.$$

Obviously,

$$(5) \quad \begin{aligned} E &= \int_0^1 F(\alpha) F(-\alpha) T(\alpha) d\alpha = \\ &= \int_0^1 \sum_{y=1}^{A(N)} e(u_y \alpha) \sum_{x=1}^{A(N)} e(-u_x \alpha) \sum_{z=1}^M e(z^2 \alpha) d\alpha = \sum_{\substack{x, y, z \\ u_y - u_x + z^2 = 0}} 1 = 0 \end{aligned}$$

since $u_1, u_2, \dots, u_{A(N)}$ is an \mathcal{A} -set.

On the other hand, we shall estimate this integral by using the Hardy—Littlewood method. For this purpose, we need some estimates for the functions $T(\alpha)$ and $F(\alpha)$.

3. In this section, we estimate the function $T(\alpha)$.

LEMMA 1. *If a, b are integers such that $a \leq b$, and β is an arbitrary real number then*

$$\left| \sum_{k=a}^b e(k\beta) \right| \leq \min \left\{ b-a+1, \frac{1}{2\|\beta\|} \right\}.$$

(For $\|\beta\|=0$, the right hand side is defined by (3).)

PROOF. Obviously,

$$\left| \sum_{k=a}^b e(k\beta) \right| \leq \sum_{k=a}^b 1 \leq b-a+1$$

for all a, b, β (where $a \leq b$).

Furthermore, for $\|\beta\| \neq 0$,

$$\begin{aligned} \left| \sum_{k=a}^b e(k\beta) \right| &= \frac{|1 - e((b-a+1)\beta)|}{|1 - e(\beta)|} \leq \frac{2}{|1 - e(\beta)|} = \\ &= \frac{2}{|e(-\beta/2) - e(\beta/2)|} = \frac{1}{|\sin \pi \beta|} = \frac{1}{\sin \pi \|\beta\|} \leq \frac{1}{\frac{2}{\pi} \cdot \pi \|\beta\|} = \frac{1}{2\|\beta\|} \end{aligned}$$

which proves Lemma 1.

LEMMA 2. Let p, q be integers and α, γ real numbers such that $q > 1, (p, q) = 1$ and

$$\left| \alpha - \frac{p}{q} \right| \leq \frac{1}{q^2}.$$

Then

$$\sum_{x=0}^{q-1} \min \left\{ q, \frac{1}{\|\gamma + \alpha x\|} \right\} < 8q \log q.$$

This lemma is identical to Theorem 44 in [1], p. 26.

LEMMA 3. Let p, q be integers and α, γ, P real numbers such that $q \geq 1, (p, q) = 1, P \geq 1$ and

$$(6) \quad \left| \alpha - \frac{p}{q} \right| \leq \frac{1}{q^2}.$$

Then

$$(7) \quad \sum_{x=0}^{q-1} \min \left\{ P, \frac{1}{\|\gamma + \alpha x\|} \right\} < 8P + 8q \log q.$$

PROOF. For $q=1$, (7) holds trivially (by $P \geq 1$).

For $q > 1$, the left hand side of (7) can be rewritten in the following way:

$$\begin{aligned} (8) \quad & \sum_{x=0}^{q-1} \min \left\{ P, \frac{1}{\|\gamma + \alpha x\|} \right\} = \\ & = \sum_{\substack{0 \leq x \leq q-1 \\ \|\gamma + \alpha x\| < \frac{1}{q}}} \min \left\{ P, \frac{1}{\|\gamma + \alpha x\|} \right\} + \sum_{\substack{0 \leq x \leq q-1 \\ \|\gamma + \alpha x\| \geq \frac{1}{q}}} \min \left\{ P, \frac{1}{\|\gamma + \alpha x\|} \right\} \cong \\ & \cong \sum_{\substack{0 \leq x \leq q-1 \\ \|\gamma + \alpha x\| < \frac{1}{q}}} P + \sum_{\substack{0 \leq x \leq q-1 \\ \|\gamma + \alpha x\| \geq \frac{1}{q}}} \frac{1}{\|\gamma + \alpha x\|} = P \sum_{\substack{0 \leq x \leq q-1 \\ \|\gamma + \alpha x\| < \frac{1}{q}}} 1 + \sum_{\substack{0 \leq x \leq q-1 \\ \|\gamma + \alpha x\| \geq \frac{1}{q}}} \min \left\{ q, \frac{1}{\|\gamma + \alpha x\|} \right\} \cong \\ & \cong P \sum_{\substack{0 \leq x \leq q-1 \\ \|\gamma + \alpha x\| < \frac{1}{q}}} 1 + \sum_{x=0}^{q-1} \min \left\{ q, \frac{1}{\|\gamma + \alpha x\|} \right\}. \end{aligned}$$

We are going to show that

$$(9) \quad \sum_{\substack{0 \leq x \leq q-1 \\ \|\gamma + \alpha x\| < \frac{1}{q}}} 1 \leq 8.$$

Let us assume indirectly that the left hand side of this inequality is ≥ 9 . This indirect assumption implies the existence of integers x_1, x_2, x_3, x_4, x_5 such that

$$(10) \quad \|\gamma + \alpha x_i\| < \frac{1}{q} \quad (\text{for } i = 1, \dots, 5)$$

and which all lie either in the interval $\left[0, \frac{q-1}{2}\right]$ or in $\left[\frac{q-1}{2}, q-1\right]$; in either case

$$(11) \quad 0 < |x_i - x_j| < \frac{q}{2} \quad (\text{for } 1 \leq i \leq j < 5).$$

By (10), there exist integers y_i and real numbers θ_i such that

$$(12) \quad \gamma + \alpha x_i = y_i + \frac{\theta_i}{q} \quad (\text{for } i = 1, \dots, 5)$$

and

$$-1 < \theta_i < 1 \quad (\text{for } i = 1, \dots, 5).$$

Using the matchbox principle, we obtain the existence of indices μ, ν such that $1 \leq \mu < \nu \leq 5$ and

$$(13) \quad |\theta_\mu - \theta_\nu| < \frac{1}{2}.$$

Writing $i = \mu$ and ν in (12), respectively, and subtracting the equalities obtained in this way, we obtain that

$$\alpha(x_\mu - x_\nu) = y_\mu - y_\nu + \frac{\theta_\mu - \theta_\nu}{q}.$$

Hence

$$(14) \quad \|\alpha(x_\mu - x_\nu)\| \leq \frac{\theta_\mu - \theta_\nu}{q} < \frac{1}{2q}$$

(by (13)).

On the other hand, we obtain with respect to (6) and (11) that

$$\begin{aligned} \|\alpha(x_\mu - x_\nu)\| &= \left\| \frac{p}{q}(x_\mu - x_\nu) + \left(\alpha - \frac{p}{q}\right)(x_\mu - x_\nu) \right\| \leq \\ &\leq \left\| \frac{p(x_\mu - x_\nu)}{q} \right\| - \left| \alpha - \frac{p}{q} \right| |x_\mu - x_\nu| > \frac{1}{q} - \frac{1}{q^2} \cdot \frac{q}{2} = \frac{1}{2q} \end{aligned}$$

in contradiction with (14), which proves (9).

(8), (9) and Lemma 2 yield (7) and Lemma 3 is proved.

LEMMA 4. Let N, p, q be integers and α a real number such that $N \geq 1$, $q \geq 1$, $(p, q) = 1$ and

$$\left| \alpha - \frac{p}{q} \right| \leq \frac{1}{q^2}.$$

Then

$$(15) \quad |T(\alpha)| = \left| \sum_{k=1}^M e(k^2 \alpha) \right| < 7 \left(\frac{M}{q^{1/2}} + (M \log q)^{1/2} + (q \log q)^{1/2} \right)$$

(where $M = \lfloor \sqrt{N} \rfloor$).

PROOF.

$$\begin{aligned}
|T(\alpha)|^2 &= T(\alpha)T(-\alpha) = \sum_{x=1}^M \sum_{y=1}^M e((x^2-y^2)\alpha) = \\
&= \sum_{x=1}^M \sum_{y=1}^M e((x-y)(x+y)\alpha) = \sum_{u=1-M}^{M-1} \sum_{y=\max\{1-u, 1\}}^{\min\{M, M-u\}} e(u(u+2y)\alpha) \cong \\
&\cong \sum_{u=1-M}^{M-1} \left| \sum_{y=\max\{1-u, 1\}}^{\min\{M, M-u\}} e(u(u+2y)\alpha) \right| = \sum_{u=1-M}^{M-1} \left| \sum_{y=\max\{1-u, 1\}}^{\min\{M, M-u\}} e(2uy\alpha) \right|.
\end{aligned}$$

To estimate the inner sum, we apply Lemma 1 with $\beta=2ux$, $a=\max\{1-u, 1\}$, $b=\min\{M, M-u\}$. Then obviously,

$$b-a = \min\{M, M-u\} - \max\{1-u, 1\} \cong M-1,$$

thus Lemma 1 yields that

$$\begin{aligned}
|T(\alpha)|^2 &\cong \sum_{u=1-M}^{M-1} \min \left\{ (M-1)+1, \frac{1}{2\|2u\alpha\|} \right\} = \\
&= \frac{1}{2} \sum_{u=1-M}^{M-1} \min \left\{ 2M, \frac{1}{\|2u\alpha\|} \right\} \cong \frac{1}{2} \sum_{u=2-2M}^{2M-2} \min \left\{ 2M, \frac{1}{\|u\alpha\|} \right\} \cong \\
&\cong \frac{1}{2} \sum_{j=0}^{\lfloor (4M-4)/q \rfloor} \sum_{v=2-2M+jq}^{2-2M+(j+1)q-1} \min \left\{ 2M, \frac{1}{\|v\alpha\|} \right\}.
\end{aligned}$$

The inner sum can be estimated by using Lemma 3 with $\gamma=(2-2M+jq)\alpha$, $P=2M$. We obtain that

$$\begin{aligned}
(16) \quad |T(\alpha)|^2 &\cong \frac{1}{2} \sum_{j=0}^{\lfloor (4M-4)/q \rfloor} (16M+8q \log q) = \\
&= \frac{1}{2} \left(\left\lfloor \frac{4M-4}{q} \right\rfloor + 1 \right) (16M+8q \log q) < \left(\frac{4M}{q} + 1 \right) (8M+4q \log q) = \\
&= 32 \frac{M^2}{q} + 8M + 16M \log q + 4q \log q.
\end{aligned}$$

For $q=1$,

$$M \cong M \cdot \frac{M}{q} = \frac{M^2}{q},$$

while for $q \cong 2$

$$M < M \log 4 = 2M \log 2 \cong 2M \log q.$$

Hence

$$M \cong \frac{M^2}{q} + 2M \log q.$$

Thus we obtain from (16) that

$$\begin{aligned} |T(\alpha)|^2 &< 32 \frac{M^2}{q} + 8 \left(\frac{M^2}{q} + 2M \log q \right) + 16M \log q + 4q \log q = \\ &= 40 \frac{M^2}{q} + 32M \log q + 4q \log q < 49 \left(\frac{M^2}{q} + M \log q + q \log q \right). \end{aligned}$$

With respect to the inequality

$$(a^2 + b^2 + c^2)^{1/2} \cong a + b + c \quad (\text{where } a, b, c \cong 0),$$

this yields (15) which proves Lemma 4.

LEMMA 5. Let N, p, q be integers and α a real number such that $(p, q) = 1$,

$$(17) \quad N \cong 3,$$

$$(18) \quad 1 \cong q \cong N^{1/2} / \log N$$

and

$$(19) \quad \left| \alpha - \frac{p}{q} \right| < \frac{1}{q^2}.$$

Then

$$|T(\alpha)| < 21 \left(\frac{N}{q} \right)^{1/2}.$$

PROOF. Applying Lemma 4, we obtain with respect to (17) and (18) that

$$\begin{aligned} |T(\alpha)| &< 7 \left(\frac{M}{q^{1/2}} + (M \log q)^{1/2} + (q \log q)^{1/2} \right) \cong \\ &\cong 7 \left(\left(\frac{N}{q} \right)^{1/2} + N^{1/4} (\log q)^{1/2} + (q \log q)^{1/2} \right) < \\ &< 7 \left(\left(\frac{N}{q} \right)^{1/2} + N^{1/4} (\log N)^{1/2} + (N^{1/2} \log N)^{1/2} \right) = \\ &= 7 \left(\frac{N}{q} \right)^{1/2} \left(1 + 2 \left(q \frac{\log N}{N^{1/2}} \right)^{1/2} \right) \cong 7 \left(\frac{N}{q} \right)^{1/2} \cdot 3 = 21 \left(\frac{N}{q} \right)^{1/2}. \end{aligned}$$

LEMMA 6. Let N, p, q be integers and α, β real numbers such that

$$(20) \quad N \cong 9,$$

$$(21) \quad (p, q) = 1,$$

$$(22) \quad 1 \cong q \cong \sqrt{N},$$

$$(23) \quad \alpha = \frac{p}{q} + \beta$$

and

$$(24) \quad \frac{\log N}{N} \cong |\beta| < \frac{1}{2q\sqrt{N}}.$$

Then

$$(25) \quad |T(\alpha)| < 30 \left(\frac{\log N}{q|\beta|} \right)^{1/2}.$$

PROOF. Let

$$Q = \left[\frac{1}{q|\beta|} \right] + 1.$$

Then obviously,

$$(26) \quad Q > \frac{1}{q|\beta|},$$

hence

$$(27) \quad |\beta| > \frac{1}{qQ}.$$

By (24),

$$(28) \quad \frac{1}{q|\beta|} > \sqrt{N}.$$

Thus

$$(29) \quad Q = \left[\frac{1}{q|\beta|} \right] + 1 \cong \frac{1}{q|\beta|} + \sqrt{N} < \frac{1}{q|\beta|} + \frac{1}{q|\beta|} = \frac{2}{q|\beta|}.$$

By Dirichlet's theorem, there exist integers r, s such that

$$(30) \quad (r, s) = 1,$$

$$(31) \quad 1 \cong s \cong Q$$

and

$$(32) \quad \left| \alpha - \frac{r}{s} \right| < \frac{1}{sQ}.$$

(31) and (32) imply that also

$$\left| \alpha - \frac{r}{s} \right| < \frac{1}{s^2}$$

holds. Thus we may apply Lemma 4 with $\frac{r}{s}$ in place of $\frac{p}{q}$. We obtain that

$$(33) \quad |T(\alpha)| < 7 \left(\frac{M}{s^{1/3}} + (M \log s)^{1/2} + (s \log s)^{1/2} \right) \cong \\ \cong 7 \left(\left(\frac{N}{s} \right)^{1/2} + N^{1/4} (\log s)^{1/2} + (s \log s)^{1/2} \right).$$

To deduce (25) from this inequality, we have to estimate s in terms of $q|\beta|$ and N , respectively.

By (29) and (31),

$$(34) \quad s \cong Q < \frac{2}{q|\beta|}$$

hence with respect to (20) and (24),

$$s < \frac{2}{q|\beta|} \cong \frac{2}{|\beta|} \cong 2 \frac{N}{\log N} < N.$$

Thus (33) implies that

$$(35) \quad |T(\alpha)| < 7 \left[\left(\frac{N}{s} \right)^{1/2} + (N^{1/4} + s^{1/2})(\log N)^{1/2} \right].$$

We are going to show that

$$(36) \quad \frac{p}{q} \neq \frac{r}{s}.$$

Let us assume indirectly that

$$(37) \quad \frac{p}{q} = \frac{r}{s}.$$

By $q \cong 1$, $s \cong 1$, (21) and (30), this implies also $q=s$. Thus in view of (23), (27) and (32)

$$\begin{aligned} \left| \frac{p}{q} - \frac{r}{s} \right| &= \left| \left(\frac{p}{q} - \alpha \right) + \left(\alpha - \frac{r}{s} \right) \right| \cong \left| \alpha - \frac{p}{q} \right| - \left| \alpha - \frac{r}{s} \right| = \\ &= |\beta| - \left| \alpha - \frac{r}{s} \right| > \frac{1}{qQ} - \frac{1}{sQ} = 0 \end{aligned}$$

in contradiction with (37), which proves (36).

(36) implies that

$$(38) \quad \left| \frac{p}{q} - \frac{r}{s} \right| = \frac{|ps - qr|}{qs} \cong \frac{1}{qs}.$$

On the other hand, with respect to (22), (23), (24), (26) and (32),

$$(39) \quad \begin{aligned} \left| \frac{p}{q} - \frac{r}{s} \right| &= \left| \left(\frac{p}{q} - \alpha \right) + \left(\alpha - \frac{r}{s} \right) \right| \cong \left| \alpha - \frac{p}{q} \right| + \left| \alpha - \frac{r}{s} \right| = \\ &= |\beta| + \left| \alpha - \frac{r}{s} \right| < |\beta| + \frac{1}{sQ} < |\beta| + \frac{q|\beta|}{s} < |\beta| + \frac{1}{s} \cdot \frac{1}{2\sqrt{N}} \cong |\beta| + \frac{1}{2sq}. \end{aligned}$$

(38) and (39) yield that

$$\frac{1}{qs} \cong |\beta| + \frac{1}{2sq}.$$

Thus by (24),

$$(40) \quad s \cong \frac{1}{2q|\beta|} > \sqrt{N}.$$

In view of (20), (34) and (40), we obtain from (35) that

$$|T(\alpha)| < 7 \left[\left(\frac{N}{s} \right)^{1/2} + 2s^{1/2}(\log N)^{1/2} \right] = 7s^{1/2}(\log N)^{1/2} \left(\frac{N^{1/2}}{s(\log N)^{1/2}} + 2 \right) <$$

$$< 7s^{1/2}(\log N)^{1/2}(1+2) = 21s^{1/2}(\log N)^{1/2} < 21 \left(\frac{2}{q|\beta|} \right)^{1/2} (\log N)^{1/2} < 30 \left(\frac{\log N}{q|\beta|} \right)^{1/2}$$

and Lemma 6 is proved.

LEMMA 7. Let N, p, q be integers, R, Q, α real numbers such that $N \geq 1, (p, q) = 1,$

$$(41) \quad 1 \leq R \leq q \leq Q,$$

$$(42) \quad \sqrt{N} \leq Q \leq N$$

and

$$(43) \quad \left| \alpha - \frac{p}{q} \right| < \frac{1}{qQ}.$$

Then

$$(44) \quad |T(\alpha)| < 7 \left(\frac{N}{R} \right)^{1/2} + 14(Q \log N)^{1/2}.$$

PROOF. (41) and (43) imply also

$$\left| \alpha - \frac{p}{q} \right| < \frac{1}{q^2}.$$

Thus with respect to (41) and (42), Lemma 4 yields that

$$\begin{aligned} |T(\alpha)| &< 7 \left(\frac{N^{1/2}}{q^{1/2}} + (N^{1/2} \log q)^{1/2} + (q \log q)^{1/2} \right) \leq \\ &\leq 7 \left(\left(\frac{N}{R} \right)^{1/2} + (Q \log N)^{1/2} + (Q \log N)^{1/2} \right) \end{aligned}$$

which proves (44).

4. In this section, we estimate the function $F(\alpha)$ by using Roth's method.

Lemmas 8 and 9 follow trivially from the definitions of the functions $A(x)$ and $a(x)$.

LEMMA 8. If x, y are positive integers such that $x \leq y$ then $A(x) \leq A(y)$.

LEMMA 9. For any positive integer $x,$

$$\frac{1}{x} \leq a(x) \leq 1.$$

For any integer b and positive integers $q, x,$ let $A_{(b,q)}(x)$ denote the greatest number of integers that can be selected from $b+q, b+2q, \dots, b+xq$ to form an \mathcal{A} -set.

LEMMA 10. For any integer b and positive integers q, x ,

$$A_{(b, q^2)}(x) = A(x).$$

PROOF. This follows trivially from the fact that the numbers $b + u_1 q^2, b + u_2 q^2, \dots, b + u_k q^2$ form an \mathcal{A} -set if and only if also the numbers u_1, u_2, \dots, u_k do.

LEMMA 11. For any positive integers x and y , we have

$$(45) \quad A(x+y) \cong A(x) + A(y),$$

$$(46) \quad A(xy) \cong xA(y),$$

$$(47) \quad a(xy) \cong a(y),$$

$$(48) \quad a(x) \cong \left(1 + \frac{y}{x}\right) a(y).$$

PROOF. Applying Lemma 10 with $b=y, q=1$, we obtain that the greatest number of integers that can be selected from $y+1, y+2, \dots, y+x$ to form an \mathcal{A} -set is $\cong A(x)$. (45) follows easily from this fact.

(46) is a consequence of (45).

(47) can be obtained from (46) by dividing by xy .

Finally, by Lemma 8 and (46),

$$A(x) \cong A\left(\left(\left[\frac{x}{y}\right] + 1\right)y\right) \cong \left(\left[\frac{x}{y}\right] + 1\right)A(y) \cong \left(\frac{x}{y} + 1\right)A(y).$$

Dividing by x , we obtain (48).

LEMMA 12. Let q, t, N be positive integers, p integer, α, β real numbers such that

$$(49) \quad \alpha - \frac{p}{q} = \beta.$$

Let

$$F_1(\alpha) = \frac{a(t)}{q^2} \left(\sum_{s=1}^{q^2} e\left(\frac{sp}{q}\right) \right) \left(\sum_{j=1}^N e(\beta j) \right),$$

so that if $(p, q)=1$ then

$$(50) \quad F_1(\alpha) = \begin{cases} a(t) \sum_{j=1}^N e(j\alpha) & \text{for } q=1 \\ 0 & \text{for } q>1 \end{cases} \quad (\text{where } (p, q)=1).$$

Then

$$(51) \quad |F(\alpha) - F_1(\alpha)| \cong (a(t) - a(N))N + 2a(t)tq^2(1 + \pi|\beta|N) = H(t, N, q, \beta).$$

PROOF. We are going to show at first that

$$(52) \quad F(\alpha) = \frac{1}{tq^2} \sum_{s=1}^{q^2} \sum_{j=1}^N \sum_{\substack{j \cong u_k < j + tq^2 \\ u_k \equiv s \pmod{q^2}}} e(\alpha u_k) + \theta(a(t)tq^2).$$

Let us investigate the coefficient of $e(\alpha u_k)$ on the right hand side.

If $tq^2 \equiv u_k \equiv N$ then we account $e(\alpha u_k)$ exactly tq^2 times, namely for the following values of j :

$$j = u_k - tq^2 + 1, \quad u_k - tq^2 + 2, \dots, u_k.$$

Thus the coefficient of $e(\alpha u_k)$ is

$$tq^2 \cdot \frac{1}{tq^2} = 1$$

in this case (and its coefficient is the same on the left hand side).

If

$$(53) \quad 1 \equiv u_k < tq^2$$

then we account $e(\alpha u_k)$ on the right of (52) for $j = 1, 2, \dots, u_k$, thus its coefficient is

$$(0 \equiv) u_k \cdot \frac{1}{tq^2} < tq^2 \cdot \frac{1}{tq^2} = 1$$

on the right and 1 on the left of (52). The numbers u_k satisfying (53) form an \mathcal{A} -set thus in view of (46) in Lemma 11, their number is

$$\equiv A(tq^2) \equiv A(t)q^2 = a(t)tq^2.$$

These facts yield that, in fact, the error term in (52) is $\theta(a(t)tq^2)$.

The term $e(\alpha u_k)$ in the inner sum in (52) can be rewritten in the following way:

$$\begin{aligned} e(\alpha u_k) &= e\left(\left(\frac{p}{q} + \beta\right)u_k\right) = e\left(\frac{pu_k}{q}\right) e(\beta u_k) = \\ &= e\left(\frac{ps}{q}\right) e(\beta j) e(\beta(u_k - j)) = e\left(\frac{ps}{q}\right) e(\beta j) (1 + \theta(2\pi|\beta(u_k - j)|)) = \\ &= e\left(\frac{ps}{q}\right) e(\beta j) + \theta(2\pi|\beta(u_k - j)|) = e\left(\frac{ps}{q}\right) e(\beta j) + \theta(2\pi|\beta|tq^2) \end{aligned}$$

since $|u_k - j| < tq^2$ in the inner sum, and

$$|e(\gamma) - 1| = |e(\gamma/2) - e(-\gamma/2)| = |2 \sin \pi\gamma| \equiv 2|\pi\gamma| = 2\pi|\gamma|$$

for any real number γ .

Thus the inner sum in (52) can be estimated in the following way:

$$(54) \quad \sum_{\substack{j \equiv u_k - j + tq^2 \\ u_k \equiv s \pmod{q^2}}} e(\alpha u_k) = \sum_{\substack{j \equiv u_k - j + tq^2 \\ u_k \equiv s \pmod{q^2}}} \left(e\left(\frac{ps}{q}\right) e(\beta j) + \theta(2\pi|\beta|tq^2) \right) = \\ = \left(e\left(\frac{ps}{q}\right) e(\beta j) + \theta(2\pi|\beta|tq^2) \right) \sum_{\substack{j \equiv u_k - j + tq^2 \\ u_k \equiv s \pmod{q^2}}} 1.$$

Let us define the integer v by

$$v < j \equiv v + q^2, \quad v \equiv s \pmod{q^2}.$$

Then by Lemma 10,

$$\sum_{\substack{j \equiv u_k < j + tq^2 \\ u_k \equiv s \pmod{q^2}}} 1 \equiv A_{(v, q^2)}(t) \equiv A(t) = a(t)t.$$

Thus if we define $D(j, t, q, s)$ by

$$\sum_{\substack{j \equiv u_k < j + tq^2 \\ u_k \equiv s \pmod{q^2}}} 1 = a(t)t - D(j, t, q, s)$$

then $D(j, t, q, s) \geq 0$. Putting this into (54):

$$\begin{aligned} \sum_{\substack{j \equiv u_k < j + tq^2 \\ u_k \equiv s \pmod{q^2}}} e(\alpha u_k) &= \left(e\left(\frac{ps}{q}\right) e(\beta j) + \theta(2\pi|\beta|tq^2) \right) (a(t)t - D(j, t, q, s)) = \\ &= e\left(\frac{ps}{q}\right) e(\beta j) (a(t)t - D(j, t, q, s)) + \theta(2\pi|\beta|a(t)t^2q^2). \end{aligned}$$

Thus (52) yields that

$$\begin{aligned} (55) \quad F(\alpha) &= \frac{1}{tq^2} \sum_{s=1}^{q^2} \sum_{j=1}^N \left\{ e\left(\frac{ps}{q}\right) e(\beta j) (a(t)t - D(j, t, q, s)) + \theta(2\pi|\beta|a(t)t^2q^2) \right\} + \\ &+ \theta(a(t)tq^2) = \frac{a(t)}{q^2} \left(\sum_{s=1}^{q^2} e\left(\frac{ps}{q}\right) \right) \left(\sum_{j=1}^N e(\beta j) \right) - \frac{1}{tq^2} \sum_{s=1}^{q^2} \sum_{j=1}^N e\left(\frac{ps}{q}\right) e(\beta j) D(j, t, q, s) + \\ &+ \theta\left(\frac{1}{tq^2} \cdot q^2 \cdot N \cdot 2\pi|\beta|a(t)t^2q^2\right) + \theta(a(t)tq^2) = \\ &= F_1(\alpha) - \frac{1}{tq^2} \sum_{s=1}^{q^2} \sum_{j=1}^N e\left(\frac{ps}{q}\right) e(\beta j) D(j, t, q, s) + \theta(2\pi|\beta|Na(t)tq^2) + \theta(a(t)tq^2). \end{aligned}$$

Putting here $\alpha = \beta = p = 0$, we obtain that

$$A(N) = a(t) \cdot N - \frac{1}{tq^2} \sum_{s=1}^{q^2} \sum_{j=1}^N D(j, t, q, s) + \theta(a(t)tq^2).$$

Hence

$$(56) \quad \frac{1}{tq^2} \sum_{s=1}^{q^2} \sum_{j=1}^N D(j, t, q, s) \equiv a(t) \cdot N - A(N) + a(t)tq^2 = \\ = (a(t) - a(N))N + a(t)tq^2.$$

With respect to (56), (55) yields that

$$\begin{aligned} |F(\alpha) - F_1(\alpha)| &\equiv \frac{1}{tq^2} \sum_{s=1}^{q^2} \sum_{j=1}^N D(j, t, q, s) + 2\pi|\beta|Na(t)tq^2 + a(t)tq^2 \equiv \\ &\equiv (a(t) - a(N))N + a(t)tq^2 + 2\pi|\beta|Na(t)tq^2 + a(t)tq^2 = \\ &= (a(t) - a(N)) + 2a(t)tq^2(1 + \pi|\beta|N) \end{aligned}$$

which proves Lemma 12.

5. In this section, we are going to deduce a functional inequality for the function $a(x)$, by investigating the integral E defined in Section 2. This inequality will contain four parameters: t, N, R, Q whose values will be fixed later.

LEMMA 13. Let t, N be positive integers, R, Q real numbers such that

$$(57) \quad N \cong e^8,$$

$$(58) \quad t/N,$$

$$(59) \quad 1 \cong R \cong N^{1/2}/\log N,$$

$$(60) \quad 2N^{1/2} < Q \cong \frac{N}{\log N}.$$

Then

$$(61) \quad \begin{aligned} a^2(t)N^{3/2} &< 1260a(t)(a(t)-a(N))N^{3/2}\log\log N + \\ &+ 12600a^2(t)tN(\log N)^{1/2}Q^{-1/2} + \\ &+ 120(a(t)-a(N))^2\{7N^{3/2}R^{3/2}\log N + 20N^2(\log N)^{1/2}Q^{-1/2}R\} + \\ &+ 26000a^2(t)t^2\{3N^{-1/2}(\log N)^3R^{11/2} + 2N^2(\log N)^{1/2}Q^{-5/2}R^3\} + \\ &+ 140a(t)\{N^{3/2}R^{-1/2} + 2NQ^{1/2}(\log N)^{1/2}\}. \end{aligned}$$

PROOF. Let us write

$$G(\alpha) = a(t) \sum_{j=1}^N e(j\alpha).$$

Then

$$E = \int_0^1 |F(\alpha)|^2 T(\alpha) d\alpha = \int_0^1 |G(\alpha)|^2 T(\alpha) d\alpha + \int_0^1 (|F(\alpha)|^2 - |G(\alpha)|^2) T(\alpha) d\alpha$$

where $E=0$ by (5). Hence

$$(62) \quad \int_0^1 |G(\alpha)|^2 T(\alpha) d\alpha = - \int_0^1 (|F(\alpha)|^2 - |G(\alpha)|^2) T(\alpha) d\alpha.$$

Here

$$(63) \quad \begin{aligned} \int_0^1 |G(\alpha)|^2 T(\alpha) d\alpha &= \int_0^1 \left(a(t) \sum_{j=1}^N e(j\alpha) \right) \left(a(t) \sum_{k=1}^N e(-k\alpha) \right) \left(\sum_{z=1}^{[N]} e(z^2\alpha) \right) d\alpha = \\ &= a^2(t) \int_0^1 \sum_{\substack{1 \leq j, k \leq N \\ 1 \leq z \leq \sqrt{N}}} e((j-k+z^2)\alpha) d\alpha = a^2(t) \sum_{\substack{j-k+z^2=0 \\ 1 \leq j, k \leq N \\ 1 \leq z \leq \sqrt{N}}} 1. \end{aligned}$$

If

$$(64) \quad 1 \cong z^2 \cong \frac{N}{2} - 1, \quad z > 0,$$

$$(65) \quad 1 \cong j \cong \frac{N}{2} + 1$$

then the numbers j , $k=j+z^2$, z satisfy the conditions

$$j-k+z^2=0, \quad 1 \leq j, k \leq N, \quad 1 \leq z \leq \sqrt{N}$$

since

$$k=j+z^2 \leq \left(\frac{N}{2}+1\right) + \left(\frac{N}{2}-1\right) = N.$$

By (57), the number of the positive integers z satisfying (64) is at least

$$\left\lfloor \sqrt{\frac{N}{2}-1} \right\rfloor \geq \left\lfloor \sqrt{\frac{N}{2}-\frac{N}{4}} \right\rfloor = \left\lfloor \frac{\sqrt{N}}{2} \right\rfloor \geq \frac{\sqrt{N}}{2} - 1 \geq \frac{\sqrt{N}}{2} - \frac{\sqrt{N}}{4} = \frac{\sqrt{N}}{4}$$

while (65) holds for $\left\lfloor \frac{N}{2} \right\rfloor + 1 > \frac{N}{2}$ integers j . Thus (63) yields that

$$(66) \quad \int_0^1 |G(\alpha)|^2 T(\alpha) d\alpha = a^2(t) \sum_{\substack{j-k+z^2=0 \\ 1 \leq j, k \leq N \\ 1 \leq z \leq \sqrt{N}}} 1 > a^2(t) \cdot \frac{\sqrt{N}}{4} \cdot \frac{N}{2} = \frac{1}{8} a^2(t) N^{3/2}.$$

To complete the proof of (61), we have to give an upper estimate for the right hand side of (62).

For $q=1, 2, \dots, [Q]$ and $p=0, 1, \dots, q-1$, let

$$I_{p,q} = \left(\frac{p}{q} - \frac{1}{pQ}, \frac{p}{q} + \frac{1}{qQ} \right)$$

and let S denote the set of those real numbers α for which

$$-\frac{1}{Q} < \alpha \leq 1 - \frac{1}{Q}$$

holds and

$$(67) \quad \alpha \notin I_{p,q} \quad \text{for } 1 \leq q \leq R, \quad 0 \leq p \leq q-1, \quad (p, q) = 1.$$

Then

$$(68) \quad \left| \int_0^1 (|F(\alpha)|^2 - |G(\alpha)|^2) T(\alpha) d\alpha \right| = \\ = \left| \int_{-1/Q}^{1-1/Q} (|F(\alpha)|^2 - |G(\alpha)|^2) T(\alpha) d\alpha \right| \leq \int_{-1/Q}^{+1/Q} \left| |F(\alpha)|^2 - |G(\alpha)|^2 \right| |T(\alpha)| d\alpha + \\ + \sum_{q=2}^{[R]} \sum_{\substack{1 \leq p \leq q-1 \\ (p,q)=1}} \int_{I_{p,q}} \left| |F(\alpha)|^2 - |G(\alpha)|^2 \right| |T(\alpha)| d\alpha + \\ + \int_S \left| |F(\alpha)|^2 - |G(\alpha)|^2 \right| |T(\alpha)| d\alpha = E_1 + E_2 + E_3.$$

Let us estimate the term E_1 at first.

For any complex numbers u, v , we have

$$(69) \quad \begin{aligned} & \left| |u|^2 - |v|^2 \right| = |u\bar{u} - v\bar{v}| = |(u-v)\bar{u} + v(\bar{u} - \bar{v})| \leq \\ & \leq |u-v|(|\bar{u}| + |\bar{v}|) = |u-v|(|u| + |v|) = |u-v|(|(u-v) + v| + |v|) \leq \\ & \leq |u-v|(|u-v| + 2|v|) = |u-v|^2 + 2|u-v||v|. \end{aligned}$$

Furthermore, applying Lemma 12 with $p=0, q=1, \alpha=\beta$, we have $F_1(\alpha) = G(\alpha)$ there, thus we obtain that

$$(70) \quad |F(\alpha) - G(\alpha)| \leq H(t, N, 1, \alpha).$$

(69) and (70) yield that

$$(71) \quad \begin{aligned} E_1 &= \int_{-1/Q}^{+1/Q} \left| |F(\alpha)|^2 - |G(\alpha)|^2 \right| |T(\alpha)| d\alpha \leq \\ &\leq \int_{-1/Q}^{+1/Q} |F(\alpha) - G(\alpha)|^2 |T(\alpha)| d\alpha + 2 \int_{-1/Q}^{+1/Q} |F(\alpha) - G(\alpha)| |G(\alpha)| |T(\alpha)| d\alpha \leq \\ &\leq \int_{-1/Q}^{+1/Q} H^2(t, N, 1, \alpha) |T(\alpha)| d\alpha + 2 \int_{-1/Q}^{+1/Q} H(t, N, 1, \alpha) |G(\alpha)| |T(\alpha)| d\alpha = E_1' + 2E_1''. \end{aligned}$$

E_1' will be estimated simultaneously with E_2 ; here we estimate only E_1'' .

The function $|G(\alpha)|$ can be estimated by using Lemma 1. Furthermore, for $|\alpha| \leq \log N/N$, we use the trivial inequality

$$(72) \quad |T(\alpha)| = \left| \sum_{z=1}^M e(z^2\alpha) \right| \leq \sum_{z=1}^M 1 = M \leq N^{1/2},$$

while for $\log N/N < |\alpha| \leq 1/Q (< 1/2\sqrt{N})$, by (60), we apply Lemma 6. In this way, we obtain that

$$(73) \quad \begin{aligned} E_1'' &< \int_{|\alpha| \leq 1/N} \left\{ (a(t) - a(N))N + 2a(t)t \left(1 + \pi \cdot \frac{1}{N} \cdot N \right) \right\} \cdot a(t)N \cdot N^{1/2} d\alpha + \\ &+ \int_{1/N < |\alpha| \leq \log N/N} \left\{ (a(t) - a(N))N + 2a(t)t(1 + \pi|\alpha|N) \right\} \cdot a(t) \frac{1}{2|\alpha|} \cdot N^{1/2} d\alpha + \\ &+ \int_{\log N/N < |\alpha| \leq 1/Q} \left\{ (a(t) - a(N))N + 2a(t)t(1 + \pi|\alpha|N) \right\} \cdot a(t) \frac{1}{2|\alpha|} \cdot 30 \left(\frac{\log N}{|\alpha|} \right)^{1/2} d\alpha < \\ &< \frac{2}{N} \left\{ a(t)(a(t) - a(N))N^{5/2} + 2a^2(t)t \cdot 5 \cdot N^{3/2} \right\} + \\ &+ \frac{1}{2} a(t)(a(t) - a(N))N^{3/2} \int_{1/N < |\alpha| \leq \log N/N} \frac{1}{|\alpha|} d\alpha + 2 \cdot \frac{\log N}{N} \cdot a^2(t)t \cdot 5 \cdot N^{3/2} + \\ &+ 15a(t)(a(t) - a(N))N(\log N)^{1/2} \int_{\log N/N < |\alpha| \leq 1/Q} \frac{1}{|\alpha|^{3/2}} d\alpha + \\ &+ 30a^2(t)t \cdot 5N(\log N)^{1/2} \int_{\log N/N < |\alpha| \leq 1/Q} \frac{1}{|\alpha|^{1/2}} d\alpha. \end{aligned}$$

Here

$$\int_{1/N < |\alpha| \leq \log N/N} \frac{1}{|\alpha|} d\alpha = 2 \log \log N,$$

$$\text{and } \int_{\log N/N < |\alpha| \leq 1/Q} \frac{1}{|\alpha|^{3/2}} d\alpha < 2 \int_{\log N/N}^{+\infty} \frac{1}{\alpha^{3/2}} d\alpha = 2 \cdot 2 \left(\frac{\log N}{N} \right)^{-1/2} = 4 \left(\frac{N}{\log N} \right)^{1/2}$$

$$\text{and } \int_{\log N/N < |\alpha| \leq 1/Q} \frac{1}{|\alpha|^{1/2}} d\alpha < 2 \int_0^{1/Q} \frac{1}{\alpha^{1/2}} d\alpha = 2 \cdot 2 \left(\frac{1}{Q} \right)^{1/2} = \frac{4}{Q^{1/2}}.$$

Thus with respect to (57), (60) and $a(t) \cong a(N)$ (by (47) and (58)),

$$(74) \quad E_1' < 2a(t)(a(t) - a(N))N^{3/2} + 20a^2(t)tN^{1/2} +$$

$$+ a(t)(a(t) - a(N))N^{3/2} \log \log N + 10a^2(t)tN^{1/2} \log N +$$

$$+ 60a(t)(a(t) - a(N))N^{3/2} + 600a^2(t)tN(\log N)^{1/2}Q^{-1/2} <$$

$$< 63a(t)(a(t) - a(N))N^{3/2} \log \log N + 30a^2(t)tN^{1/2} \log N \left\{ 1 + 20 \left(\frac{N}{Q \log N} \right)^{1/2} \right\} \cong$$

$$\cong 63a(t)(a(t) - a(N))N^{3/2} \log \log N + 630a^2(t)tN(\log N)^{1/2}Q^{-1/2}.$$

Now we are going to estimate $E_1' + E_2$.

If $2 \cong q \cong Q$, $1 \cong p \cong q-1$ then $\alpha \in I_{p,q}$ implies that

$$\|\alpha\| \cong \frac{1}{q} - \frac{1}{qQ} \cong \frac{1}{q} - \frac{1}{2q} = \frac{1}{2q}.$$

Thus for $2 \cong q \cong Q$, $1 \cong p \cong q-1$ and $(p, q) = 1$, Lemmas 1 and 12 (where $F_1(\alpha) = 0$ in this case) and the trivial inequality (72) yield that

$$\int_{I_{p,q}} \left| |F(\alpha)|^2 - |G(\alpha)|^2 \right| |T(\alpha)| d\alpha \cong \int_{I_{p,q}} |F(\alpha)|^2 |T(\alpha)| d\alpha + \int_{I_{p,q}} |G(\alpha)|^2 |T(\alpha)| d\alpha \cong$$

$$\cong \int_{I_{p,q}} |F(\alpha)|^2 |T(\alpha)| d\alpha + \int_{I_{p,q}} a^2(t) \frac{1}{2} (2q)^2 N^{1/2} d\alpha =$$

$$= \int_{I_{p,q}} |F(\alpha)|^2 |T(\alpha)| d\alpha + \frac{1}{2qQ} \cdot 2a^2(t)q^2 N^{1/2} \cong$$

$$\cong \int_{-1/qQ}^{+1/qQ} H^2(t, N, q, \beta) \left| T\left(\frac{p}{q} + \beta\right) \right| d\beta + a^2(t)N^{1/2}.$$

Hence

$$(75) \quad E_1' + E_2 \cong \int_{-1/Q}^{+1/Q} H^2(t, N, 1, \alpha) |T(\alpha)| d\alpha +$$

$$+ \sum_{q=2}^{[R]} \sum_{\substack{1 \cong p \cong q-1 \\ (p, q) = 1}} \left\{ \int_{-1/qQ}^{+1/qQ} H^2(t, N, q, \beta) \left| T\left(\frac{p}{q} + \beta\right) \right| d\beta + a^2(t)N^{1/2} \right\} \cong$$

$$\cong \sum_{q=1}^{[R]} \sum_{\substack{0 \cong p \cong q-1 \\ (p, q) = 1}} \int_{-1/qQ}^{+1/qQ} H^2(t, N, q, \beta) \left| T\left(\frac{p}{q} + \beta\right) \right| d\beta + a^2(t)R^2 N^{1/2}.$$

To estimate $T\left(\frac{p}{q} + \beta\right)$, we use Lemmas 5 and 6 for $|\beta| \leq \log N/N$ and $\log N/N < |\beta| \leq 1/qQ$, respectively. We obtain with respect to (57), (59) and (60) that (for $q \equiv R$, $(p, q) = 1$)

$$\begin{aligned} & \int_{-1/qQ}^{+1/qQ} H^2(t, N, q, \beta) \left| T\left(\frac{p}{q} + \beta\right) \right| d\beta < \\ & < \int_{|\beta| \leq \log N/N} \{2(a(t) - a(N))^2 N^2 + 16a^2(t) t^2 q^4 (1 + \pi^2 \beta^2 N^2)\} 21 \left(\frac{N}{q}\right)^{1/2} d\beta + \\ & + \int_{\log N/N < |\beta| \leq 1/qQ} \{2(a(t) - a(N))^2 N^2 + 16a^2(t) t^2 q^4 (1 + \pi^2 \beta^2 N^2)\} 30 \left(\frac{\log N}{q|\beta|}\right)^{1/2} d\beta < \\ & < 2 \frac{\log N}{N} \left\{ 2(a(t) - a(N))^2 N^2 + 16a^2(t) t^2 q^4 \left(1 + \pi^2 \frac{\log^2 N}{N^2} N^2\right) \right\} 21 \left(\frac{N}{q}\right)^{1/2} + \\ & + 60(a(t) - a(N))^2 N^2 (\log N)^{1/2} q^{-1/2} \int_{\log N/N < |\beta| \leq 1/qQ} |\beta|^{-1/2} d\beta + \\ & + \int_{\log N/N < |\beta| \leq 1/qQ} 480a^2(t) t^2 q^4 \cdot 11\beta^2 N^2 \cdot \left(\frac{\log N}{q|\beta|}\right)^{1/2} d\beta < \\ & < 84(a(t) - a(N))^2 N^{3/2} \log N \cdot q^{-1/2} + 32 \cdot \log N \cdot N^{-1/2} a^2(t) t^2 q^{7/2} \cdot 11 \cdot \log^2 N \cdot 21 + \\ & + 120(a(t) - a(N))^2 N^2 (\log N)^{1/2} q^{-1/2} \int_0^{1/qQ} \beta^{-1/2} d\beta + \\ & + 10560a^2(t) t^2 q^{7/2} N^2 (\log N)^{1/2} \int_0^{1/qQ} \beta^{3/2} d\beta = \\ & = 84(a(t) - a(N))^2 N^{3/2} \log N \cdot q^{-1/2} + 7392a^2(t) t^2 N^{-1/2} (\log N)^2 q^{7/2} + \\ & + 120(a(t) - a(N))^2 N^2 (\log N)^{1/2} q^{-1/2} \cdot 2(qQ)^{-1/2} + \\ & + 10560a^2(t) t^2 q^{7/2} N^2 (\log N)^{1/2} \cdot \frac{2}{5} (qQ)^{-5/2} = \\ & = (a(t) - a(N))^2 \{84N^{3/2} \log N \cdot q^{-1/2} + 240N^2 (\log N)^{1/2} Q^{-1/2} q^{-1}\} + \\ & + a^2(t) t^2 \{7392N^{-1/2} (\log N)^2 q^{7/2} + 4224N^2 (\log N)^{1/2} Q^{-5/2} q\} \end{aligned}$$

since

$$(x+y)^2 \leq 2x^2 + 2y^2$$

for any real numbers x, y . Thus (75) yields with respect to (57) and (59) that
(76)

$$\begin{aligned} E_1' + E_2 &< \sum_{q=1}^{[R]} \sum_{p=0}^{q-1} \{ (a(t) - a(N))^2 (84N^{3/2} \log N \cdot q^{-1/2} + 240N^2 (\log N)^{1/2} Q^{-1/2} q^{-1}) + \\ &+ a^2(t) t^2 (7392N^{-1/2} (\log N)^3 q^{7/2} + 4224N^2 (\log N)^{1/2} Q^{-5/2} q) \} + a^2(t) R^2 N^{1/2} < \\ &< \sum_{q=1}^{[R]} \{ (a(t) - a(N))^2 (84N^{3/2} \log N \cdot q^{1/2} + 240N^2 (\log N)^{1/2} Q^{-1/2}) + \\ &+ a^2(t) t^2 (7392N^{-1/2} (\log N)^3 q^{9/2} + 4224N^2 (\log N)^{1/2} Q^{-5/2} q^2) \} + \\ &+ a^2(t) (N^{1/2} / \log N)^2 N^{1/2} \cong \\ &\cong (a(t) - a(N))^2 \{ 84N^{3/2} R^{3/2} \log N + 240N^2 (\log N)^{1/2} Q^{-1/2} R \} + \\ &+ a^2(t) t^2 \{ 7392N^{-1/2} (\log N)^3 R^{11/2} + 4224N^2 (\log N)^{1/2} Q^{-5/2} R^3 \} + \frac{1}{64} a^2(t) N^{3/2}. \end{aligned}$$

Finally, in order to estimate E_3 , we use Lemma 7. Namely, if $\alpha \in S$ then there exist integers p, q such that

$$1 \cong q \cong Q, \quad 0 \cong p \cong q-1, \quad (p, q) = 1$$

and

$$\left| \alpha - \frac{p}{q} \right| < \frac{1}{qQ};$$

by (67), q satisfies also $R < q$. Thus (41) and (43) in Lemma 7 hold (and also (42) holds by (60)). Hence, Lemma 7 yields that

$$\sup_{\alpha \in S} |T(\alpha)| \cong 7 \left(\frac{N}{R} \right)^{1/2} + 14(Q \log N)^{1/2}.$$

Thus we obtain applying Parseval's formula that

$$\begin{aligned} (77) \quad E_3 &= \int_S | |F(\alpha)|^2 - |G(\alpha)|^2 | |T(\alpha)| d\alpha \cong \\ &\cong \sup_{\alpha \in S} |T(\alpha)| \left(\int_S |F(\alpha)|^2 d\alpha + \int_S |G(\alpha)|^2 d\alpha \right) < \\ &< \{ 7N^{1/2} R^{-1/2} + 14Q^{1/2} (\log N)^{1/2} \} \left(\int_0^1 |F(x)|^2 dx + \int_0^1 |G(x)|^2 dx \right) = \\ &= \{ 7N^{1/2} R^{-1/2} + 14Q^{1/2} (\log N)^{1/2} \} (A(N) + a^2(t) N) = \\ &= \{ 7N^{1/2} R^{-1/2} + 14Q^{1/2} (\log N)^{1/2} \} (a(N) N + a^2(t) N) \cong \\ &\cong \{ 7N^{1/2} R^{-1/2} + 14Q^{1/2} (\log N)^{1/2} \} (a(t) N + a(t) N) = \\ &= a(t) \{ 14N^{3/2} R^{-1/2} + 28NQ^{1/2} (\log N)^{1/2} \} \end{aligned}$$

by Lemma 9 and since $a(N) \cong a(t)$ by (47) in Lemma 11 and (58).

Collecting the results (62), (66), (68), (71), (74), (76) and (77) together, we obtain that

$$\begin{aligned} \frac{1}{8} a^2(t) N^{3/2} &< \int_0^1 |G(x)|^2 T(x) dx \cong E_1 + E_2 + E_3 \cong \\ &\cong (E_1' + 2E_1'') + E_2 + E_3 = 2E_1'' + (E_1' + E_2) + E_3 < \\ &< 126a(t)(a(t) - a(N)) N^{3/2} \log \log N + 1260a^2(t) tN (\log N)^{1/2} Q^{-1/2} + \\ &+ (a(t) - a(N))^2 \{84N^{3/2} R^{3/2} \log N + 240N^2 (\log N)^{1/2} Q^{-1/2} R\} + \\ &+ a^2(t) t^2 \{7392N^{-1/2} (\log N)^3 R^{11/2} + 4224N^2 (\log N)^{1/2} Q^{-5/2} R^3\} + \\ &+ \frac{1}{64} a^2(t) N^{3/2} + a(t) \{14N^{3/2} R^{-1/2} + 28NQ^{1/2} (\log N)^{1/2}\}. \end{aligned}$$

Subtracting $\frac{1}{64} a^2(t) N^{3/2}$ and then multiplying by

$$\left(\frac{1}{8} - \frac{1}{64}\right)^{-1} = \left(\frac{7}{64}\right)^{-1} = \frac{64}{7} < 10,$$

we obtain that

$$\begin{aligned} a^2(t) N^{3/2} &< 1260a(t)(a(t) - a(N))^{3/2} \log \log N + 12600a^2(t) tN (\log N)^{1/2} Q^{-1/2} + \\ &+ 120(a(t) - a(N))^2 \{7N^{3/2} R^{3/2} \log N + 20N^2 (\log N)^{1/2} Q^{-1/2} R\} + \\ &+ 26000a^2(t) t^2 \{3N^{-1/2} (\log N)^3 R^{11/2} + 2N^2 (\log N)^{1/2} Q^{-5/2} R^3\} + \\ &+ 140a(t) \{N^{3/2} R^{-1/2} + 2NQ^{1/2} (\log N)^{1/2}\} \end{aligned}$$

which proves Lemma 13.

6. In this section, we are going to simplify the functional inequality given in Lemma 13. It can be shown that we obtain the best possible estimate for $a(x)$ in the case when the order of magnitude of the product QR is $N/\log N$; thus we choose

$$R = \frac{N}{Q} \cdot \frac{1}{\log N}.$$

Furthermore, we put

$$r = \frac{N}{t}, \quad s = \frac{N}{Q}.$$

Then

$$R = \frac{s}{\log N}.$$

The inequalities (59) and (60) can be rewritten in form

$$1 \cong \frac{s}{\log N} \cong \frac{N^{1/2}}{\log N},$$

(78)

$$\log N \cong s \cong N^{1/2}$$

and

$$(79) \quad \begin{aligned} 2N^{1/2} &< \frac{N}{s} \cong \frac{N}{\log N}, \\ \frac{1}{2}N^{1/2} &> s \cong \log N, \end{aligned}$$

respectively. (79) implies (78) thus it suffices to assume that (79) holds. Finally, if we divide (61) by $N^{3/2}$ then we obtain that

$$\begin{aligned} a^2(t) &< 1260a(t)(a(t) - a(N)) \log \log N + 12600a^2(t)r^{-1}s^{1/2}(\log N)^{1/2} + \\ &+ 120(a(t) - a(N))^2 \{7s^{3/2}(\log N)^{-1/2} + 20s^{3/2}(\log N)^{-1/2}\} + \\ &+ 26000a^2(t)r^{-2} \{3s^{11/2}(\log N)^{-5/2} + 2s^{11/2}(\log N)^{-5/2}\} + \\ &+ 140a(t) \{s^{-1/2}(\log N)^{1/2} + 2s^{-1/2}(\log N)^{1/2}\} < \\ &< 1260a(t)(a(t) - a(N)) \log \log N + \\ &+ 130000a^2(t) \{r^{-1}s^{1/2}(\log N)^{1/2} + r^{-2}s^{11/2}(\log N)^{-5/2}\} + \\ &+ 3240(a(t) - a(N))^2 s^{3/2}(\log N)^{-1/2} + 420a(t)s^{-1/2}(\log N)^{1/2}. \end{aligned}$$

Thus we have proved the following

LEMMA 14. Let t, N, r be positive integers, s a real number such that

$$(80) \quad N \cong e^8,$$

$$(81) \quad N = tr,$$

$$(82) \quad \log N \cong s < \frac{1}{2}N^{1/2}.$$

Then

$$(83) \quad \begin{aligned} a^2(t) &< 1260a(t)(a(t) - a(N)) \log \log N + \\ &+ 130000a^2(t) \{r^{-1}s^{1/2}(\log N)^{1/2} + r^{-2}s^{11/2}(\log N)^{-5/2}\} + \\ &+ 3240(a(t) - a(N))^2 s^{3/2}(\log N)^{-1/2} + 420a(t)s^{-1/2}(\log N)^{1/2}. \end{aligned}$$

7. In this section, we are going to complete the proof of our theorem by showing that the functional inequality given in Lemma 14 implies (2).

Let us put

$$\varphi(x) = \frac{(\log \log x)^{2/3}}{(\log x)^{1/3}}$$

for $x \cong 3$. Furthermore, for $L=3, 4, \dots$, let us define the sequence $\mathcal{B}(L) = \{b_1, b_2, \dots\}$ by the following recursion: let $b_1=L$, and for $k=1, 2, \dots$, let

$$b_{k+1} = b_k [4 \cdot 10^{22} (\varphi(b_k))^{-11/2} (\log b_k)^{3/2}].$$

Obviously, $(\varphi(x))^{-1} \rightarrow +\infty$ as $x \rightarrow +\infty$. Thus there exists a real number X_0

such that $(\varphi(x))^{-1} > 1$ for $x \geq X_0$. Let us put $L_1 = \max\{X_0, 3\}$. Then we obtain (by straight induction) that for $L \geq L_1$,

$$(84) \quad \frac{b_{k+1}}{b_k} = [4 \cdot 10^{22} (\varphi(b_k))^{-11/2} (\log b_k)^{3/2}] > [4 \cdot 10^{22} \cdot 1 \cdot 1] = 4 \cdot 10^{22} > 1.$$

Hence

$$(85) \quad L = b_1 < b_2 < b_3 < \dots \quad (\text{for } L \geq L_1).$$

We are going to show (by straight induction, using Lemma 14) that if L is large enough then for $k=1, 2, \dots$,

$$(86) \quad a(b_k) \cong \frac{1}{\varphi(L)} \cdot \varphi(b_k).$$

For $k=1$, the right hand side of (86) is

$$\frac{1}{\varphi(L)} \cdot \varphi(b_1) = \frac{1}{\varphi(L)} \cdot \varphi(b_1) = \frac{1}{\varphi(L)} \cdot \varphi(L) = 1$$

thus in this case, (86) holds trivially by Lemma 9.

Let us suppose now that (86) holds for some positive integer k . We have to show that this implies also

$$(87) \quad a(b_{k+1}) \cong \frac{1}{\varphi(L)} \varphi(b_{k+1}).$$

Let us suppose indirectly that

$$(88) \quad a(b_{k+1}) > \frac{1}{\varphi(L)} \varphi(b_{k+1}).$$

By the construction of the sequence $\mathcal{B}(L)$,

$$(89) \quad b_k/b_{k+1}.$$

Hence,

$$(90) \quad a(b_k) \cong a(b_{k+1})$$

by (47) in Lemma 11.

We are going to show that for sufficiently large L , Lemma 14 is applicable with $t = b_k$, $N = b_{k+1}$,

$$r = N/t = b_{k+1}/b_k = [4 \cdot 10^{22} (\varphi(b_k))^{-11/2} (\log b_k)^{3/2}],$$

$$s = 5 \cdot 10^6 (a(b_k))^{-2} \log b_{k+1} = 5 \cdot 10^6 (a(t))^{-2} \log N.$$

Then by (85),

$$(91) \quad N > t \cong L \quad (\text{for } L \geq L_1).$$

Thus (80) holds for $L \geq \max\{e^8, L_1\}$.

(81) holds by the definitions of N, t, r and $\mathcal{B}(L)$.

By Lemma 9,

$$s = 5 \cdot 10^6 (a(b_k))^{-2} \log N \cong 5 \cdot 10^6 \log N > \log N.$$

Finally, we have to prove that

$$s < \frac{1}{2} N^{1/2}.$$

With respect to (84), (88) and (89),

$$\begin{aligned} (92) \quad s &= 5 \cdot 10^6 (a(b_k))^{-2} \log b_{k+1} \cong \\ &\cong 5 \cdot 10^6 (a(b_{k+1}))^{-2} \log b_{k+1} \cong 5 \cdot 10^6 \left(\frac{1}{\varphi(L)} \varphi(b_{k+1}) \right)^{-2} \log b_{k+1} \cong \\ &\cong 5 \cdot 10^6 (\varphi(b_{k+1}))^{-2} \log b_{k+1} = 5 \cdot 10^6 (\varphi(N))^{-2} \log N = 5 \cdot 10^6 \frac{(\log N)^{5/3}}{(\log \log N)^{4/3}} \end{aligned}$$

for $L \cong L_1 \cong X_0$. Thus it suffices to show that

$$5 \cdot 10^6 \frac{(\log N)^{5/3}}{(\log \log N)^{4/3}} < \frac{1}{2} N^{1/2}.$$

But this holds trivially for large N , i.e. for $L \cong L_2$ (in view of (91)).

Thus Lemma 14 is applicable and it yields that (83) holds. To deduce a contradiction from (83), we have to estimate r and $a(t) - a(N)$ (in terms of $a(t)$ and N).

Obviously,

$$\begin{aligned} (93) \quad r &= \frac{N}{t} = \frac{b_{k+1}}{b_k} = [4 \cdot 10^{22} (\varphi(b_k))^{-11/2} (\log b_k)^{3/2}] = \\ &= [4 \cdot 10^{22} (\varphi(t))^{-11/2} (\log t)^{3/2}] = \left[4 \cdot 10^{22} \left\{ \frac{(\log \log t)^{2/3}}{(\log t)^{1/3}} \right\}^{-11/2} (\log t)^{3/2} \right] = \\ &= \left[4 \cdot 10^{22} \frac{(\log t)^{10/3}}{(\log \log t)^{11/3}} \right] < 4 \cdot 10^{22} \frac{(\log t)^{10/3}}{(\log \log t)^{11/3}} \end{aligned}$$

and with respect to (84),

$$(94) \quad r = \frac{N}{t} = \left[4 \cdot 10^{22} \frac{(\log t)^{10/3}}{(\log \log t)^{11/3}} \right] > 2 \cdot 10^{22} \frac{(\log t)^{10/3}}{(\log \log t)^{11/3}} \quad (\text{for } L \cong L_2).$$

In view of (91), (93) and (94) imply that for $L \cong L_3$,

$$\begin{aligned} (95) \quad (0 <) \log r &= \log N - \log t = \\ &= \frac{10}{3} \log \log t + \theta(1) \cdot \log(4 \cdot 10^{22}) + \theta \left(\frac{11}{3} \log \log \log t \right) < \\ &< 4 \log \log t \quad (\text{for } L \cong L_3) \end{aligned}$$

hence for $L \cong L_4$,

$$\begin{aligned} (96) \quad \log t &> \log N - 4 \log \log t > \log N - 4 \log \log N > \\ &> \frac{1}{2} \log N \quad (\text{for } L \cong L_4) \end{aligned}$$

and for $L \cong L_5$,

$$(97) \quad \log \log t > \log \left(\frac{1}{2} \log N \right) = \log \log N - \log 2 > \frac{1}{2} \log \log N \quad (\text{for } L \cong L_5).$$

(86), (88), (91), (95), (96) and (97) yield that for $L \cong L_6$,

$$(98) \quad \begin{aligned} a(t) - a(N) &= a(t) \left(1 - \frac{a(N)}{a(t)} \right) < a(t) \left(1 - \frac{(\varphi(L))^{-1} \varphi(N)}{(\varphi(L))^{-1} \varphi(t)} \right) = \\ &= a(t) \frac{\varphi(t) - \varphi(N)}{\varphi(t)} = a(t) \frac{(\log t)^{1/3}}{(\log \log t)^{2/3}} \left(\frac{(\log \log t)^{2/3}}{(\log t)^{1/3}} - \frac{(\log \log N)^{2/3}}{(\log N)^{1/3}} \right) < \\ &< a(t) \frac{(\log t)^{1/3}}{\left(\frac{1}{2} \log \log N \right)^{2/3}} \left(\frac{(\log \log N)^{2/3}}{(\log t)^{1/3}} - \frac{(\log \log N)^{2/3}}{(\log N)^{1/3}} \right) < \\ &< a(t) \frac{2(\log t)^{1/3}}{(\log \log N)^{2/3}} (\log \log N)^{2/3} \frac{(\log N)^{1/3} - (\log t)^{1/3}}{(\log t)^{1/3} (\log N)^{1/3}} = \\ &= 2a(t) \frac{\log N - \log t}{(\log N)^{1/3} ((\log N)^{2/3} + (\log N \log t)^{1/3} + (\log t)^{2/3})} < \\ &< 2a(t) \frac{4 \log \log t}{(\log N)^{1/3} \cdot 3 (\log t)^{2/3}} < \frac{8}{3} a(t) \frac{\log \log N}{(\log N)^{1/3} \left(\frac{1}{2} \log N \right)^{2/3}} < \\ &< 6a(t) \frac{\log \log N}{\log N}. \end{aligned}$$

By (88), (90), (91), (92), (94), (96) and (98), (83) implies that for $L \cong L_7$,

$$\begin{aligned} a^2(t) &< 1260a(t) \cdot 6a(t) \frac{\log \log N}{\log N} \cdot \log \log N + \\ &+ 130000a^2(t) \left\{ 2^{-1} \cdot 10^{-22} \frac{(\log \log t)^{11/3}}{(\log t)^{10/3}} \cdot \left(5 \cdot 10^6 \frac{(\log N)^{5/3}}{(\log \log N)^{4/3}} \right)^{1/2} \cdot (\log N)^{1/2} + \right. \\ &\quad \left. + \left(2^{-1} 10^{-22} \frac{(\log \log t)^{11/3}}{(\log t)^{10/3}} \right)^2 \left(5 \cdot 10^6 \frac{(\log N)^{5/3}}{(\log \log N)^{4/3}} \right)^{11/2} \cdot (\log N)^{-5/2} \right\} + \\ &\quad + 3240 \left(6a(t) \frac{\log \log N}{\log N} \right)^2 (5 \cdot 10^6 (a(t))^{-2} \log N)^{3/2} (\log N)^{-1/2} + \\ &\quad - 420a(t) (5 \cdot 10^6 (a(t))^{-2} \log N)^{-1/2} (\log N)^{1/2} < \frac{7560 (\log \log N)^2}{\log N} a^2(t) + \\ &\quad + 130 \cdot 10^3 a^2(t) \left\{ 2^{-1} \cdot 10^{-22} \frac{(\log \log N)^{11/3}}{\left(\frac{1}{2} \log N \right)^{10/3}} \cdot 3 \cdot 10^3 \frac{(\log N)^{4/3}}{(\log \log N)^{2/3}} + \right. \end{aligned}$$

$$\begin{aligned}
& + 2^{-2} \cdot 10^{-44} \frac{(\log \log N)^{22/3}}{\left(\frac{1}{2} \log N\right)^{20/3}} \cdot 5^{1/2} \cdot 5^5 \cdot 10^{33} \frac{(\log N)^{20/3}}{(\log \log N)^{20/3}} \Bigg\} + \\
& + 4 \cdot 10^3 \cdot 36 \cdot a^2(t) \frac{(\log \log N)^2}{(\log N)^2} \cdot 125^{1/2} \cdot 10^9 (a(t))^{-3} \log N + \\
& + (21^2 \cdot 2^2 \cdot 10^2 / 500 \cdot 2^2 \cdot 10^2 \cdot 25)^{1/2} a^2(t) < \\
& < \frac{1}{10} a^2(t) + a^2(t) \left\{ 130 \cdot 10^{-16} \cdot 2^{7/3} \cdot 3 \cdot \frac{(\log \log N)^2}{(\log N)^2} + 130 \cdot 10^{-8} \cdot 2^{-2} \cdot 2^7 \cdot 3 \cdot 5^5 \right\} + \\
& + 144 \cdot 10^{12} \cdot 12 a^2(t) \frac{(\log \log N)^2}{\log N} \left(\frac{1}{\varphi(L)} \varphi(N) \right)^{-3} + \left(\frac{441}{500} \cdot \frac{1}{25} \right)^{1/2} a^2(t) < \\
& < \frac{1}{10} a^2(t) + a^2(t) \left\{ \frac{1}{10} + 390 \cdot 10^{-9} \right\} + 2 \cdot 10^{15} a^2(t) \varphi^3(L) + \frac{1}{5} a^2(t) < \\
& < \frac{1}{10} a^2(t) + a^2(t) \left(\frac{1}{10} + \frac{2}{5} \right) + \frac{1}{5} a^2(t) + \frac{1}{5} a^2(t) = 2a^2(t)
\end{aligned}$$

provided that

$$2 \cdot 10^{15} \varphi^3(L) < \frac{1}{5}, \quad \varphi(L) < 10^{-16/3}$$

but this holds for $L \cong L_8$.

Thus for large L , the indirect assumption (88) leads to the contradiction $a^2(t) < a^2(t)$ which proves (86).

Finally, if x is a positive integer satisfying $x \cong b_1 = L$, then there exists a positive integer k such that $b_k \cong x < b_{k+1}$. By (48) in Lemma 11, (86) implies that

$$\begin{aligned}
(99) \quad a(x) & \cong \left(1 + \frac{b_k}{x} \right) a(b_k) \cong 2 \cdot \frac{1}{\varphi(L)} \varphi(b_k) \cong \\
& \cong 2 \cdot \frac{1}{\varphi(L)} \cdot \frac{(\log \log b_k)^{2/3}}{(\log b_k)^{1/3}} \cong 2 \cdot \frac{1}{\varphi(L)} \cdot \frac{(\log \log x)^{2/3}}{(\log x)^{1/3}}.
\end{aligned}$$

With respect to (96),

$$\log b_k > \frac{1}{2} \log b_{k+1} > \frac{1}{2} \log x.$$

Thus we obtain from (99) that

$$(100) \quad a(x) \cong \frac{2}{\varphi(L)} \cdot \frac{(\log \log x)^{2/3}}{\left(\frac{1}{2} \log x\right)^{1/3}} \cong \frac{4}{\varphi(L)} \cdot \frac{(\log \log x)^{2/3}}{(\log x)^{1/3}} \quad (\text{for } x \cong L)$$

which completes the proof of our theorem.

We remark that L can be chosen as the least positive integer L satisfying $L \in \{e^s, L_1, L_2, \dots, L_8\}$. All the constants L_1, L_2, \dots, L_8 are explicitly computable thus also the constants in (100) are explicitly computable.

8. In Part II of this series, we will give a lower estimate for $a(x)$.

References

- [1] E. LANDAU, *Über einige neuere Fortschritte der additiven Zahlentheorie*, Cambridge University Press (1937).
- [2] K. F. ROTH, Sur quelques ensembles d'entiers, *C. R. Acad. Sci. Paris*, **234** (1952), 388—390.
- [3] K. F. ROTH, On certain sets of integers, *J. London Math. Soc.*, **28** (1953), 104—109.

(Received January 12, 1977)

MATHEMATICAL INSTITUTE
OF THE HUNGARIAN ACADEMY OF SCIENCES
1053 BUDAPEST, REALTANODA U. 13—15.