

Keep Our Future Generation Alive:


Reinforcing Routine HIV Testing & Treatment Among Children in Zambia

Key Messages

- Only 3% of children visiting clinics are tested for HIV, despite Ministry of Health recommendations to test and treat ALL children.
- If unidentified and left untreated, 75% of HIV positive children die by the age of 5 years.
- Reinforcement of routine HIV testing and treatment of children will increase testing up to 90 - 95%, and importantly reduce mortality by 52%.
- Adding Integrated Primary School Screening will reach 5-10 year olds who were not previously tested.



The Problem




Source: CIDRZ Annual Progress Reports 2006-2015

Zambia has made great strides in tackling the HIV/AIDS epidemic, initiating 58% of HIV infected people on treatment¹, prescribing more effective drugs, and delivering these services “closer to home”. Over the past 10 years the prevention-of-mother-to-child-transmission (PMTCT) of HIV has changed from single dose anti-retroviral prevention to full Anti-Retroviral Treatment (ART), reducing the percentage of HIV infected children being born of HIV positive mothers from 7% to 2%. **However, despite these efforts, there are approximately 36,000 HIV positive children not being identified through HIV testing annually, according to national estimate¹.**

To address this child testing gap a team from the Centre for Infectious Disease Research in Zambia (CIDRZ) developed a model to estimate the number of unidentified positive children in the catchment population of 338 CIDRZ-supported facilities in the Lusaka, Western, and Eastern Provinces of the Zambia.

The model results indicated that from 2006 through to the end of 2015, there were 30,203 HIV positive children that had died from HIV-related conditions. In addition, there was an estimated 14,348 HIV-infected children that remained untested, unidentified, and therefore untreated.

When HIV-Infected Children are Left Undiagnosed and Untreated



Keep Our Future Generation Alive

Recent literature from sub-Saharan African indicates that 75% of those unidentified HIV infected children will die by the age of 5, while up to 80% will die by the age of 10, if they remain without HIV treatment².

Since 2008, Zambia has recommended routine HIV testing at immunization clinics for those children born to HIV-positive mothers who attended antenatal clinics. This has increased the testing of HIV-exposed infants to 98%. But after the HIV-exposed child reaches one year of age they no longer receive regular testing. Children, of mothers who did not attend antenatal clinics and have unknown HIV status, are missed and do not receive routine HIV testing.

Though the Ministry of Health has promoted Provider Initiated Testing and Counseling (PITC) in Out-Patient departments, the uptake of child testing is very low at only 3% compared to the 40% of adult testing found in most clinics. Healthcare staff are reluctant to ask permission to test a child, especially when the accompanying adult is not the parent. Non-parent caregivers are also reluctant to give permission for testing. Mothers tend to refuse having their child being tested if they don't have the father's or the spouse's permission. This results in an "Opt-In" approach that presents many missed opportunities for child testing.

How can we find un-identified HIV positive children who will die if they don't receive ART?

<i>No Intervention in 2015</i>				<i>Reinforcing Routine HIV Testing in 2015</i>		
Age (yrs)	NEW on ART	NOT on ART	Died	NEW on ART	NOT on ART	Died
1	132	915	1081	1265	425	439
2	324	1411	317	1415	538	98
3	195	2154	239	1055	1379	153
4	124	1869	221	860	1219	135
5	107	1323	159	635	858	95
6	116	1360	94	638	876	56
7	117	1253	87	601	804	51
8	151	1570	110	760	1007	64
9	115	1283	89	609	825	53
10	119	1209	85	589	775	49
Total	1500	14348	2483	8429	8707	1195

Calculated Estimated Impact of 1 Year of Option 1 in 338 Facilities in 2015: 6,929 more children on treatment, and 1,288 (52%) deaths avoided.

Policy Options

In order to find all children living with HIV, reinforcement of routine HIV testing with community sensitization on the importance of HIV testing in children is essential. Proposed policy options to achieve these include: (1) Reinforcing Routine HIV testing; (2) Introducing Pre- and Primary School Screening Drives; and (3) Introducing Primary School Entry Screening.

1. Reinforcing Routine HIV Testing Strategy

WHAT: Routinely test all children for HIV, regardless of their age and condition as long as they visit a health facility with adequate information provided to the caregiver.

WHY: 80% of children up to 2 years of age, and 50% of children up to 5 years of age visit a health facility, but currently only 3 out of every 100 children are tested for HIV in Outpatient Departments. Routine testing removes the responsibility for the decision of testing from healthcare workers and parents/caregivers. According to literature in Zimbabwe³ and Tanzania⁴ this strategy has increased child testing up to 90%. **In our model, applying reinforcing routine HIV testing in 2015 for 0-10 year olds would have identified an additional 39% untested HIV-positive, and would have prevented 52% of child deaths.**

FEASIBILITY: MEDIUM to HIGH This strategy builds on the government's decision to identify HIV positive children. It will require a reinforced **legal framework**, community sensitization, placement of more trained counselors, and an increase in HIV test kits.

2. Introducing Screening Drives at Pre- and Primary Schools

WHAT: Conduct integrated screening drives for Pre-School and Primary School children, assessing development, eyes, ears, and dental, immunization status, and testing for malaria, TB and HIV.

WHY: 80% of children in Zambia attend Pre or Primary school. This strategy will find the 5 to 10 year old HIV positive children that have been missed during usual health services. **In our model, applying School Screening Drive in 2015 for school-goers would have identified, 50% of untested HIV-positive children (5-10-year-old), and will reduce 5-10-year child deaths by 54%.** Additional benefits of school screening include reducing absenteeism and improve school performance⁵.

FEASIBILITY: HIGH This strategy will require community sensitization, funds, transport, and human resource, including orientation in school screening for nurses in collaboration with Ministry of Education and other key stakeholders.

3. Introducing Primary School Entry Screening

WHAT: Conduct integrated screening for new children entering Primary School, assessing development, eyes, ears, and dental, immunization status and test for malaria, TB and HIV before the child is enrolled.

WHY: 80% of children in Zambia attend Pre- and Primary School. **In our model, this strategy will identify 35% of the 5-year-old HIV positive children who were missed in the health facilities, and will prevent 40% of death among 5 year olds.** Additional benefits of school screening include reducing absenteeism and improve school performance.

FEASIBILITY: HIGH This strategy builds on health services already available at the clinic for children. Children can visit the clinic for school entry testing. Nurses will require orientation on screening activities, and collaboration will be needed with the Ministry of Education and other key stakeholders.


Keep Our Future Generation Alive

ESTIMATED COSTS BY OPTION	Option 1 Routine Testing	Option 2 Pre & Primary School Drives	Option 3 Pre School Entry Screening
Positive Children Found in 1 yr	6,929	4,311	528
Lives Saved in 1 yr	1,288	336	64
Annual Testing Cost (testing, HR, training, community sensitisation)	\$1,289,722	\$2,375,951	\$493,476
Testing Cost Per Positive Child	\$186	\$551	\$934
Annual Treatment Cost (testing, HR, training, community sensitisation)	\$1,216,060	\$1,133,911	\$138,967
Treatment Cost Per Positive Child	\$176	\$263	\$263
Political Feasibility			
Operational Feasibility			

Recommendations

Reinforcing Routine HIV testing is the most cost effective and feasible option to increase HIV testing among children with unknown status, as recommended by the WHO⁶ and the CDC⁷. Implementation of routine HIV testing will identify the majority of HIV-positive children in Zambia, reduce HIV-related mortality and promote long-term control of the epidemic.

Routine Testing Policy Will Save Children's Lives



In addition to a legal framework and collaboration amongst stakeholders this strategy will require:

Additional test kits, more trained counsellors, and a robust referral system to link children to treatment.

Ministry of Health to intensify community sensitization and trainings for healthcare providers.

Ministry of Finance to ensure sufficient resources so all identified children can be placed on treatment.

Additional school screening will identify children who were missed at the health institutions.



References:

1. Spectrum Zambia 2013/2014, <http://www.avenirhealth.org/software-spectrum.php>
2. Renaud Becquet, 'Children who acquire HIV inf perinatally are at higher risk of early death than those acquiring inf through breastmilk: a meta-analysis', source PLoS One, 2012
3. Rashida Abas Ferrand, Jamilah Meghji, The effectiveness of Routine Opt-Out HIV testing for children in Harare, Zimbabwe, Aquire Immune Deficiency Syndrome, 2016
4. Kathy Baisley, Aoife Doyl, Uptake of voluntary counselling and testing among young people participating in an HIV prevention trial: comparison of opt-out and Opt-in strategies, Plos One, 2012
5. Tsitsi Bandason, Lisa Langhaug, Burden of HIV among primary school children and feasibility of primary school-linked HIV testing in Harare, Zimbabwe: A mixed methods study, Aids Care, 2014
6. World Health Organization May 2007, Provider-Initiated Testing and Counseling in Health Facilities
7. CDC 2006, Revised Recommendation for HIV testing of adults, adolescents and pregnant women in Health Care setting: home/HIV/AIDS Basics/Prevention: HIV Testing: Opt-Out Testing
8. Zambian Demographic Health Survey 2013/2014
9. Nigel Rollins, Mary Mahy, Renaud Becquet, 'Estimates of peripartum and postnatal mother to child transmission probabilities of HIV for use in Spectrum and other population-based models', UNAIDS report 2012.