

Unusual Eruptions Associated with *Mycoplasma pneumoniae* Respiratory Infections: Review of the Literature

Luisa Terraneo^a Sebastiano A.G. Lava^{a,b} Pietro Camozzi^a Lorenzo Zgraggen^{a,b}
Giacomo D. Simonetti^a Mario G. Bianchetti^a Gregorio P. Milani^c

^aPediatric Department of Southern Switzerland, Bellinzona, and ^bUniversity Children's Hospital Bern and University of Bern, Bern, Switzerland; ^cFoundation IRCCS Ca' Granda, Ospedale Maggiore Policlinico, Pediatric Emergency Department, Milan, Italy

Key Words

Erythema nodosum · Gianotti-Crosti syndrome · Henoch-Schönlein syndrome · Mucositis · *Mycoplasma pneumoniae* · Papular acrodermatitis · Pityriasis lichenoides et varioliformis acuta Mucha-Habermann · Pityriasis rosea Gibert · Varicella-like eruption

Mycoplasma infections. **Conclusion:** This review implies that *M. pneumoniae* may cause, in addition to erythematous maculopapular (or urticarial) eruptions and erythema multiforme, Fuchs' syndrome and varicella-like eruptions. Furthermore, there is an intriguing link with leukocytoclastic vasculitides or erythema nodosum that deserves further investigation.

© 2015 S. Karger AG, Basel

Abstract

Background: Maculopapular or urticarial eruptions and erythema multiforme sometimes occur in patients affected with *Mycoplasma pneumoniae* respiratory infections. Further eruptions have also been reported. **Objective:** To review the literature addressing *M. pneumoniae* respiratory infection and rather unusual eruptions. **Methods:** Computer-based search in the US National Library of Medicine database as well as in the search engine Google. **Results:** We found a possible relationship between *M. pneumoniae* infection and Fuchs' syndrome ($n = 37$), varicella-like eruptions ($n = 8$), Henoch-Schönlein syndrome and further leukocytoclastic vasculitides ($n = 21$) and erythema nodosum ($n = 11$). A temporal relationship was also observed with 2 cases of Gianotti-Crosti syndrome. Finally, there exists reasonable evidence that pityriasis rosea Gibert and pityriasis lichenoides et varioliformis acuta Mucha-Habermann are not associated with

Introduction

Skin eruptions sometimes occur in patients affected with *Mycoplasma pneumoniae* respiratory infections [1]. The most usual eruption is a mild erythematous maculopapular (or urticarial) rash. Targetoid lesions, the hallmark of erythema multiforme, mostly associated with erosions or bullae involving the oral, ocular or anogenital mucosae, are a further common manifestation [2]. Obviously, similar eruptions can also be caused by the antimicrobials prescribed to treat this infection.

Further eruptions have also been observed in patients with *M. pneumoniae* respiratory infections. Since textbooks and reviews only marginally mention these unusual associations, we analyzed the available literature.

Data Assessment

Between October and December 2014, we performed a computer-based search in the National Library of Medicine database as well as in the search engine Google. Articles published after 1960 were considered, which address a possible link between *M. pneumoniae* respiratory infections and erythema nodosum, Fuchs' syndrome, Gianotti-Crosti syndrome, Henoch-Schönlein syndrome, pityriasis lichenoides et varioliformis acuta Mucha-Habermann, pityriasis rosea Gibert or varicella-like eruptions. For this purpose 'Mycoplasma pneumoniae' and each of the aforementioned conditions (and their synonyms) together with the Boolean operator 'AND' were utilized. Pertinent secondary references were also selected. Reports published in languages other than English, French, German, Italian, Portuguese, or Spanish were not included. Exclusively cases other than erythematous maculopapular and urticarial eruptions or classical erythema multiforme associated with *M. pneumoniae* infections were considered for the analysis. In retained cases, diagnosis of respiratory *M. pneumoniae* infection was based both on a distinctive clinical presentation and an appropriate laboratory testing [3]. Cases with eruptions possibly due both to Mycoplasma as well as to an already recognized cause (e.g. erythema nodosum with clinical and laboratory findings consistent with both a streptococcal and a *M. pneumoniae* infection) were excluded [4].

Furthermore, to address their possible quantitative relationship with *M. pneumoniae* infections, case series dealing with Henoch-Schönlein syndrome, erythema nodosum or Gianotti-Crosti syndrome published in English after 1980 were also reviewed.

Using the described strategy, 94 reports were retained for the final analysis [4–97]: 1 in German [23], 1 in Italian [38], 3 in French [46, 51, 92], 4 in Spanish [20, 24, 50, 81] and the remaining 85 in English.

The Fisher test was used to compare dichotomous variables. Statistical significance was assigned at $p < 0.05$.

Results

The characteristics of 77 patients with Fuchs' syndrome, varicella-like eruption, vasculitis and erythema nodosum associated with *M. pneumoniae* respiratory infection appear in table 1. Fuchs' syndrome (and varicella-like eruption) more frequently affected male subjects than erythema nodosum.

Fuchs' Syndrome (Atypical Erythema Multiforme Major)

The characteristics of erythema multiforme major, sometimes referred to as Stevens-Johnson syndrome, comprise both targetoid cutaneous eruptions and mucosal erythema, erosions or ulcers involving ≥ 2 different sites, including the oral region. Erythema multiforme major sometimes presents exclusively with mucosal involvement and is termed atypical erythema multiforme major or Fuchs' syndrome, because this German author

Table 1. Characteristics of patients with Fuchs' syndrome, varicella-like eruption, vasculitis and erythema nodosum associated with *M. pneumoniae* respiratory infection

	Fuchs' syndrome	Varicella-like eruption	Vasculitis	Erythema nodosum
Patients, n	37	8	21	11
Age, years				
Childhood	21	5	14	8
Adulthood	16	3	7	3
Gender				
Male	32 ^a	6 ^a	12	2 ^b
Female	5 ^a	2 ^a	9	7 ^b

Number of patients.

^a p < 0.01 vs. patients with erythema nodosum.

^b Information not available in 2 cases.

first reported the condition as 'herpes oris conjunctivae' approximately 150 years ago [98]. Our review disclosed 37 patients affected with Mycoplasma respiratory infection and oral mucositis [5–35]. In addition, ocular involvement was noted in 36, genital in 26 (22 male and 4 female subjects) and anal in 3 cases.

Varicella-Like Eruption

Eight cases of Mycoplasma infection developed vesicular lesions surrounded by an erythematous halo resembling varicella [36–41]. In these patients, the Tzanck preparation or the virus culture had been used to exclude varicella. Like in patients with atypical erythema multiforme major, lesions of the oral and ocular mucosae were noted in 5 of the 8 cases. Genital lesions were observed uniquely in 2 males.

Henoch-Schönlein Syndrome and Further Vasculitides

Leukocytoclastic vasculitides classically present as palpable purpura in the context of Henoch-Schönlein syndrome. Less commonly, they present with targetoid lesions and nonpitting edema in the context of acute hemorrhagic edema of young children, which is considered the infantile variant of Henoch-Schönlein syndrome, or with urticarial plaques in the context of urticarial vasculitis. Our review disclosed 21 cases of leukocytoclastic vasculitides associated with *M. pneumoniae* infection: 18 cases of Henoch-Schönlein syndrome, 1 case of acute hemorrhagic edema and 2 cases of urticarial vasculitis [42–62].

We found 14 case series addressing Henoch-Schönlein syndrome in a total of 2,186 cases published in English-

language journals between 1984 and 2014 [63–76]. None of the reports included cases associated with *M. pneumoniae* respiratory infection. It is true, however, that none of the articles specifically investigated this association. A systematic review dealing with acute hemorrhagic edema of young children published a few years ago did not find any further association with infections caused by *M. pneumoniae* [77].

Erythema Nodosum

Erythema nodosum is characterized by painful erythematous nodules located on the anterior surface of the legs. Our review disclosed 11 erythema nodosum cases associated with *M. pneumoniae* infection [4, 54, 58, 78–83]. No more than 1 case of erythema nodosum associated with *M. pneumoniae* respiratory infection was noted in 9 case series [4, 79, 84–90] including a total of 654 erythema nodosum cases not addressing this possible association. However, 4 cases were noted in 3 case series [79, 80, 91] addressing the mentioned association in a total of 77 erythema nodosum cases.

Other Possible Associations

Papular acrodermatitis Gianotti-Crosti presents with acral papular eruptions and usually occurs in association with a viral illness, most commonly hepatitis B virus, Epstein-Barr virus or picornaviruses. The association between Gianotti-Crosti syndrome and *M. pneumoniae* infection was observed in a 4-year-old girl [92] and in a 44-year-old woman [93]. A systematic review of the infections underlying this condition did not disclose any further case of Mycoplasma infection associated with Gianotti-Crosti syndrome [94].

Finally, there exists reasonable evidence that pityriasis rosea Gibert [95, 96] and pityriasis lichenoides et varioliformis acuta Mucha-Habermann [97] are not associated with Mycoplasma infections.

Discussion

There is a recognized causal relationship between *M. pneumoniae* respiratory infections and erythematous maculopapular (or urticarial) eruptions or erythema multiforme major [2]. To our knowledge, this is the first comprehensive review addressing a possible relationship with further skin eruptions. The survey focused on a possible relationship between *M. pneumoniae* infections and erythema nodosum, Fuchs' syndrome, Gianotti-Crosti syndrome, vasculitides such as Henoch-Schönlein syn-

Table 2. Skin lesions that have been associated with *M. pneumoniae* respiratory infections in the literature

Recognized causal relationship

- Erythematous maculopapular (or urticarial) lesions
- Erythema multiforme (both minor and major form)
- Fuchs' syndrome
- Varicella-like skin eruptions

Not well recognized causal relationship

- Vasculitides (most frequently Henoch-Schönlein syndrome)
- Erythema nodosum

drome, pityriasis rosea Gibert, pityriasis lichenoides et varioliformis acuta Mucha-Habermann and varicella-like eruptions (table 2).

Fuchs' syndrome is a rare form of erythema multiforme major that presents exclusively with mucosal involvement. Likewise, in patients with *M. pneumoniae* infections, varicella-like eruptions are usually associated with a mucous membrane involvement that resembles Fuchs' syndrome. These observations suggest that, like for classical erythema multiforme [2], the correlation between Mycoplasma and Fuchs' syndrome or varicella-like eruptions is causal.

Many cases of Henoch-Schönlein syndrome are preceded by a respiratory illness [99]. In this survey, a relationship between Mycoplasma infection and Henoch-Schönlein syndrome or other leukocytoclastic vasculitides was documented in no more than 21 cases. Erythema nodosum is traditionally associated with inflammatory bowel disease and with mycobacterial and streptococcal infections [100]. However, most cases are caused by different respiratory infections. Our survey shows a possible relationship between erythema nodosum and *M. pneumoniae* infection in 11 cases. Many cases of Henoch-Schönlein syndrome or erythema nodosum are preceded by symptoms and signs consistent with a mycoplasmal infection [99, 100]. Nonetheless, testing for *M. pneumoniae* is usually performed neither in Henoch-Schönlein syndrome nor in erythema nodosum. It is therefore concluded that these possible associations deserve further inquiry.

Gianotti-Crosti syndrome is usually associated with viral infections such as hepatitis B or infectious mononucleosis. Since a relationship between Mycoplasma infection and this condition has been documented exclusively in 2 cases, we assume that the association is accidental. A possible relationship between Mycoplasma infections and pityriasis rosea Gibert or pityriasis lichen-

noides et varioliformis acuta Mucha-Habermann has previously been debated, but recent evidence strongly argues against such an association.

The link between *M. pneumoniae* and skin lesions deserves some discussion. *M. pneumoniae* has sometimes been isolated from targetoid lesions, suggesting that in some instances they represent extrapulmonary spread of the infection [2]. Alternatively, skin lesions result from similarities shared between macromolecules found on Mycoplasma and in host tissue that trigger an immune response [2].

It is traditionally assumed that macrolides, tetracyclines or fluoroquinolones speed recovery in Mycoplasma respiratory infection but do not shorten the course of mucocutaneous manifestations [2, 3]. Recent data indicate that there is no clear evidence that antibiotic treatment of mycoplasmal pneumonia is effective [101].

References

- 1 Ludlam GB, Bridges JB: Association of Stevens-Johnson syndrome with antibody for *Mycoplasma pneumoniae*. Lancet 1964;283: 958–959.
- 2 Schalock PC, Dinulos JG: *Mycoplasma pneumoniae*-induced cutaneous disease. Int J Dermatol 2009;48:673–681.
- 3 Atkinson TP, Waites KB: *Mycoplasma pneumoniae* infections in childhood. Pediatr Infect Dis J 2014;33:92–94.
- 4 Aydin-Teke T, Tanir G, Bayhan GI, Metin O, Oz N: Erythema nodosum in children: evaluation of 39 patients. Turk J Pediatr 2014;56: 144–149.
- 5 Sieber OF Jr, John TJ, Fulginiti VA, Overholt EC: Stevens-Johnson syndrome associated with *Mycoplasma pneumoniae* infection. JAMA 1967;200:79–81.
- 6 Alter SJ, Stringer B: *Mycoplasma pneumoniae* infections associated with severe mucositis. Clin Pediatr (Phila) 1990;29:602–604.
- 7 Kirke S, Powell FC: Mucosal erosions and a cough. Ir Med J 2003;96:245.
- 8 Vanfleteren I, Van Gysel D, De Brandt C: Stevens-Johnson syndrome: a diagnostic challenge in the absence of skin lesions. Pediatr Dermatol 2003;20:52–56.
- 9 Schalock PC, Dinulos JG: *Mycoplasma pneumoniae*-induced Stevens-Johnson syndrome without skin lesions: fact or fiction? J Am Acad Dermatol 2005;52:312–315.
- 10 De Ru DH, Sukhai RN: Stevens-Johnson syndrome. Eur J Pediatr 2007;166:1303–1304.
- 11 Fearon D, Hesketh EL, Mitchell AE, Grimwood K: *Mycoplasma pneumoniae* infection complicated by pneumomediastinum and severe mucositis. J Paediatr Child Health 2007; 43:403–405.
- 12 Latsch K, Girsick HJ, Abele-Horn M: Stevens-Johnson syndrome without skin lesions. J Med Microbiol 2007;56:1696–1699.
- 13 Ravin KA, Rappaport LD, Zuckerbraun NS, Wadowsky RM, Wald ER, Michaels MM: *Mycoplasma pneumoniae* and atypical Stevens-Johnson syndrome: a case series. Pediatrics 2007;119:e1002–e1005.
- 14 Birch J, Chamlin S, Duerst R, Jacobsohn D: *Mycoplasma pneumoniae* and atypical Stevens-Johnson syndrome in a hematopoietic stem cell transplant recipient. Pediatr Blood Cancer 2008;50:1278–1279.
- 15 Figueira-Coelho J, Lourenço S, Pires AC, Mendonça P, Malhado JA: *Mycoplasma pneumoniae*-associated mucositis with minimal skin manifestations. Am J Clin Dermatol 2008;9:399–403.
- 16 Hillebrand-Haverkort ME, Budding AE, bij de Vaate LA, van Agtmael MA: *Mycoplasma pneumoniae* infection with incomplete Stevens-Johnson syndrome. Lancet Infect Dis 2008;8:586–587.
- 17 Li K, Haber RM: Stevens-Johnson syndrome without skin lesions (Fuchs syndrome): a literature review of adult cases with Mycoplasma cause. Arch Dermatol 2008;148:963–964.
- 18 Sendi P, Gruber P, Lepère F, Schiller P, Zimmerli W: *Mycoplasma pneumoniae* infection complicated by severe mucocutaneous lesions. Lancet Infect Dis 2008;8:268.
- 19 Walicka M, Majsterek M, Rakowska A, Słowińska M, Sicińska J, Górska B, Ptasińska M, Rudnicka L, Marcinowska-Suchowierska E: *Mycoplasma pneumoniae*-induced pneumonia with Stevens-Johnson syndrome of acute atypical course. Pol Arch Med Wewn 2008;118:449–453.
- 20 Artés Figueres M, Oltra Benavent M, Fernández Calatayud A, Revert Gomar M: Mucositis grave inducida por *Mycoplasma pneumoniae*. An Pediatr (Barc) 2009;71:573–574.
- 21 Choi S-H, Lee Y-M, Rha Y-H: *Mycoplasma pneumoniae*-induced Stevens-Johnson syndrome without skin manifestations. Korean J Pediatr 2009;52:247–250.
- 22 Havliza K, Jakob A, Rompel R: Erythema multiforme majus (Fuchs syndrome) associated with *Mycoplasma pneumoniae* infection in two patients. J Dtsch Dermatol Ges 2009;7: 445–448.
- 23 Jakob M, Stuhrmann N, Jordan J, Bootz F, Schröck A: Stomatitis, Cheilitis und Konjunktivitis nach grippalem Infekt. HNO 2009; 57:914–917.
- 24 Villarroel J, Bustamante MC, Denegri M, Pérez L: Manifestaciones muco-cutáneas de la infección por *Mycoplasma pneumoniae*: presentación de cuatro casos. Rev Chilena Infectol 2009;26:457–463.
- 25 Bressan S, Mion T, Andreola B, Bisogno G, Da Dalt L: Severe *Mycoplasma pneumoniae*-associated mucositis treated with immunoglobulins. Acta Paediatr 2011;100:e238–e240.
- 26 McGouran DC, Petterson T, McLaren JM, Wolbinski MP: Mucositis, conjunctivitis but no rash – the ‘Atypical Stevens-Johnson syndrome’. Acute Med 2011;10:81–82.
- 27 Meyer Sauteur PM, Gansser-Kälin U, Lautenschlager S, Goetschel P: Fuchs syndrome associated with *Mycoplasma pneumoniae* (Stevens-Johnson syndrome without skin lesions). Pediatr Dermatol 2011;28:474–476.
- 28 Ramasamy A, Patel C, Conlon C: Incomplete Stevens-Johnson syndrome secondary to atypical pneumonia. BMJ Case Rep 2011 DOI: 10.1136/bcr.08.2011.4568.

The results of our review must be viewed with an understanding of the inherent limitations of the analysis, which is based on the scanty available literature.

In conclusion, this review of the literature implies that *M. pneumoniae* may cause, in addition to erythematous maculopapular (or urticarial) eruptions and erythema multiforme, Fuchs’ syndrome and varicella-like eruptions. Furthermore, the link between this microorganism and vasculitides or erythema nodosum is intriguing and needs further investigations.

Disclosure Statement

The authors have no conflict of interest to declare.

- 29 Schomburg J, Vogel M: A 12-year-old boy with severe mucositis: extrapulmonary manifestation of *Mycoplasma pneumoniae* infection. *Klin Padiatr* 2012;224:94–95.
- 30 Hochreiter D, Jackson JM, Shetty AK: Fever, severe mucositis, and conjunctivitis in a 15-year-old male. *Clin Pediatr (Phila)* 2012; 51:1103–1105.
- 31 Trapp LW, Schrantz SJ, Joseph-Griffin MA, Hageman JR, Waskow SE: A 13-year-old boy with pharyngitis, oral ulcers, and dehydration. *Mycoplasma pneumoniae*-associated mucositis. *Pediatr Ann* 2013;42:148–150.
- 32 Yachoui R, Kolasinski SL, Feinstein DE: *Mycoplasma pneumoniae* with atypical Stevens-Johnson syndrome: a diagnostic challenge. *Case Rep Infect Dis* 2013;2013:457161.
- 33 Varghese C, Sharain K, Skalski J, Ramar K: *Mycoplasma pneumoniae*-associated mucositis. *BMJ Case Rep* 2014 DOI: 10.1136/bcr-2014-203795.
- 34 Šternberský J, Tichý M: Fuchs' syndrome (Stevens-Johnson syndrome without skin involvement) in an adult male – a case report and general characteristics of the sporadically diagnosed disease. *Acta Dermatovenerol Croat* 2014;22:284–287.
- 35 Vujić I, Shroff A, Grzelka M, Posch C, Monshī B, Sanlorenzo M, Ortiz-Urda S, Rappersberger K: *Mycoplasma pneumoniae*-associated mucositis – case report and systematic review of literature. *J Eur Acad Dermatol Venereol* 2015;29:595–598.
- 36 Teisch JA, Shapiro L, Walzer RA: Vesiculopustular eruption with mycoplasma infection. *JAMA* 1970;211:1694–1697.
- 37 Lascari AD, Garfunkel JM, Mauro DJ: Varicella-like rash associated with Mycoplasma infection. *Am J Dis Child* 1974;128:254–255.
- 38 Ulivelli A, Cocchi P, Paoletti A: Rash varicelliforme associato a infezione da micoplasma. *Minerva Pediatr* 1976;28:32–35.
- 39 Chusid MJ, Lachmann BS, Lazerson J: Severe *Mycoplasma pneumoniae* and vesicular eruption in SC hemoglobinopathy. *J Pediatr* 1978;93:449–451.
- 40 Sequeira W, Jones E, Bronson DM: *Mycoplasma pneumoniae* infection with arthritis and a varicella-like eruption. *JAMA* 1981;246: 1936–1937.
- 41 Winnock T, Wang J, Suys E, De Coninck A, Roseeuw D: Vesiculopustular eruption associated with *Mycoplasma pneumoniae* pneumopathy. *Dermatology* 1996;192:73–74.
- 42 Sussman M, Jones JH, Almeida JD, Lachmann PJ: Deficiency of the second component of complement associated with anaphylactoid purpura and presence of mycoplasma in the serum. *Clin Exp Immunol* 1973;14: 531–539.
- 43 Liew SW, Kessel I: Mycoplasmal pneumonia preceding Henoch-Schönlein purpura. *Arch Dis Child* 1974;49:912–913.
- 44 Steare SE, Wiselka MJ, Kurinczuk JJ, Nicholson KG: *Mycoplasma pneumoniae* infection associated with Henoch-Schönlein purpura. *J Infect* 1988;16:305–307.
- 45 Van Bever HP, Van Doorn JW, Demey HE: Adult respiratory distress syndrome associated with *Mycoplasma pneumoniae* infection. *Eur J Pediatr* 1992;151:227–228.
- 46 Callens A, Machet MC, Vaillant L, Pelucio-Lopes C, Machet L, Barruet K, Lorette G: Purpura rhumatoïde de l'adulte et infection à mycoplasme. *Ann Med Interne (Paris)* 1993;144: 297–299.
- 47 Perez C, Mendoza H, Hernandez R, Valcayo A, Guardo R: Leukocytoclastic vasculitis and polyarthritides associated with *Mycoplasma pneumoniae* infection. *Clin Infect Dis* 199;25: 154–155.
- 48 Kaneko K, Fujinaga S, Ohtomo Y, Nagaoka R, Obinata K, Yamashiro Y: *Mycoplasma pneumoniae*-associated Henoch-Schönlein purpura nephritis. *Pediatr Nephrol* 1999;13: 1000–1001.
- 49 Perez C, Montes M: Cutaneous leukocytoclastic vasculitis and encephalitis associated with *Mycoplasma pneumoniae* infection. *Arch Intern Med* 2002;162:352–354.
- 50 Jover F, Cuadrado JM, Ivars J, Merino J: Vasculitis urticariforme e infección por *Mycoplasma pneumoniae*. *Enferm Infect Microbiol Clin* 2003;21:215–219.
- 51 Orlandini V, Dega H, Dubertret L: Vasculite cutanée révélant une infection à *Mycoplasma pneumoniae*. *Presse Med* 2004;33: 1365–1366.
- 52 Timitilli A, Di Rocco M, Nattero G, Tacchella A, Giacchino R: Unusual manifestations of infections due to *Mycoplasma pneumoniae* in children. *Infez Med* 2004;12:113–117.
- 53 Greco F, Sorge A, Salvo V, Sorge G: Cutaneous vasculitis associated with *Mycoplasma pneumoniae* infection: case report and literature review. *Clin Pediatr (Phila)* 2007;46:451–453.
- 54 Kano Y, Mitsuyama Y, Hirahara K, Shiohara T: *Mycoplasma pneumoniae* infection-induced erythema nodosum, anaphylactoid purpura, and acute urticaria in 3 people in a single family. *J Am Acad Dermatol* 2007;57(2 suppl):S33–S35.
- 55 Fukuda S, Takahashi T, Kumori K, Takahashi Y, Yasuda K, Kasai T, Yamaguchi S: Idiopathic testicular infarction in a boy initially suspected to have acute epididymo-orchitis associated with mycoplasma infection and Henoch-Schönlein purpura. *J Pediatr Urol* 2009; 5:68–71.
- 56 Lim CSH, Lim SL: Henoch-Schönlein purpura associated with *Mycoplasma pneumoniae* infection. *Cutis* 2011;87:273–276.
- 57 Dua J, Nandagudi A, Sutcliffe N: *Mycoplasma pneumoniae* infection associated with urticarial vasculitis mimicking adult-onset Still's disease. *Rheumatol Int* 2012;32:4053–4056.
- 58 Shimizu M, Hamaguchi Y, Matsushita T, Sakakibara Y, Yachie A: Sequentially appearing erythema nodosum, erythema multiforme and Henoch-Schönlein purpura in a patient with *Mycoplasma pneumoniae* infection: a case report. *J Med Case Rep* 2012;6:398.
- 59 Trčko K, Marko PB, Miljković J: Leukocytoclastic vasculitis induced by *Mycoplasma pneumoniae* infection. *Acta Dermatovenerol Croat* 2012;20:119–122.
- 60 Yiallouros P, Moustaki M, Voutsioti A, Sharifi F, Karpathios T: Association of *Mycoplasma pneumoniae* infection with Henoch-Schönlein purpura. *Prague Med Rep* 2013; 114:177–179.
- 61 Di Lernia V: *Mycoplasma pneumoniae*: an aetiological agent of acute haemorrhagic oedema of infancy. *Australas J Dermatol* 2014; 55:e69–e70.
- 62 Hu P, Guan Y, Lu L: Henoch-Schönlein purpura triggered by *Mycoplasma pneumoniae* in a female infant. *Kaohsiung J Med Sci* 2015;31: 163–164.
- 63 Saulsbury FT: Henoch-Schönlein purpura. *Pediatr Dermatol* 1984;1:195–201.
- 64 Kumar L, Singh S, Goraya JS, Uppal B, Kakkar S, Walker R, Sehgal S: Henoch-Schönlein purpura: the Chandigarh experience. *Indian Pediatr* 1998;35:19–25.
- 65 Nussinovitch M, Prais D, Finkelstein Y, Varsano I: Cutaneous manifestations of Henoch-Schönlein purpura in young children. *Pediatr Dermatol* 1998;15:426–428.
- 66 Saulsbury FT: Henoch-Schönlein purpura in children. Report of 100 patients and review of the literature. *Medicine (Baltimore)* 1999;78: 395–409.
- 67 Calviño MC, Llorca J, García-Porrúa C, Fernández-Iglesias JL, Rodríguez-Ledo P, González-Gay MA: Henoch-Schönlein purpura in children from northwestern Spain: a 20-year epidemiologic and clinical study. *Medicine (Baltimore)* 2001;80:279–290.
- 68 Trapani S, Micheli A, Grisolía F, Resti M, Chiappini E, Falcini F, De Martino M: Henoch Schönlein purpura in childhood: epidemiological and clinical analysis of 150 cases over a 5-year period and review of literature. *Semin Arthritis Rheum* 2005;35:143–153.
- 69 Cakir M, Orhan F, Mungan I, Sonmez FM, Aslan Y, Kalyoncu M, Erduran E, Gedik Y, Okten A: Henoch-Schönlein purpura in north-eastern Turkey. *Ann Trop Paediatr* 2006;26:59–65.
- 70 Nong BR, Huang YF, Chuang CM, Liu CC, Hsieh KS: Fifteen-year experience of children with Henoch-Schönlein purpura in southern Taiwan, 1991–2005. *J Microbiol Immunol Infect* 2007;40:371–376.
- 71 Peru H, Soylemezoglu O, Bakkaloglu SA, Elmas S, Bozkaya D, Elmali AM, Kara F, Buyan N, Hasanoglu E: Henoch Schönlein purpura in childhood: clinical analysis of 254 cases over a 3-year period. *Clin Rheumatol* 2008;27: 1087–1092.
- 72 Anil M, Aksu N, Kara OD, Bal A, Anil AB, Yavascan O, Un B: Henoch-Schönlein purpura in children from western Turkey: a retrospective analysis of 430 cases. *Turk J Pediatr* 2009;51:429–436.
- 73 Hung SP, Yang YH, Lin YT, Wang LC, Lee JH, Chiang BL: Clinical manifestations and outcomes of Henoch-Schönlein purpura: comparison between adults and children. *Pediatr Neonatol* 2009;50:162–168.

- 74 Watson L, Richardson AR, Holt RC, Jones CA, Beresford MW: Henoch Schölein purpura – a 5-year review and proposed pathway. *PLoS One* 2012;7:e29512.
- 75 Chen O, Zhu XB, Ren P, Wang YB, Sun RP, Wei DE: Henoch Schölein purpura in children: clinical analysis of 120 cases. *Afr Health Sci* 2013;13:94–99.
- 76 Calvo-Río V, Loricera J, Mata C, Martín L, Ortiz-Sanjuán F, Alvarez L, González-Vela MC, González-Lamuño D, Rueda-Gotor J, Fernández-Llaca H, González-López MA, Armesto S, Peiró E, Arias M, González-Gay MA, Blanco R: Henoch-Schönlein purpura in northern Spain: clinical spectrum of the disease in 417 patients from a single center. *Medicine (Baltimore)* 2014;93:106–113.
- 77 Fiore E, Rizzi M, Simonetti GD, Garzoni L, Bianchetti MG, Bettinelli A: Acute hemorrhagic edema of young children: a concise narrative review. *Eur J Pediatr* 2011;170: 1507–1511.
- 78 Teyssandier R, Guidet B, Pinta P, Offenstadt G: Pneumopathie à *Mycoplasma pneumoniae* avec anémie grave et érythème noueux. *Presse Med* 1985;14:1613.
- 79 Cribier B, Caille A, Heid E, Grosshans E: Erythema nodosum and associated diseases. A study of 129 cases. *Int J Dermatol* 1998;37: 667–672.
- 80 Kakourou T, Drosatou P, Psychou F, Aroni K, Nicolaïdou P: Erythema nodosum in children: a prospective study. *J Am Acad Dermatol* 2001;44:17–21.
- 81 Rodríguez Blanco MA, Martín Morales JM, Casas Fernández L: Nódulos eritematovioláceos y neumonía. *An Esp Pediatr* 2002;57: 279–280.
- 82 Kano Y, Mitsuyama Y, Hirahara K, Shiohara T: *Mycoplasma pneumoniae* infection-induced erythema nodosum, anaphylactoid purpura, and acute urticaria in 3 people in a single family. *J Am Acad Dermatol* 2007; 57(suppl):S33–S35.
- 83 Greco F, Catania R, Pira AL, Saporito M, Scalora L, Aguglia MG, Smilari P, Sorge G: Erythema nodosum and *Mycoplasma pneumoniae* infections in childhood: further observations in two patients and a literature review. *J Clin Med Res* 2015;7:274–277.
- 84 Labbé L, Perel Y, Maleville J, Taïeb A: Erythema nodosum in children: a study of 27 patients. *Pediatr Dermatol* 1996;13:447–450.
- 85 Hassink RI, Pasquini-Egli CE, Jacomella V, Laux-End R, Bianchetti MG: Conditions currently associated with erythema nodosum in Swiss children. *Eur J Pediatr* 1997;156:851–853.
- 86 García-Porrúa C, González-Gay MA, Vázquez-Caruncho M, López-Lazaro L, Luero M, Fernández ML, Alvarez-Ferreira J, Pujol RM: Erythema nodosum: etiologic and predictive factors in a defined population. *Arthritis Rheum* 2000;43:584–592 (Erratum in: *Arthritis Rheum* 2000;43:1061).
- 87 Garty BZ, Poznanski O: Erythema nodosum in Israeli children. *Isr Med Assoc J* 2000;2: 145–146.
- 88 Psychos DN, Voulgari PV, Skopouli FN, Drosos AA, Moutsopoulos HM: Erythema nodosum: the underlying conditions. *Clin Rheumatol* 2000;19:212–216.
- 89 Tay YK: Erythema nodosum in Singapore. *Clin Exp Dermatol* 2000;25:377–380.
- 90 Mert A, Kumbasar H, Ozaras R, Erten S, Tasli L, Tabak F, Ozturk R: Erythema nodosum: an evaluation of 100 cases. *Clin Exp Rheumatol* 2007;25:563–570.
- 91 Cengiz AB, Kara A, Kanra G, Seçmeer G, Ceyhan M: Erythema nodosum in childhood: evaluation of ten patients. *Turk J Pediatr* 2006;48:38–42.
- 92 Angoulvant N, Grézard P, Wolf F, Truchot F, Marcilly MC, Perrot H: Infection aiguë à *Mycoplasma pneumoniae*: nouvelle cause de syndrome de Gianotti Crosti. *Presse Med* 2000;29:1287.
- 93 Manoharan S, Muir J, Williamson R: Gianotti-Crosti syndrome in an adult following recent *Mycoplasma pneumoniae* infection. *Australas J Dermatol* 2005;46:106–109.
- 94 Brandt O, Abeck D, Gianotti R, Burgdorf W: Gianotti-Crosti syndrome. *J Am Acad Dermatol* 2006;54:136–145.
- 95 Chuh AA, Chan HH: Prospective case-control study of chlamydia, legionella and mycoplasma infections in patients with pityriasis rosea. *Eur J Dermatol* 2002;12:170–173.
- 96 Chuh A, Chan H, Zawar V: Pityriasis rosea – evidence for and against an infectious aetiology. *Epidemiol Infect* 2004;132:381–390.
- 97 Fernandes NF, Rozdeba PJ, Schwartz RA, Kihiczak G, Lambert WC: Pityriasis lichenoides et varioliformis acuta: a disease spectrum. *Int J Dermatol* 2010;49:257–261.
- 98 Fuchs E: Herpes oris conjunctivae. *Klin Monbl Augenheilkd* 1876;14:333–351.
- 99 Ting TV: Diagnosis and management of cutaneous vasculitis in children. *Pediatr Clin North Am* 2014;61:321–346.
- 100 Requena L, Yus ES: Erythema nodosum. *Dermatol Clin* 2008;26:425–438.
- 101 Biondi E, McCulloh R, Alverson B, Klein A, Dixon A: Treatment of mycoplasma pneumonia: a systematic review. *Pediatrics* 2014; 133:1081–1090.