



Original Article

Bladder management and urological complications in patients with chronic spinal cord injuries in Taiwan



Sheng-Fu Chen, Yuan-Hong Jiang, Jia-Fong Jhang, Cheng-Ling Lee, Hann-Chorng Kuo*

Department of Urology, Buddhist Tzu Chi General Hospital and Tzu Chi University, Hualien, Taiwan

ARTICLE INFO

Article history:

Received 20 August 2013

Received in revised form

30 August 2013

Accepted 6 September 2013

Keywords:

Detrusor overactivity

Lower urinary tract dysfunction

Neurogenic bladder

Urological complication

ABSTRACT

Objectives: Neurogenic voiding dysfunction (NVD) results in lower urinary tract symptoms and upper urinary tract complications. Management of urinary tract dysfunction in patients with spinal cord injury (SCI) must be based on patient needs and urological complications. This study surveyed the bladder management and urological complications in patients with SCI in Taiwan.

Materials and methods: A total of 894 patients with SCI were surveyed throughout Taiwan over a period of 5 years (2007–2011). All patients received neurological and urological examinations, renal sonography, bladder sonography, and urinalysis. They were further requested to report urinary tract infections (UTI), voiding conditions, and bladder management in the past 3 years. The bladder management and urological complications were analyzed based on different SCI levels and duration of disease.

Results: Among all patients, 39.7% voided spontaneously or by reflex, 23.8% used percussion to void, 20.9% voided by abdominal pressure, 18.1% used clean intermittent catheterization (CIC), and 22.9% had indwelling catheters or cystostomy. Detrusor sphincter dyssynergia (DSD) was noted in 39.7% of patients and autonomic dysreflexia (AD) in 19.9%. UTI was noted in 483 (54%) patients, hydronephrosis in 110 (12.3%) patients, and severe urinary incontinence in 257 (28.7%) patients. UTI occurred significantly more often in patients without, than with normal voiding. CIC was more frequently used in patients with duration of SCI < 1 year, but the rate of CIC significantly decreased as the duration of SCI increased.

Conclusion: The rates of urological complications in patients with chronic SCI remained high in this survey in Taiwan. Patients with a duration of SCI > 5 years chose indwelling catheters or a cystostomy more often than CIC.

Copyright © 2013, Buddhist Compassion Relief Tzu Chi Foundation. Published by Elsevier Taiwan LLC. All rights reserved.

1. Introduction

Neurogenic voiding dysfunction (NVD) includes dysfunction of the urinary bladder and urethra, due to central nervous system or peripheral neurogenic lesions. Cerebrovascular accidents, Parkinson's disease, cerebral palsy, multiple sclerosis, transverse myelitis, and spinal cord lesions can result in NVD [1]. Among all types of NVD, spinal cord injury (SCI) can lead to the most difficult to manage voiding disorders. A significant association between the level of injury and the type of voiding dysfunction has been noted in patients with a single level of SCI. Management of the urinary tract in patients with SCI must be based on urodynamic findings rather than inferences from neurologic evaluation [2].

Conflicts of interest: none.

* Corresponding author. Department of Urology, Buddhist Tzu Chi General Hospital, 707, Section 3, Chung-Yang Road, Hualien, Taiwan. Tel.: +886 3 8561825x2117; fax: +886 3 8560794.

E-mail address: hck@tzuchi.com.tw (H.-C. Kuo).

The main problems of NVD are failure to store due to detrusor overactivity (DO) or urethral incompetence, and failure to empty due to detrusor areflexia, bladder neck dysfunction, or detrusor sphincter dyssynergia (DSD), and combined failure to store and empty due to DSD or DO and impaired contractility [3]. The priorities in the management of NVD should be preservation of renal function, freedom from urinary tract infection (UTI), and efficient bladder emptying [3]. It is important to screen patients at high risk, including those with complete neurological lesions, cervical spinal cord paraplegia, prolonged indwelling catheters, high detrusor leak-point pressure, presence of DSD and autonomic dysreflexia (AD), large postvoid residual, and vesicoureteral reflux [4]. Correction of urological complications and improving the quality of life in urination are the two main goals in the management of NVD. In addition, individualized treatment strategies for each patient should be carefully evaluated.

The worldwide incidence of SCI reported in the literature ranges from 12.1 to 57.8 per million [5]. Bladder dysfunction occurs

Table 1
The level of spinal cord injury and vesicourethral dysfunction.

Level of SCI	N	Normal	Detrusor areflexia	Detrusor hyperreflexia	DSD	Autonomic dysreflexia
Total	894	115 (12.9)	232 (26.0)	534 (59.7)	355 (39.7)	178 (19.9)
Cervical	395	49 (12.6)	21 (5.4)	320 (82.1)	238 (61)	151 (38.7)
Thoracic	344	23 (6.9)	131 (39.1)	187 (55.8)	109 (32.5)	26 (7.8)
Lumbar	148	38 (26.8)	79 (55.6)	26 (18.3)	8 (5.6)	1 (0.7)
Sacral	7	5 (71.4)	1 (14.3)	1 (14.3)	0	0
<i>p</i>		<0.001	<0.001	<0.001	<0.001	<0.001

Data are presented as *n* (%).

DSD = detrusor sphincter dyssynergia; SCI = spinal cord injury.

depending on the level and location of the SCI [6]. One study of neurogenic DO (NDO) in SCI in Taiwan from 2006 to 2008 revealed that the prevalence rate over the 3 years was 855 per million, and the incidence rate was 241 per million person-years, similar to a previous study and much higher than in other countries [7]. Eighty-two percent of all patients with SCI and NDO used catheterization with (46%) or without (36%) pharmacological treatment.

We previously reported a survey of urological complications in SCI patients in Taiwan from 1995 to 1998 [8]. In that report, the overall complication rate was 40%. This study reported a recent survey of the voiding management and urological complications in patients with chronic SCI in Taiwan from 2007 through to 2011.

2. Materials and methods

Beginning in 2007, the authors conducted a cross national survey of SCI patients in Taiwan. We performed health education and urological examinations for SCI patients in 14 counties of Taiwan, in conjunction with the SCI association of each county. All patients in the 28 surveys were visited and only the data from the first visit were collected in this study.

During the SCI survey, all patients received neurological and urological examinations, including a physical examination, renal sonography, bladder sonography, and urinalysis. The level of SCI was classified as cervical, thoracic, lumbar, and sacral based on the history and neurological examination. Patients' voiding dysfunction was classified as normal, reflex, detrusor areflexia, detrusor hyperreflexia, or DSD, according to the history and neurological examination. If patients had symptoms of headache, hypertension, and sweating at full bladder, they were considered to have AD.

Current bladder management was also recorded including normal spontaneous voiding, reflex voiding, percussion, abdominal pressure to void, clean intermittent catheterization (CIC)/clean intermittent self-catheterization (CISC), indwelling urethral catheter, cystostomy, or urinary diversion with a neobladder or ileostoma. Urological complications included symptomatic UTIs, hydronephrosis, renal stones, contracted kidneys, trabeculated bladder, and bladder stones. Symptomatic UTI was defined if

patients had documented acute pyelonephritis, acute cystitis, or urosepsis during the previous 3 years. Hydronephrosis was defined if renal sonography disclosed dilatation of the renal pelvis and calyces. The patients were further requested to report episodes of UTI, voiding conditions, and bladder management in the past 3 years.

The types of bladder management and urological complications were analyzed according to SCI level and disease duration. Episodes of UTI were further analyzed according to the type of bladder management. For all analyses, the variables are presented as patient number and percentage of the group. Categorical data were analyzed using the Chi-square test with Fisher's exact probability test as appropriate. A *p* value of <0.05 was considered statistically significant. The statistics program used for the analysis was SPSS version 17.0 (SPSS, Chicago, IL, USA) for Windows.

3. Results

A total of 894 patients with SCI completed the survey and urological examinations. Their mean age was 43.4 ± 13.3 years (range, 14–80 years). Among the patients, 395 (44.2%) had cervical SCI, 344 (38.5%) had thoracic SCI, 148 (16.6%) had lumbar SCI, and 7 (0.8%) had sacral SCI. No significant difference was noted in the mean age and sex distribution among different levels of SCI. The duration of SCI was <1 year in 77 patients, ≥ 1 –10 years in 391 patients, 11–20 years in 284 patients, and ≥ 20 years in 142 patients.

The vesicourethral dysfunction at different levels of SCI is shown in Table 1. Detrusor hyperreflexia was noted significantly more in patients with cervical and thoracic SCI than in those with lumbar or sacral SCI, whereas significantly more patients with lumbar SCI had detrusor areflexia. DSD was noted in 61% of patients and AD was noted in 38.7% of patients with cervical SCI.

Table 2 shows the bladder management in patients with different levels of SCI. Among all patients, 18.2% used CIC or CISC to empty the bladder, 11.4% used indwelling Foley catheters, and 11.5% had a cystostomy. Patients with SCI for <5 years used CIC/CISC more often than an indwelling catheter to empty the bladder, but indwelling catheters or cystostomies were used more often by those with SCI for >5 years (Fig. 1). The rate ratios of CIC/CISC to indwelling catheter/cystostomy were 3.31 for a duration of SCI of <1 year, 1.35 for 2–5 years, 0.75 for 6–10 years, 1.0 for 11–15 years, 0.59 for 16–20 years, and 0.68 for ≥ 20 years. Indwelling catheters/cystostomies were chosen because of frequent UTIs and hydronephrosis ($n = 72$, 35.1%), convenience in daily working life ($n = 65$, 31.7%), tetraplegia and lack of a care giver ($n = 42$, 20.5%), and severe urinary incontinence ($n = 26$, 12.7%).

We also surveyed urological complications in SCI patients in the past 3 years. Symptomatic UTI (>1 episode per year) was reported by 54.0% of patients, hydronephrosis was detected in 12.3%, severe urinary incontinence in 28.7%, and urolithiasis in 4.0%. The occurrence of symptomatic UTI ($p = 0.343$) and hydronephrosis

Table 2
Bladder management in patients with different levels of spinal cord injury.

SCI level	N	Normal/reflex voiding	Percussion to void	Abdominal pressure	CIC/CISC	Foley catheter	Cystostomy	Neobladder
Total	894	355 (39.7)	213 (23.8)	187 (20.9)	163 (18.2)	102 (11.4)	103 (11.5)	7 (0.8)
Cervical	395	191 (48.4)	109 (27.6)	31 (7.8)	66 (16.7)	42 (10.6)	55 (13.9)	5 (1.3)
Thoracic	344	102 (29.7)	79 (23.0)	94 (27.3)	73 (21.2)	51 (14.8)	31 (9.0)	2 (0.6)
Lumbar	148	57 (38.5)	23 (15.5)	61 (41.2)	22 (14.9)	9 (6.1)	17 (11.5)	0
Sacral	7	5 (71.4)	2 (28.6)	1 (14.3)	2 (28.6)	0	0	0
<i>p</i>		<0.001	0.030	<0.001	0.230	0.026	0.152	0.460

Data are presented as *n* (%).

Patient may use more than one bladder management.

CIC/CISC = clean intermittent catheterization/clean intermittent self-catheterization; SCI = spinal cord injury.




Fig. 1. Use of clean intermittent catheterization/clean intermittent self-catheterization (CIC/CISC) and cystostomy/indwelling catheters by patients with different durations of spinal cord injury (SCI).

($p = 0.058$) was not significantly different among different levels of SCI, but severe urinary incontinence was noted more often in patients with cervical and thoracic SCI than lumbar SCI ($p < 0.001$). Urolithiasis was more frequent in patients with sacral SCI than other levels of SCI ($p < 0.001$; Table 3). There were no significant differences in the occurrence rate of urological complications among different durations of SCI (Table 4).

UTI was noted in 238 (32.5%) of the 732 patients with available urinalysis results. UTI occurred significantly less often in patients with normal voiding than in those with other methods of bladder management ($p = 0.001$). However, there was no significant difference in the rate of UTI among patients with different methods of bladder management (Table 5).

4. Discussion

This large cohort survey reveals that urological complication rates remain high in patients with chronic SCI in Taiwan. Symptomatic UTI, hydronephrosis, and severe urinary incontinence were highly prevalent in SCI patients. UTI occurred more frequently in patients without normal voiding, and SCI patients preferred indwelling catheters or a cystostomy rather than CIC or CISC as the duration of SCI increased over 5 years.

Conservative management is the mainstay of urological treatment for NVD. Patients can be instructed to void by abdominal stimulation, the Crede maneuver, or abdominal straining. If they do not have a balanced bladder after training, CIC from a caregiver or CISC is necessary. Spontaneous voiding with and without triggered voiding and/or bladder expression has proven to be less safe except in well-defined patients with regular urological follow-up [9]. In SCI patients with DSD, long-term antimuscarinic therapy, in conjunction with CIC/CIS, might be necessary to avoid persistently

Table 3
Urological complications in different levels of spinal cord injury.

SCI level	N	Symptomatic UTI >1/y	Hydronephrosis	Severe urinary incontinence	Urolithiasis
Total	894	483 (54.0)	110 (12.3)	257 (28.7)	36 (4.0)
Cervical	395	210 (53.2)	46 (11.6)	118 (29.9)	8 (2.0)
Thoracic	344	184 (53.5)	53 (15.4)	115 (33.4)	14 (4.7)
Lumbar	148	87 (58.8)	11 (7.4)	22 (14.9)	12 (8.1)
Sacral	7	2 (28.6)	0	2 (28.6)	2 (28.6)
<i>p</i>		0.343	0.058	<0.001	<0.001

Data are presented as *n* (%).

SCI = spinal cord injury; UTI = urinary tract infection.

Table 4

Occurrence of symptomatic urinary tract infection (UTI) (>1/year) and hydronephrosis in spinal cord injury of different durations.

Duration of SCI	N	Symptomatic UTI > 1/year	Hydronephrosis	Severe urinary incontinence	Urolithiasis
Total	894	483 (54.0)	110 (12.3)	257 (28.7)	36 (4.0)
<1 y	77	40 (51.9)	7 (9.1)	23 (29.9)	4 (5.2)
1–10 y	391	216 (55.2)	52 (13.3)	110 (28.1)	9 (2.3)
11–20 y	284	152 (53.5)	34 (12.0)	81 (28.5)	19 (6.7)
>20 y	142	75 (52.8)	17 (12.0)	43 (30.3)	4 (2.8)
<i>p</i>		0.923	0.786	0.962	0.029

Data are presented as *n* (%).

SCI = spinal cord injury; UTI = urinary tract infection.

elevated intravesical pressure, which jeopardizes upper tract function [10–12].

Long-term indwelling catheters should be avoided except for patients with tetraplegia, or those who are bedbound, in whom an indwelling urethral catheter or suprapubic cystostomy may be an alternative [13,14]. However, because of a lack of public supportive facilities and Medicare, patients with chronic SCI cannot use CIC/CISC, and they often choose a more convenient method of bladder management. During the first few years after SCI, more patients used CIC/CISC because they were instructed to do so in the hospital. However, when they returned to the community with a long duration of SCI, they needed a more convenient method for daily life and to prevent urological complications. Nevertheless, UTI did not increase in patients using indwelling catheters or a cystostomy in comparison with other methods of bladder management.

CIC/CISC has been documented as the standard treatment for patients who are unable to empty the bladder [15]. CIC is a superior method for preserving bladder compliance and preventing upper tract complications associated with low compliance [16]. However, patients should be well instructed in the technique and risks of CIC/CISC. Less frequent catheterization results in higher catheterization volumes and a higher risk of UTI. Insufficient patient education and the inherent greater risk of UTI in patients with NVD are contributing factors [15,17].

Correction of urological complications and improving quality of life in urination are the two main goals for management of NVD. The level of SCI and the type of voiding dysfunction are closely associated, but not absolutely correlated [2,18]. Some patients with more than one lesion may also have different estimated voiding dysfunctions. It is important to screen SCI patients at high risk. When the detrusor leak point pressure is >40 cmH₂O, the upper tract is endangered [19,20]. Our study did not find differences in symptomatic UTI and hydronephrosis among different SCI levels or different disease durations. Therefore, close follow-up of UTI and the renal and bladder condition are important for patients with all levels of SCI and any disease duration.

In this study, the rate of UTI in patients who used CIC/CISC was not lower than that in those with other methods of bladder management, indicating that patients' education in the correct technique for CIC/CISC is not adequate. The other possible reason is a lack of annual monitoring of the bladder condition and resetting of a suitable and correct CIC program. Although patients with NVD may be properly diagnosed and treated in the first few years, all patients should receive life-long surveillance to prevent development of urological complications and undesired lower urinary tract symptoms [4,21,22].

Renal function is an important issue in the management of NVD, especially in patients with chronic SCI. Patients with DSD, low bladder compliance, and high intravesical pressure at end-bladder filling, may be at high risk of renal failure. The incidence of chronic renal disease is

Table 5
Relationship between bladder management and urinalysis results at spinal cord injury survey.

UTI	Normal voiding	Reflex voiding	Percussion to void	Abdominal pressure	CIC/CISC	Foley catheter	Cystostomy
Normal	112 (84.2)	97 (62.2)	47 (64.4)	72 (64.9)	44 (58.7)	63 (67.0)	48 (62.3)
UTI	21 (15.8)	59 (37.8)	26 (35.6)	39 (35.1)	31 (41.3)	31 (33.0)	29 (37.7)
Total	133	156	73	111	75	94	77

Data are presented as *n* (%).

CIC/CISC = clean intermittent catheterization/clean intermittent self-catheterization; UTI = urinary tract infection.

higher in those with paraplegia than in the healthy population [23]. Bladder management affects bladder compliance and changes in compliance with time. Patients with DSD who are currently using an indwelling catheter, performing CIC, or voiding spontaneously, should be monitored annually to prevent renal failure.

The results of this study revealed that 59.7% of SCI patients had detrusor hyperreflexia, 39.7% had DSD, and 19.9% had AD, suggesting that a high percentage of SCI patients are at high risk of upper tract deterioration. It is not surprising to know that 54% of patients in this study had frequent UTIs and 12.3% had hydro-nephrosis. Furthermore, the rates of frequent UTI and hydro-nephrosis did not change with disease duration, indicating that urological complications might occur at any stage after SCI. Therefore, patients with NVD should be regularly followed up for lower urinary tract dysfunction and any urological complication should be adequately treated.

5. Conclusion

In this large cohort SCI survey in Taiwan, urological complications in patients with chronic SCI remained high. Patients using methods of bladder management other than normal voiding had a significantly higher rate of UTI than those who voided normally. Patients with SCI > 5 years preferred indwelling catheters or cystostomies rather than CIC. Improvement in patient education, regular urinary tract surveillance, and correct medical treatment are mandatory to decrease the rate of urological complications and improve the quality of life of patients with chronic SCI.

References

- Abrams P, Andersson KE, Birder L, Brubaker L, Cardozo L, Chapple C, et al. Fourth International Consultation on Incontinence Recommendations of the International Scientific Committee: evaluation and treatment of urinary incontinence, pelvic organ prolapse, and fecal incontinence. *Neurourol Urodyn* 2010;29:213–40.
- Weld KJ, Dmochowski RR. Association of level of injury and bladder behavior in patients with post-traumatic spinal cord injury. *Urology* 2000;55:490–4.
- Kuo HC. Quality of life after active urological management of chronic spinal cord injury in eastern Taiwan. *Eur Urol* 1998;34:37–46.
- Stöhrer M, Kramer G, Löchner-Ernst D, Goepel M, Noll F, Rübber H. Diagnosis and treatment of bladder dysfunction in spinal cord injury patients. *Eur Urol Update Series* 1994;3:170–5.
- Van den Berg ME, Castellote JM, Mahillo-Fernandez I, de Pedro-Cuesta J. Incidence of spinal cord injury worldwide: a systematic review. *Neuro-epidemiology* 2010;34:184–92.
- Jeong SJ, Cho SY, Oh SJ. Spinal cord/brain injury and the neurogenic bladder. *Urol Clin North Am* 2010;37:537–46.
- Lai CC, Kao YH, Kuo HC, Ng K, Cheng E. Prevalence of neurogenic detrusor overactivity associated with spinal cord injury in Taiwan. *ICS Annual Meeting Proceedings* 2012:358 (Abstract).
- Kuo HC. Investigation of the voiding management of chronic spinal cord injured patients in Taiwan. *Tzu Chi Med J* 1999;11:69–78.
- Wyndaele JJ, Madersbacher H, Kovindha A. Conservative treatment of the neuropathic bladder in spinal cord injured patients. *Spinal Cord* 2001;39:294–300.
- Gajewski JB, Awad SA. Oxybutynin versus propanteline in patients with multiple sclerosis and detrusor hyperreflexia. *J Urol* 1986;135:966–8.
- Stöhrer M, Madersbacher H, Richter R, Wehnert J, Dreikorn K. Efficacy and safety of propiverine in SCI patients suffering from detrusor hyperreflexia—a double-blind, placebo-controlled clinical trial. *Spinal Cord* 1999;37:196–200.
- Thuroff JW, Bunke B, Ebner A, Faber P, de Geeter P, Hannappel J, et al. Randomized, double-blind, multicenter trial on treatment of frequency, urgency and incontinence related to detrusor hyperactivity: oxybutynin versus propanteline versus placebo. *J Urol* 1991;145:813–6.
- Weld KJ, Dmochowski RR. Effect of bladder management on urological complications in spinal cord injured patients. *J Urol* 2000;163:768–72.
- Mitsui T, Minami K, Furuno T, Morita H, Koyanagi T. Is suprapubic cystostomy an optimal urinary management in high quadriplegics? A comparative study of suprapubic cystostomy and clean intermittent catheterization. *Eur Urol* 2000;38:434–8.
- Madersbacher H, Wyndaele JJ, Igawa Y, Chancellor M, Chartier-Kastler E, Kovindha A. Conservative management in neuropathic urinary incontinence. In: Abrams P, Khoury S, Wein A, editors. *Incontinence*. 2nd ed. Plymouth: Health Publication Ltd; 2002, p. 697–754.
- Weld KJ, Graney MJ, Dmochowski RR. Difference in bladder compliance with time and associations of bladder management with compliance in spinal cord injured patients. *J Urol* 2000;163:1228–33.
- Wyndaele JJ. Intermittent catheterization: which is the optimal technique? *Spinal Cord* 2002;40:432–7.
- Weld KJ, Graney MJ, Dmochowski RR. Clinical significance of detrusor sphincter dyssynergia type in patients with post-traumatic spinal cord injury. *Urology* 2000;56:565–8.
- McGuire EJ, Cespedes RD, O'Connell HE. Leak-point pressures. *Urol Clin North Am* 1996;23:253–62.
- Stöhrer M, Goepel M, Kondo A, Kramer G, Madersbacher H, Millard R, et al. The standardization of terminology in neurogenic lower urinary tract dysfunction with suggestions for diagnostic procedures. *Neurourol Urodyn* 1999;18:139–58.
- Nijman RJ. Neurogenic and non-neurogenic bladder dysfunction. *Curr Opin Urol* 2001;11:577–83.
- Kachourbos MJ, Creasey GH. Health promotion in motion: improving quality of life for persons with neurogenic bladder and bowel using assistive technology. *SCI Nurs* 2000;17:125–9.
- Lawrenson R, Wyndaele JJ, Vlachonikolis I, Farmer C, Glickman S. Renal failure in patients with neurogenic lower urinary tract dysfunction. *Neuro-epidemiology* 2001;20:138–43.