



Contents lists available at ScienceDirect

## Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb



Document heading doi:10.1016/S2221-1691(14)60218-5 © 2014 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

# Lupeol: An antioxidant triterpene in *Ficus pseudopalma* Blanco (Moraceae)

Librado A Santiago<sup>1,2,3\*</sup>, Anna Beatriz R Mayor<sup>1,3</sup>

<sup>1</sup>Research Center for the Natural and Applied Sciences, University of Santo Tomas, Manila, Philippines

<sup>2</sup>Department of Biochemistry, Faculty of Pharmacy, University of Santo Tomas, Manila, Philippines

<sup>3</sup>The Graduate School, University of Santo Tomas, Manila, Philippines

## PEER REVIEW

### Peer reviewer

Roldan M. de Guia, Joint Research Division, Molecular Metabolic Control (A170)–German Cancer Research Center (DKFZ), Heidelberg, Center for Molecular Biology (ZMBH), University of Heidelberg, Heidelberg University Hospital, Im Neuenheimer Feld 280, 69120 Heidelberg, Germany.  
Tel: +49-6221-42-3597  
E-mail: r.deguia@dkfz-heidelberg.de

### Comments

This is an interesting research work in which authors have demonstrated the free-radical scavenging activity of the leaf extracts of *F. pseudopalma* Blanco (Moraceae) using biochemical tests. Lupeol was identified as one of the possible active components based on chromatographic analysis and infrared spectroscopy.

Details on Page 117

## ABSTRACT

**Objective:** To assess the antioxidant activity of *Ficus pseudopalma* Blanco (Moraceae) (*F. pseudopalma*) and characterize the active components present in it.

**Methods:** Column chromatography of crude ethanol leaf extract of *F. pseudopalma* was performed and seven fractions were obtained, labeled as F1, F2, F3, F4, F5, F6, F7. DPPH, FRAP, Griess, Fenton and superoxide radical scavenging assays were performed to assess the antioxidant ability of the fractions. Thin layer chromatography (TLC), high performance liquid chromatography and Fourier transfer infrared spectroscopy (FTIR) were performed to identify and characterize the bioactive component present in each fractions of *F. pseudopalma*.

**Results:** DPPH and FRAP assay showed that F5, F6 and F7 exhibited the good proton accepting ability and reducing power as compared to the other fractions. All fractions exhibited a good nitric oxide radical scavenging activity wherein F1, F2 and F3 showed the highest inhibition. However, all of the fractions exhibited a stimulatory activity on hydroxyl and superoxide radicals. Lupeol matched one of the spots on the thin layer chromatography chromatogram of the fractions. Linear gradient high performance liquid chromatography and spiking of lupeol with the fraction revealed the presence of 5.84 mg/L lupeol in F6. Infrared spectra of the fractions revealed the presence of C–C, OH, aromatic C=C and C=O groups.

**Conclusions:** The identified lupeol in *F. pseudopalma* may be responsible for the exhibited antioxidant property of the plant. Furthermore, knowing the antioxidant capability of the plant, *F. pseudopalma* can be developed into products which can help prevent the occurrence of oxidative stress related diseases.

### KEYWORDS

Antioxidant, Oxidative stress, Free radical scavenging, Reducing power, Phenolic acids, *Ficus pseudopalma* Blanco

## 1. Introduction

Physiologically, normal metabolic processes of the body produce significant amounts of reactive oxygen species (ROS). The damaging effects brought by these ROS are being counteracted by the cellular antioxidant defense system of the body which consist of enzymatic and non-enzymatic components[1]. However, at certain conditions,

oxidative stress is triggered due to the imbalance between the production of ROS and the antioxidant systems of the body[2,3]. Oxidative stress are usually related to high risk health conditions such as cardiovascular disease, neurodegenerative diseases, diabetes, cancer and inflammation[4–6].

Henceforth, potential antioxidative agents are being utilized and tested in order to alleviate the occurrence of

\*Corresponding author: Librado A Santiago, Research Center for the Natural and Applied Sciences, University of Santo Tomas, España Blvd. Manila, Philippines.

Tel: +632 406-1611 loc. 4053

Fax: +632 731-4031

E-mail: librado\_santiago@yahoo.com

Foundation Project: Supported by Grants-in-aid of the Philippine Council for Health Research and Development, Department of Science and Technology (Grant no. FP 120023).

Article history:

Received 12 Nov 2013

Received in revised form 23 Nov, 2nd revised form 28 Nov, 3rd revised form 5 Dec 2013

Accepted 13 Jan 2014

Available online 28 Feb 2014

such health conditions. In this regard, phytochemicals from various natural products had become the subject of most drug development researches.

Plants are the most commonly known sources of natural antioxidants which includes ascorbate, tocopherols, polyphenolic compounds and terpenoids[7,8]. The numerous health benefits that people obtained from natural products is not only related to their antioxidative components but also to some of their phytochemical contents which work hand in hand in order to elicit functions that may specifically be beneficial in the treatment of certain diseases[8].

*Ficus pseudopalma* Blanco (*F. pseudopalma*), an endemic medicinal plant in Philippines, was reported to have an antioxidant activity[9,10]. As presented in a study, *F. pseudopalma* contains terpenoids and sterols, which includes,  $\alpha$ -amyrin acetate,  $\beta$ -amyrin acetate, ursenone and lupeol acetate[11], and they were reported to have antioxidant properties[12]. Terpenoids are among the commonly useful phytochemicals isolated from plants. They have various biological functions that help in maintaining good health. A study conducted by Goto *et al.* (2010) relating to peroxisome proliferator-activated receptors revealed that terpenoids can regulate the activity of peroxisome proliferator-activated receptors so as to maintain the energy homeostasis in the body and manage obesity induced metabolic disorders including type 2 diabetes, hyperlipidemia, insulin resistance and cardiovascular disease[13].

Lupeol is one of the identified compounds in *F. pseudopalma* which has several biological activities[11]. These activities were described in some review articles wherein according to them lupeol has anti-inflammatory, antimicrobial, antiprotozoal and anti-tumor activities[14,15]. In this regard, lupeol can also be used as a nutraceutical and chemopreventive agent[15].

According to Stuart, *F. pseudopalma* is used as herbal medicine for the treatment of kidney stones and diabetes[16]. The plant also exhibits no toxicity to tested female Sprague Dawley rats at dose of 2000 mg/kg body weight[17], which is why the young shoots and leaves can be eaten as salad and mixed with other vegetables. The anti-urolithiatic property of the plant was verified with the study conducted by Acosta *et al.* (2013) which reported that the crude dichloromethane extract of *F. pseudopalma* significantly decreased the creatinine and urine oxalates levels of ethylene-glycol induced male Sprague Dawley rats[17], which supports the ethnobotanical use of *F. pseudopalma* as an agent that cure kidney stones.

Considering the health benefits one can derive from *F. pseudopalma*, its biological, biochemical and pharmacological studies were scarce. Moreover, chemical profiling of the antioxidative components present in the plant is not yet been fully elucidated. Hence, the plant remains underutilized and insignificant. Therefore this study was undertaken to back up its ethnobotanical uses while intensively evaluating the antioxidative components of *F. pseudopalma* as well as assessing its potential role as a powerful antioxidant and a scavenger of free radicals.

## 2. Materials and methods

Fresh leaves of *F. pseudopalma* Blanco were collected from Brgy. San Jose, Pili, Camarines Sur, Philippines. All reagents and solvents used were either analytical or high performance liquid chromatography (HPLC) grade and were obtained from Sigma–Aldrich Co., Singapore.

### 2.1. Plant preparation and extraction

Plant preparation and extraction was done accordingly as what were described[9]. In brief, air-dried leaves were ground to fine powder using Wiley mill and sieved in 20 mm mesh size. The powder sample was kept in a clean, dried, well-sealed amber glass container to protect it from sunlight and contamination.

About 1 kg powdered leaves were soaked with 2.5 L 95% ethanol in a percolator for 4 d while changing the extracting solvent every after 24 h. The collected ethanol extract was then concentrated using the rotary evaporator (Eyela, USA) at 40 °C until syrupy consistency was obtained. The syrupy extract was further evaporated until it became dry. The air-dried crude extract was weighed, obtaining 4.572% yield and then kept in amber-colored container under 0 °C until use.

### 2.2. Column chromatography

Different factions from the crude ethanolic leaf extract was obtained through sequential elution column chromatography. Silica gel G–50, 100–200 mesh (Hi-media, India) was used as the stationary phase. Five grams of silica gel was added with 10 mL of hexane. The mixture was then transferred to a column and was allowed to stabilize for 20 min. The crude ethanolic leaf extract of *F. pseudoplama* (1.5 g) was dissolved in 2 mL 95% ethyl alcohol to form a slurry. After the stabilization of the column, the crude ethanol leaf extract slurry was loaded to the column and was continuously eluted with the mobile phase (hexane, ethyl acetate, acetone and methanol). Approximately 10 mL of eluents were collected in pre-weighed vials. A total of seven fractions were collected and were labeled accordingly: hexane (F1), hexane : ethyl acetate (F2), ethyl acetate (F3), ethyl acetate : acetone (F4), acetone (F5), acetone : methanol (F6) and methanol (F7) fractions. These fractions were concentrated by removing the solvent through rotary evaporator (Eyela, USA). The fractionated extracts that were obtained were kept in a refrigerator at –20 °C until use.

### 2.3. Micro-scale antioxidant tests

Procedures for the antioxidant tests were performed according to the discussed procedures[9].

#### 2.3.1. DPPH radical scavenging

The hydrogen donating ability of each fractions was evaluated using the stable DPPH radical. Each fractions

were prepared in ethanol at different concentrations. Ten microliter of each fractions at various concentrations were loaded in a 96–well microplate (triplicate). Then 140  $\mu\text{L}$  of  $6.58 \times 10^{-5}$  mol/L DPPH solution was added to each well. The microplate was incubated for 30 min at 25 °C in the dark. The absorbance of the fractions were measured at 517 nm. The free radical scavenging activity was expressed as the percentage inhibition of free radical by each fraction.

### 2.3.2. FRAP assay

The reducing power can be measured through the intensity of the Prussian blue color that results from the direct reduction of potassium ferricyanide [ $\text{K}_3\text{Fe}_2(\text{CN})_6$ ] to potassium ferrocyanide [ $\text{K}_4\text{Fe}(\text{CN})_6$ ][18].

Different concentrations of each fraction (40  $\mu\text{L}$ ) were mixed with 1.0 mol/L hydrochloric acid (100  $\mu\text{L}$ ), 1% sodium dodecylsulphate (20  $\mu\text{L}$ ) and 1% potassium ferricyanide (30  $\mu\text{L}$ ) in an Eppendorf tube. The mixtures were incubated at 50 °C for 20 min. Then, 30  $\mu\text{L}$  of 0.1% ferric chloride was added to each tubes. Aliquots of the mixtures were transferred to a 96–well microplate and absorbance was read at 750 nm.

### 2.3.3. Nitric oxide radical scavenging

The ability of the fractions to scavenge nitric oxide radical was assessed through the Griess reaction[19]. One hundred microliters of each fractions of the ethanolic leaf extract of *F. pseudopalma*, at different concentrations, were mixed with 10 mmol/L sodium nitroprusside (400  $\mu\text{L}$ ) and phosphate buffered saline, pH 7.4 (100  $\mu\text{L}$ ). The mixture was incubated for 150 min at 25 °C. After incubation, 100  $\mu\text{L}$  of each mixture was transferred to a new tube and was added with 0.33% sulfanilamide (200  $\mu\text{L}$ ). The resulting mixture is incubated for 5 min at 25 °C. Then 0.1% naphthylethylenediamine (200  $\mu\text{L}$ ) was added to each tubes. The tubes were incubated for another 30 min 25°C. An aliquot of 250  $\mu\text{L}$  of the resulting mixture was transferred to a 96–well microplate in triplicate and was read at 540 nm.

### 2.3.4. Hydroxyl radical scavenging

The method used for the scavenging of the highly reactive hydroxyl radical was determined by Valko M, *et al.* with slight modification[20]. Before starting, Fenton reagent—a mixture of 2.5 mL 0.1 mmol/L  $\text{FeCl}_3$ , 0.625 mL 1.5%  $\text{H}_2\text{O}_2$  and 1.88 mL 0.0029% EDTA—was prepared. Briefly, 50  $\mu\text{L}$  of varied concentration of sample was loaded in each designated plates followed by the addition of 200  $\mu\text{L}$  Fenton reagent. The absorbance of each wells were read a 500 nm as an optimized wavelength of the blank away from the original 288 nm.

### 2.3.5. Superoxide radical scavenging

Five microliter of each sample was loaded in designated wells. After, 50  $\mu\text{L}$  73  $\mu\text{mol/L}$  NADH, 50  $\mu\text{L}$  156  $\mu\text{mol/L}$  nitrobluetetrazolium and 50  $\mu\text{L}$  60  $\mu\text{mol/L}$  phenazine methosulfate was introduced to the sample loaded wells. To complete the reaction, the mixture was incubated for 5 min at 25 °C and was immediately read at 560 nm[18].

## 2.4. Structural evaluation of the active antioxidant components

### 2.4.1. Thin layer chromatography (TLC)

Thin layer chromatography was performed to identify the components present in each fractions of *F. pseudopalma*. In a TLC plate pre-coated with silica gel (6 cm $\times$ 8 cm), each fractions were applied in small spots. The TLC chamber was developed using ethyl acetate: toluene: methanol (6:3:1) and was used as the solvent system.

### 2.4.2. Infrared spectroscopy

Attenuated total reflectance–Fourier transform infrared spectroscopy was performed using Shimadzu IR–Prestige 21 in order to analyze the functional group present on the seven fractions obtained from the crude ethanolic leaf extract of *F. pseudopalma*.

### 2.4.3. HPLC analysis

#### 2.4.3.1. Standard preparation

Standard quercetin (24.0 mg), rutin (40.0 mg), gallic acid (22.0 mg) and lupeol (2.7 mg) were accurately weighed and dissolved in vacuum–filtered methanol (HPLC grade) to obtain a stock solution. These solutions were further diluted to 6.67 mg/mL, 2.4 mg/mL, 4.4 mg/mL and 2.7 mg/mL, respectively before the chromatographic analysis.

#### 2.4.3.2. Sample preparation

Seven fractions from the crude ethanolic leaf extract of *F. pseudopalma*, labeled as F1, F2, F3, F4, F5, F6, and F7, were prepared and dissolved in vacuum–filtered methanol (HPLC grade). Each solutions were filtered using 0.22  $\mu\text{m}$  membrane filter before injecting to the sample port.

#### 2.4.3.3. Apparatus and chromatographic condition

The components present in each fractions were separated using reverse–phase C18 column. The mobile phase was comprised of 70% methanol (A), phosphate buffer, pH 7.2 (B) and ultrapure water (C). Linear elution of the solvent system was programmed at 4% B and 96% C for the first 2 min; 50% A, 1% B and 49% C for 2 to 6 min; 80% A, 2% B and 18% C for 6 to 26 min; 50% A, 1% B and 49% C for 26–30 min; and the last 5 min for 4% B and 96% C. The mobile phase was delivered at 1 mL/min with detection wavelength at 280 nm.

Sixty microliters of each fractions and standard solutions were introduced to the column. The chromatographic peaks obtained of each fractions were compared to that of the peaks obtained by the standard solutions. Spiking was performed in order to confirm the presence of each standard used on the fractions of *F. pseudopalma*.

## 2.5. Statistical analysis

Mean $\pm$ SEM were used to summarize the data gathered from the experiment. Single–factor analysis of variance (ANOVA) was used to determine if there is a significant difference in the mean percentage inhibition on antioxidant assays (DPPH,

nitric oxide, hydroxide and superoxide), mean percentage reducing power using FRAP assay. Tukey's HSD was used for post-hoc analyses.

All the statistical tests were performed using Graphpad's Prism 5.0, and SAS 9.0.  $P$ -values less than 0.05 indicate significant difference.

### 3. Results

#### 3.1. Free radical scavenging activity of the fractions of *F. pseudopalma*

##### 3.1.1. DPPH radical scavenging activity

DPPH assay of the fractions from the ethanolic leaf extract of *F. pseudopalma* showed that F1 ( $P=0.006$ ), F2 ( $P=0.008$ ) and F3 ( $P<0.001$ ) has a significantly least mean percentage of inhibition of DPPH radical compared to the other fractions (Figure 1). The scavenging activity of the F4 is not dependent on the concentration of the fraction ( $P=0.324$ ) as indicated by its mean percentage inhibition. Moreover, the increase in DPPH scavenging activity is observed in the polar fractions of *F. pseudopalma*, F5 ( $P=0.003$ ), F6 ( $P<0.001$ ), F7 ( $P<0.001$ ) as shown by their mean percentage inhibition. In connection to this, the highest concentration of the indicated fractions has demonstrated the best DPPH scavenging activity ( $P<0.05$ ). The activity of each fractions was compared to the standard ascorbic acid, which exhibits a good scavenging activity ( $P=0.004$ ).

##### 3.1.2. Ferric reducing power

The ferric reducing power of the fractions of *F. pseudopalma*

were evaluated using FRAP assay which utilizes the reduction of  $Fe^{2+}$  ions to  $Fe^{3+}$  to form a Prussian blue complex.

As illustrated in Figure 2, the ferric reducing power of almost all the fractions increases with concentrations ( $P>0.05$ ). Significant differences in the mean percentage reducing power were observed between the concentrations of each fractions. As shown, F1 ( $P=0.003$ ) and F2 ( $P<0.001$ ) has exhibited the least ferric reducing power. Whereas, F3 ( $P<0.001$ ) and F4 ( $P<0.001$ ) has almost the same activity.

Polar fractions of *F. pseudopalma*, F5 ( $P<0.001$ ), F6 ( $P<0.001$ ) and F7 ( $P<0.001$ ), had demonstrate the highest mean percentage of reducing power. All fractions were compared to ascorbic acid ( $P=0.029$ ), which exhibited a good reducing power.

##### 3.1.3. Nitric oxide scavenging activity

Nitric oxide scavenging ability of the fractions of *F. pseudopalma* was determined using Griess' reaction. As shown in Figure 3, significant difference ( $P<0.05$ ) was observed between the activity of each fractions. The activity of F1 ( $P=0.043$ ), F2 ( $P<0.001$ ) and F3 ( $P<0.001$ ) showed the highest scavenging ability compared to the other fractions. This is followed by F5 ( $P<0.001$ ), F6 ( $P=0.216$ ) and F7 ( $P<0.001$ ). Among the fractions, F4 ( $P<0.001$ ) has the lowest scavenging activity. The scavenging ability of all the fractions was compared to ascorbic acid ( $P<0.001$ ) which elicit a concentration-dependent inhibition of nitric oxide radical production.

##### 3.1.4. Hydroxyl radical scavenging activity

Hydroxyl radicals ( $\cdot OH$ ) is among the free radicals that are physiologically produced by the body through its metabolic processes[21]. In the assay, hydroxyl radical was produced by




Figure 1. Proton-donating ability of the fractions obtained from the ethanolic leaf extract of *F. pseudopalma*. Analysis was performed in triplicates ( $n=3$ ,  $P>0.05$ ).






Figure 2. Ferric reducing activity of the fractions obtained from the ethanolic leaf extract of *F. pseudopalma*. Analysis was performed in triplicates ( $n=3$ ,  $P>0.05$ ).




Figure 3. Concentration-dependent inhibition of NO· by the fractions obtained from the ethanolic leaf extract of *F. pseudopalma*. Analysis was performed in triplicates ( $n=3$ ,  $P>0.05$ ).

the reaction of H<sub>2</sub>O<sub>2</sub> with EDTA-bound Fe<sup>2+</sup>.

Almost all the fractions of *F. pseudopalma* were able to stimulate the production of hydroxyl radicals (Figure 4). Significant differences were observed on scavenging activity of each fractions ( $P<0.05$ ). Hexane ( $P<0.001$ ), hexane: ethyl acetate ( $P<0.001$ ) and ethyl acetate ( $P<0.001$ ) fractions showed the highest stimulation of hydroxyl radicals. Semi-polar and polar fractions, F4 ( $P<0.001$ ), F5 ( $P<0.001$ ), F6 ( $P<0.001$ ) and F7 ( $P=0.496$ ), has low stimulatory activity towards hydroxyl radicals as compared to the first three fractions mentioned as indicated by their mean percentage inhibition. In addition to that, difference in the concentration of F7 does not affect its activity towards scavenging of hydroxyl radicals.

### 3.1.5. Superoxide radical scavenging activity

Another common oxygen radical in the body is superoxide radicals ( $\cdot\text{O}_2^-$ ) [21]. It is produced with the  $\cdot\text{OH}$  and continue to a cascade of free radical formation in the body. In this experiment,  $\cdot\text{O}_2^-$  are produced from phenazine methosulfate-NADH and direct reduction of nitrobluetetrazolium.

As demonstrated in Figure 5, there is a significant difference in the superoxide radical scavenging ability of each fractions ( $P>0.05$ ). The nonpolar fractions such as hexane ( $F_{5,12}=3.648$ ,  $P=0.031$ ), hexane: ethyl acetate ( $P=0.069$ ) and ethyl acetate ( $P=0.001$ ) exhibited the highest stimulatory effect towards superoxide radicals. They were followed by the semi-polar and polar fractions: F4 ( $P<0.001$ ), F5 ( $P=0.040$ ), F6 ( $P=0.019$ ) and F7 ( $P<0.001$ ), which have lower stimulatory effect as compared to the first three fractions that were

mentioned.

On the other hand, ascorbic acid ( $P<0.001$ ) was able to inhibit the production of superoxide radical as opposed to the exhibited activities of the fractions of *F. pseudopalma*.

## 3.2. Structural evaluation of the active antioxidant components

### 3.2.1. Thin layer chromatography (TLC)

The TLC chromatogram of the fractions revealed that fractions 1, 2 and 3 have the most number of components and one of which corresponds to lupeol ( $R_f=0.831$ ). Other fractions showed lesser number of components. In order to further verify the results and confirm the compounds present in the fractions of *F. pseudopalma*, infrared spectroscopy and HPLC analysis was performed.

### 3.2.2. Infrared spectroscopy

Infrared spectra of the fractions obtained from *F. pseudopalma* revealed the presence of alkane and alcohol functional group (Figures 6 and 7). The sharp peak for the IR spectra of each fraction with values ranging from 2800–3000  $\text{cm}^{-1}$  corresponds to a C–H stretch functional group of alkane. Terminal alkene group ( $=\text{C}-\text{H}$ ) was observed at hexane: ethyl acetate (50: 50) and ethyl acetate fractions. Of the seven spectra, methanol fraction showed a more prominent broad peak that corresponds to an O–H stretch (3300–3400  $\text{cm}^{-1}$ ) of phenol or alcohol functional group. Aromatic C=C was also shown in the spectra of all the fractions. Other functional groups that were assumed to be present in each fraction are




Figure 4. Concentration-dependent stimulation of  $\cdot\text{OH}$  by the fractions obtained from the ethanolic leaf extract of *F. pseudopalma*. Analysis was performed in triplicates ( $n=3, P>0.05$ ).




Figure 5. Concentration-dependent stimulation of  $\cdot\text{O}_2$  by the fractions obtained from the ethanolic leaf extract of *F. pseudopalma*. Analysis was performed in triplicates ( $n=3, P>0.05$ ).





Figure 6. Infrared spectra of the 7 fractions from the crude ethanolic leaf extract of *F. pseudopalma*.

A: Hexane fraction; B: Hexane: ethyl acetate fraction; C: Ethyl acetate fraction; D: Ethyl acetate: acetone fraction; E: Acetone fraction; F: Acetone: methanol fraction; G: Methanol fraction .



**Figure 7.** HPLC chromatogram of the crude ethanolic leaf extract of *F. pseudopalma*. A: hexane fraction; B: hexane:ethyl acetate fraction; C: ethyl acetate fraction; D: ethyl acetate: acetone fraction; E: acetone fraction; F: acetone:methanol fraction; G: methanol fraction . The numbers correspond to the major peaks observed with their corresponding retention time.

**Table 1**

Summary of the infrared spectra of each fractions of *F. pseudopalma*.

Chemicals	Infrared spectra						
	Hexane	Hexane: ethyl acetate	Ethyl acetate	Ethyl acetate:acetone	Acetone	Acetone: methanol	Methanol
Alkane C–H	2853.81	2853.81	2854.77	2852.84	2854.77	2853.81	2930.00
	2924.21	2924.21	2926.14	2925.17	2925.17	2924.21	
	2955.07	2954.11					
Alkene =C–H	–	3099.74	3038.02	–	–	–	–
C=C	1647.28	1651.14	1651.14	1641.49	1653.07	1651.14	1640.53
Alcohol C–O	1668.50	1666.57	1668.50	1649.21		1665.60	
	1046.43	1047.39	1043.53	1641.49	1032.93	1050.29	1030.03
O–H	3365.93	3371.71	3359.18	3382.32	3334.10	3302.27	3295.52
	3369.79	3385.22			3344.71	3312.88	
	3383.29	3393.90				3564.60	
	3499.02	3544.35			3354.35	3585.82	
	3544.35	3565.57					
	3564.60						
Ether C–O	1168.91	1068.61	–	1073.43	1146.73	1050.29	–
						1144.80	
Aldehyde Aromatic/Conjugated C=O Saturated C=O	–	1708.04	–	–	–	–	–
	1732.15	1733.12	–	–	–	1734.08	–
C–H	2727.46	2728.43					
Ketone Aromatic/Conjugated C=O Cyclic C=O Saturated C=O	–	–	–	–	–	1683.93	–
	–	–	–	–	–	1716.72	–
	1714.79	–	–	–	–	–	–
Arene C–H	821.71	819.78	819.78	3009.08	817.85	817.85	771.56
		836.18					817.85
Aromatic Ring	1454.39	1454.39	1453.43	1458.25	1456.32	1455.35	1507.43
	1540.23	1508.40	1514.19	1507.43	1653.07	1622.20	1520.94
	1605.81	1541.19	1519.01	1521.90	1702.25	1636.67	1543.12
	1647.28	1604.84	1544.08	1539.26		1651.14	1558.55
	1668.50	1651.14	1555.66	1556.62		1665.60	1622.20
	1714.79	1666.57	1610.63	1641.49		1683.93	1637.64
	1732.15	1708.04	1624.13	1649.21		1700.32	1640.53
		17333.12	1651.14	1684.89		1716.72	
		1668.50	1694.54		1734.08		
			1699.36				


summarized in Table 1.

Spectral analysis of the fractions help in the determination of the active compounds present in each fraction that contribute to their antioxidant activity.

### 3.2.3. HPLC analysis

High pressure liquid chromatography was performed in order to further confirm the active antioxidative components of each fractions from the crude ethanolic leaf extract of *F. pseudopalma*.

Chromatograms of all the fractions were compared to different standards which includes quercetin, rutin, gallic acid and lupeol. However, only lupeol was observed on the chromatogram of F6. This finding was further confirmed through spiking wherein the standard lupeol was mixed with the F6 solution and was injected to the HPLC unit. The chromatogram of the spiked F6 (Figure 8) showed that one of the peaks had increased in height and area, which correspond mainly to the presence of lupeol. Quantitative analysis revealed that 5.84 mg/L lupeol was contained in F6 using a standard curve.



**Figure 8.** HPLC chromatogram of F6.

(A) is compared to the chromatogram of F6 spiked with lupeol (B).

## 4. Discussion

The DPPH and FRAP assay are general tests used to evaluate the antioxidant capacity of substance. The DPPH radical is commonly used for fast evaluation of the antioxidant property of a given compound[22]. The dark purple solution of DPPH is turned to yellow when the DPPH radical subsequently receive an electron or a hydrogen radical from an antioxidant[23]. This change in color can be measured spectrophotometrically at 517 nm. Moreover, the antioxidant activity can be also linked

to the reducing power of a bioactive compound[24]. In this assay, the ability of each fractions from the ethanolic leaf extract of *F. pseudopalma* to directly reduce of  $Fe^{3+}$  to  $Fe^{2+}$  was monitored by measuring the increase in absorbance through the formation of Pearl's Prussian blue complex at 750 nm[18].

As indicated in the results, most of the polar fractions of *F. pseudopalma* were scavengers of DPPH radicals and has a better reducing power compared to that of the nonpolar and semi-polar fractions. The electron donating property of each of the fractions can potentially neutralize the action of free radicals. In relation to this, the reducing power exhibited by each of the fractions can serve as an indicator of their antioxidant potential.

Specific antioxidant tests were also performed in order to evaluate the influence of the fractions from *F. pseudopalma* on the activity of free radicals that are physiologically produced by the body. Nitric oxide free radical ( $NO\cdot$ ) is a biologically active specie that can readily react with other free radicals inside the body which in turn mediate specific biological effects depending on the local chemical environment[25].  $NO\cdot$  is usually linked in inflammation[21] and carcinogenesis[26]. Overproduction of  $NO\cdot$  by nitric oxide synthase is commonly observed in patients with rheumatoid arthritis and osteoarthritis[21].

Hydroxyl radicals and superoxide radicals are among the free radicals that are physiologically produced by the body through its metabolic processes[27]. The immune system utilize these reactive species in order to fight the invading pathogens so as to maintain the homeostasis inside the body.

Generally, free radicals do not only cause harmful effects in the body. In fact, free radicals have been also linked to the killing of cancer cells. Cancer cells are normally anaerobic cells which survive in a less oxygenated environment. Exposure of cancer cells to oxidative stress would lead to growth inhibition or even death.

The antioxidant activity of several plant extracts are usually related to the phytochemicals that were isolated from them. Some of the known antioxidant agents are the phenolic acids, flavonoids and terpenoids. In order to determine the active antioxidative component of *F. pseudopalma*, the fractions of the ethanolic leaf extract were subjected to TLC, high pressure liquid chromatography and infrared spectroscopy to verify the structure of the compound present.

The TLC chromatogram showed that lupeol was present in F1, F2, F3 and F4. However, HPLC analysis revealed that 5.84 mg/L lupeol was present in the acetone: methanol (50: 50) (retention time=9.00 min). Lupeol is a pentacyclic triterpenes that exhibit antioxidative activity through direct scavenging of free radicals and protects membrane permeability[14]. This property can be related to its hepatoprotective and anticancer properties. In one particular study, lupeol compound isolated from *Verpis punctata* exhibited cytotoxicity ( $IC_{50}$ =26.4 $\mu$ g/mL) on A2780 human ovarian cancer cell line[28].

Additionally, lupeol was also reported to have cardioprotective activity due to its ability to ameliorate lipidemic-oxidative abnormalities in the early stage of hypocholesterolemic atherosclerosis in rats[29]. Other findings



showed that lupeol provided 34.4% protection against *in vitro* low density lipoprotein oxidation and hypotensive activity that makes it a preventive agent against cardiac disorder<sup>[30,31]</sup>.

Another reported biological activity of lupeol is hepatoprotections. In one related study, lupeol showed some effectiveness in lessening the action of aflatoxin B1, which is a fungal derived toxin<sup>[32]</sup> that is considered to be the most potent of the mycotoxins<sup>[33]</sup> which can cause acute hepatotoxicity and liver carcinoma in exposed animal. In relation to that, it was observed that lupeol can substantially normalized degenerative alterations in the hepatocytes with granular cytoplasm<sup>[34]</sup>. Additionally, lupeol treatment induced growth inhibition and apoptosis in hepatocellular carcinoma SMMC7721 cells by down-regulation of the death receptor 3 expression<sup>[35]</sup>.

Detection of this compound in *F. pseudopalma* supports the antioxidant activity exhibited by the plant. In addition to that, lupeol may also be the one that contribute to the anticancer property of the plant. A study showed that the ethanolic extract of *F. pseudopalma* demonstrated a cytotoxic effect against HepG2 cells that was comparable to the activity of curcumin<sup>[10]</sup>.

Based on the aforementioned information, lupeol is indeed one of the most valuable phytochemical constituents of the plant, where several benefits can be derived that include cardioprotection, hepatoprotection and can help in the prevention of cancer progression.

Being one of the local medicinal plant in Philippines which has only meager studies, it is noteworthy to know other uses of *F. pseudopalma* Blanco (Moraceae). Identification of lupeol in *F. pseudopalma* supports the exhibited antioxidant activity of the plant which can be used to avoid the prevalence of several diseases related to oxidative stress.

Since little is known about the specific chemical constituents of the plant, further analysis such as liquid chromatography coupled with mass spectroscopy (LC-MS) can be performed in order to elucidate and further confirm the existence of other phytochemicals that may contribute to its pharmaceutical function.

### Conflict of interest statement

We declare that we have no conflict of interest.

### Acknowledgements

The authors would like to extend their gratitude to the Grants-in-Aid of the Philippine Council for Health Research and Development, Department of Science and Technology for providing the financial support for the study with Grant no. FP 120023. The authors are also in gratitude for the Research Center for the Natural and Applied Sciences (RCNAS) of the University of Santo Tomas for the assistance in conducting the analytical experiments included in the study.

### Comments

#### Background

The damaging effects of imbalances in the oxidative state of the cell could trigger development and exacerbation of cardiovascular diseases, hepatocellular maladies and cancer. The biochemical and instrumental identification of lupeol, a biologically active triterpene, in the Philippine-endemic *F. pseudopalma* Blanco (Moraceae) has provided insights into the potential use of phytochemicals as antioxidants.

#### Research frontiers

The current research work presents biochemical characterization of antioxidant properties and structural evaluation of phytochemicals, lupeol in particular, present in leaf extracts of *F. pseudopalma* Blanco (Moraceae).

#### Related reports

The present work complements the NMR instrumentation done by Ragasa CY *et al.* in 2009 on the leaf extracts of *F. pseudopalma* Blanco (Moraceae). Furthermore, the paper supports previous reports on the potential biological and pharmacological property of lupeol and the plant as a whole.

#### Innovations and breakthroughs

*F. pseudopalma* Blanco (Moraceae) commonly known as the Philippine fig or niog-niogan in some parts of the country, is being used as an ornamental or medicinal plant. In the present study, the authors showed the antioxidant properties of different organic solvent fractions of the leaf of the plant eventually identifying lupeol as one of the active components.

#### Applications

The demonstrated free-radical scavenging activity of the different leaf extracts of *F. pseudopalma* Blanco (Moraceae) as well as the identification of lupeol as an active component, support the prospective application as an anti-cancer and cardio/hepatoprotective agent.

#### Peer review

This is an interesting research work in which authors have demonstrated the free-radical scavenging activity of the leaf extracts of *F. pseudopalma* Blanco (Moraceae) using biochemical tests. Lupeol was identified as one of the possible active components based on chromatographic analysis and infrared spectroscopy. These significant results have shown the importance of phytochemical screening of endemic flora that potentially harbors pharmacologically promising drug candidates.

### References

- [1] Hopps E, Noto D, Caimi G, Averna MR. A novel component of

- metabolic syndrome: the oxidative stress. *Nutr Metab Cardiovasc Dis* 2010; **20**(1): 72–77.
- [2] Wiernsperger NF. Oxidative stress as a therapeutic target in diabetes: revisiting the controversy. *Diabetes Metab* 2008; **29**(6): 579–585.
- [3] Pepe H, Balci SS, Revan S, Akalin PP, Kurtoglu F. Comparison of oxidative stress and antioxidant capacity before and after running exercises in both sexes. *Gen Med* 2009; **6**(4): 587–595.
- [4] Fearon IM, Faux SP. Oxidative stress and cardiovascular disease: Novel tools give free radical insight. *J Mol Cell Cardiol* 2009; **47**(3): 372–381.
- [5] Pashkow FJ. Oxidative stress and inflammation in heart disease: do antioxidants have a role in treatment and/or prevention? *Int J Inflamm* 2011; doi: 10.4061/2011/514623.
- [6] Fang YZ, Yang S, Wu G. Free radicals, antioxidants and nutrition. *Nutrition* 2002; **18**(10): 872–879.
- [7] Grassmann J. Terpenoids as plant antioxidants. *Vitam Horm* 2005; **72**: 505–535.
- [8] Dimitrios B. Sources of natural phenolics and antioxidants. *Trends Food Sci Technol* 2006; **17**: 506–512.
- [9] Santiago LA, Valerio VLM. Assessment of antioxidant activities of crude ethanolic leaf extract of *Ficus pseudopalma* Blanco (Moraceae). *Int J Pharm Front Res* 2013; **3**(1): 15–25.
- [10] Bueno PRP, Buno CBM, Santos DLM, Santiago LA. Antioxidant activity of *Ficus pseudopalma* Blanco and its cytotoxic effect on hepatocellular carcinoma and peripheral blood mononuclear cells. *Curr Res Bio Pharm Sci* 2013; **2**(2): 14–21.
- [11] Ragasa CY, Tsai P, Shen CC. Terpenoids and sterols from the endemic and endangered Philippine trees, *Ficus pseudopalma* and *Ficus ulmifolia*. *Philipp J Sci* 2009; **138**(2): 205–209.
- [12] Topcu G, Ertas A, Kolak U, Ozturk M, Ulubelen A. Antioxidant activity tests on novel triterpenoids from *Salvia macrochlamys*. *ARKIVOC* 2007; **7**: 195–208.
- [13] Goto T, Takahashi N, Hirai S, Kawada T. Various terpenoids derived from herbal and dietary plants function as PPAR modulators and regulate carbohydrate and lipid metabolism. *PPAR Res* 2010; doi: 10.1155/2010/483958.
- [14] Wal P, Wal A, Sharma G, Rai AK. Biological activities of lupeol. *Syst Rev Pharm* 2011; **2**(2): 96–103.
- [15] Gallo MBC, Sarachine MJ. Biological activities of lupeol. *Int J Biomed Pharm Sci* 2009; **3**(1): 46–66.
- [16] Stuart G. Niyog–niyogan, *Quisqualis indica* Linn. Yesterday, today and tomorrow. Philippines: StuartXchange; 2011. [Online] Available from: <http://www.stuartxchange.org/niyog–niyogan.html>. [Accessed on September 23, 2012]
- [17] Acosta CJH, Macabeo APH., Santiago LA. Assessment of the antiuro lithiatic and antioxidant properties of *Ficus pseudopalma* Blanco leaves (Moraceae). *Curr Res Bio Pharm Sci* 2013; **2**(2): 22–30.
- [18] Ahmed AS, Elgorashi EE, Moodley N, McGaw LJ, Naido V, Eloff JN. The antimicrobial, antioxidative, anti–inflammatory activity and cytotoxicity of different fractions of four South African *Bauhinia* species used traditionally to treat diarrhea. *J Ethnopharmacol* 2012; **143**(3): 826–839.
- [19] Hazra B, Biswas S, Mandal N. Antioxidant and free radical scavenging activity of *Spondias pinnata*. *BMC Complement Altern Med* 2008; **8**: 63.
- [20] Valko M, Morris H, Cronin MT. Metals, toxicity and oxidative stress. *Curr Med Chem* 2005; **12**: 1161–1208.
- [21] Murrell GA, Dolan MM, Jang D, Szabo C, Warren RF, Hannafin JA. Nitric oxide: an important articular free radical. *J Bone Joint Surg Am* 1996; **78**: 265–274.
- [22] Bozin B, Mimica–Dukic N, Samojlik I, Goran A, Igic R. Phenolics as antioxidants in garlic (*Allium sativum* L., Alliaceae). *Food Chem* 2008; **111**: 925–929.
- [23] Shukla S, Mehta A, John J, Singh S, Mehta P, Vyas SP. Antioxidant activity and total phenolic content of ethanolic extract of *Caesalpinia bonducella* seeds. *Food Chem Toxicol* 2009; **47**: 1848–1851.
- [24] Siddhuraju P, Mohan PS, Becker K. Studies on the antioxidant activity of Indian Laburnum (*Cassia fistula* L.): a preliminary assessment of crude extracts from stem bark, leaves, flowers and fruit pulp. *Food Chem* 2002; **79**: 61–67.
- [25] Mayer O Jr, Simon J, Holubec L, Pikner R, Trefil L. Folate co–administration improves the effectiveness of fenofibrate to decrease the lipoprotein oxidation and endothelial dysfunction surrogates. *Physiol Res* 2006; **55**: 475–481.
- [26] Hofseth LJ, Saito S, Hussain SP, Espey MG, Miranda KM, Araki Y, et al. Nitric oxide–induced cellular stress and p53 activation in chronic inflammation. *Proc Natl Acad Sci U S A* 2003; **100**(1): 143–148.
- [27] Wong CC, Li HB, Cheng KW, Chen F. A systematic survey of antioxidant activity of 30 Chinese medicinal plants using the ferric reducing antioxidant power assay. *Food Chem* 2006; **97**: 705–711.
- [28] Chaturvedula VS, Schilling JK, Miller JS, Andriantsiferana R, Rasamison VE, Kingston DG. New cytotoxic terpenoids from the wood of *Vepris punctata* from the Madagascar rainforest. *J Nat Prod* 2004; **67**(5): 895–898.
- [29] Sudhakar V, Ashokkumar S, Varalakshmi P. Effect of lupeol and lupeol linolate on lipemic–hepato cellular aberrations in rats fed a high cholesterol diet. *Mol Nutr Food Res* 2006; **50**(12): 1212–1219.
- [30] Andrikopoulos NK, Kaliora AC, Assimopolou AN, Papapeorgiou VP. Biological activity of some naturally occurring resins, gums and pigments against *in vitro* LDL oxidation. *Phytother Res* 2003; **7**: 501–507.
- [31] Saleem M, Adhami VM, Siddiqui IA, Mukhtar H. Tea beverage in chemoprevention of prostate cancer: a mini–review. *Nutr Cancer* 2003; **47**: 13–23.
- [32] Preetha SP, Kannappan M, Selvakumar E, Nagaraj M, Varalakshmi P. Lupeol ameliorates aflatoxin B1–induced peroxidative hepatic damage in rats. *Comp Biochem Physiol C Toxicol Pharmacol* 2006; **143**: 333–339.
- [33] Luyendyk JP, Copple B, Barton CC, Ganey PE, Roth RA. Augmentation of aflatoxin B1 hepatotoxicity by endotoxin: involvement of endothelium and coagulation system. *Toxicol Sci* 2003; **72**(1): 171–181.
- [34] Akcam M, Artan R, Yilmaz A, Ozdem S, Gelen T, Naziroglu M. Caffeic acid phenethyl ester modulates aflatoxin B1–induced hepatotoxicity in rats. *Cell Biochem Funct* 2013; doi: 10.1002/cbf.2957.
- [35] Zhang L, Zhang Y, Zhang L, Yang X, Lv Z. Lupeol, a dietary triterpene, inhibited growth and induced apoptosis through down–regulation of DR3 in SMMC7721 cells. *Cancer Invest* 2009; **27**: 163–170.