

Elsevier Editorial System(tm) for Journal of Cleaner Production
Manuscript Draft

Manuscript Number: JCLEPRO-D-14-01991R3

Title: Life Cycle Assessment of multiyear peach production

Article Type: Original Research Paper

Corresponding Author: Ms. Elisabet Vinyes,

Corresponding Author's Institution: Sostenipra.Research Group. Universitat Autònoma de Barcelona (UAB).

First Author: Elisabet Vinyes

Order of Authors: Elisabet Vinyes; Carles M Gasol; Luis Asin; Simó Alegre; Pere Muñoz

Abstract: Considering that peach is a significant fruit in the Mediterranean countries, and most publications on environmental impacts of fruit productions are based on one single productive year, this study attempts to perform an environmental analysis of peach production using Life Cycle Assessment, in order to provide new information on peaches, and also introduce a multiyear perspective analysis to identify the variability of the environmental impacts related to annual orchard yield and weather variations. The system studied is a peach orchard (*Prunus persica* L.) with integrated agricultural practices. The study analyses the cultivation period, as well as the impact of the initial orchard establishment tasks (soil preparation and planting). Data used have been collected directly from an orchard located in the North East of Spain, and covers 15 years of real production. The functional unit adopted was the cultivation of 1kg of peach. Four scenarios have been considered according to the different yield periods of the peach fruit tree: Growth, Low, High and Multiyear. The results of the study reveals that, depending on production scenario considered, the results per kg of peach can vary between 7% and 69% depending on the environmental indicator. If the impact of initial orchard establishment tasks (soil preparation and planting) is not included in the quantification, then 5% of total emissions may be overlooked, but sometimes a lack of data makes it difficult to include these stages. Caution should be taken when the functional unit is related to mass and only when a single year of production is studied, because unproductive years increase impacts on value per functional unit, whereas over-productive years decrease them. According to variability of the results obtained, multiyear approach should be considered in crops with an average life time of twenty years or higher. The present study can be considered a useful methodological framework for providing a deeper understanding of the key environmental impact issues related to fruit production based on peach case study, and how to avoid multiple interpretation of results associated to reporting annual environmental impact variations.

Highlights

- This work is based on multiyear perspective analysis.
- It includes the impact of the initial orchard establishment tasks.
- It analyses the variability of the environmental impacts related to annual orchard yield.
- Data used were collected directly from a real orchard.

Life Cycle Assessment of multiyear peach production

Vinyes Elisabet^{1*}, Gasol Carles M.^{1,2} Asin Luis³, Alegre Simó³, Muñoz Pere^{1,4}

¹Sostenipra Research Group. Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain

²Inèdit Innovació s.l. Research Park of UAB. UAB Campus. 08193 Bellaterra(Spain)

³Institute for Research and Technology in Food and Agriculture (IRTA), 25198 Lleida, Spain

^{1,4}Institute for Research and Technology in Food and Agriculture (IRTA), 08348 Cabriels, Barcelona, Spain

* **Corresponding author:** elisabet.vinyes@gmail.com

Key words: LCA, environmental impacts, integrated fruit production, sustainable farming.

ABSTRACT.

Considering that peach is a significant fruit in the Mediterranean countries, and most publications on environmental impacts of fruit productions are based on one single productive year, this study attempts to perform an environmental analysis of peach production using Life Cycle Assessment, in order to provide new information on peaches, and also introduce a multiyear perspective analysis to identify the variability of the environmental impacts related to annual orchard yield and weather variations. The system studied is a peach orchard (*Prunus persica L.*) with integrated agricultural practices. The study analyses the cultivation period, as well as the impact of the initial orchard establishment tasks (soil preparation and planting). Data used have been collected directly from an orchard located in the North East of Spain, and covers 15 years of real production. The functional unit adopted was the cultivation of 1kg of peach. Four scenarios have been considered according to the different yield periods of the peach fruit tree: *Growth, Low, High and Multiyear*. The results of the study reveals that, depending on production scenario considered, the results per kg of peach can vary between 7% and 69% depending on the environmental indicator. If the impact of initial orchard establishment tasks (soil preparation and planting) is not included in the quantification, then 5% of total emissions may be overlooked, but sometimes a lack of data makes it difficult to include these stages. Caution should be taken when the functional unit is related to mass and only when a single year of production is studied, because unproductive years increase impacts on value per functional unit, whereas over-productive years decrease them. According to variability of the results obtained, multiyear approach should be considered in crops with an average life time of twenty years or higher. The present study can be considered a useful methodological framework for providing a deeper understanding of the key environmental impact issues related to fruit production based on peach case study, and how to avoid multiple interpretation of results associated to reporting annual environmental impact variations.

1. INTRODUCTION

The agricultural sector has changed in Europe over the last ten years, from traditional practices to more intensive methods, in order to increase the productivity of the plantations, and as a response to the growing demand of an increasing population. As a consequence of the increase of intensive methods, food production has become an important contribution to the depletion of natural resources and climate change (Martínez-Blanco et al., 2011^a). The IPCC Climate Change Synthesis Report 2007 estimates that the direct impacts of agriculture contribute about 13.5% of global anthropogenic GHG emissions. Europe is currently encouraging farmers to practice more sustainable agriculture in order to meet all the needs of society: environmental, social and economic (European Commission 2012). Even so, to promote environmental friendly agricultural production it is essential for farmers to identify the causes of environmental impacts of their production systems.

A few years ago, the main concern in the food industry was safety, but recently it is becoming conscious of the environmental repercussions of their products, and is attempting to open new horizons towards sustainable production. Consumers are also increasingly aware of the environmental performance of the food products they buy, and this is reflected in their purchasing decisions. To develop a proper environmental management for industries and farmers, it is essential to know the main environmental indicators of their products: emissions, energy and water consumption, waste generation, efficiency, etc. It can also give an environmental added value to their product, at the same time that it provides valuable information for consumers (Environdec, 2012).

Table 1 shows a review of some publications about environmental impacts of fruit production. The literature review was carried out from papers in international journals and conference proceedings. The review covered all main aspects for conducting environmental analyses of fruit production systems, giving preference to the agricultural stage. The information was collected from two main approaches: LCA and agricultural aspects. For LCA approach the following items were considered: functional unit (FU), system boundaries, environmental impact assessment method, initial stages consideration, and cultivation period considered. For the agricultural approach, the country of the study, and fruit variety were taken into account.

Most of reviewed studies only consider one productive year, and initial stages of orchard establishment (soil preparation and planting) are not included. The application of environmental assessment methods in the fruit sector is conventionally divided into a field phase and a retail phase (considering a spatial time of one productive year). Although there are important differences in the environmental impacts in the field phase, a major part of the impacts is related to the distribution chain in the retail phase, mainly due to the cooling (Cerutti et al., 2013^a). Another important aspect to be considered is that some resources are used annually, whilst others are present during the whole lifetime of the orchard (Mila i Canals and Polo, 2003).

<<Table 1. Environmental fruit studies published in last 10 years.>>

1
2
3
4
5
6
7
8
9 According Table 1, Life Cycle Assessment (LCA) is one of the most used standardized methodologies
10 (ISO14040) for estimating the environmental burdens linked to fruit production, and it has shown to be
11 an effective mechanism to report environmental performance in the food and beverage sector in general
12 (Vazquez-Rowe et al., 2012). However there are a limited number of fruit crop LCA studies, and they still
13 do not present enough environmental information, the impacts are partially analysed, and the existing
14 studies mainly focus on one productive year, when the life span of fruit crop plantations range from 20 to
15 60 years. Quantitative environmental assessment methodologies such as LCA require significant time and
16 resource inputs during the data acquisition and life cycle inventory (LCI) phase. Approaches to
17 streamlining the LCI data collection process without degrading data quality are therefore required, and is
18 especially true for agricultural products (Bellon-Maurel et al., 2014).The main reason that may explain
19 this is due to environmental and energy aspects for the development of fruit crops were not taken in
20 account by farmers during the last decades, so there is no available data and the existing information that
21 can be found are not reliable data. In recent times, after the emergence of new private sustainability
22 standards such as: Global Gap, SAGP Guidelines, SAI initiative, etc.) and the growing competitiveness in
23 the private markets, all the actors involved in fruit production showed much interest in environmental
24 impacts that their products generate, and became aware of the need to collect much more reliable data to
25 improve the quality, the availability and the temporality of these, in order to develop environmental
26 studies with a high quality and rigorousness. Nonetheless, in the European context, the EU Framework
27 Programme for Research and Innovation has been developed: Horizon 2020, which encourages
28 companies to develop more sustainable strategies in order to reduce the environmental impact of their
29 companies and the use of resources. For the fruit sector in Mediterranean countries, the peach is an
30 important product. The main producers of peaches in Europe are: Italy, Spain, Greece and France, all
31 together produce 42% of the world production. According to FAO statistics (2011) the largest peach and
32 nectarine producer is Italy with 1,474,337 tonnes, followed by Spain 1,129,300 tonnes, Greece 810,000
33 and France 313,300. While Italy stands out as the largest producer, Spain is the major exporter, due to its
34 early season harvest, lower production costs, and varietal renewal using higher quality varieties. Greece is
35 the major EU peach processor. Spain is the second peach and nectarine producer in the European Union,
36 and ranks third in the world after China and Italy (European Commission 2012). This study will analyse
37 the region of Catalonia, located in the North East of Spain, and is the region with the second highest
38 peach production in Spain, with a production of 367,887 tonnes/year ($t\ y^{-1}$) and a cultivated area of
39 11,299 ha, which is 35% of the total Spanish peach production, and 26.4% of the total fruit cultivated area
40 (MAGRAMA 2012).

41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58 The aim of this work is to calculate the environmental impact of a multiyear peach production system in
59 the North East of Spain, using LCA methodology. Data used in the study have been directly collected
60
61
62
63
64
65

1 from an experimental orchard, and fifteen years of production have been considered. This study is a peach
2 analysis based on a multiyear system experiment, which allows working with a reliable and high-quality
3 experimental dataset and supports to highlight the importance of the multiyear approach in order to
4 reduce variability and underestimated environmental impacts. The results also may be useful to identify
5 the hot spots of peach production in its agriculture stage, and provide new inventory data and results of
6 Mediterranean peach fruit.
7
8
9

10 11 **2. MATERIALS AND METHODS.** 12

13 The environmental analysis of multiyear peach production will be performed using LCA methodology
14 according to Standard ISO 14040:2006.
15

16 **2.1. Life Cycle Assessment** 17

18 LCA is defined by ISO 14040 as the compilation and evaluation of the inputs, outputs and potential
19 environmental impacts of a product system throughout its life cycle. The end results are dependent on the
20 systems boundaries and the functional unit (FU), which is the unit to which the results of the LCA are
21 related, and used further for the communication of the LCA results.
22
23

24 The impact categories considered in this study on the performance of LCA are mid-point characterization
25 factors, and the calculation method used is Recipe Midpoint H. In this method, the environmental flows
26 were assigned by multiplying them by the corresponding characterization factor for different impact
27 categories. Calculations were performed using Simapro 7.3.3 software, together with ecoinvent database
28 3.0.
29
30

31 In order to obtain scientific, verified, and comparable information about the environmental performance
32 of the products, the results will be reported according to product category rules (PCR) procedures that
33 have been developed in accordance with ISO 14025. Given that the scope of the study is fruit, the PCR
34 model chosen will be: Fruits and nuts.
35
36
37

38 According to Recipe Midpoint (H) characterization method, the following environmental impact
39 indicators were considered in the study:
40
41

- 42 – Climate Change (CCH) expressed in kg CO₂ eq
 - 43 – Ozone Depletion (ODP) expressed in kg CFC-11 eq
 - 44 – Photochemical oxidant formation (PHO) expressed in kg NMVOC
 - 45 – Terrestrial Acidification (TAC) expressed in kg SO₂ eq
 - 46 – Freshwater Eutrophication (FEU) expressed in kg P eq
 - 47 – Marine eutrophication (MEU) expressed in kg N eq
 - 48 – Agricultural land occupation (ALO) expressed in m²a
 - 49 – Urban land occupation (ULO) expressed in m²a
 - 50 – Natural land transformation (NLT) expressed in m²
 - 51 – Water depletion (WDP) expressed in m³
 - 52 – Metal depletion (MDP) expressed in kg Feeq
 - 53 – Fossil depletion (FDP) expressed in kg oil eq
 - 54 – Ecotoxicity (Etox) expressed in CTUe
 - 55 – Demand for non-renewable energy resources (NRE) expressed in MJ-eq
- 56
57
58
59
60
61
62
63
64
65

1 The indicators chosen are midpoint indicators. The classification and characterization stages were carried
2 out excluding normalization, in order to avoid subjectivity in the analysis. The quantification of the
3 ecotoxicity (Etox) indicator has been done according the USEtox model using Simapro software. USEtox
4 is a model based on scientific consensus, providing midpoint characterization factors for human and
5 freshwater eco-toxicological impacts of chemicals in life cycle impact assessment, developed under the
6 United Nations Environment Program (UNEP) and the Society for Environmental Toxicology and
7 Chemistry, (SETAC) Life Cycle Initiative (UNEP-SETAC toxicity model, 2008).
8

9 **2.2 Functional unit.**

10 The choice of system boundaries, the definition of functional unit (FU), and allocation procedures plays
11 an important role in the LCA of food products. The most commonly used functional unit is based on
12 mass, but recently there are more ways of expressing the functional unit for food products, such as energy
13 balance or protein content (Schau et al., 2008), and nutritional value (Martínez Blanco et al., 2011^b). A
14 mass based functional unit is adequate when only analysing the agricultural stages of the life cycle of
15 fruit for descriptive purposes (Milà i Canals et al., 2006). According to fruit and nuts PCR, and in order to
16 compare the results according to yield variations over the years, a mass based functional unit has been
17 chosen, so the FU was defined as a “cultivation of 1kg of peach. Considering that the function of the
18 orchard is to produce, the fact of using an FU based on a mass unit also allows reflecting seasonal
19 variability. A functional unit based on hectares could hide the variability of results.
20

21 **2.3 Experimental orchard design.**

22 The demonstrative plantation studied is a peach orchard of *Prunus persica L.* It has an area of 1 hectare
23 and is located in Gimenells (Lleida) in the North East of Spain. According to a USDA soil Survey (1999),
24 the soil texture is loamy, and its physical proprieties are favourable for the root development, with fruit
25 cultivation being possible. The orchard has a planting frame of 4x5m, the plot is designed with 15 rows of
26 20 trees (300trees/ha). Peach cultivation is done using integrated fruit production according to the
27 European Integrated Farming Framework 2010. Integrated Production is defined as a system of
28 agricultural production of quality food, using methods that respect the environment and human health in
29 order to obtain high quality products, minimize the use of agrochemicals, optimize production methods
30 and reduce waste. The average yield of the orchard is 36,148 kg· ha⁻¹ ±10%. Maximum production was
31 achieved in year eight, with 48,350 kg· ha⁻¹, and the minimum was achieved in the second year, with a
32 production of 18,745 kg· ha⁻¹
33

34 **2.4 System description.**

35 The study only focuses on the cultivation period, but it also includes the initial establishment tasks: soil
36 preparation and plantation. The nursery stage has been excluded, mainly due to the lack of reliable data
37 regarding this phase of peach-growing. According to Vazquez-Rowe et al. 2012, given the longevity of
38 the crops, and the small percentage of annual tree replacement, its exclusion should not significantly
39 affect the final results. The agricultural stages considered in this study are: soil preparation and plantation,
40 fertilization, irrigation, pest management, weeds mowing, pruning, and harvest. Post-harvest operations
41 (storage, processing, packaging, and commercialization) are not included. Figure 1 shows the boundaries
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62

1 of the system studied. Different management tasks are involved depending on the agricultural stage. Soil
2 preparation and plantation tasks were performed mechanically with tractors (these tasks are only done
3 once, at the beginning of the orchard when trees are planted). Agrochemicals were applied using a tractor
4 and a sprayer. Fertilizers were applied through the irrigation system with electric pumps (fertigation).
5 Pruning was done manually, and the wood was crushed with a tractor implement and incorporated into
6 the soil (it is considered as a soil structuring effect). The emissions related to pruning waste degradation
7 have not been considered. It has been assumed that biogenic carbon emitted, as well as the biogenic
8 carbon fixed from the biomass of the tree or the peach, was not taken into account as a potential global
9 warming indicator according to the ISO14067/ PAS2050-1. For this case study, the information about the
10 effect of the pruning waste has on the soil was not available, however it is an interesting effect that will be
11 considered for further research. Weed mowing was done with a tractor implement. Harvesting was done
12 manually, but required a self-driven platform and a tractor to transport the fruit to the storehouse.
13 Considering the characteristics of the system studied and all the tasks involved, the following inputs have
14 been taken into account to make the inventory: production of fertilizers and their application to the field,
15 pest management substances manufacture and their application (fungicides and insecticides), machinery
16 manufacture and implements used with their transport to the orchard, water use, energy use (from
17 irrigation pumps and inputs manufacturing).

18 <<Figure 1. System boundaries. >>

19 2.5 Inventory.

20 Experimental data have been directly obtained from the Catalan Research Institute of Food and
21 Agriculture Technology (IRTA) orchards. The data used to make the inventory cover 15 years of real
22 production. Table 2 shows the inventory considered in the study, according to the FU described in Section
23 2.2 and also in Figure 1. As the FU chosen is related to the kg of fruit produced, four different production
24 scenarios have been considered related to the different production periods of the fruit tree: *Growth*, *Low*,
25 *High* and *Multiyear*, in order to analyse the variation of environmental results according to the orchard
26 yield, all four scenarios are based on real data. The *Growth* scenario covers between years 1 and 3, when
27 the orchard starts to produce fruit but has not yet reached full production. *Low* scenario covers when the
28 fruit tree starts to go into full production between years 4 and 5. *High* scenario is when the orchard
29 reaches the maximum fruit yield, around year 7. *Multiyear* scenario is the average of 15 years production.
30 The production considered for the Growth scenario is 18,745 kg ha⁻¹, for Low it is 31,625 kg ha⁻¹, for
31 High is 48,350 kg ha⁻¹, and the production considered for Multiyear is 36,280 kg ha⁻¹. The production
32 ranges chosen are in line with the fruit growing study reported by Iglesias (2013). These scenarios are
33 proposed to quantify the variability of LCA indicators depending on the consideration of a single year
34 (productive or not) or a multiyear approach.

35 << Table 2. Inventory for FU and selected production scenarios (FU=1kg of peach). >>

2.5.1. Data assumptions.

Some assumptions have been made in order to optimize the calculation and the application of the methodologies:

- **Change Land Use:** It was assumed that the land occupied is arable and that it had been used for agriculture for a long time. Therefore, no impacts caused by land transformation were taken into account as the plot has been an orchard for more than 25 years (ISO14067).

- **Agrochemicals substances** (insecticides and fungicides). The active ingredients of the pesticides used have been taken into account. Individual pesticide production data were not available, thus the generic pesticide process from ecoinvent database v3.0 has been chosen (ecoinvent Centre).

- **Machinery:** Experimental data of the number of machinery operations and the working hours for running the machines and their implements have been used to quantify the amount of machinery input per kg of fruit. Emissions of machinery production and diesel consumed for machinery operations have also been taken from ecoinvent database v.3.0 (Ecoinvent Centre).

- **Electricity:**

The electricity consumed for irrigation pumps was known, and the impact of generation and distribution of electricity demand was estimated using information from ecoinvent database v3.0, according to the Spanish electricity mix of low voltage (ecoinvent Centre).

- **Irrigation water:** The orchard studied was irrigated with electric pumps, and the water came from a Catalan public irrigation canal (Catalonia-Aragon).

- **Fertilizer emissions:** It has been considered the emissions of fertilizer production and the emissions of fertilizers used have been taken into account. Nitrogen (N_2O), phosphorus (P_2O_5) and potassium (K_2O) emissions were modelled according to the literature (Bentrup 2001; Audsley 1997). As regards diffuse emissions, according Audsley (1997), it was assumed as 2% of NH_3 volatilization for simple nutrient fertilizer (ammonium nitrate), and 4% for multinutrient fertilizer (NPK). NO_x emissions were assumed as 10% of the N_2O emissions. The N_2O emission factor assumed for all fertilizers is 1.25% of N addition (Bentrup, 2001).

- **Transport of input materials and substances to the orchard:** It was assumed that the vehicle used to transport the materials and substances from the production plant to the local point of sale was a 7.5 t lorry, and the distance covered was 150km. The vehicle considered to deliver the materials from the regional cooperative to the plantation was a small van<3.5t and the distance, 15km.

- **Carbon sequestration:** there is a lack of knowledge on specific topics, and in particular a lack of inventories to estimate carbon sequestration (Alaphilipe et al., 2012). No specific land and biomass carbon sequestrations were taken into account in this work, as the soil carbon content remained constant during the years of the study, and there was no change in the use of the land. Biogenic carbon has not been considered as either kidnapped or as issued, because is for temporary short chain.

3. RESULTS AND DISCUSSION.

The results obtained are based on the FU (production of 1kilogram of peach) and according to the impact indicators defined in Section 2.1. Results and discussion are organized into three sections: 3.1 LCA results, 3.2 Agricultural stages impact contribution, 3.3 Annual evolution of impacts

3.1 LCA results

Figure 2 shows relative percentage values of all impact categories analysed according to scenarios considered. Large differences between 30-50% were identified in the environmental impact values depending on the production scenario (Growth, Low, High and Multiyear). *High* scenario presents the lowest impact in relative percentage in all impact categories, because the maximum yield is achieved in this scenario, thus the amount kg of peach produced is higher than other scenarios (48,350 kg.ha⁻¹), so the impact associated per kg of fruit becomes lower. *Growth* scenario has the highest relative percentage in all impact categories, as the kg of fruit produced is very low (18,745 kg.ha⁻¹), thus the impact associated per kg of fruit produced becomes the highest.

<< **Figure 2.** Relative impact values per FU depending on the production scenarios considered. >>

3.2 Agricultural stages impact contribution

Table 3 shows the results of total impact values and agricultural stages contribution expressed in % for each impact category studied, and according to the production scenario (*Growth, Low, High and Multiyear*). Depending on the scenario considered, the percentage contribution of each impact category can vary between 7%-69%. Impact values per FU for all agricultural stages studied become higher if they are calculated according to *Growth* and *Low* scenarios because the kg of peach produced in this scenario are lower. In the impact categories required in Fruit and nuts PCR a range of variance can be observed in the results: NRE 8-52%, GWP 11-50%, WDP 14-52% and Etox 12-69%.

Of the all agricultural stages considered in the study, fertigation is the stage that presents the highest contribution percentage in 10 of the 14 impact categories studied for all scenarios, with a maximum contribution of 99.93% in WDP category (*Growth* scenario) and a minimum contribution of 45.66% in FDP category (*Multiyear* scenario). Pest management presents the highest contribution in the 4 remaining impact categories, with a maximum of 64.50% in Etox (*Growth* scenario) and a minimum of 47.22% in ODP. Fertilizer dosages are calculated according to the requirements of the fruit tree, and the maximum potential yield expected for each year according to the variation in annual conditions (age of the orchard, soil analysis results and climatic conditions); thus, every year a maxim value of fertilizers dosage is defined for the orchard, because once the maximum dose is exceeded it does not guarantee higher yield.

If the soil preparation and planting stages are not included in the calculation, then the 5% of total emissions can be overlooked, considering that the life span of peach plantation of 15 years. However they are distributed over the fifteen years of production (these tasks are only done once, at the beginning of the orchard when trees are planted). Cerutti (2013^a) reported that orchards are perennial and biological systems, and these two characteristics add complexity to the modelling of fruit systems. But if the productive period alone is considered, the environmental impacts of the final product are underestimated considerably. The longer the life span of the peach crop plantation, the lower is the contribution of initial

1 agricultural stages such as soil preparation and planting. Milà I Canals (2006) stress that it is important to
2 consider the nursery in environmental impact assessments, but a lack of data makes this difficult.
3 According to Cerutti (2013^a), the nursery where orchard seedlings are produced should be considered an
4 upstream process delivering grafted plants to the orchards, and the impact during this stage should be
5 included in assessments of fruit production systems, even if impacts are spread over the lifetime of the
6 orchard. This has not been included in this study due to the lack of available experimental information.
7

8
9 Results obtained were compared with other publications of fruit production with similar agricultural
10 stages and life span such as: apple, orange and peach production. The impact percentage contribution of
11 the agricultural stages considered in the study are consistent with the results described for other fruit
12 studies described in Table 1 such as Sanjuán (2005), Coltro (2009), Clasadonte (2010^{a,b}) and Alaphilipe
13 (2012). It should be mentioned that the aforementioned publications only considered productive periods
14 for one year. The existing studies for peach fruit are related to Eco indicators, so it has not been possible
15 to validate and compare the total impact values obtained in this study because there are no existing peach
16 studies focused on mid-point characterization factors.
17

18
19 Once the environmental impact of the different agricultural stages is evaluated, and taking into
20 consideration that fertigation is the stage that presents the highest contribution percentage in 10 of the 14
21 impact categories studied for all scenarios (this is because manufacturing of fertilizers have a significant
22 impact), it would be interesting for farmers to try and choose another kind of fertilizer with low
23 environmental impact and encourage them to try to adjust the application dose, with better the monitoring
24 of nutrients contents in soil and crop. Other important recommendations for farmers, in order to promote
25 orchard better environmental performance are: the orchard design (trees orientation, planting frame,
26 irrigation system) and geographic location, in order to reduce water consumption and pesticides use.
27
28
29
30
31
32
33
34
35

36 <<Table 3. Total and relative percentage impact values per FU depending on the production scenarios considered in
37 this study.>>
38

39 **3.3 Annual evolution of impacts**

40 This section describes the annual evolution of the impact categories recommended for Fruit and Nuts
41 PCR: a) Non-renewable resources, b) Global warming potential, c) Water depletion, and d) Ecotoxicity
42 during the 15 years of life span of peach orchard.
43
44

45 From year 1 to year 4 orchard is not very productive because the fruit tree is getting stabilized, from year
46 5 it starts to come into full production. Between years 6 to 12 it reaches the stable production. From year
47 13 to 15 the orchard begins to lose efficiency until year 15 when it ceases to be productive and acquires
48 the finest of its useful life. Note that in the first year (year 1) there is no fruit production, so results are
49 discussed from second year; maximum production was achieved in the year 8 with 48,350 kg· ha⁻¹ and
50 the minimum was achieved in the year 2, with 18,745 kg· ha⁻¹. Note that annual yield variability not only
51 depends on the orchard age, it also depends on the meteorological conditions, so the fact of considering a
52 range of years allows these variables to be reflected on the yield. The climatic conditions are an important
53 fact to be considered in agricultural practices when annual variation is studied, because weather has an
54 effect on many aspects of fruit production such as: yield, water requirements, pest management and weed
55
56
57
58
59
60
61
62
63
64
65

1 mowing. Thus, it is important to have meteorological inventory with local data evolution, in order to
2 evaluate the multiyear approach. The fact of studying 15 years of real production, means that the
3 variability of the climatic conditions are more reflected in the results than if we only had studied a single
4 year of production. Note that the yield variations not only depend on the orchard age, it also strongly
5 depends on other variables, such as fruit variety, geographic location, planting frame, climatic conditions,
6 irrigation system, fertilizer supply and pest control optimization. Therefore, it is important to have data
7 related to these other variables and for as long a period of time as possible to as to reflect them in the
8 results.
9
10

11 12 13 14 **a) Non-renewable energy resources (NRE)**

15 Figure 3a represents the MJ.ha⁻¹ consumed for the inputs considered during 15 years. It can also can be
16 observed The evolution of MJ per kg of peach per year (dashed line) can also be observed in the Graph.
17 As regards the MJ required to produce 1 kg of peach, it can be observed that during the second year the
18 value is very high around 4 MJ·kg⁻¹, but from the second year onwards fruit yield per hectare begins to
19 increase, then the MJ. kg⁻¹ decreases, reaching a minimum value in year 8 of around 1.8 MJ·kg⁻¹ (in this
20 year the maximum yield is achieved). Concerning the values of the MJ.ha⁻¹ for inputs considered, it can
21 be observed that they are inversely proportional to the kg produced. Fertilizers are the input with the
22 highest contribution in this category during the fifteen years considered, due to the energy needed for
23 their manufacturing. In the second year the MJ.ha⁻¹ has a minimum of 3.8 E+04 MJ.ha⁻¹, as fruit trees are
24 not very productive low fertilization is needed, just the dose to promote the correct development of the
25 crop but no the fruit. On the other hand during years 8-10 the maximum yield is achieved, so more
26 fertilization is needed to promote fruit development, in consequence the MJ.ha⁻¹ consumed has a
27 maximum value 6.5 E+04 MJ.ha⁻¹. Between years 10 to 12 the MJ.ha⁻¹ tends to stabilize because as the kg
28 produced becomes stable so fertilizers dose also becomes stable. In years 14 and 15 the MJ.ha⁻¹
29 consumed decreases, because the orchard starts to become unproductive, so less agricultural tasks are
30 invested due to fruit yield decrease.
31
32
33
34
35
36
37
38
39
40
41

42 **b) Climate change (CCH)**

43 Figure 3b quantifies the emissions in terms of kg CO_{2eq}.ha⁻¹ for the inputs considered during 15 years, and
44 also illustrates the evolution of the kg CO_{2eq} per kg of peach and year. Regarding to the kg CO_{2eq} emitted
45 to produce 1 kg of peach it can be observed that during the second year the maximum value is achieved
46 0.37 kg CO_{2eq}·kg⁻¹, but from the second year onwards fruit yield begins to increase, then the kgCO_{2eq}·kg⁻¹
47 decreases getting the minimum value in year 8 around 0.16 kgCO_{2eq}·kg⁻¹. When fruit yield begins to be
48 constant then kg CO_{2eq} per kg of peach value decreases and then tends to stabilize (0.15-0.20 kgCO_{2eq}·kg⁻¹).
49 About the values of kg CO_{2eq}.ha⁻¹, once more fertilizers are the input with the highest contribution for
50 the fifteen years considered, due mainly to the high emissions related to their manufacturing process. The
51 maxim value is achieved in year 8 with 6E+03 kg CO_{2eq}.ha⁻¹. This year is when more fertilization is
52 needed to promote fruit development, and more machinery hours and more diesel consumption is required
53
54
55
56
57
58
59
60
61
62
63
64
65

to collect the kg of fruit produced; and on the contrary the minimum is in year 2 with $3.5E+03 \text{ kg CO}_{2\text{eq}}\cdot\text{ha}^{-1}$ when less fertilization, machinery and diesel are involved.

c) Water Depletion (WDP)

Figure 3c represents the $\text{m}^3\cdot\text{ha}^{-1}$ consumed for the inputs considered during 15 years, and the m^3 per kg of peach and year. It can be observed that the maximum m^3 consumed to produce 1 kg of peach is in the second year, $0.5 \text{ kg CO}_{2\text{eq}}\cdot\text{kg}^{-1}$, then as fruit yield begins to increase, the $\text{m}^3\cdot\text{kg}^{-1}$ decreases, reaching the minimum value in year 8 of around $0.2 \text{ kgCO}_{2\text{eq}}\cdot\text{kg}^{-1}$. As regards the inputs considered, in this category, water has the highest contribution during the fifteen years, due to the amount of water needed for the irrigation. The minimum value of $\text{m}^3\cdot\text{ha}^{-1}$ is achieved in the second year with $1.6E+03 \text{ m}^3\cdot\text{ha}^{-1}$ and this is also the year when low irrigation is required because tree requirements are lower due to it not being a full productive year. The maximum value is achieved in year 8 with $1.8E+04 \text{ m}^3\cdot\text{ha}^{-1}$, as this is the year when high irrigation is needed to help to promote fruit progress. Between years 6-13 the quantification of $\text{m}^3\cdot\text{ha}^{-1}$ tends to be stable because the yield is stable, and as agricultural management tasks are constant, it starts to decrease at years 14-15, when the yield and water requirements become lower.

d) Ecotoxicity (Ecotox)

Figure 3d represents ecotoxicity impact in $\text{CTUe}\cdot\text{ha}^{-1}$ for the inputs considered for 15 years, and also $\text{CTUe}\cdot\text{kg}^{-1}$ per kg of peach and year. In this category, once more, the maximum impact to produce 1 kg of peach is in achieved in the second year, $1.E^{+04} \text{ CTUe}\cdot\text{kg}^{-1}$, and reaches a minimum value in year 8, $4.E^{+03} \text{ CTUe}\cdot\text{kg}^{-1}$. As regards the inputs considered in the values of $\text{CTUe}\cdot\text{ha}^{-1}$, diesel consumption has the highest contribution during the fifteen years, due mainly to the high ecotoxicity of some of its chemical components. The maxim value of $\text{CTUe}\cdot\text{ha}^{-1}$ is achieved in year 8 with $1.2E+08 \text{ CTUe}\cdot\text{ha}^{-1}$. In this year more machinery hours and more diesel consumption are required to collect the kg of fruit produced, and on the contrary the minimum is in year 2 with $1E+08 \text{ CTUe}\cdot\text{ha}^{-1}$ when less machinery and diesel are required because fruit yield is very low. During the intermediate years the quantification of $\text{CTUe}\cdot\text{ha}^{-1}$ tends to be stable, and it starts to decrease at years 14-15 when the yield decreases.

<<Figure 3. Evolution PCR impact values per FU over the years.>

4. CONCLUSIONS

The results of the study reveal that when the FU is related to mass units, using different production scenarios can generate a variation in the environmental impact results of between 7% and 69%, depending on the impact indicator. Therefore, caution should be taken when the FU is related to mass and only a single year is studied, because in the years that the yield is low the impact values per FU increase. On the contrary, in the years that the yield is high, the impact values per FU decrease; thus, depending on the year chosen for the fruit studies, the results can be biased. Geographic location of fruit orchards is also an important aspect to be considered in the data collection phase, because in temperate areas orchards reach maturity as early as two years after the plantation, and reach full production from the fifth year, which can significantly affect yield average, depending on the amount of years taken into account. Yield variability not only depends on the orchard age, it also strongly depends on other variables such as fruit variety, geographic location, planting frame, climatic conditions, irrigation system, fertilizers supply and pest control optimization. So it is important, when the UF is related to yield, to have data related to other variables and for many periods of time as possible, to reflect these variables in the results. According to the results obtained in this study in crops with an average life time of twenty years or longer, a multiyear approach is strongly recommended when the functional unit is related to kg produced.

In agricultural stages contribution, fertigation has the highest contribution in all impact categories studied, followed by pest management. This is because manufacturing of fertilizers and pesticides have a significant impact. Weed mowing, pruning and harvest impacts are mainly due to the use of machinery, and their involvement in the cultivation process is more sporadic than fertigation. As regards initial orchard establishment tasks (soil preparation and planting), if they are not included in the impacts quantification, 5% of total emissions may be overlooked. Sometimes a lack of data makes it difficult to inventory and include these stages. On the other hand, it is essential to encourage the farmers to try to choose another kind of fertilizer with low environmental impact, and encourage them to try to regulate the application dose, and improve the monitoring of nutrients contents in soil and crop. They should also consider the orchard design and its geographic location to promote a better orchard environmental performance.

This study contributes to complete the fruit LCA literature and provides new information for peach analysis, as well as introducing a multiyear perspective analysis to identify the variability of results related to annual yield conditionings and climatic conditions. The results may be useful to identify the hot spots of peach production, in order to identify the stages with higher impact and obtaining more environmentally friendly fruit practices. The study also provides new inventory data and results on the Mediterranean peach fruit. This work also provides an additional methodological perspective. Although LCA is a useful tool for estimating the impact associated with a product or process, there are still some issues to be resolved regarding to the quality of environmental impact databases and data available because sometimes, due the need to work with generic data, as in the case of pesticides or fertilizers, it may vary the results.

To complete this study, systems boundaries will be further extended to embrace the whole life of peach production, from plant production and plantation to final consumer disposal, in order to estimate the overall impact.

41 **Acknowledgments** We thank the staff members of IRTA experimental Station of Lleida unit in charge of
42 the management of the orchards for their helpful comments on an earlier draft of the manuscript.
43 Constructive comments from reviewers have improved the clarity of the paper and are also
44 acknowledged.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

45 5. REFERENCES

- 46 Alaphilippe A., Simon S., Brun L., Hayer F., Gaillard G., 2012. Life cycle analysis reveals higher
47 agroecological benefits of organic and low-input apple production. *Agron. Sustain. Dev.* DOI
48 [10.1007/s13593-012-0124-7](https://doi.org/10.1007/s13593-012-0124-7).
49
- 50 Assomela (Associazione Italiana Produttori di Meleapple), 2012. Apple produced in the Region of
51 Trentino Alto Adige Italy. PCR:2012:07 version 1.0 of 23/08/2012. Environdec (2012)
52 <http://www.environdec.com> (Accessed 01.06.14).
53
- 54 Audsley E., 1997. Harmonization of environmental life cycle assessment for agriculture. European
55 Commission DG VI Agriculture. Final Report Concerted Action AIR 3-CT94-2028.
56
- 57 Beccali M., Iudicello M., Mistretta M., 2010. Life cycle assessment of Italian citrus-based products.
58 Sensitivity analysis and improvement scenarios. *Journal of Environmental Management*.
59 DOI:[10.1016/j.jenvman.2010.02.028](https://doi.org/10.1016/j.jenvman.2010.02.028)
60
- 61 Brentrup F., Küsters J., Kuhlmann H., Lammel J., 2001. Application of the life cycle assessment
62 methodology to agricultural production: an example of sugar beet production with different forms of
63 nitrogen fertilizers. *European Journal of Agronomy* 14 (3). 221–233.
64
- 65 Bellon-Maurel V., Short M.D., Roux P., Schulz M., Peters G.M (2014). Streamlining life cycle inventory
66 data generation in agriculture using traceability data and information and communication technologies e
67 part I: concepts and technical basis. *Journal of Cleaner Production* 69 (2014) 60e66 DOI:
68 <http://dx.doi.org/10.1016/j.jclepro.2014.01.079>
69
- 70 Cerutti AK., Bagliani M., Beccaro GL., Bounous G., 2010. Application of Ecological Footprint Analysis
71 on nectarine production: methodological issues and results from a case study in Italy. *Journal of Cleaner*
72 *Production* 18 (2010) 771–776. DOI:[10.1016/j.jclepro.2010.01.009](https://doi.org/10.1016/j.jclepro.2010.01.009)
73
- 74 Cerutti AK., Beccaro GL., Bruun S., Bosco S., Donno D., Notarnicola B., Bounous G., 2013^a. LCA
75 application in the fruit sector: State of the art and recommendations for environmental declarations of
76 fruit products, *Journal of Cleaner Production*. DOI: [10.1016/j.jclepro.2013.09.017](https://doi.org/10.1016/j.jclepro.2013.09.017)
77
- 78 Cerutti A., Bruun S., Donno D., Beccaro G., Bounous G., 2013^b. Environmental sustainability of
79 traditional foods: the case of ancient apple cultivars in Northern Italy. *Journal of Cleaner Production*,
80 52:245–252.
81
- 82 Clasadonte MT., Matarazzo A., Ingrao C., 2010a. Life cycle assessment of Sicilian peach sector.
83 Notarnicola, B., Settanni, E., Tassielli, G., Giungato, P. (Eds.), *Proceedings of LCA food 2010*. Bari, pp.
84 295-300.
85
- 86 Clasadonte, MT., Lo Giudice A., Ingrao C., 2010b. Life Cycle Assessment of the Sicilian citrus fruit
87 field. *Proceedings of LCA food 2010*. Bari, pp. 301-306.
88
- 89 Coltro L., Mourad A.L., Kletecke L.M., Mendonça T.A., Germer S., 2009. Assessing the environmental
90 profile of orange production in Brazil. *Int. J. of LCA*.14 (7).pp 656-664. DOI [10.1007/s11367-009-0097-1](https://doi.org/10.1007/s11367-009-0097-1)
91
- 92 Ecoinvent centre. Technical Documentation of the ecoinvent Database. Available at:
93 <http://ecoinvent.org/database> (Accessed 12.10.14).
94
- 95 European Integrated Farming Framework: A European Definition and Characterization of Integrated
96 Farming as Guideline for Sustainable Development of Agriculture, revised version August 2010.
97 European Initiative for Sustainable Development in Agriculture.
98
- 99 European Commission Agriculture and rural development publications (2012). Available at:
100 <http://ec.europa.eu/agriculture> (Accessed 12.04.14).
101
- 102 Environdec Organization, 2012. The International EPD^R System- a Communications tool for International
103 Markets <http://www.environdec.com> (Accessed 20.07.14).
104

105 FAO (Food and Agriculture Organization of the United Nations) Annual statistics,
106 2011.<http://faostat.fao.org/site/339/default.aspx> (Accessed 27.03.14).
107

108 Global Gap (EU working group). Available at: ([http://www.globalgap.org/uk_en/what-we-do/the-gg-
109 system/gg-farm-assurers/](http://www.globalgap.org/uk_en/what-we-do/the-gg-system/gg-farm-assurers/)) (Accessed 01.12.14).
110

111 ISO 14040:2006 - Environmental management - Life cycle assessment - Principles and framework.
112 International Standard Organization. Available at: <http://www.iso.org/iso/home.htm> (Accessed 16.05.14).
113

114 ISO 14025:2006 - Environmental labels and declarations-Type III environmental declarations- Principles
115 and procedures. International Standard Organization. Available at: <http://www.iso.org/iso/home.htm>
116 (Accessed 16.05.14).
117

118 ISO14047:2012 - Environmental management - Life cycle assessment - Illustrative examples on how to
119 apply ISO 14044 to impact assessment situations. Available at: <http://www.iso.org/iso/home.htm>
120 (Accessed 16.05.14).
121

122 IPCC Climate Change, 2007. Synthesis Report. Contribution of Working Groups I. II and III to the
123 Fourth Assessment Report of the Intergovernmental Panel on Climate Change .International Panel on
124 Climate Change. Geneva. Available at: <http://www.ipcc.ch/> (Accessed 28.02.14).
125

126 Ingwersen W., 2012. Life cycle assessment of fresh pineapple from Costa Rica. Journal of Cleaner
127 Production 35 (2012) 152e163.DOI:[10.1016/j.jclepro.2012.05.035](https://doi.org/10.1016/j.jclepro.2012.05.035)
128

129 Iglesias I., 2013. Peach production in Spain: current situation and trends, from production to
130 consumption.Proceedings of the 4th Conference, Innovations in Fruit Growing, 75-98. Ed.: D. Milatović.
131 Belgrad (Serbia)
132

133 Liu Y., Langer V., Høgh-Jensen H., Egelyng H., 2010. Life Cycle Assessment of fossil energy use and
134 greenhouse gas emissions in Chinese pear production. Journal of Cleaner Production Volume 18.Issue 14.
135 September 2010. Pages 1423-1430.
136 DOI:[10.1016/j.jclepro.2010.05.025](https://doi.org/10.1016/j.jclepro.2010.05.025).
137

138 MAGRAMA. Spanish Ministry of Environment and Rural and Marine Affairs, 2010 Practical guide to
139 rational crop fertilization in Spain.
140 Available at: <http://www.magrama.gob.es/ca/agricultura/publicaciones> (Accessed 11.05.14).
141

142 Martínez-Blanco J., Muñoz P., Antón A., Rieradevall J., 2011^a. Assessment of tomato Mediterranean
143 production in open-field and in standard multi-tunnel greenhouse with compost or mineral fertilizers from
144 an agricultural and environmental standpoint. Journal of Cleaner Production 19 (2011) 985e997.
145 DOI:[10.1016/j.jclepro.2010.11.018](https://doi.org/10.1016/j.jclepro.2010.11.018)
146

147 Martinez-Blanco, J.; Muñoz, P.; Antón, A.; Rieradevall, J.; Castellari, M., 2011^b. Comparing nutritional
148 value and yield as functional units in the environmental assessment of horticultural production with
149 organic or mineral fertilization. International Journal of Life Cycle Assessment 16 :12-26
150

151 McLaren SJ., Hume A., Mithraratne N., 2010. Carbon management for the primary agricultural sector in
152 New Zealand: case studies for the pipfruit and kiwifruit industries. Proceedings of LCA food 2010, pp.
153 293-298.
154

155 Mila i Canals L., Clemente Polo G., 2003. Life cycle assessment of fruit production. In: Mattsson, B.,
156 Sonesson, U. (Eds.), Environmentally friendly food processing. Woodhead Publishing Limited and CRC
157 Press LLC, Cambridge and Boca Raton, 2003, 29–53.
158

159 Mila i Canals L., Burni GM., Cowell SJ., 2006. Evaluation of the environmental impacts of apple
160 production using Life Cycle Assessment (LCA): case study in New Zealand. Agriculture Ecosystems and
161 Environment 114. 226–238.DOI:[10.1016/j.agee.2005.10.023](https://doi.org/10.1016/j.agee.2005.10.023)
162

163 Mouron P., Nemecek T., Scholz R., Weber O., 2006. Management influence on environmental impacts in
164 an apple production system on Swiss fruit farms: Combining Life Cycle. Assessment with statistical risk
165 assessment/Agriculture. Ecosystems and Environment 114 (2006) 311–322.
166 DOI:[10.1016/j.agee.2005.11.020](https://doi.org/10.1016/j.agee.2005.11.020)
167

168 Ntiamoah A., Afrane G., 2008. Environmental impacts of cocoa production and processing in Ghana: life
169 cycle assessment approach. Journal of Cleaner Production 16 (2008) 1735e1740.
170 DOI:[10.1016/j.jclepro.2007.11.004](https://doi.org/10.1016/j.jclepro.2007.11.004)
171

172 (SAGP) Sustainable Agriculture Guidelines Principles. The Coca-Cola Company. Available at:
173 [http://www.coca-colacompany.com/press-center/company-statements/sustainable-agriculture-guiding-](http://www.coca-colacompany.com/press-center/company-statements/sustainable-agriculture-guiding-principles)
174 [principles](http://www.coca-colacompany.com/press-center/company-statements/sustainable-agriculture-guiding-principles) (Accessed 21.12.14).
175

176 (SAI) Sustainable Agriculture Initiative. the global food and drink industry initiative for sustainable
177 agriculture. Available at: <http://www.saipatform.org/> (Accessed 01.12.14).
178

179 Sanjuan N., Climent M., Dominguez A., Girona F., 2005. LCA of integrated orange production in the
180 Comunidad valenciana (Spain). International Journal of agriculture resources. Governance and ecology.4
181 (2). pp. 163-177. Poster International Conference of biocitrics. November 2005. Gandia. Spain.
182

183 Schau EM., Fet AM., 2008. LCA Studies of food products as background for environmental product
184 declarations. International Journal of LCA 13 (3). 255e264.DOI:[10.1065/lca2007.12.372](https://doi.org/10.1065/lca2007.12.372)
185

186 Sim S., Barry M., Clift R., Cowell SJ., 2007. The relative importance of transport in determining an
187 appropriate sustainability strategy for food sourcing. The International Journal of Life Cycle Assessment.
188 Springer Berlin / Heidelberg. 12, 422-431.
189

190 UNEP-SETAC toxicity model- USEtox: recommended characterisation factors for human toxicity and
191 freshwater ecotoxicity, 2008. International Journal Of Life Cycle Assessment, vol. 13, p. 532-546.
192

193 USETOX official USEtox model documentation <http://www.usetox.org> (Accessed 01.09.14).
194

195 USDA Soil Survey Staff. 1999. Soil Taxonomy, 2nd ed. USDA, Natural Resources Conservation Service,
196 Agricultural Handbook No. 436, U.S. Gov. Print. Office, Washington, D C. 869 pp. (afegir a les
197 references).
198

199 Vazquez-Row I., Villanueva Rey P., Moreira MT., Feijoo G., 2012. Environmental analysis of Ribeiro
200 wine from a timeline perspective: Harvest matters when reporting environmental impacts. Journal of
201 Environmental Management 98 (2012) 73e83
202

203 Williams A., Pell E., Webb J., Moorhouse E., Audsley E., 2008. Strawberry and tomato production for
204 the UK compared between the UK and Spain. Proceedings of the 6th International Conference on LCA in
205 the Agri-Food Sector, pp. 254-262.
206

207 Zeus Kiwi. Greece EPD PCR 2011:02 UN CPC 01342 Version 1.0 kiwi fruit. Approval date: 3 January
208 2012. Available at: <http://www.zeuskiwi.gr> (Accessed 26.07.14).
209
210
211
212
213
214
215
216
217

Table 1. Environmental fruit studies published in last 10 years.

Fruit	Country	Tool	Method	Boundaries	FU	Cultivation Period	Initial stages	References.
EUROPE								
Apple	Switzerland	LCA	SALCA v.31	Production	ha, \$	4 years	No	Mouron et al. 2006
Apple	France	LCA	CML EDIP97 IPCC 2007	Production	ha	1 year	No	Alaphilipe et al. 2012
Apple	Italy	LCA	EDIP	Production	kg	1 year	yes	Cerutti et al. 2013 ^b
Apple	Italy	LCA	CML01	Production & supply chain	kg	1 year	yes	Assomela 2012
Kiwi	Greece	LCA	CML01	Production & supply chain	kg	1 year	yes	Zeus 2012
Citrus (products)	Italy	LCA	IPCC GWP100 CML01	Production & processing	kg	1 year	No	Beccali et al. 2010
Orange	Spain	LCA	CML	Production	t	1 year	No	Sanjuán et al. 2005
Orange	Italy	LCA	Impact 2002+	Production & processing	kg	1 year	No	Clasadonte et al. 2010 ^b
Peach	Italy	LCA	Impact 2002+	Production & processing	kg	1 year	No	Clasadonte et al. 2010 ^a
Nectarine	Italy	EF	Eco indicator 99	Production	gha t ⁻¹	1 year	No	Cerutti et al. 2010
Strawberry	Uk, Spain	Literature	IPCC 2007	Production & processing	kg	1 year	No	Williams et al. 2008
OTHER COUNTRIES								
Apple	New Zealand	LCA	EDIP97	Production & processing	t	2 years	yes	Milà Canals et al. 2006
Apple	Brazil, Uk	LCA	CML01 Baseline 2000	supply chain	t	1 year	yes	Sim et al. 2007
Cacao	Ghana	LCA	CML01	Production & processing	kg	1 year	No	Ntiamoah et al. 2008
Pear	China	LCA	IPCC 2007	Production & processing	t	1 year	No	Liu et al. 2010
Orange	Brazil	LCA	EMS4 PIRA	Production & processing	kg	1 year	No	Coltro et al. 2009
Pineapple	Costa-Rica	LCA	PAS,2050 USEtox TRACI	Production & processing	kg	1 year	No	Ingwersen 2012
kiwi	New Zeland	CF	PAS 2050	Production & supply chain	kg	6 years	yes	Mc. Laren et al 2010

Table 2. Inventory for FU and selected production scenarios (FU=1kg of peach).

Production scenarios		Growth	Low prod	High prod	Multiyear*
INPUTS					
<i>From the technosphere</i>					
Energy inputs					
	Units				
Diesel	g	8.49	6.11	4.48	6.04
Electricity	kwh	0.06	0.04	0.03	0.04
Transport	tkm	1.2	1.24	1.31	1.28
Materials inputs					
<i>Fertilizers:</i>					
K ₂ O	g	4.91	3.64	2.38	3.10
N ₂ O	g	4.96	3.68	2.40	3.14
P ₂ O ₅	g	0.67	0.51	0.33	0.43
<i>Agrochemicals:</i>					
fungicides (generic)	g	2.25	1.33	0.87	1.18
insecticides (generic)	g	0.56	0.33	0.22	0.30
Machinery:					0
Use	g	0.76	0.55	0.40	0.54
Accessories	g	0.05	0.04	0.04	0.05
Water	l	347.43	253.36	165.72	216.93
OUTPUTS					
<i>To the technosphere</i>					
Peach	kg	18,745	31,625	48,350	36,280
<i>Emissions to the atmosphere</i>					
Diesel:					
CO ₂	g	25.95	18.67	13.71	18.47
SO ₂	mg	4.98	6.04	4.44	5.98
VOC	mg	18.46	28.72	21.08	28.41
NOX	mg	169.28	268.67	197.23	265.79
NH ₃	mg	0.07	0.12	0.09	0.12
CH ₄	mg	0.42	0.77	0.57	0.76
N ₂ O	mg	0.41	0.72	0.53	0.71
Fertilizers:					
N ₂ O	mg	6.04E-03	4.48E-03	1.32E-04	5.20E-03
NH ₃	mg	1.27E-02	9.41E-03	9.41E-03	1.09E-02

*Average production

Table 3. Total and relative percentage impact values per FU depending on the production scenarios considered in this study.

Impact category		Soil prep	Planting	Fertigation	Pest m.	Pruning	Moving	Harvest	TOTAL
Units		% total	% total	% total	% total	% total	% total	% total	
<i>Growth</i>									
kg CO2 eq	CCH	0.42	0.58	70.22	21.66	0.86	1.71	4.54	2.50E-01
kg CFC-11 eq	ODP	0.46	0.57	43.89	47.22	0.95	1.89	5.02	2.81E-08
kg NMVOC	PHO	1.91	2.27	0.00	64.47	3.78	7.55	20.02	3.44E+04
kg SO2 eq	TAC	0.52	0.65	67.14	23.10	1.04	2.07	5.49	1.64E-03
kg P eq	FEU	0.15	0.23	66.06	30.48	0.37	0.74	1.97	5.65E-05
kg N eq	MEU	0.44	0.53	40.33	51.45	0.87	1.75	4.63	1.12E-04
m2a	ALO	0.12	0.20	79.31	17.71	0.32	0.64	1.70	2.33E-03
m2a	ULO	0.04	0.05	96.12	3.05	0.09	0.18	0.47	5.45E-03
m2	NLT	0.73	0.89	56.47	29.68	1.47	2.95	7.81	6.87E-05
m3	WDP	0.00	0.00	99.93	0.06	0.00	0.00	0.01	3.21E-01
kg Fe eq	MDP	0.34	0.60	61.84	29.30	0.96	1.90	5.06	1.52E-02
kg oil eq	FDP	0.71	1.16	48.64	37.39	1.46	2.92	7.73	5.21E-02
CTUe	Etox	2.42	0.94	0.00	64.50	3.77	7.55	20.00	5.98E+03
MJ	NRE	0.86	1.76	55.94	30.58	1.41	2.81	7.46	2.38E+00
<i>Low</i>									
kg CO2 eq	CCH	0.34	0.47	70.74	20.23	0.70	1.39	6.15	1.83E-01
kg CFC-11 eq	ODP	0.37	0.46	39.81	50.39	0.76	1.48	6.73	2.08E-08
kg NMVOC	PHO	1.68	2.00	0.00	56.75	3.33	6.65	29.60	2.31E+04
kg SO2 eq	TAC	0.42	0.53	68.16	20.85	0.85	1.68	7.51	1.19E-03
kg P eq	FEU	0.11	0.18	64.37	31.97	0.29	0.58	2.49	4.28E-05
kg N eq	MEU	0.33	0.41	38.20	53.17	0.66	1.33	5.90	8.75E-05
m2a	ALO	0.10	0.16	77.97	18.80	0.26	0.51	2.20	1.72E-03
m2a	ULO	0.03	0.04	96.03	3.06	0.07	0.14	0.62	3.98E-03
m2	NLT	0.61	0.73	57.81	26.39	1.22	2.39	10.85	4.91E-05
m3	WDP	0.00	0.00	99.93	0.06	0.00	0.00	0.01	2.34E-01
kg Fe eq	MDP	0.27	0.48	61.48	29.16	0.77	1.52	6.31	1.12E-02
kg oil eq	FDP	0.58	0.95	49.11	35.29	1.19	2.33	10.54	3.78E-02
CTUe	Etox	1.96	1.43	0.00	52.44	3.07	6.13	34.97	4.36E+03
MJ	NRE	0.69	0.75	55.48	29.75	1.13	2.21	9.98	1.76E+00
<i>High</i>									
kg CO2 eq	CCH	0.32	0.45	67.87	21.60	0.75	1.41	7.60	1.25E-01
kg CFC-11 eq	ODP	0.33	0.40	35.19	54.42	0.67	1.34	7.65	1.42E-08
kg NMVOC	PHO	1.56	1.84	0.00	52.39	3.07	6.14	35.00	1.07E+04
kg SO2 eq	TAC	0.41	0.51	65.64	21.70	0.82	1.63	9.29	7.32E-04
kg P eq	FEU	0.11	0.17	59.83	36.01	0.27	0.54	3.08	2.92E-05
kg N eq	MEU	0.29	0.36	33.44	57.53	0.58	1.16	6.63	6.11E-05
m2a	ALO	0.09	0.15	74.47	21.74	0.25	0.49	2.80	1.14E-03
m2a	ULO	0.03	0.04	95.37	3.54	0.07	0.14	0.81	2.60E-03
m2	NLT	0.58	0.70	55.23	26.67	1.17	2.34	13.32	2.91E-05
m3	WDP	0.00	0.00	99.91	0.07	0.00	0.00	0.01	1.53E-01
kg Fe eq	MDP	0.25	0.45	57.22	31.81	0.72	1.42	8.13	7.24E-03
kg oil eq	FDP	0.54	0.89	45.82	36.74	1.11	2.22	12.68	2.31E-02

CTUe	Etox	1.96	1.43	0.00	52.45	3.07	6.13	34.96	1.86E+03
MJ	NRE	0.65	0.70	51.64	31.84	1.06	2.12	11.99	1.09E+00

Multiyear

kg CO2 eq	CCH	0.36	0.50	68.09	21.73	0.80	1.53	6.98	1.62E-01
kg CFC-11 eq	ODP	0.36	0.36	36.38	53.57	0.74	1.48	7.02	2.00E-08
kg NMVOC	PHO	1.69	1.69	0.00	54.60	3.34	6.68	31.69	2.16E+04
kg SO2 eq	TAC	0.45	0.45	65.48	22.30	0.90	1.79	8.51	1.05E-03
kg P eq	FEU	0.12	0.12	60.58	35.39	0.30	0.60	2.83	3.90E-05
kg N eq	MEU	0.33	0.33	34.17	56.91	0.66	1.31	6.22	8.32E-05
m2a	ALO	0.10	0.10	75.08	21.28	0.27	0.54	2.56	1.54E-03
m2a	ULO	0.04	0.04	95.42	3.53	0.08	0.15	0.73	3.44E-03
m2	NLT	0.63	0.63	54.68	28.16	1.26	2.52	11.99	4.46E-05
m3	WDP	0.00	0.00	99.92	0.07	0.00	0.00	0.01	2.01E-01
kg Fe eq	MDP	0.28	0.28	57.69	31.75	0.79	1.57	7.42	1.03E-02
kg oil eq	FDP	0.59	0.59	45.66	37.67	1.21	2.42	11.49	3.49E-02
CTUe	Etox	2.09	1.52	0.00	55.09	3.26	6.52	31.51	3.85E+03
MJ	NRE	0.71	0.71	51.44	32.49	1.23	2.48	10.90	1.62E+00

Figure 1. System boundaries for the production of 1 kg of peach






Figure 2. Relative impact values per FU depending on the production scenarios considered




CCH = climate change; ODP = ozone depletion; PHO = photochemical oxidant formation; TAC = terrestrial acidification; FEU = freshwater eutrophication; MEU = marine eutrophication; ALO = agricultural land occupation; ULO = urban land occupation; NLT = natural land transformation; WDP = water depletion; MDP = metal depletion; FDP= fossil depletion; Etox = Ecotoxicity; NRE demand for non-renewable energy resources.

Figure 3. Evolution PCR impact values per FU over the years


3a) NRE= non renewable energy



3b)CCH= climate change



3c) WDP= water depletion



3d) Ecotox= ecotoxicity

