

Letter to the Editor

Nomenclature for members of the expansin superfamily of genes and proteins

Hans Kende^{1,*}, Kent J. Bradford², David A. Brummell³, Hyung-Taeg Cho⁴, Daniel J. Cosgrove⁵, Andrew J. Fleming⁶, Chris Gehring⁷, Yi Lee⁸, Simon McQueen-Mason⁹, Jocelyn K.C. Rose¹⁰ and Laurentius A.C.J. Voesenek¹¹

¹MSU-DOE Plant Research Laboratory, Michigan State University, East Lansing, MI 48824, USA (*author for correspondence; e-mail hkende@msu.edu); ²Seed Biotechnology Center, University of California, Davis CA 95616, USA; ³Crop and Food Research, Private Bag 11600, Palmerston North, 5301, New Zealand; ⁴School of Biosciences and Biotechnology, Chungnam National University, Daejeon 305-764, Republic of Korea; ⁵Department of Biology, 208 Mueller Laboratory, Pennsylvania State University, University Park, PA 16802, USA; ⁶Department of Animal and Plant Sciences, University of Sheffield, Western Bank, Sheffield S10 2TN, UK; ⁷University of the Western Cape, Department of Biotechnology, Private Bag X17, Bellville 7535, South Africa; ⁸Department of Tobacco Science, Chungbuk National University, 48 Gaesin-dong Hungduk-ku, Chongju 361-763, Republic of Korea; ⁹Biology Department, University of York, PO Box 373, York, YO10 5YW, UK; ¹⁰Department of Plant Biology, Cornell University, Ithaca, NY 14853, USA; ¹¹Plant Ecophysiology, Utrecht University, Sorbonnelaan 16, 3584 CA Utrecht, The Netherlands

Expansins were discovered as proteinaceous factors that have the capacity to induce extensibility and stress relaxation of plant cell walls in a pH-dependent manner (McQueen-Mason *et al.*, 1992). Originally, two such proteins were isolated from cucumber cell walls. After the two genes encoding these expansins were cloned and sequenced (Shcherban *et al.*, 1995), searches of the growing EST and genomic databases showed that expansins are encoded by large multigene families, and it was found that they occur in all land plants examined, ranging from bryophytes to angiosperms (for a review, see Li and McQueen-Mason, 2003). Evidence from work with transgenic plants indicates that endogenous expansins are not only involved in regulating growth (Cho and Cosgrove, 2000; Choi *et al.*, 2003; Zenoni *et al.*, 2004), but also developmental processes such as initiation of leaves (Pien *et al.*, 2001), abscission (Cho and Cosgrove, 2000), and softening of tomato fruits (Brummell *et al.*, 1999). After the initial discovery of the expansins, Cosgrove *et al.* (1997) found that group I allergens of grass pollen have regions of significant amino acid similarity to expansins and a high level of predicted secondary structure identity. Extracts of maize pollen and purified maize group I pollen allergen

applied to grass cell walls *in vitro* were shown to have, indeed, expansin activity. Group I pollen allergens from other grass species likewise exhibited wall extension activity (Li and Cosgrove, 2001). Transcripts encoding proteins similar to pollen allergens were also found in vegetative tissues, and it was suggested that these proteins also act as cell wall-loosening factors (Cosgrove *et al.*, 1997).

The discovery of the pollen allergen-type expansins necessitated the introduction of a nomenclature that differentiated between the two expansin families. Expansins that are phylogenetically related to the originally discovered cucumber expansins were called α -expansins, and expansins that are phylogenetically related to the group-I grass pollen allergens were designated β -expansins (for reviews see Lee *et al.*, 2001; Li and McQueen-Mason, 2003).

Further database searches revealed the existence of plant genes that encode proteins with distant homology to α - and β -expansins. These proteins have been called expansin-like and expansin-related (Lee *et al.*, 2001), or were classified as β 2- and β 3-expansins (Li *et al.*, 2002, Li and McQueen-Mason, 2003). Both α - and β -expansins consist of two domains: an N-terminal domain I

that is distantly related to the catalytic domain of glycoside hydrolase family-45 (GH45) and a C-terminal domain II that is distantly related to group-2 grass pollen allergens (Cosgrove, 2000). Plants also have single-domain proteins that are distantly related to expansin domain I, and these have been named p12 proteins (Ceccardi *et al.*, 1998), plant natriuretic peptides (Gehring and Irving, 2003) and γ -expansins (Li *et al.*, 2003). Moreover, grasses have a group of small proteins homologous to expansin domain II. These proteins, which have long been known as group-2/3 grass pollen allergens (Ansari *et al.*, 1989), were recently designated δ -expansins (Li and McQueen-Mason 2003). Meanwhile, it also has become evident that the GH45-like domain is found in diverse proteins from many organisms other than plants, such as *Dictyostelium* (Li *et al.*, 2002), mussels (Xu *et al.*, 2001), fungi (Li *et al.*, 2002; Saloheimo *et al.*, 2002), nematodes (Qin *et al.*, 2004), endosymbiotic protists in the termite hindgut (Ohtoko *et al.*, 2000), and bacteria (Nembaware *et al.*, 2004; D.J. Cosgrove and J. Sampedro Jimenez, unpublished data). This multitude of expansins and proteins with homologies to one or both expansin domains called for a unified and agreed-upon nomenclature.

General rules

The expansin gene and protein nomenclature will follow the convention established for *Arabidopsis* (Meinke and Koornneef, 1997). Wild-type alleles are abbreviated with three italicized capital letters followed by a qualifying letter, for example, *EXPA* for a wild-type α -expansin gene. Symbols for mutant alleles are written in lower-case and italicized letters (*expA*). Different wild-type genes with the same gene symbol are assigned numbers (*EXPA1*, *EXPA2*, etc.). The numbering of genes will be in chronological order of their discovery. In cases of duplicate genes encoding identical proteins, the genes will be given the same number followed by a lower case letter (*a*, *b*, *c*, and so on). Mutant alleles of a gene are assigned numbers connected with a hyphen to the gene symbol. Accordingly, mutant allele 1 of *EXPA1* is designated *expA1-1*. The abbreviations for protein products of wild-type and mutant alleles correspond to the respective gene symbols but are written in non-italicized let-

ters (*EXPA1* or *expA1-1*). The species designation is given by the initial of the genus and species, such as *AtEXPA1* for *Arabidopsis thaliana* α -expansin gene 1. Potential conflicts arising from two species having the same initials should be resolved by adding one more differentiating letter to the genus or species abbreviation. Under any circumstance, the full gene name must be spelled out at first mention, for example, *Arabidopsis thaliana EXPANSIN A1*.

Nomenclature of expansin and expansin-like genes and proteins

The expansin and expansin-like designations are reserved for genes and their protein products that contain both domain I (the GH45-like domain) and domain II (similar to group-2 grass pollen allergens). Proteins having only one of these domains but not the other will not be classified as expansins.

The classification of expansin and expansin-like genes is based on their phylogenetic relationship. Phylogenetic evidence indicates that all plant expansin and expansin-like genes belong to the same superfamily of genes with a common ancestor (Lee *et al.*, 2001; Li and McQueen-Mason, 2003). Figure 1 shows a phylogenetic tree of the expansin superfamily with representative sequences selected to illustrate the structure of the tree. Four protein families are currently recognized in plants and designated *EXPA*, *EXPB*, *EXLA* and *EXLB* (Table 1). Multiple members of the *EXPA* and *EXPB* families have been shown to possess in vitro expansin activity, namely the capacity to induce rapid extension or stress relaxation of isolated cell walls placed under uniaxial tension. The two expansin-like families, designated *EXLA* (*EXPANSIN-LIKE A*) and *EXLB* (*EXPANSIN-LIKE B*), possess both expansin domains, but their amino acid sequences are very divergent from *EXPA* and *EXPB*. No biological or biochemical function has yet been established for any member of the *EXLA* or *EXLB* families.

If any new gene families within the expansin superfamily of plants are discovered, they will be given the symbols *EXPC*, *EXPD*, or *EXLC*, *EXLD*, etc., as appropriate, following consultations with other workers in the expansin field via the expansin web site (see below). The symbol *EXLX* may be




Figure 1. Phylogenetic tree of expansin protein sequences from *Arabidopsis thaliana* with additional EXLA and EXLB sequences from other species to balance the tree. This is a neighbor-joining tree constructed using the methods and sequences recommended at the website <http://www.bio.psu.edu/expansins/naming.htm>. The additional EXLA and EXLB sequences are either from rice (Os) or from EST assemblies generated at www.tigr.org for cotton (cot), tomato (tom), alfalfa (medi) and lettuce (let). The TIGR EST assembly code is indicated in each name, e.g., TC88001.

Table 1. Revised names and symbols of expansin genes.

Original name	Former symbol ^a	New symbol	New name
α -Expansin	<i>EXP</i> or <i>EXPα1</i>	<i>EXPA</i>	<i>EXPANSIN A</i>
β -Expansin	<i>EXPB</i> or <i>EXPβ1</i>	<i>EXPB</i>	<i>EXPANSIN B</i>
Expansin-like	<i>EXPL</i> or <i>EXPβ2</i>	<i>EXLA</i>	<i>EXPANSIN-LIKE A</i>
Expansin-related	<i>EXPR</i> or <i>EXPβ3</i>	<i>EXLB</i>	<i>EXPANSIN-LIKE B</i>
Various	<i>EXP</i> or none	<i>EXLX</i>	<i>EXPANSIN-LIKE X</i>

^aAs listed on the web site <http://www.bio.psu.edu/expansins/arabidopsis.htm> and in Li *et al.* (2003).

used for genes that fulfill the following criteria: (i) they occur in non-plant organisms; (ii) their protein products have homologies to both expansin domains; and (iii) they do not fall within the established expansin gene families. The symbol *EXLX* does not represent a monophyletic gene family but is rather a temporary polyphyletic 'catch-all' group of expansin-like sequences that do not belong to the recognized expansin gene families. These genes will probably be renamed once the function of their protein products has been identified.

Web site

The web site <http://www.bio.psu.edu/expansins/>, currently maintained by Daniel Cosgrove, Pennsylvania State University, contains recent and archival information on expansins, including a summary of the nomenclature rules and a diagnostic tool to place newly identified expansins on the expansin phylogenetic tree. It lists the newly accepted symbols for all *Arabidopsis* and rice expansin genes, together with previously used symbols. This website is also the clearing house for questions concerning expansins, such as assignment of genus and species abbreviations when the initials are identical in two plant species.

References

- Ansari, A.A., Shenbagamurthi, P. and Marsh, D.G. 1989. Complete primary structure of a *Lolium perenne* (perennial rye grass) pollen allergen, Lol p III: comparison with known Lol p I and II sequences. *Biochemistry* 28: 8665–8670.
- Brummell, D.A., Harpster, M.H., Civeello, P.M., Palys, J.M., Bennett, A.B. and Dunsmuir, P. 1999. Modification of expansin protein abundance in tomato fruit alters softening and cell wall polymer metabolism during ripening. *Plant Cell* 11: 2203–2216.

- Ceccardi, T.L., Barthe, G.A. and Derrick, K.S. 1998. A novel protein associated with citrus blight has sequence similarities to expansin. *Plant Mol. Biol.* 38: 775–783.
- Cho, H.-T. and Cosgrove, D.J. 2000. Altered expression of expansin modulates leaf growth and pedicel abscission in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* 97: 9783–9788.
- Choi, D., Lee Y., Cho, H.-T. and Kende, H. 2003. Regulation of expansin gene expression affects growth and development in transgenic rice plants. *Plant Cell* 15: 1386–1398.
- Cosgrove, D.J. 2000. Loosening of plant cell walls by expansins. *Nature* 407: 321–326.
- Cosgrove, D.J., Bedinger, P. and Durachko, D.M. 1997. Group I allergens of grass pollen as cell wall-loosening agents. *Proc. Natl. Acad. Sci. USA* 94: 6559–6564.
- Gehring, C.A. and Irving, H.R. 2003. Natriuretic peptides—a class of heterologous molecules in plants. *Int. J. Biochem. Cell Biol.* 35: 1318–1322.
- Lee, Y., Choi, D. and Kende, H. 2001. Expansins: ever-expanding numbers and functions. *Curr. Opin. Plant Biol.* 4: 527–532.
- Li, L.-C. and Cosgrove, D.J. 2001. Grass group I pollen allergens (β -expansins) lack proteinase activity and do not cause wall loosening via proteolysis. *Eur. J. Biochem.* 268: 4217–4226.
- Li, Y., Darley, C.P., Ongaro, V., Fleming, A., Schipper, O., Baldauf, S.L. and McQueen-Mason, S.J. 2002. Plant expansins are a complex multigene family with an ancient evolutionary origin. *Plant Physiol.* 128: 854–864.
- Li, Y. and McQueen-Mason, S. 2003. Expansins and cell growth. *Curr. Opin. Plant Biol.* 6: 603–610.
- Meinke, D. and Koornneef, M. 1997. Community standards for *Arabidopsis* genetics. *Plant J.* 12: 247–253.
- McQueen-Mason, S.J., Durachko, D.M. and Cosgrove, D.J. 1992. Two endogenous proteins that induce cell wall expansion in plants. *Plant Cell* 4: 1425–1433.
- Nembaware, V., Seoighe, C., Sayed, M. and Gehring, C. 2004. A plant natriuretic peptide-like gene in the bacterial pathogen *Xanthomonas axonopodis* may induce hyperhydration in the plant host: a hypothesis of molecular mimicry. *BMC Evol. Biol.* 4: 10.
- Pien, S., Wyrzykowska, J., McQueen-Mason, S.J., Smart, C. and Fleming, A.J. 2001. Local expression of expansin induces the entire process of leaf development and modifies leaf shape. *Proc. Natl. Acad. Sci. USA* 98: 11812–11817.
- Qin, L., Kudla, U., Roze, E.H.A., Goverse, A., Popeijus, H., Nieuwland, H.O., Jones, J.T., Schots, A., Smant, G., Bakker, J. and Helder, J. 2004. A nematode expansin acting on plants. *Nature* 427: 30.
- Saloheimo, M., Paloheimo, M., Hakola, S., Pere, J., Swanson, B., Nyyssonen, E., Bhatia, A., Ward, M. and Penttila, M. 2002. Swollenin, a *Trichoderma reesei* protein with sequence similarity to the plant expansins, exhibits disruption activity on cellulosic materials. *Eur. J. Biochem.* 269: 4202–4211.
- Shcherban, T.Y., Shi, J., Durachko, D.M., Guiltinan, M.J., McQueen-Mason, S.J., Shieh, M. and Cosgrove, D.J. 1995. Molecular cloning and sequence analysis of expansins—a highly conserved, multigene family of proteins that mediate cell wall extension in plants. *Proc. Natl. Acad. Sci. USA* 92: 9245–9249.
- Xu, B., Janson, J.C. and Sellos, D. 2001. Cloning and sequencing of a molluscan endo- β -1,4-glucanase gene from the blue mussel, *Mytilus edulis*. *Eur. J. Biochem.* 268: 3718–3727.
- Zenoni, S., Reale L., Torielli, G.B., Lanfaloni, L., Porceddu, A., Ferrarini, A., Moretti, C., Zamboni, A., Speghini, A., Ferranti, F. and Pezzotti, M. 2004. Downregulation of the *Petunia hybrida* α -expansin gene *PhEXPI* reduces the amount of crystalline cellulose in cell walls and leads to phenotypic changes in petal limbs. *Plant Cell* 16: 295–308.