

UC Davis

Recent Work

Title

The effect of road kills on amphibian populations

Permalink

<https://escholarship.org/uc/item/0mf9859v>

Authors

Hels, Tove
Buchwald,, Erik

Publication Date

2001-09-24

THE EFFECT OF ROAD KILLS ON AMPHIBIAN POPULATIONS

Reprinted from Biological Conservation 99(3), Hels T. and Buchwald E. The effect of road kills on amphibian populations, p. 331-340, Copyright 2001, with permission from Elsevier Science

Tove Hels, Danish Forest and Landscape Research Institute, Department of Park and Landscape, Hørsholm Kongevej 11, DK-2970 Hørsholm, Denmark, Email: the@fsl.dk,
Phone: + 45 45 76 32 00, Fax: + 45 45 76 32 33

Erik Buchwald, The National Forest and Nature Agency
Haraldsgade 53, DK-2100 Copenhagen, Denmark, Email: ecb@sns.dk

Abstract: The diurnal movement patterns of *Triturus vulgaris*, *T. cristatus*, *Pelobates fuscus*, *Bufo bufo*, *Rana temporaria*, and *R. arvalis* were investigated during five breeding seasons (1994-1998). Two main questions were addressed: 1) What is the probability of an individual amphibian getting killed when crossing the road? and 2) What fraction of the amphibian populations gets killed by traffic? The rate of movement of 203 adult amphibians was recorded. Information on traffic loads was provided, and mortality risk was calculated depending on traffic loads and movement rate. The probability of getting killed ranged from 0.34 to 0.61 when crossing a road with a traffic load of 3,207 vehicles/day, and from 0.89 to 0.98 when crossing a motorway. The number of amphibians killed on the road was estimated by systematic counts. Population sizes were estimated for all ponds within 250m of the relevant highway stretch. Results indicate that about 10% of the adult population of *P. fuscus* and brown frogs (*R. temporaria* and *R. arvalis*) were killed annually by traffic at this site.

Introduction

The decline of amphibian populations throughout the world is a well-established fact that has received a lot of attention during the last ten years (Blaustein and Wake 1990; Wyman 1990; Blaustein et al. 1994). Several factors have been proposed, but there seems to be a consensus about the fact that human activities are responsible for most of the declines. Road mortality is one factor which is potentially important but has received little attention (but see Fahrig et al. 1995). Traffic may be destructive to animal populations in two ways: directly, in the sense of actually killing individuals and indirectly, by fragmenting a population's habitat (Mader 1984; Mader et al. 1990; Andrews 1990; Groot Bruinderink and Hazebroek 1996; Reed et al. 1996). Fragmentation in turn may lead to isolation of populations which again may result in a reduced population size and an increased stochastic risk of extinction (Bennett 1990). As Fahrig et al. (1995) point out, traffic intensity throughout the world has increased in the last two decades and this goes for Denmark too (Anonymous 1998). Thus, it is likely that the toll of animal lives taken by traffic has increased accordingly.

Several studies have quantified road kills of many different taxa, e.g. toads (van Gelder 1973; Cooke 1995), birds, mammals, amphibians, and reptiles (Hansen 1982; Fuellhaas et al. 1989), butterflies (Munguira and Thomas 1992), snakes (Rosen and Lowe 1994), mammals, birds, and reptiles (Drews 1995), deer and other ungulates (Romin and Bissonette 1996; Groot Bruinderink and Hazebroek 1996). Due to their activity pattern, population structure, and preferred habitats, aquatic breeding amphibians are more vulnerable to traffic mortality than most other species. If they have to cross a road to get from their hibernation site to the breeding pond, or if a road runs through their terrestrial habitat, it may pose a serious threat to the population. Few studies, however, have related the number of road-killed individuals to the size of the total population, and as Huijser and Bergers (1997) mention, and Mallick et al. (1998) infer, a species often found killed on roads may simply reflect the presence of large thriving populations.

It is even more uncommon in the literature to relate road kills to the spatial organisation of the population. Vos and Chardon (1998), however, demonstrated a significant negative effect of road density on the occupation probability of ponds by moor frogs (*Rana arvalis*) in the Netherlands.

Most studies regularly count road kills from slow-moving vehicles (Hansen 1982; Drews 1995; Rosen and Lowe 1994; Mallick et al. 1998), or by foot (Fuellhaas et al. 1989; Munguira and Thomas 1992). These assume that every victim is observed, which may be true for large conspicuous mammals, but is certainly not true for small animals. If an estimate of the total number of animals killed on a road in a given period of time is needed, one must quantitatively compensate for the number of animals that disappear from the road between censuses. Although often noted in the above-mentioned studies, only Munguira and Thomas (1992) attempt to make this compensation.

The present study aimed at quantifying road kills in populations of six amphibian species: common newt (*Triturus vulgaris*), crested newt (*Triturus cristatus*), spadefoot toad (*Pelobates fuscus*), common toad (*Bufo bufo*), moor frog (*Rana arvalis*) and common frog (*Rana temporaria*). During a period of five years (1994-1998) the population sizes were estimated and road kills quantified (including corrections for animals not observed). The problem of quantifying road kills was approached from two different angles:

by establishing a relationship between the probability of getting killed by crossing the road and (a) velocity of the animal, (b) diurnal activity pattern, (c) traffic intensity.

by identifying the proportion of the populations killed on the road, and assessing the importance to the probability of population persistence.

Methods

Study site

The study site is located on the peninsula of Djursland, northern Denmark (56°26'N, 10°34'E). It is situated next to a two-lane road connecting two towns. The road is about 8m wide, with a traffic intensity of c. 3,200 vehicles per 24 hours. The speed limit is 80 km/h but the usual vehicle speed is 10-20 km/h higher. The landscape is dominated by intensively cultivated fields (mostly barley), with small woods and farms here and there. Ponds are quite numerous in the area; most of them are of natural origin (glacial formations), although some are artificially dug (peat bogs, gravel pits).

Population size estimates

Population sizes of *P. fuscus*, *R. temporaria* and *R. arvalis* were estimated in 1996 and 1997 for all ponds within a distance of 1,000m from the relevant road stretch ($n = 14$ ponds). Five of the ponds were completely encircled by drift fences in both years. Estimates for the two *Rana* species were based on egg-clumps counted in 1996 and 1997, which corresponded very well with the number of females and males known to have entered the ponds by the pitfall traps. Moreover, the number of calling males was found to correspond well with the number of males known to be present in the pond. The number of egg clumps and the number of calling males were then used as a basis for an estimate of the population size in the ponds that were not fenced, assuming one egg clump per female and a sex ratio of unity. Population sizes of *T. vulgaris*, *T. cristatus* and *B. bufo* were estimated in the five fenced ponds only. Here, all adults of *P. fuscus* were caught by the fence in pitfall traps and were individually marked in both years. In the nine unfenced ponds, *P. fuscus* was assumed to be absent because there were no males calling either year. For *B. bufo*, the number of calling and/or visible males was assessed at the peak of the calling season and used as a basis of a population estimate.

Velocities and activity patterns of amphibians crossing roads

In order to establish a relationship between the probability for an amphibian of getting killed by crossing the road, data on velocity and diurnal activity pattern of the amphibians, and vehicle intensity, and diurnal

variation, were needed. The velocity of adults of the six naturally occurring amphibian species was recorded during their spring migration to the breeding ponds (i.e. before spawning), as well as during their summer movements (in August) in 1996-1998. Some of the animals were spontaneous migrators, i.e., they were discovered on their way to the breeding pond (and their movement speed recorded directly), whereas some of the animals were caught in pitfall traps by the drift fences. They were then taken to a nearby paved area and released. Time spent and distance moved were then recorded from the time the animal started moving until it left the paved area. Pitfall traps were used to describe the amphibian diurnal activity pattern by emptying the traps at regular intervals during each 24-hour period (April-May 1996 and 1997).

Figures on traffic intensity (number of vehicles per 24 hours, and frequency distribution during a 24-hour period), and the number and frequency of different vehicle types were provided by the Danish Road Directorate. Traffic intensity was recorded as the mean number of vehicles on the road in each one-hour interval in the months of April, May, June, and August 1996 (where amphibian activity and road kills were recorded).

Monitoring of road kills

A 600m stretch of the road was monitored for road-killed adult amphibians every morning at dawn, in order to minimise the number of corpses removed by daytime scavengers (routine monitoring). Both sides of the road were carefully examined (one side at a time) by foot. All road victims were removed after recording in order to avoid double counts. This took place during the breeding period, and in late summer: 30 March - 31 May and 27 July - 1 September, 1995; 11 April - 2 June and 1 August - 2 September, 1996; 25 March - 2 June and 29 July - 1 September, 1997.

In addition to the routine monitoring, 19 control monitorings were conducted during peak spring migration (20-27 April, 1996, and 31 March - 28 April, 1997), to assess the efficiency of our method of monitoring by foot. The control monitorings were spread evenly around the clock. Each road victim was recorded and the site carefully but invisibly marked, so as not to influence the probability of its recording during the routine monitoring. During control monitorings, victims were not removed from the road, so victims not recorded by the following routine monitoring must have disappeared or been missed. A few control monitorings were undertaken as direct continuations of the routine monitoring to check the number of extant victims missed.

The efficiency of monitoring was calculated using basic mark-recapture theory (Lincoln index, e.g. Begon 1979). Let:

R = number of victims recorded by routine monitorings only, C = number of victims recorded by control monitorings only, B = number of victims recorded by routine *and* control monitorings, N = total number of victims.

Assuming that we are dealing with the same population of road kills for both routine and control monitoring and that the probability of missing an amphibian during routine monitoring is independent of the probability of missing it during control monitoring, then

$$\frac{B}{B + R} = \frac{B + C}{N} \quad (1)$$

and by rearranging (1), we get an estimate of the total number of victims (\hat{N}):

$$\hat{N} = \frac{(B + C)(B + R)}{B} \quad (2)$$

We define efficiency of routine monitorings (E) as the fraction of all victims found by routine monitorings:

$$E = \frac{R + B}{N} \quad (3)$$

Replacing N with the expression in (2), we get:

$$E = \frac{(R + B)B}{(B + C)(B + R)} = \frac{B}{B + C} \quad (4)$$

Substituting the expression for E in (3) into (4) yields:

$$\hat{N} = \frac{1}{E} (R + B) \quad (5)$$

Consequently, $F = 1/E$ is the factor to be multiplied by the number of road victims found by routine monitorings to get the estimated total number of road victims.

Model for probability of getting killed when crossing the road

The probability of surviving one road crossing ($P(surv)$) is:

$$P(surv) = \frac{1}{\pi} \int_{\alpha=-\pi/2}^{\alpha=\pi/2} e^{-\frac{Na}{v \cos \alpha}} d\alpha \quad (\text{cf. eq. 10, Appendix})$$

where N = number of cars passing per time unit, a = killing width of car, v = velocity of animal and α = angle of road crossing.

This expression averages the survival probability for all possible crossing angles. Note that according to the expression, survival probability decreases exponentially with increasing traffic intensity (N), and increases exponentially with velocity of the animal. $\alpha = 0$ corresponds to perpendicular road crossing; in this case, survival probability is at its maximum value. As the crossing angle deviates from perpendicular, $\cos \alpha$ – and thereby survival probability – decreases.

Our calculation is based on the fact that amphibians get killed if they are hit, even if only partly, by a wheel but usually not if they remain still under a passing vehicle (pers.obs.). Therefore, we calculated the killing width of vehicles (a) as twice the width of a tyre plus twice the body length of the species in question, assuming that the front and rear wheels traverse exactly the same part of the road. a was calculated as a weighted average of all vehicles occurring on this particular road: 72% cars (< 2 tonnes), 18% vans (between 2 and 3.5 tonnes) and 10% trucks (> 3.5 tonnes, 2% with single wheels and 8% with twin wheels) (Møller, pers.com.). Tyre widths for cars, vans and trucks were 0.22 m, 0.24m and 0.38 m/0.64m (single wheels/twin wheels), respectively (Møller, pers. com.). Anurans usually jump when passed by heavy vehicles (> 3.5 tonnes) (pers. obs.), so for such traffic twice the length of a jump was added to the width of the wheels instead of twice the length of the body. Finally, for all vehicles, a was increased by 5% which is an assessed fraction of vehicles that kill by their wind speed alone rather than by hitting the animals. a is thus proportional to the body length of the amphibians and ranges from 0.71m (*P. fuscus*) to 0.77m (*Triturus* spp.).

Results

Velocities, activity patterns and probability of getting killed by road crossing

We recorded velocities of 203 adult amphibians (Table 1): 185 were recorded during their spring migration (31 March - 10 June) and 18 during their summer movements (30 July - 24 August). All velocities were recorded between 2000 and 0230, reflecting the peak activity period. There was no significant difference between velocities of spontaneously and non-spontaneously moving individuals (Table 1). Because of this non-significant difference, we did not distinguish between spontaneous and non-spontaneous movers in the following analyses. Also, despite the low number of amphibians moving in summer, there was no significant difference between movement rates of adult amphibian individuals moving in spring and summer.

Table 1

Mean speed and distance moved for the six naturally occurring amphibian species in the study area. Sp: undisturbed, spontaneous movement across pavement. Non-sp: movement after relocation to metalled road. P: P-value for H0: movement speeds equal for spontaneous and non-spontaneous moving amphibians (Kruskal-Wallis test).

Species	Number of records		Mean distance moved (m \pm SD)			Mean speed (m/min \pm SD)			
	sp	non-sp	P	sp	non-sp	all	Sp	non-sp	all
<i>Triturus vulgaris</i>	6	19	0.52	1.37(0.39)	1.28(0.65)	1.30(0.59)	0.69(0.49)	0.51(0.25)	0.55(0.32)
<i>Triturus cristatus</i>	0	7	-	-	1.66(0.81)	1.66(0.81)	-	0.99(1.01)	0.99(1.01)
<i>Pelobates fuscus</i>	8	40	0.18	2.81(3.04)	2.33(1.01)	2.41(1.50)	0.97(0.84)	1.50(0.99)	1.41(0.98)
<i>Bufo bufo</i>	38	0	-	3.63(2.67)	-	3.63(2.67)	0.93(0.82)	-	0.93(0.82)
<i>Rana temporaria</i>	11	19	0.06	2.77(2.03)	3.41(1.03)	3.18(1.48)	1.51(2.39)	2.39(1.94)	2.07(2.12)
<i>Rana arvalis</i>	20	35	0.59	5.29(3.05)	2.60(1.77)	3.58(2.64)	2.51(3.06)	1.76(2.18)	2.03(2.54)
Total	83	120	-	3.67(2.81)	2.37(1.38)	2.90(2.17)	1.37(1.93)	1.53(1.62)	1.46(1.75)

In the period of investigation, the sun set between 2000 and 2130 and rose between 0530 and 0700 and the activity patterns of the investigated amphibian species were concentrated at night (Fig. 1). It is clear, however, that the time of peak activity differed between species: most *R. temporaria* and *R. arvalis* were active soon after sunset whereas most *B. bufo* were active between 2200 and 2300. The two *Triturus* species and *P. fuscus* were later still, the latter with a distinct activity peak around midnight to 0200. There was a small rush-hour peak of traffic intensity around 0700 and a large peak around 1500. From 1500 to 0200, traffic intensity decreased steadily, reaching a minimum value of nine vehicles per hour in the middle of the night.




Fig. 1. Diurnal activity patterns of the six species of amphibians investigated and diurnal variation in vehicle intensity. Traffic data for April and May 1996.

According to equation 12 (located in Appendix A), the probability of getting killed increases to a maximum value with increasing traffic intensity. We investigated this for different velocities of amphibians, representative of the species in the study area, assuming perpendicular road crossing (Fig. 2) and for perpendicular road crossing in contrast to road crossing with all possible angles (Fig. 3). In order to investigate the significance of velocity only, α was set to 0.74 (mean of α) in both Fig. 2 and Fig. 3. Up to a traffic intensity of 625 vehicles/hour (15,000 vehicles/day), corresponding to a busy road, the velocity of the animals has a large influence on the probability of getting killed. Above this traffic intensity, the probability of getting killed during a road crossing is very close to 1 for all amphibian species investigated, whatever their velocity (within the range investigated).




Fig. 2. Probability of getting killed for one individual of different species on the road, as a function of its velocity

and traffic intensity, following the model: $P(\text{death}) = 1 - \frac{1}{\pi} \int_{\alpha=-\pi/2}^{\alpha=\pi/2} e^{\frac{-Na}{v \cos \alpha}} d\alpha$, and assuming

perpendicular road crossing ($\alpha = 0$). Velocities representative for different amphibian species are shown, together with velocities representative for hedgehog and hare. A traffic intensity of 3,200 vehicles per 24 hours corresponds to the road investigated.




Fig. 3. The effect on probability of getting killed by crossing the road randomly (*all angles*) compared to perpendicular road crossing (*perpendicular*) for three representative velocities of amphibians. 0.50 m/min corresponds to slow moving amphibians (mainly *Triturus* species), 1.50 m/min corresponds to *P. fuscus*, and 9.00 m/min is the velocity of the fastest moving *Rana temporaria* and *R. arvalis*.

We also calculated the probability of getting killed at different traffic intensities for velocities representative for hedgehog (*Erinaceus europaeus*, 45 m/min) and hare (*Lepus europaeus*, 120 m/min)(pers.obs). For these two species, *a* was set to 2.0m (total vehicle width), since, because of their size, they are killed by any part of a vehicle, not just the tyres. At these velocities, the probability of getting killed is far lower than for any of the amphibians considered (Fig. 2).

The angle of crossing clearly has an effect on the probability of getting killed (Fig. 3). The difference in probability of getting killed by perpendicular and random road crossing is most pronounced at medium probabilities of getting killed – for slow movers this corresponds to a traffic intensity of about 1,500 to 3,000 vehicles per 24 hours.

Diurnal variation in probability of death from crossing the road

In order to estimate diurnal variation in the probability of getting killed, species were grouped taxonomically, i.e., *Triturus* species, *Rana* species, *B. bufo*, *P. fuscus*. For any taxonomic group (k) at any time interval (j), the probability of getting killed ($P_{j,k}(\text{death})$) was calculated as the mean probability of getting killed for all individuals in that particular group. Assuming perpendicular road crossing, equation (9) (cf. Appendix) reduces to:

$$P_{j,k}(\text{death}) = \frac{\sum_{i=1}^{n(k)} \left(1 - e^{-\frac{aN_j}{v_{i(k)}}} \right)}{n(k)}, \quad (13)$$

where $n(k)$ is the total number of individuals in the taxonomic group in question (k), $v_{i(k)}$ denotes the velocity of one individual in group (k), and a is species specific. $P(\text{death})$ was then multiplied by the fraction of amphibians in this taxonomic group that are known from the recordings of diurnal activity pattern to be active in this particular time interval. Diurnal variation in the actual probability of getting killed was thus calculated, given the particular activity pattern and velocities of amphibians recorded, and the traffic intensity pattern. Diurnal variation in the probability of getting killed by a single road crossing not only reflects diurnal variation in traffic intensity (Fig. 4) but more particularly, the diurnal movement pattern of the species, and to a lesser extent the velocity of the species. The probability of getting killed is very small in the daytime with a small increase before dawn, reflecting the early rush-hour peak in traffic intensity late at night when the amphibians are still active. However, the probability of getting killed reaches a peak just after sunset, owing to the activity pattern of the amphibians (Fig. 1).




Fig. 4. Diurnal variation in probability of getting killed for the six amphibians species investigated. Probability of getting killed has been weighted by the movement pattern of the species.

Finally, the overall probability of getting killed by a single road crossing was found by summing up probabilities of getting killed for each of the 24 hour-long intervals. This probability was also extrapolated to other traffic intensities (Fig. 5). Again, the difference in levels of probability of getting killed reflects the velocity and the diurnal activity pattern of the species. The slow-moving salamanders face the highest probability of getting killed and the fast-moving *Rana* species face a somewhat lower risk. The diurnal activity pattern for *P. fuscus* is complementary to that of vehicles and lowers the probability of getting killed considerably.




Fig. 5. Probability of getting killed for one individual of different amphibian species on the road, as a function of traffic intensity. Probability of getting killed has been weighted by movement pattern of the amphibian species and diurnal variation in traffic intensity.

The efficiency of recording road deaths

The efficiency of monitoring road victims by foot was estimated with the expression derived in the Methods section (Table 2). The two *Rana* species were pooled since they are hard to distinguish as road kills. *Triturus vulgaris* and *T. cristatus* were pooled because of low numbers and taxonomic similarity.

Table 2

The efficiency of monitoring road victims by foot

The efficiency (E^*) with standard error is given as a fraction of road victims discovered out of the total (unknown) number (N).

R = number of victims recorded by routine monitorings only

C = number of victims recorded by control monitorings only

B = number of victims recorded by routine *and* control monitorings

The numbers are sums of 16 routine monitorings and 19 control monitorings, all by foot

F is the factor that converts, by multiplication, the number of victims found by routine monitorings ($R + B$) into the estimated total number of victims (\hat{N}).

	<i>Triturus vulgaris and Triturus cristatus</i>	<i>Pelobates fuscus</i>	<i>Bufo bufo</i>	<i>Rana temporaria and Rana arvalis</i>
R	5	2	4	29
C	14	2	7	73
B	1	4	8	34
Total = $R + C + B$	20	8	19	136
\hat{N}	90.0	9.0	22.5	198.3
E^*	0.067	0.667	0.533	0.318
SE	0.065	0.192	0.129	0.045
$F = 1/E^*$	15.00	1.50	1.88	3.15

Monitoring road victims by foot was surprisingly inefficient, ranging from about 7% to 67% of the road victims discovered (Table 2). The efficiency of foot monitoring was highest for *P. fuscus* and *B. bufo* – species that are believed to stay on the road for some time after getting killed due to their relatively tough skin and unpalatability, while only about one third of the brown frogs (*R. temporaria* and *R. arvalis*), and about seven percent of the salamanders were discovered by foot.

Impact of road deaths on populations

In 1996, anuran adult population sizes in the area were estimated to be 1,075 *Rana temporaria*, 3,309 *Rana arvalis*, and 265 *Pelobates fuscus*. In 1997, the figures were 425 *R. temporaria*, 1,680 *R. arvalis*, and 439 *P. fuscus*.

The estimated total number of adult amphibians killed was found by extrapolating the results from the period of road kill monitoring (April, May, and August) to the assumed whole active season (1 April – 15 October), assuming that the three months of monitoring are representative (Table 3). Finally, the estimated fraction of road-killed adults in 1996 and 1997 was calculated.

Table 3

The number of road victims and the estimated fraction of the adult populations killed on the road, found by foot-monitoring in 1996 (85 days) and 1997 (102 days), extrapolated to whole active seasons (1 April – 15 October).

The ranges of the total number killed are calculated from estimates of monitoring efficiency (i.e. $E^* \pm 2$ times SE (cf. Table 2). As E^* minus 2 times SE for *Triturus vulgaris* and *T. cristatus* is below zero, only minimum values of total number killed are given. Population estimates for *T. vulgaris* and *T. cristatus* and for *Bufo bufo* are incomplete, and therefore the fraction of adult populations killed is not estimated for these three species.

	<i>T. vulgaris and T. cristatus</i>	<i>P. fuscus</i>	<i>B. bufo</i>	<i>R. temporaria and R. arvalis</i>
1996				
Observed victims	17	8	20	91
Estimated victims ^a	86.29-	7.61-28.27	25.28-72.73	223.04-399.13
Total estimate for active season	201.02-	17.73-65.85	58.90-169.42	519.55-929.73
Fraction of adult population	-	0.07-0.25	-	0.11-0.21
1997				
Observed victims	11	13	14	30
Estimated victims ^a	55.88-	12.37-45.94	17.70-50.91	73.53-131.58
Total estimate for active season	108.47-	24.01-89.17	34.36-98.82	142.73-255.42
Fraction of adult population	-	0.05-0.20	-	0.07-0.12

Discussion

Probability of getting killed on the road

Formally expressing the probability for an animal to get killed by a single road crossing has been attempted before (Heine 1987). However, Heine's equation suffers from the logic shortcoming that high vehicle intensities and/or slow-moving animals result in negative values of survival probability – values that are then truncated at zero. van Langevelde and Jaarsma (1997) overcome this by turning the equation (very similar to that of Heine 1987), into an exponential expression, allowing the probability of getting killed by road crossing to approach 1 asymptotically for high traffic intensities and/or slow-moving animals. Their equation is very similar to ours, except that van Langevelde and Jaarsma consider the entire paved width to be effective in killing whereas we operate with a narrower killing width of vehicles (a). Since van Langevelde and Jaarsma mainly consider large animals, i.e., roe deer (*Capreolus capreolus*), foxes (*Vulpes vulpes*), and mustelids, their assumption is a

realistic approximation, but this does not apply to small animals that may remain still under a passing vehicle without getting hurt. It is also clear from our study that the actual distribution of crossing angles at a site has a large effect on the probability of getting killed; the effect is most pronounced for fast-moving animals, and for intermediate values of probabilities of getting killed. The actual distribution of crossing angles at a site is expected to vary considerably between sites, and in this work, we did not record crossing angles in a systematic way. Therefore, we mainly considered perpendicular road crossing, and consequently it needs to be stressed that our calculations of probabilities of getting killed by road crossing must be considered minimum values. Where amphibians have a fixed route to-and-from spawning sites, they may be undeterred by low-to-medium traffic intensity (i.e., below 12,000 vehicles per 24 hours). Mortality on this type of road may therefore be higher than predicted from traffic intensity alone. However, this is only speculative, and we recommend investigations on actual crossing angles.

The width (a) of vehicles that kills, was calculated as a weighted average of the vehicles on this particular road. The proportion of vehicles is likely to vary somewhat diurnally, seasonally, and with the day of the week. Because there is no existing data on this variation, we did not include it in our model. We did, however, tentatively increase and decrease the proportion of trucks by 5% and change the proportion of cars accordingly. The resulting probability of getting killed was increased and decreased by up to 5%, respectively. The change in probability was largest on roads with low traffic intensity, and for species with low overall probability of getting killed. Thus, it seems that in situations where traffic is intense, the results of the model are most reliable; for busy roads (15,000+ vehicles per 24 hours), with a 5% change in proportion of vehicles as described above, the change in model results for *T. vulgaris*, *T. cristatus*, *B. bufo*, *R. temporaria*, and *R. arvalis* was 1% or less.

Clearly, the three most important factors determining species vulnerability to road mortality are velocity of the species and diurnal movement pattern of the species and the vehicles. As a logical consequence of this, the most vulnerable species are day-active, slow-moving species. Velocity as an important factor has been stressed by several authors (e.g. Heine 1987; Rosen and Lowe 1994; Schlupp and Podlousky 1994; van Langevelde and Jaarsma 1997), but none considers the diurnal movement pattern of the animals and the vehicles (movement patterns are discussed by Rosen and Lowe (1994), but on a seasonal basis only).

The road investigated has a large diurnal variation in traffic intensity (high traffic intensity in the daytime dropping to almost zero at night), and this pattern reduces the vulnerability of nocturnal amphibians considerably. Other types of roads (e.g., motorways) may have diurnal variation in traffic intensity quite different from this one. In assessing the overall vulnerability of a species to traffic, the frequency of road crossing has to be included as well. *P. fuscus* may have a low frequency of road crossing whereas the badger (*Meles meles*), for example, is vulnerable to traffic mortality because of its frequent road crossings (Verboom, pers. com.).

Assessing the number of road kills

Despite earlier attempts to quantify road killed amphibians and reptiles, none of the authors have quantified the efficiency of their estimated number of road kills. Göransson et al. (1978) developed an expression to calculate the efficiency of foot monitoring. Their equation calibrates the efficiency of routine and control monitorings against all victims found. In contrast with our approach, Göransson et al. do not consider the unknown total number of victims killed on the road, i.e., the ones found plus the ones missed, and they thereby tend to overestimate monitoring efficiency by underestimating the total number of animals killed on the road.

Road-killed amphibians do not remain on the road for long; they are eaten by scavengers or are, especially in rainy weather, obliterated by being repeatedly run over by cars. They presumably have a shorter duration than mammals, for example, whose size and furry skin make them visible for a longer time. Our calculations confirm that many amphibians are missed even when patrolling the road by foot once every 24 hours: only 7% (*Triturus vulgaris* and *T. cristatus*) to 67% (*Pelobates fuscus*) of the total number of road victims are found. Patrolling by

car, which is done by most authors, is far less efficient. Thus, uncorrected road kill estimates are highly unreliable.

One basic assumption of estimating the size of a population using a Lincoln index is that the population is closed. This assumption is clearly violated in the case of estimating the total number of road-killed amphibians. On the road, new victims are continuously added to the "population," and others are removed by scavengers. Violations of this assumption tend to underestimate the victims found both at the routine and the control monitoring, and thereby overestimate the total number of amphibians killed. Another basic assumption is the one of independence in observing/missing a road victim during routine and control monitorings, respectively. There may be a slight bias in that very obvious roadkills are more likely to be recorded, but to all intents and purposes, the assumption is met.

Population size estimates

Our estimates of the population size of *P. fuscus* in the five fenced ponds are fairly accurate since they are based on marked individuals. The estimate of the population size of the two *Rana* species is not as accurate as the one for *P. fuscus*, partly because the brown frogs were not individually marked and partly because a larger fraction of the population bred in the ponds that were not fenced, compared to the *P. fuscus* population. The population estimates of *B. bufo* are very inaccurate, since the main part of the population bred in other ponds than the five fenced ones. They are therefore omitted from the following calculations together with *Triturus vulgaris* and *T. cristatus* and all data from 1995, where population sizes were estimated in the five fenced ponds only and thus do not form a basis of a total population estimate.

Our calculation of the fraction of adult amphibians killed on the road is based on the assumption that we have monitored the entire (meta)population affected by the highway. We chose 1,000m from the highway to be the upper limit of movement (in the sense that all populations within 1,000m from the highway are believed to be affected by its presence. In the literature, exact data on amphibian movement range are scarce). This assumption is in reasonable accordance with existing data, as *P. fuscus* is recorded to move a maximum of 1,200m between hibernation site and breeding pond (Nöllert 1990). Moreover, *P. fuscus* seems to be philopatric to its native pond with few adults changing breeding pond from year to year, which makes it reasonable to assume that we have monitored the entire metapopulation affected by road mortality. The two *Rana* species were pooled when estimating the fraction of adults killed on the road, partly because they are difficult to distinguish as road victims and their egg-clumps cannot be distinguished with certainty (Fog et al. 1997). Haapanen (1970) found that the maximum distance moved between years by *R. arvalis* and *R. temporaria* was 350 and 600m, respectively. Despite the accordance between our assumptions and existing data on amphibian movement range, it is clear that more data on the subject are needed until a firm fraction of amphibians killed can be established. Results should therefore be regarded as preliminary, although within the right range.

The impact of road kills on the populations

The road mortality estimates from this study cannot be extrapolated to other populations, but the equation, relating probability of getting killed for one individual on the road to the crossing angle, the velocity of the animal and the traffic intensity, is directly applicable to other populations, other geographical areas, and even to other species. The use of the equation is restricted, however, to species that do not behave intelligently towards traffic (e.g., stay on the side of the road until no vehicles are present). It is still useful to consider whether an annual mortality of up to 25% and 21% of the reproductively active adult population of *P. fuscus* and *R. temporaria/R. arvalis*, respectively, would have a significant effect on the population size (note that the annual mortality range is underestimated since it does not take into account the standard errors of the population estimates). That is, does road mortality constitute an additive or a compensatory mortality effect? For anuran adults in general, density independent mortality factors seem to be most important whereas for larvae, both density dependent and density independent mortality factors seem to be important (Duellman and Trueb 1994, and references therein). If the population in question is mainly regulated by density independent

mechanisms, such as climatic variability, road mortality will be an additive effect and is thus important as a population-regulating factor. If, however, the population is regulated by density dependent factors (mainly intraspecific competition among the larvae, cf. e.g. Wilbur 1972, 1977), road mortality will be compensated for by higher larval survival and the impact on the population will not be very large. This particular *P. fuscus* population is probably regulated in the larval state by density dependent mechanisms (Hels unpublished), and the road mortality is therefore expected to have no large regulating effect. If traffic intensity continues to increase, however, increased road mortality may eventually reduce the population to a level where its reproductive output is too small to reach the carrying capacities of the breeding ponds. This in turn may drive the population down to a level where demographic stochastic processes become important for the survival of the population. Finally, it should be noted that road mortality may be even more serious to the juveniles in the population since they are slow movers. This is a field where more investigations are needed, since very little is known about the movement ranges of juvenile amphibians.

Acknowledgements: We thank Niels F. Christiansen, The Danish Road Directorate, for supplying us with data on traffic intensity, Jesper Bruun-Schmidt for allowing us to use unpublished data, Gösta Nachman, Chris Topping, Jana Verboom and W. A. Scott for valuable comments on earlier versions of the manuscript, and Tove Petersen for improving the language. This study was supported by a grant from The Danish Road Directorate to TH.

Biographical Sketch: Dr. Tove Hels, an amphibian population ecologist, is a senior advisor and publications editor for the Danish Forest and Landscape Research Institute. She has published scientific and popular papers on the conflict between wildlife and traffic, and on amphibian ecology. Tove is also the main advisor on amphibians in the Danish Ministry of Environment. Her Ph.D. was earned from Copenhagen University.

References

- Andrews, A. 1990. Fragmentation of Habitat by Roads and Utility Corridors: A Review. *Australian Zoologist*, 26(3&4), 130-141.
- Anonymous 1998. *Trafikrapport 1996* (Traffic Report 1996, in Danish). The Road Directorate, Copenhagen, Denmark. 1-48.
- Begon, M. 1979. *Investigating Animal Abundance*. Edward Arnold Publishers Ltd., London.
- Bennett, A.F. 1990. *Habitat Corridors: Their Role in Wildlife Management and Conservation*, Department of Conservation and Environment, Melbourne, Australia. 1-37.
- Blaustein, A.R. and Wake, D.B. 1990. Declining Amphibian Populations: A Global Phenomenon? *Trends in Ecology and Evolution*, 5(7), 203-204.
- Blaustein, A.R., D.B. Wake, and W.P. Sousa 1994. Amphibian Declines: Judging Stability, Persistence, and Susceptibility of Populations to Local and Global Extinctions. *Conservation Biology*, 8(1), 60-71.
- Cooke, A.S. 1995. Road mortality of common toads (*Bufo bufo*) near a breeding site, 1974-1994. *Amphibia-Reptilia*, 16, 87-90.
- Drews, C. 1995. Road kills of animals by public traffic in Mikumi National Park, Tanzania, with notes on baboon mortality. *African Journal of Ecology*, 33, 89-100.
- Duellman, W.E. and L. Trueb 1994. *Biology of Amphibians*. The Johns Hopkins Press Ltd., London.
- Fahrig, L., J.H. Pedlar, S.E. Pope, P.D. Taylor, and J.F. Wegner 1995. Effect of Road Traffic on Amphibian Density. *Biological Conservation*, 73, 177-182.
- Fog, K., A. Schmedes, and D. Rosenørn de Lasson 1997. *Nordens padder og krybdyr* (Amphibians and Reptiles of the Nordic Countries, in Danish). GAD, Copenhagen.
- Fuellhaas, U., C. Klemp, A. Kordes, H. Ottersberg, M. Pirmann, A. Thiessen, C. Tschoetschel, and H. Zucchi 1989. Untersuchungen zum Strassentod von Vögeln, Säugetieren, Amphibien und Reptilien. *Beiträge zur Naturkunde Niedersachsens*, 42, 129-147.

- Groot Bruinderink, G.W.T.A. and E. Hazebroek 1996. Ungulate Traffic Collisions in Europe. *Conservation Biology*, 10(4), 1059-1067.
- Göransson, G., J. Karlson, and A. Lindgren 1978. *Vägers inverkan på omgivande natur II. Fauna* (Influence of roads on the surrounding nature II. Fauna, in Swedish). Statens Naturvårdsverk, Lund, Sweden. pp. 3-124.
- Haapanen, A. 1970. Site tenacity of the common frog (*Rana temporaria* L.) and the moor frog (*R. arvalis* Nilss.) *Ann. Zool. Fennici*, 7, 61-66.
- Hansen, L. 1982. Trafikdræbte dyr i Danmark (Road kills in Denmark, in Danish). *Dansk Ornitologisk Forenings Tidsskrift*, 76, 97-110.
- Heine, G. 1987. Einfache Mess- und Rechenmethode zur Ermittlung der Überlebenschance wandernder Amphibien beim Überqueren von Strassen. *Beih. Veröff. Naturschutz Landschaftspflege Bad.-Württ.*, 41, 473-479.
- Huijser, M.P. and P.J.M. Bergers 1997. The mortality rate in a hedgehog population: The relative importance of road kills. In *Habitat Fragmentation & Infrastructure*, Proceedings of the international conference on habitat fragmentation, infrastructure and the role of ecological engineering, September 1995, The Netherlands, ed. Canters, K. pp. 98-104. Published by Ministry of Transport, Public Works and Water Management, The Netherlands.
- Mader, H.-J. 1984. Animal Habitat Isolation by Roads and Agricultural Fields. *Biological Conservation*, 29, 81-96.
- Mader, H.-J., C. Schell, and P. Kornacker 1990. Linear Barriers to Arthropod Movements in the Landscape. *Biological Conservation*, 54, 209-222.
- Mallick, S.A., G.J. Hocking, and M.M. Driessen 1998. Road-kills of the eastern barred bandicoot (*Perameles gunnii*) in Tasmania: an index of abundance. *Wildlife Research*, 25, 139-145.
- Munguira, M.L. and J.A. Thomas 1992. Use of road verges by butterfly and burnet populations, and the effect of roads on adult dispersal and mortality. *Journal of Applied Ecology*, 29, 316-329.
- Nöllert, A. 1990. *Die Knoblauchkröte*, 2nd ed. A. Ziemsen Verlag, Wittenberg Lutherstadt.
- Reed, R.A., Johnson-Barnard, J., and W.L. Baker 1996. Contribution of Roads to Forest Fragmentation in the Rocky Mountains. *Conservation Biology*, 10(4), 1098-1106.
- Romin, L.A. and J.A. Bissonette 1996. Deer-vehicle collisions: status of state monitoring activities and mitigation efforts. *Wildlife Society Bulletin*, 24(2), 276-283.
- Rosen, P.C. and C.H. Lowe 1994. Highway Mortality of Snakes in the Sonoran Desert of Southern Arizona. *Biological Conservation*, 68, 143-148.
- Schlupp, I. and R. Podlucky 1994. Changes in Breeding Site Fidelity: A Combined Study of Conservation and Behaviour in the Common toad *Bufo bufo*. *Biological Conservation*, 69, 285-291.
- van Gelder, J.J. 1973. A Quantitative Approach to the Mortality Resulting from Traffic in a Population of *Bufo bufo* L. *Oecologia* (Berlin), 13, 93-95.
- van Langevelde, F. and Jaarsma, C.F. 1997. Habitat fragmentation, the role of minor rural roads and their traversability. In *Habitat Fragmentation & Infrastructure*, Proceedings of the international conference on habitat fragmentation, infrastructure and the role of ecological engineering, September 1995, The Netherlands, ed. Canters, K. pp. 171-182. Published by Ministry of Transport, Public Works and Water Management, The Netherlands.
- Vos, C.C. and J. P. Chardon 1998. Effects of habitat fragmentation and road density on the distribution pattern of the moor frog, *Rana arvalis*. *Journal of Applied Ecology*, 35, 44-56.

- Wilbur, H.M. 1972. Competition, predation, and the structure of the *Ambystoma-Rana sylvatica* community. *Ecology*, 53, 3-21.
- Wilbur, H.M. 1977. Density-dependent aspects of growth and metamorphosis in *Bufo americanus*. *Ecology*, 58, 196-200.
- Wyman, R.L. 1990. What's Happening to the Amphibians? *Conservation Biology*, 4(4), 350-352.

Appendix A

A FORMAL EXPRESSION FOR AN ANIMAL TO GET KILLED WHEN CROSSING THE ROAD

The probability of an amphibian surviving t time units on the road is denoted p . The change in p during one time step (dt) becomes:

$$dp = -\lambda p dt, \quad (6)$$

where λ is a positive constant depending on:

N : mean number of vehicles passing per time unit (the actual number is assumed to be Poisson distributed with mean N),

$\frac{a}{L}$

L : the probability of getting hit, where a is the width of vehicles that actually kills, and L = width of the road.

The change in p thus becomes:

$$dp = -\frac{Na}{L} p dt \quad (7)$$

Integration of (6) yields:

$$p = e^{-\lambda t}. \quad (8)$$

Since $t = \frac{s}{v}$, where s = distance moved to cross the road and v = velocity of the animal in question,

equation (8) can be rewritten as:

$$p = e^{-\frac{\lambda}{v}s} = e^{-\frac{Na}{Lv}s}. \quad (9)$$

s depends on the angle of crossing the road. Crossing angle is denoted α , $\alpha = 0$ corresponds to

perpendicular road crossing, which implies that the following condition must be satisfied for the animal to

cross the road: $-\frac{\pi}{2} < \alpha < \frac{\pi}{2}$. Consequently, the distance moved by crossing the road becomes $s = \frac{L}{\cos \alpha}$

(note that for perpendicular road crossing: $s = L$). Mean probability of surviving one road crossing ($P(surv)$), including all possible crossing angles, becomes:

$$P(surv) = \frac{1}{\pi} \int_{\alpha=-\pi/2}^{\alpha=\pi/2} e^{-\frac{Na}{v \cos \alpha}} d\alpha \quad (10)$$

and the mean probability of getting killed ($P(death)$) consequently becomes:

$$P(death) = 1 - \frac{1}{\pi} \int_{\alpha=-\pi/2}^{\alpha=\pi/2} e^{-\frac{Na}{v \cos \alpha}} d\alpha, \quad (11)$$

which can be extended to include variation in animal velocity by weighting Eq. (11) with the density function of velocities ($p(v)$), i.e.

$$P(death) = 1 - \frac{1}{\pi} \int_{v=0}^{\infty} p(v) \int_{\alpha=-\pi/2}^{\pi/2} e^{-\frac{Na}{v \cos \alpha}} d\alpha dv. \quad (12)$$