

**Table P-6a. Age-adjusted annualized rate (per 1,000 population) (with standard errors) of medically consulted injury and poisoning episodes, by external cause and selected characteristics: United States, 2014**

Selected characteristic	Fall <sup>1</sup>	Struck by a person or an object <sup>1</sup>	Transportation <sup>1</sup>	Overexertion <sup>1</sup>	Cutting or piercing instruments <sup>1</sup>	Other causes (injury) <sup>1</sup>	Poisoning <sup>1</sup>
Total	46.88 (2.68)	13.62 (1.44)	10.62 (1.19)	11.78 (1.25)	10.25 (1.19)	30.56 (2.21)	1.99 (0.55)
<b>Sex</b>							
Male	42.45 (3.53)	18.51 (2.54)	10.67 (1.61)	11.20 (1.77)	12.87 (1.92)	34.59 (3.22)	*1.55 (0.72)
Female	50.95 (4.09)	9.07 (1.51)	10.64 (1.69)	12.21 (1.69)	7.77 (1.55)	26.55 (2.90)	*2.46 (0.85)
<b>Age</b>							
Under 12 years	46.84 (6.20)	15.04 (3.42)	*3.54 (1.63)	*	*3.48 (1.50)	25.16 (5.10)	*
12-17 years	69.03 (15.99)	26.38 (6.44)	21.28 (5.97)	*18.36 (6.10)	*6.89 (2.83)	53.16 (9.86)	*
18-44 years	26.11 (3.02)	10.20 (2.30)	12.75 (2.08)	13.72 (2.11)	15.20 (2.57)	28.95 (3.46)	*1.79 (0.88)
45-64 years	45.14 (5.06)	12.04 (2.45)	9.79 (2.27)	13.37 (2.78)	9.03 (2.17)	31.31 (4.30)	*
65-74 years	71.14 (11.22)	*10.02 (3.83)	*7.83 (3.22)	*7.72 (3.55)	*11.92 (4.11)	24.05 (6.85)	*
75 years and over	130.85 (17.47)	*23.47 (8.12)	*	*13.45 (5.37)	*	28.64 (8.26)	-
<b>Race</b>							
One race <sup>2</sup>	47.40 (2.74)	13.19 (1.45)	10.48 (1.20)	11.49 (1.26)	10.17 (1.20)	30.53 (2.25)	2.04 (0.57)
White	50.57 (3.24)	14.05 (1.70)	9.93 (1.33)	12.28 (1.48)	10.84 (1.44)	30.63 (2.52)	*2.32 (0.69)
Black or African American	32.20 (5.42)	*10.64 (3.20)	14.81 (3.59)	*7.98 (2.74)	*8.94 (3.00)	30.45 (5.64)	*
American Indian or Alaska Native	*54.33 (20.99)	*	*	*	-	78.62 (20.80)	*
Asian	30.29 (8.24)	*	*	*10.22 (4.24)	*8.14 (3.52)	*20.40 (9.31)	-
Native Hawaiian or Other Pacific Islander	*	-	-	-	-	-	-
Two or more races <sup>3</sup>	*33.32 (13.36)	*24.28 (9.45)	*	*28.21 (13.40)	*	*28.93 (11.78)	-
Black or African American, white	*	*	*	*	-	*	-
American Indian or Alaska Native, white	*	*	*	*	*	*	-
<b>Hispanic or Latino origin<sup>4</sup> and race</b>							
Hispanic or Latino	32.50 (5.33)	5.99 (1.63)	7.66 (1.92)	5.62 (1.58)	*5.15 (1.75)	14.78 (3.61)	*
Mexican or Mexican American	30.67 (7.85)	8.75 (2.58)	*5.50 (1.85)	*3.61 (1.53)	*4.30 (2.11)	13.42 (3.46)	*
Not Hispanic or Latino	50.57 (3.15)	15.26 (1.73)	11.38 (1.41)	13.00 (1.48)	11.36 (1.43)	34.07 (2.61)	2.45 (0.69)
White, single race	56.36 (4.00)	15.92 (2.15)	10.70 (1.63)	13.95 (1.83)	12.79 (1.83)	34.92 (3.15)	*2.96 (0.91)
Black or African American, single race	32.79 (5.65)	*11.30 (3.40)	15.26 (3.80)	*8.52 (2.93)	*9.39 (3.15)	31.14 (5.91)	*
<b>Education<sup>5</sup></b>							
Less than a high school diploma	58.51 (10.18)	*	*11.12 (3.57)	*6.70 (3.20)	*12.29 (4.07)	*7.00 (2.36)	-
High school diploma or GED <sup>6</sup>	44.16 (5.88)	9.84 (2.55)	9.84 (2.85)	10.78 (2.54)	13.20 (3.66)	28.05 (4.63)	*4.96 (2.34)
Some college	55.66 (6.80)	14.40 (3.61)	9.73 (2.80)	14.11 (3.05)	16.09 (3.65)	38.60 (5.94)	*
Bachelor's degree or higher	40.20 (5.50)	16.20 (3.85)	10.50 (2.59)	17.89 (3.24)	8.74 (2.48)	36.73 (5.43)	*
<b>Family income<sup>7</sup></b>							
Less than \$35,000	57.61 (5.16)	13.60 (2.60)	11.87 (2.48)	11.78 (2.37)	5.96 (1.66)	34.26 (4.37)	*5.21 (1.78)
\$35,000 or more	43.10 (3.42)	14.24 (2.06)	9.82 (1.63)	13.66 (1.88)	11.53 (1.67)	30.89 (2.98)	*1.07 (0.51)
\$35,000-\$49,999	42.47 (6.79)	*10.33 (3.27)	16.75 (4.58)	15.27 (4.18)	12.99 (3.85)	41.69 (8.23)	*
\$50,000-\$74,999	34.63 (6.11)	*11.50 (3.81)	*10.45 (3.52)	15.94 (4.15)	*10.97 (3.41)	24.49 (4.25)	*
\$75,000-\$99,999	49.01 (8.81)	*14.39 (4.57)	*	*9.70 (3.16)	13.07 (3.90)	25.57 (6.39)	-
\$100,000 or more	46.71 (6.38)	17.55 (3.92)	9.05 (2.41)	13.90 (3.45)	10.47 (2.83)	33.59 (5.68)	*
<b>Poverty status<sup>8</sup></b>							
Poor	52.57 (7.43)	15.07 (3.76)	14.19 (3.71)	*11.97 (3.72)	*3.89 (1.55)	32.02 (5.56)	*
Near poor	57.85 (6.63)	10.66 (2.39)	9.69 (2.50)	11.66 (2.80)	13.12 (3.18)	33.56 (5.33)	*
Not poor	44.21 (3.43)	14.93 (2.14)	11.00 (1.68)	13.63 (1.89)	9.66 (1.50)	30.62 (2.99)	*1.73 (0.72)

**Table P-6a. Age-adjusted annualized rate (per 1,000 population) (with standard errors) of medically consulted injury and poisoning episodes, by external cause and selected characteristics: United States, 2014**

Selected characteristic	Fall <sup>1</sup>	Struck by a person or an object <sup>1</sup>	Transportation <sup>1</sup>	Overexertion <sup>1</sup>	Cutting or piercing instruments <sup>1</sup>	Other causes (injury) <sup>1</sup>	Poisoning <sup>1</sup>
Health insurance coverage <sup>9</sup>							
Under 65 years:							
Private	40.36 (3.97)	13.48 (2.02)	9.26 (1.57)	14.94 (1.96)	10.68 (1.72)	30.87 (3.20)	*1.78 (0.75)
Medicaid	63.47 (8.43)	12.15 (3.18)	18.08 (4.05)	*6.26 (2.27)	*12.49 (4.07)	34.20 (5.62)	*
Other	58.41 (16.91)	*42.67 (21.22)	*	*	*	31.23 (8.83)	-
Uninsured	21.39 (4.71)	*16.24 (7.41)	11.71 (3.06)	*6.68 (2.44)	*12.88 (4.72)	*31.25 (11.33)	*
65 years and over:							
Private	93.35 (15.45)	*12.03 (4.80)	*	*8.59 (3.21)	*6.34 (3.12)	23.18 (6.40)	*
Medicare and Medicaid	215.98 (58.02)	*	*	*	*	*	-
Medicare only	91.02 (16.46)	*20.29 (8.11)	*8.23 (4.09)	*	*12.11 (5.23)	*33.66 (11.32)	*
Other	*92.13 (33.19)	*	*	*	*	*	*
Uninsured	*	-	-	-	-	-	-
Place of residence <sup>10</sup>							
Large MSA	41.83 (3.82)	12.46 (2.00)	8.24 (1.43)	8.83 (1.50)	9.12 (1.59)	29.28 (3.03)	*2.11 (0.82)
Small MSA	47.54 (4.22)	18.23 (2.73)	13.11 (2.31)	18.21 (2.73)	10.89 (2.25)	29.98 (3.73)	*1.95 (0.86)
Not in MSA	62.27 (7.04)	7.40 (2.02)	15.15 (3.99)	*9.03 (2.84)	12.73 (3.10)	36.15 (6.02)	*
Region							
Northeast	43.30 (6.31)	*11.12 (3.76)	11.62 (3.01)	10.30 (2.84)	12.19 (3.43)	30.80 (5.63)	*
Midwest	56.19 (7.21)	14.01 (2.68)	12.35 (2.93)	12.15 (2.86)	10.92 (2.78)	35.46 (5.29)	*4.98 (1.93)
South	47.60 (3.96)	14.52 (2.55)	10.59 (1.93)	10.67 (1.95)	9.98 (1.79)	26.86 (3.42)	*
West	38.78 (4.58)	13.55 (2.72)	8.18 (1.90)	14.33 (2.75)	8.76 (2.15)	32.30 (4.29)	*

\* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) greater than 30% and less than or equal to 50% and should be used with caution. Data not shown have an RSE greater than 50%.

- Quantity zero.

<sup>1</sup>Based on the questions, "Of the [number] times that [person] was injured, how many of those times was the injury serious enough that a medical professional was consulted?" and "Of the [number] times that [person] was poisoned, how many of those times was the poisoning serious enough that a medical professional was consulted?" Injury and poisoning episodes are classified according to the external cause of the episode using categories based on the International Classification of Diseases, 9th revision, Clinical Modification (ICD-9-CM) external cause codes (E codes). The category "Transportation" includes motor vehicle, bicycle, motorcycle, pedestrian, train, boat, and airplane. The category "Other" includes fire/burn/scald-related, animal or insect bite, machinery, other (not specified) and unknown causes. The category "Poisoning" excludes food poisoning and allergic reaction. Rates of episodes have been annualized in this table. Estimates from this report should not be compared with estimates from 2003 or earlier.

<sup>2</sup>Refers to persons who indicated only a single race group, including those of Hispanic or Latino origin.

<sup>3</sup>Refers to persons who indicated more than one race group, including those of Hispanic or Latino origin. Only two combinations of multiple race groups are shown due to small sample sizes for other combinations.

<sup>4</sup>Refers to persons who are of Hispanic or Latino origin and may be of any race or combination of races. "Not Hispanic or Latino" refers to persons who are not of Hispanic or Latino origin, regardless of race.

<sup>5</sup>Shown only for persons aged 25 and over. Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using four age groups: 25-44, 45-64, 65-74, and 75 and over.

<sup>6</sup>GED is General Educational Development high school equivalency diploma.

<sup>7</sup>Includes persons who reported a dollar amount or who would not provide a dollar amount but provided an income interval.

<sup>8</sup>"Poor" persons are defined as having income below the poverty threshold. "Near poor" persons have incomes of 100% to less than 200% of the poverty threshold. "Not poor" persons have incomes that are 200% of the poverty threshold or greater.

<sup>9</sup>Based on a hierarchy of mutually exclusive categories. Persons with more than one type of health insurance were assigned to the first appropriate category in the hierarchy. "Uninsured" includes persons who had no coverage as well as those who had only Indian Health Service coverage or had only a private plan that paid for one type of service such as accidents or dental care.

<sup>10</sup>MSA is metropolitan statistical area. Large MSAs have a population of 1 million or more; small MSAs have a population of less than 1 million. "Not in MSA" consists of persons not living in a metropolitan statistical area.

NOTES: Estimates are based on household interviews of a sample of the civilian noninstitutionalized population. This table is based on responses about all persons in the family. Data came from the Person file and the Injury and Poisoning Episode file and were weighted using the Person weight. Unknowns for the columns were excluded from the denominator when calculating rates. Includes other races not shown separately and persons with unknown education, family income, poverty status, and health insurance characteristics. Unless otherwise specified, estimates are age adjusted using the projected 2000 U.S. population as the standard population and using six age groups: 0-11, 12-17, 18-44, 45-64, 65-74, and 75 and over. Estimates for age groups are not age adjusted. For more information on the data source, methods, and definitions used for this table, refer to Technical Notes for Summary Health Statistics Tables: National Health Interview Survey (available from <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>). Suggested citation: Adams PF, Benson V. Tables of Summary Health Statistics for the U.S. Population: 2014 National Health Interview Survey, 2015. Available from: <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, 2014.

**Table P-6b. Annualized frequency (number in thousands) of medically consulted injury and poisoning episodes, by external cause and selected characteristics: United States, 2014**

Selected characteristic	Struck by a person or an object <sup>1</sup>				Cutting or piercing instruments <sup>1</sup>	Other causes (injury) <sup>1</sup>	Poisoning <sup>1</sup>
	Fall <sup>1</sup>	Transportation <sup>1</sup>	Overexertion <sup>1</sup>				
Total	14,963	4,228	3,293	3,699	3,191	9,546	629
Sex							
Male	6,542	2,733	1,624	1,698	2,003	5,244	*247
Female	8,421	1,495	1,669	2,001	1,189	4,301	*382
Age							
Under 12 years	2,269	729	*172	*	*168	1,219	*
12-17 years	1,723	658	531	*458	*172	1,327	*
18-44 years	2,928	1,144	1,429	1,538	1,705	3,247	*201
45-64 years	3,728	994	809	1,104	746	2,586	*
65-74 years	1,865	*263	*205	*202	*312	630	*
75 years and over	2,450	*439	*	*252	*	536	-
Race							
One race <sup>2</sup>	14,784	4,009	3,177	3,535	3,108	9,324	629
White	12,821	3,412	2,420	3,026	2,615	7,520	573
Black or African American	1,280	452	593	*328	*347	1,184	*
American Indian or Alaska Native	*165	*	*	*	-	*297	*
Asian	478	*	*	*179	*146	*323	-
Native Hawaiian or Other Pacific Islander	*	-	-	-	-	-	-
Two or more races <sup>3</sup>	*179	*219	*116	*164	*	*222	-
Black or African American, white	*	*	*	*	-	*	-
American Indian or Alaska Native, white	*	*	*	*	*	*	-
Hispanic or Latino origin <sup>4</sup> and race							
Hispanic or Latino	1,551	336	416	317	*279	839	*
Mexican or Mexican American	785	304	*185	*128	*154	483	*
Not Hispanic or Latino	13,412	3,892	2,878	3,382	2,912	8,706	627
White, single race	11,461	3,080	2,064	2,745	2,424	6,775	571
Black or African American, single race	1,228	452	574	*328	*347	1,141	*
Education <sup>5</sup>							
Less than a high school diploma	1,760	*	*291	*194	*351	*190	-
High school diploma or GED <sup>6</sup>	2,697	608	548	575	680	1,508	*256
Some college	3,144	820	534	796	894	2,240	*
Bachelor's degree or higher	2,490	947	733	1,161	565	2,368	*
Family income <sup>7</sup>							
Less than \$35,000	5,294	1,147	967	1,008	502	2,828	*438
\$35,000 or more	7,461	2,530	1,762	2,492	2,074	5,431	*191
\$35,000-\$49,999	1,447	*347	581	539	439	1,370	*
\$50,000-\$74,999	1,559	*488	*416	702	*508	1,078	*
\$75,000-\$99,999	1,442	*470	*	*319	*436	825	-
\$100,000 or more	3,014	1,225	665	933	692	2,158	*
Poverty status <sup>8</sup>							
Poor	2,108	706	578	*474	*189	1,357	*
Near poor	3,434	618	535	658	747	1,909	*
Not poor	7,957	2,638	1,987	2,496	1,877	5,484	*297

**Table P-6b. Annualized frequency (number in thousands) of medically consulted injury and poisoning episodes, by external cause and selected characteristics: United States, 2014**

Selected characteristic	Fall <sup>1</sup>	Struck by a person or an object <sup>1</sup>	Transportation <sup>1</sup>	Overexertion <sup>1</sup>	Cutting or piercing instruments <sup>1</sup>	Other causes (injury) <sup>1</sup>	Poisoning <sup>1</sup>
Health insurance coverage <sup>9</sup>							
Under 65 years:							
Private	6,530	2,214	1,577	2,578	1,836	5,217	*272
Medicaid	2,671	643	771	*283	489	1,785	*
Other	710	*306	*	*	*	*490	-
Uninsured	738	*363	535	*308	437	861	*
65 years and over:							
Private	1,933	*266	*	*182	*	494	*
Medicare and Medicaid	647	*	*	*	*	*	-
Medicare only	1,391	*305	*	*	*203	*541	*
Other	*323	*	*	*	*	*	*
Uninsured	*	-	-	-	-	-	-
Place of residence <sup>10</sup>							
Large MSA	7,009	2,103	1,429	1,499	1,543	5,012	*354
Small MSA	4,873	1,776	1,236	1,769	1,071	2,947	*205
Not in MSA	3,080	349	628	*432	577	1,587	*
Region							
Northeast	2,453	*550	627	581	605	1,593	*
Midwest	4,046	1,030	862	876	775	2,459	*346
South	5,649	1,681	1,231	1,217	1,200	3,131	*
West	2,814	968	573	1,025	611	2,362	*164

\* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) greater than 30% and less than or equal to 50% and should be used with caution. Data not shown have an RSE greater than 50%.

- Quantity zero.

<sup>1</sup>Based on the questions, "Of the [number] times that [person] was injured, how many of those times was the injury serious enough that a medical professional was consulted?" and "Of the [number] times that [person] was poisoned, how many of those times was the poisoning serious enough that a medical professional was consulted?" Injury and poisoning episodes are classified according to the external cause of the episode using categories based on the International Classification of Diseases, 9th revision, Clinical Modification (ICD-9-CM) external cause codes (E codes). The category "Transportation" includes motor vehicle, bicycle, motorcycle, pedestrian, train, boat, and airplane. The category "Other" includes fire/burn/scald-related, animal or insect bite, machinery, other (not specified) and unknown causes. The category "Poisoning" excludes food poisoning and allergic reaction. Counts of episodes have been annualized in this table. Estimates from this report should not be compared with estimates from 2003 or earlier.

<sup>2</sup>Refers to persons who indicated only a single race group, including those of Hispanic or Latino origin.

<sup>3</sup>Refers to persons who indicated more than one race group, including those of Hispanic or Latino origin. Only two combinations of multiple race groups are shown due to small sample sizes for other combinations.

<sup>4</sup>Refers to persons who are of Hispanic or Latino origin and may be of any race or combination of races. "Not Hispanic or Latino" refers to persons who are not of Hispanic or Latino origin, regardless of race.

<sup>5</sup>Shown only for persons aged 25 and over.

<sup>6</sup>GED is General Educational Development high school equivalency diploma.

<sup>7</sup>Includes persons who reported a dollar amount or who would not provide a dollar amount but provided an income interval.

<sup>8</sup>"Poor" persons are defined as having income below the poverty threshold. "Near poor" persons have incomes of 100% to less than 200% of the poverty threshold. "Not poor" persons have incomes that are 200% of the poverty threshold or greater.

<sup>9</sup>Based on a hierarchy of mutually exclusive categories. Persons with more than one type of health insurance were assigned to the first appropriate category in the hierarchy. "Uninsured" includes persons who had no coverage as well as those who had only Indian Health Service coverage or had only a private plan that paid for one type of service such as accidents or dental care.

<sup>10</sup>MSA is metropolitan statistical area. Large MSAs have a population of 1 million or more; small MSAs have a population of less than 1 million. "Not in MSA" consists of persons not living in a metropolitan statistical area.

NOTES: Estimates are based on household interviews of a sample of the civilian noninstitutionalized population. This table is based on responses about all persons in the family. Data came from the Person file and the Injury and Poisoning Episode file and were weighted using the Person weight. Unknowns for the columns were excluded from the frequencies. Includes other races not shown separately and persons with unknown education, family income, poverty status, and health insurance characteristics. For more information on the data source, methods, and definitions used for this table, refer to Technical Notes for Summary Health Statistics Tables: National Health Interview Survey (available from <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>). Suggested citation: Adams PF, Benson V. Tables of Summary Health Statistics for the U.S. Population: 2014 National Health Interview Survey, 2015. Available from: <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, 2014.

**Table P-6c. Crude annualized rate (per 1,000 population) (with standard errors) of medically consulted injury and poisoning episodes, by external cause and selected characteristics: United States, 2014**

Selected characteristic	Fall <sup>1</sup>	Struck by a person or an object <sup>1</sup>	Transportation <sup>1</sup>	Overexertion <sup>1</sup>	Cutting or piercing instruments <sup>1</sup>	Other causes (injury) <sup>1</sup>	Poisoning <sup>1</sup>
Total	47.79 (2.73)	13.50 (1.40)	10.52 (1.18)	11.82 (1.25)	10.19 (1.16)	30.49 (2.19)	2.01 (0.55)
Sex							
Male	42.75 (3.60)	17.86 (2.40)	10.62 (1.57)	11.10 (1.75)	13.09 (1.93)	34.27 (3.18)	*1.61 (0.73)
Female	52.61 (4.05)	9.34 (1.54)	10.43 (1.65)	12.50 (1.70)	7.43 (1.46)	26.87 (2.91)	*2.39 (0.82)
Age							
Under 12 years	46.84 (6.20)	15.04 (3.42)	*3.54 (1.63)	*	*3.48 (1.50)	25.16 (5.10)	*
12-17 years	69.03 (15.99)	26.38 (6.44)	21.28 (5.97)	*18.36 (6.10)	*6.89 (2.83)	53.16 (9.86)	*
18-44 years	26.11 (3.02)	10.20 (2.30)	12.75 (2.08)	13.72 (2.11)	15.20 (2.57)	28.95 (3.46)	*1.79 (0.88)
45-64 years	45.14 (5.06)	12.04 (2.45)	9.79 (2.27)	13.37 (2.78)	9.03 (2.17)	31.31 (4.30)	*
65-74 years	71.14 (11.22)	*10.02 (3.83)	*7.83 (3.22)	*7.72 (3.55)	*11.92 (4.11)	24.05 (6.85)	*
75 years and over	130.85 (17.47)	*23.47 (8.12)	*	*13.45 (5.37)	*	28.64 (8.26)	-
Race							
One race <sup>2</sup>	48.31 (2.77)	13.10 (1.41)	10.38 (1.19)	11.55 (1.26)	10.16 (1.18)	30.47 (2.22)	2.06 (0.56)
White	52.38 (3.30)	13.94 (1.65)	9.89 (1.32)	12.36 (1.48)	10.68 (1.37)	30.72 (2.51)	2.34 (0.68)
Black or African American	31.73 (5.28)	11.21 (3.34)	14.69 (3.56)	*8.12 (2.75)	*8.61 (2.88)	29.34 (5.40)	*
American Indian or Alaska Native	*55.15 (21.37)	*	*	*	-	*99.20 (32.03)	*
Asian	27.40 (7.69)	*	*	*10.27 (4.30)	*8.37 (3.60)	*18.49 (8.09)	-
Native Hawaiian or Other Pacific Islander	*	-	-	-	-	-	-
Two or more races <sup>3</sup>	*25.33 (8.59)	*30.99 (11.47)	*16.45 (7.96)	*23.27 (10.57)	*	*31.45 (11.95)	-
Black or African American, white	*	*	*	*	-	*	-
American Indian or Alaska Native, white	*	*	*	*	*	*	-
Hispanic or Latino origin <sup>4</sup> and race							
Hispanic or Latino	28.48 (4.24)	6.16 (1.69)	7.63 (1.90)	5.82 (1.65)	*5.13 (1.72)	15.41 (3.39)	*
Mexican or Mexican American	22.54 (4.93)	8.72 (2.50)	*5.32 (1.80)	*3.66 (1.54)	*4.42 (2.13)	13.86 (3.27)	*
Not Hispanic or Latino	51.86 (3.17)	15.05 (1.66)	11.13 (1.38)	13.08 (1.46)	11.26 (1.37)	33.66 (2.53)	2.43 (0.67)
White, single race	58.50 (3.95)	15.72 (2.02)	10.54 (1.58)	14.01 (1.78)	12.37 (1.68)	34.58 (3.01)	2.92 (0.86)
Black or African American, single race	32.36 (5.52)	11.91 (3.55)	15.13 (3.76)	*8.63 (2.91)	*9.15 (3.06)	30.07 (5.68)	*
Education <sup>5</sup>							
Less than a high school diploma	64.79 (10.73)	*	*10.72 (3.42)	*7.13 (3.33)	*12.93 (4.17)	*6.99 (2.49)	-
High school diploma or GED <sup>6</sup>	49.32 (6.61)	11.12 (2.97)	10.02 (2.89)	10.52 (2.36)	12.44 (3.31)	27.58 (4.61)	*4.68 (2.12)
Some college	54.48 (6.60)	14.20 (3.40)	9.26 (2.64)	13.80 (2.98)	15.49 (3.46)	38.81 (5.86)	*
Bachelor's degree or higher	37.73 (5.01)	14.35 (3.27)	11.11 (2.74)	17.59 (3.13)	8.57 (2.41)	35.89 (5.23)	*
Family income <sup>7</sup>							
Less than \$35,000	61.77 (5.53)	13.38 (2.53)	11.28 (2.33)	11.76 (2.32)	5.86 (1.61)	33.00 (4.17)	*5.11 (1.71)
\$35,000 or more	41.01 (3.21)	13.90 (1.93)	9.69 (1.56)	13.70 (1.87)	11.40 (1.64)	29.85 (2.83)	*1.05 (0.49)
\$35,000-\$49,999	42.70 (6.83)	*10.25 (3.26)	17.14 (4.68)	15.91 (4.34)	12.94 (3.82)	40.44 (7.97)	*
\$50,000-\$74,999	35.28 (6.34)	*11.04 (3.49)	*9.41 (3.03)	15.88 (4.07)	*11.50 (3.55)	24.40 (4.18)	*
\$75,000-\$99,999	42.70 (7.13)	*13.92 (4.40)	*	*9.45 (3.03)	*12.91 (3.91)	24.43 (6.06)	-
\$100,000 or more	42.99 (5.45)	17.47 (3.70)	9.49 (2.57)	13.30 (3.05)	9.86 (2.56)	30.78 (4.62)	*
Poverty status <sup>8</sup>							
Poor	47.10 (6.14)	15.78 (3.91)	12.91 (3.31)	*10.59 (3.28)	*4.22 (1.62)	30.31 (5.07)	*
Near poor	59.99 (6.82)	10.79 (2.40)	9.34 (2.41)	11.50 (2.79)	13.05 (3.12)	33.34 (5.20)	*
Not poor	43.36 (3.34)	14.38 (1.99)	10.83 (1.63)	13.60 (1.85)	10.23 (1.56)	29.88 (2.84)	*1.62 (0.65)

**Table P-6c. Crude annualized rate (per 1,000 population) (with standard errors) of medically consulted injury and poisoning episodes, by external cause and selected characteristics: United States, 2014**

Selected characteristic	Fall <sup>1</sup>	Struck by a person or an object <sup>1</sup>	Transportation <sup>1</sup>	Overexertion <sup>1</sup>	Cutting or piercing instruments <sup>1</sup>	Other causes (injury) <sup>1</sup>	Poisoning <sup>1</sup>
Health insurance coverage <sup>9</sup>							
Under 65 years:							
Private	38.70 (3.64)	13.12 (1.91)	9.35 (1.55)	15.28 (2.00)	10.88 (1.73)	30.91 (3.11)	*1.61 (0.66)
Medicaid	52.88 (6.24)	12.72 (2.90)	15.26 (3.13)	*5.60 (2.06)	9.67 (2.81)	35.35 (5.22)	*
Other	67.50 (17.14)	*29.05 (13.25)	*	*	*	46.58 (13.69)	-
Uninsured	20.91 (4.39)	*10.28 (3.64)	15.18 (3.96)	*8.74 (3.26)	12.38 (3.66)	24.39 (6.80)	*
65 years and over:							
Private	87.80 (14.60)	*12.09 (4.82)	*	*8.26 (3.08)	*	22.43 (6.07)	*
Medicare and Medicaid	219.98 (59.70)	*	*	*	*	*	-
Medicare only	87.68 (15.76)	*19.21 (7.52)	*	*	*12.80 (5.55)	*34.13 (11.56)	*
Other	*92.70 (31.83)	*	*	*	*	*	*
Uninsured	*	-	-	-	-	-	-
Place of residence <sup>10</sup>							
Large MSA	41.15 (3.70)	12.34 (1.95)	8.39 (1.45)	8.80 (1.46)	9.06 (1.56)	29.42 (3.03)	*2.08 (0.81)
Small MSA	49.94 (4.58)	18.20 (2.73)	12.67 (2.22)	18.13 (2.75)	10.97 (2.26)	30.20 (3.72)	*2.10 (0.92)
Not in MSA	68.17 (7.82)	7.73 (2.27)	13.90 (3.67)	*9.55 (3.00)	12.78 (2.89)	35.12 (5.78)	*
Region							
Northeast	46.25 (6.84)	*10.36 (3.40)	11.82 (3.14)	10.96 (2.72)	11.41 (3.05)	30.03 (5.29)	*
Midwest	57.22 (7.23)	14.56 (2.81)	12.20 (2.86)	12.38 (2.88)	10.96 (2.77)	34.78 (5.17)	*4.90 (1.92)
South	47.99 (3.99)	14.28 (2.46)	10.46 (1.90)	10.34 (1.93)	10.20 (1.80)	26.60 (3.37)	*
West	39.29 (4.68)	13.51 (2.69)	8.00 (1.84)	14.32 (2.75)	8.53 (2.07)	32.99 (4.43)	*2.29 (1.14)

\* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) greater than 30% and less than or equal to 50% and should be used with caution. Data not shown have an RSE greater than 50%.

- Quantity zero.

<sup>1</sup>Based on the questions, "Of the [number] times that [person] was injured, how many of those times was the injury serious enough that a medical professional was consulted?" and "Of the [number] times that [person] was poisoned, how many of those times was the poisoning serious enough that a medical professional was consulted?" Injury and poisoning episodes are classified according to the external cause of the episode using categories based on the International Classification of Diseases, 9th revision, Clinical Modification (ICD-9-CM) external cause codes (E codes). The category "Transportation" includes motor vehicle, bicycle, motorcycle, pedestrian, train, boat, and airplane. The category "Other" includes fire/burn/scald-related, animal or insect bite, machinery, other (not specified) and unknown causes. The category "Poisoning" excludes food poisoning and allergic reaction. Rates of episodes have been annualized in this table. Estimates from this report should not be compared with estimates from 2003 or earlier.

<sup>2</sup>Refers to persons who indicated only a single race group, including those of Hispanic or Latino origin.

<sup>3</sup>Refers to persons who indicated more than one race group, including those of Hispanic or Latino origin. Only two combinations of multiple race groups are shown due to small sample sizes for other combinations.

<sup>4</sup>Refers to persons who are of Hispanic or Latino origin and may be of any race or combination of races. "Not Hispanic or Latino" refers to persons who are not of Hispanic or Latino origin, regardless of race.

<sup>5</sup>Shown only for persons aged 25 and over.

<sup>6</sup>GED is General Educational Development high school equivalency diploma.

<sup>7</sup>Includes persons who reported a dollar amount or who would not provide a dollar amount but provided an income interval.

<sup>8</sup>"Poor" persons are defined as having income below the poverty threshold. "Near poor" persons have incomes of 100% to less than 200% of the poverty threshold. "Not poor" persons have incomes that are 200% of the poverty threshold or greater.

<sup>9</sup>Based on a hierarchy of mutually exclusive categories. Persons with more than one type of health insurance were assigned to the first appropriate category in the hierarchy. "Uninsured" includes persons who had no coverage as well as those who had only Indian Health Service coverage or had only a private plan that paid for one type of service such as accidents or dental care.

<sup>10</sup>MSA is metropolitan statistical area. Large MSAs have a population of 1 million or more; small MSAs have a population of less than 1 million. "Not in MSA" consists of persons not living in a metropolitan statistical area.

NOTES: Estimates are based on household interviews of a sample of the civilian noninstitutionalized population. This table is based on responses about all persons in the family. Data came from the Person file and the Injury and Poisoning Episode file and were weighted using the Person weight. Unknowns for the columns were excluded from the denominator when calculating rates. Includes other races not shown separately and persons with unknown education, family income, poverty status, and health insurance characteristics. For more information on the data source, methods, and definitions used for this table, refer to Technical Notes for Summary Health Statistics Tables: National Health Interview Survey (available from <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>). Suggested citation: Adams PF, Benson V. Tables of Summary Health Statistics for the U.S. Population: 2014 National Health Interview Survey, 2015. Available from: <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey, 2014.