

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Temporal variability of soil management effects on soil hydrological properties, runoff and erosion at the field scale in a hillslope vineyard, North-West Italy**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1619505> since 2018-02-28T12:40:14Z

*Published version:*

DOI:10.1016/j.still.2016.07.017

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Manuscript Number: STILL-15-700R3

Title: Temporal variability of soil management effects on soil hydrological properties, runoff and erosion at the field scale in a hillslope vineyard, North-West Italy

Article Type: Research paper

Section/Category: Coupled Soil & Tillage Processes

Keywords: Soil management; vineyards; soil hydrological properties

Corresponding Author: Dr. Marcella Biddoccu, M.Sc.Eng.

Corresponding Author's Institution: Institute for Agricultural and Earthmoving Machines (IMAMOTER), National Research Council of Italy (CNR)

First Author: Marcella Biddoccu, M.Sc.Eng.

Order of Authors: Marcella Biddoccu, M.Sc.Eng.; Stefano Ferraris; Andrea Pitacco; Eugenio Cavallo, PhD

Abstract: Soil management in vineyard inter-rows has a great influence on soil hydraulic conductivity and bulk density, and, consequently, on runoff and soil erosion processes at the field scale. The maintenance of bare soil in vineyard inter-rows with tillage, as well as the tractor traffic, are known to expose the soil to compaction, reduction of soil water holding capacity and increase of runoff and erosion formation. The use of grass cover is one of the most common and effective practices in order to reduce such threats. It is therefore important to relate rainfall characteristics, soil properties and response in terms of runoff and soil erosion, from yearly to seasonal and to single event temporal scales.

The objective of this work is to quantify the temporal variability of the effects of two different kind of inter-row management on soil hydrological properties, runoff and erosion in vineyards. For this reason two vineyard field-scale plots in the Alto Monferrato vine-growing area (Piedmont, NW Italy) were monitored in two years. The inter-rows were managed with conventional tillage (CT) and grass cover (GC), respectively. Fifteen series of infiltration tests were carried out during a 2-year period of observation (October 2012 to November 2014). In order to take into account the effect of tractors traffic, the tests were done on the track, and outside the track. Furthermore, a dataset of 29 rainfall-runoff events covering a wide range of topsoil characteristics was collected in the two plots, along with soil water content and runoff discharge monitoring, and determination of sediment yield in case of erosive events. An optical disdrometer installed in the plots provided also 1-min rainfall intensity data. In summer, just one month after tillage, CT soil showed very low hydraulic conductivity, so storms were able to cause Hortonian runoff and soil losses up to 5.7 Mg ha<sup>-1</sup>. In autumn and winter very high saturation-excess runoff was observed in CT, that reached 83% of the precipitation. Runoff in the grass cover plot was mainly due to saturation of the topsoil, and the annual reduction of

runoff in the GC plot was about 63%. Soil erosion up to 1.2 Mg ha<sup>-1</sup> in a single event was observed in the GC vineyard in winter. In each year of observation, most of the erosion occurred during a single event, while the total annual erosion was up to 9 times higher in the CT treatment than in the GC.

*Consiglio Nazionale delle Ricerche*

**IMAMOTER**

Istituto per le Macchine Agricole e Movimento Terra

Marcella Biddoccu  
Institute for Agricultural and Earthmoving Machines  
Italian National Research Council  
Strada delle Cacce, 73  
10135 Torino, Italy  
m.biddoccu@ima.to.cnr.it  
+39 011 39 77 723

June 23<sup>th</sup>, 2016

Dear editorial board of Soil and Tillage Research,

Please find enclosed the new revised version of the manuscript entitled "Temporal and soil management effects on hydrological properties, runoff and erosion at field scale in a hillslope vineyard, North-West Italy", to be submitted as a research paper to Soil and Tillage Research for consideration of publication. We followed indications of editor in preparing this new version of the manuscript. All co-authors have seen and agree with the contents of the manuscript. We certify that the submission is original work and is not under review at any other publication.

In this manuscript, we reported the results of a research evaluating the temporal and soil management effects on soil hydrological properties in two vineyard field-scale plots (Piedmont, North-West Italy), which inter-rows were managed with grass cover and conventional tilled, respectively. Furthermore, the study was addressed to identify correlations between rainfall characteristics, soil properties and field-scale response in terms of runoff and soil erosion, at event temporal scale. During a 2-years period of observation, several series of infiltration tests were carried out, and a dataset of 29

rainfall-runoff events covering a wide range of topsoil characteristics was collected in the two plots, along with soil water content monitoring, measurements and sampling of runoff in order to determine the sediment yield.

The results highlighted the positive effect of grass cover in favoring water infiltration, reducing runoff and soil erosion throughout the year, compared with the conventional tillage management, with greatest effectiveness in summer. The annual reduction was greater than 63% and up to 90%, respectively for runoff and soil erosion. Only saturation-excess runoff was observed in the grassed vineyard. The highest runoff rates and soil losses were measured in the vineyard managed with conventional tillage even some weeks after the execution of tillage. In summer and early autumn, the tilled soil showed the lowest hydraulic conductivity, so summer storm were able to generate hortonian runoff and high soil losses, up to 5.7 Mg ha<sup>-1</sup> soil erosion for a single event in the period of observation. In autumn and winter, the wettest seasons, despite the autumn tillage, very high saturation-excess runoff was observed, so the winter runoff reached 83% of the precipitation.

We believe that our research could be of interest to the readers of Soil and Tillage Research and we hope that the editorial board will agree on the interest of this paper.

Thank you for your consideration.

Sincerely yours,

Marcella Biddoccu

on behalf of the authors.

Corresponding author: Marcella Biddoccu, Institute for Agricultural and Earthmoving Machines (IMAMOTER), National Research Council, 10135 Torino, Strada delle Cacce, 73, Italy

m.biddoccu@ima.to.cnr.it, Ph: +390113977723

June, 23<sup>th</sup>, 2016

Dear editor,

The authors thanks for your comments and suggestion to further improve the manuscript. We reviewed the manuscript taking in account them.

The conclusion section was revised and shortened to be more

Apart from the editorial comments, which were solved, the answers to the specific comments are described below.

Finally, the text was revised in English by a native speaker.

Reviewers' comments:

Line 4 should read: "... are known to expose ...".

L. 6 should read: "practices".

L. 9: spelling of "scales".

L. 9: Somewhere here, the objective of this study should be provided before you tell what was done in the study.

**Answer: the objective is now provided in L10-11**

L. 16: spelling of "in".

L. 17 should read: "... just one month after tillage ...".

L. 18: spelling of "Hortonian" as this is based on a name.

L. 27 should read: "... one of the land uses ...".

L. 34-40: This sentence is grammatically incomplete.

L. 47 should read: "... and other land uses ...".

L. 58 should read: "... at yearly or multi-annual scales ...".

L. 59: Spelling of "Gómez".

L. 63 should read: "... runoff and infiltration at the field-scale ...".

L. 71 should read: "... objectives ...".

L. 92 should read: "The soil has been managed ...".

L. 96 should read: "... on the soil surface."

L. 100: spelling of "Glyphosate".

L. 121 should read: "... recorded at 10-min intervals ...".

L 123 should read: "... data have been obtained from ...".

L. 132 should read: "To obtain the sediment yield from each erosive event ...".

L. 135-136: How were the TM sensors calibrated?

**Answer: The TM sensors were gravimetrically calibrated (L136)**

L. 138 should read: "... in the 2-year period of observations, ...".

L. 140 should read: "... on the same date ...".

L. 149 should read: "... that slope does not affect the measurements significantly."

L. 161 should read: "Rainfall events occurring after August 2013 for which precipitation was recorded at 10-min intervals, were ...".

L. 166 should read: "... were computed ...".

L. 170-171: "Soil characteristics (Kfs, SWCs, BD) for NT and T positions, were associated at each event."

I do not understand this statement. What do you mean by "associated". Do not erase this statement, clarify it.

**Answer: Each rainfall event was analyzed considering its characteristics and the "soil properties (Kfs, SWCs, BD) measured in T and NT position in the closest date (L172-173).**

L. 174: spelling of "Hortonian".

L. 178-179: "the lower mean values between Kfs(T) and Kfs(NT) and between SWCs(T) and SWCs(NT) were chosen."

This statement is not clear. What do you mean by lower mean values, and were chosen for what?

**Answer: The paragraph was modified, to clarify this point (L178-188). Each rainfall event was characterized by values of Kfs and SWCs measured in T and NT. The lower value between Kfs(T) and Kfs(NT) was chosen as reference Kfs to compare the time series of rainfall intensity during the event. The lower between SWCs(T) and SWCs(NT) was chosen as reference SWCs to compare the SWC time series, to complete the event analysis in order to detect runoff generation and the type of runoff.**

L. 181 should read: "reached" and "remained".

L. 197: I can't make sense of "both in T than in NT position". Do you mean "both in T and in NT position"?

**Answer: it was corrected throughout the text**

L. 200 should read: "... for both treatments."

L. 202 should read: "... some days after tillage."

L. 212 should be "cumulative" instead of "cumulated".

L. 216 should read: "The 40% and 35% of annual precipitation were recorded ...".

L. 218 should read: "... in both plots."

L. 244 should read: "... 36 mm of rain fell with the highest ...".

L. 246 should read: "5.6" (decimal point).

L. 252-253 should read: "... with a higher determination coefficient ...".

L. 255 should read: "... at 7-day intervals ...">

L. 264 should read: "... are displayed in Figs. 3 and 4."

L. 270-271: Similar as above, I can't make sense of "both from CT than from GC". Please clarify.

L. 271 should read: "... with a maximum 10-min rainfall intensity...".

L. 275 should read: "... did not overcome ...".

L. 282 should read: "... thus runoff caused by infiltration excess ...".

L. 284 should read: "... shows a rainfall occurring in autumn ...".

L. 284 should read: "... the 10-minute rainfall intensity ...".

L. 288 should read: "... near the 140 ...".

L. 289-290: I could not grasp the sense of this sentence, but did you intend to say "Erosion detected in the two plots was nearly 40 times higher in CT than in GC."?

**Answer: L295-296 "Erosion was detected in the two plots, and in CT it was nearly 40 times higher than in GC"**

L. 294-205 should read: "Runoff appeared in the first hours of the event ...".

L. 296 should read: "Within a few hours ...".

L. 297: Replace "first" with "upper".

L. 304: Wghat do you mean by "this kind of saturation runoff events"? Clarify.

L. 307 "Hortonian".

L. 308-309 should read: "... and rainfall depth from 14 to 36 mm."

L. 311: "Hortonian".

L. 312 should read: "Runoff occurred for 50% of the observations ...".

L. 317: "Hortonian".

L. 319 should read: "whereas 50% of the rainfall events produced Hortonian ...".



L. 320 should read: "... were examined in the following."

L. 323 "Hortonian".

L. 324: "... runoff in CT, but no runoff in GC."

L. 324: "Hortonian".

L. 325 should read: "... with a depth ranging from ...".

L. 326 should read: "... in both plots."

L. 329: "Hortonian".

L. 331: "Hortonian".

L. 332: "Hortonian".

L. 335 should read: "... in GC ...".

L. 338 should read: "With a few exceptions ...".

L. 338-339 should read: "... so soil water saturation was reached sooner than in the T position."

L. 344 should read: "vineyard."

L. 346-347 should read: "The coefficients of variation ...".

L. 348 should read: "During most of the sampling dates ...".

L. 349: Replace "exclusion" with "exception".

L. 357 should read: "during summer and autumn, a finding that was particularly evident ...".

L. 364 should read: "In 2014, the CT topsoil showed higher sand content ...".

L. 369-370 should read: "... could also be related to these differences ...".

L. 371: There is no "role" on something. I suggest to say: "Impact of soil management, soil properties and rainfall on runoff".

**Answer: the section title was changed (L381)**

L. 372 should read: "... was usually 2 to 3.6 times higher ...".

L. 374: should read: "... the highest runoff was observed during ...".

L. 375: Do you mean "... both in CT and in GC."?

L. 375-376: "The highest differences in runoff between CT and GC occurred in ...".

L. 376 should read: "... when the grass cover was higher."

L. 377 should read: "... and runoff coefficients were observed in winter ...".

L. 378 should read: "... when snowfall was followed by rainfall."

L. 378-379 should read: "... whereas it was 28 % in GC."

L. 383: replace "have" with "had".

L. 385: I could not understand this statement. Instead of "which was originated" do you mean "that was generated"?

**Answer: it was corrected (L395)**

L. 387 should read: "... despite Kfs showing the highest mean values ...".

L. 397: Do you mean "... at 7-day intervals."?

L. 400: This statement seems to be erroneous: "when the some precipitation was recorded in the previous 7 days". Do you mean "when the same precipitation" or "when some precipitation"?

**Answer: it was corrected (L411) "when some precipitation was recorded in the previous 7 days"**

L. 403 should read "rainfall-runoff event analysis".

L. 404 should read: "... runoff was caused by saturation of the topsoil ...".

L. 407 should read: "... of a structural crust ...".

L. 408: "cumulative" instead of "cumulated".

L. 409 should read: "factors".

L. 416: "Hortonian".

L. 421: Replace "Role" with "Impact".

L. 422 should read: "Sediment yields of ...".

L. 423 should read: "... was close to 7.4 ...".

L. 424: Replace "yearly" with "annual".

L. 435 should read: "... were due to the variability of rainfall ...".

L. 441: Replace "where" with "when".

L. 446 should read: "These results ...".

L. 446: Spelling of "Gómez".

L. 449: I was not sure what you meant with "and then with 7-days antecedent precipitation". Did you mean "... and is also related with the 7-day antecedent precipitation"?

**Answer: it was corrected in "7-day antecedent precipitation" all over in the text**

L. 458 should read "... was observed."

L. 462 should read: "... over a 5-minute period."

L. 464-466: This sentence is grammatically incomplete.

L. 469 should read: "... to those of Gómez et al. ...".

L. 482: "Hortonian".

L. 483 should read: "Due to high compaction after grape harvest, the worst conditions for infiltration were found before tillage in CT and also in winter for GC."

L. 489: Replace "in case of" with "during".

## Highlights

- Soil moisture, runoff, soil erosion were monitored in vineyard field-scale plots
- Temporal and soil management effects on soil hydrological properties were evaluated
- Summer storms caused hortonian runoff and high soil losses just a month after tillage
- Highest runoff rates were observed in late autumn and winter in tilled vineyard
- High runoff was due to soil saturation in the wet seasons

**Title page**

Title:

Temporal variability of soil management effects on soil hydrological properties, runoff and erosion at the field scale in a hillslope vineyard, North-West Italy

Authors and affiliations:

Marcella Biddoccu<sup>1,2\*</sup>, Stefano Ferraris<sup>2,4</sup>, Andrea Pitacco<sup>3</sup>, Eugenio Cavallo<sup>1</sup>

<sup>1</sup>Institute for Agricultural and Earthmoving Machines (IMAMOTER), National Research Council, 10135 Torino, Strada delle Cacce, 73, Italy

<sup>2</sup>Interuniversity Department of Regional and Urban Studies and Planning (DIST), Politecnico di Torino and Università di Torino, 10125 Torino, Castello del Valentino, Viale Mattioli, 39, Italy

<sup>3</sup>Università degli Studi di Padova, Department of Agronomy, Food, Natural Resources, Animals and Environment (DAFNAE), Padova, Italy

<sup>4</sup> Istituto di Geoscienze e Georisorse, CNR, Pisa, Italy

\*Corresponding author: [m.biddoccu@ima.to.cnr.it](mailto:m.biddoccu@ima.to.cnr.it), Ph: +390113977723

1 **Abstract**

2 Soil management in vineyard inter-rows has a great influence on soil hydraulic conductivity and  
3 bulk density, and, consequently, on runoff and soil erosion processes at the field scale. The  
4 maintenance of bare soil in vineyard inter-rows with tillage, as well as the tractor traffic, are known  
5 to expose the soil to compaction, reduction of soil water holding capacity and increase of runoff and  
6 erosion formation. The use of grass cover is one of the most common and effective practices in  
7 order to reduce such threats. It is therefore important to relate rainfall characteristics, soil properties  
8 and response in terms of runoff and soil erosion, from yearly to seasonal and to single event  
9 temporal scales.

10 The objective of this work is to quantify the temporal variability of the effects of two different kind  
11 of inter-row management on soil hydrological properties, runoff and erosion in vineyards. For this  
12 reason two vineyard field-scale plots in the Alto Monferrato vine-growing area (Piedmont, NW  
13 Italy) were monitored in two years. The inter-rows were managed with conventional tillage (CT)  
14 and grass cover (GC), respectively. Fifteen series of infiltration tests were carried out during a 2-  
15 year period of observation (October 2012 to November 2014). In order to take into account the  
16 effect of tractors traffic, the tests were done on the track, and outside the track. Furthermore, a  
17 dataset of 29 rainfall-runoff events covering a wide range of topsoil characteristics was collected in  
18 the two plots, along with soil water content and runoff discharge monitoring, and determination of  
19 sediment yield in case of erosive events. An optical disdrometer installed in the plots provided also  
20 1-min rainfall intensity data. In summer, just one month after tillage, CT soil showed very low  
21 hydraulic conductivity, so storms were able to cause Hortonian runoff and soil losses up to 5.7 Mg  
22 ha<sup>-1</sup>. In autumn and winter very high saturation-excess runoff was observed in CT, that reached  
23 83% of the precipitation. Runoff in the grass cover plot was mainly due to saturation of the topsoil,  
24 and the annual reduction of runoff in the GC plot was about 63%. Soil erosion up to 1.2 Mg ha<sup>-1</sup> in  
25 a single event was observed in the GC vineyard in winter. In each year of observation, most of the

26 erosion occurred during a single event, while the total annual erosion was up to 9 times higher in  
27 the CT treatment than in the GC.

## 28 **1. Introduction**

29 Grapevine cultivation represents one of the land uses for which higher runoff rates and sediment  
30 losses are observed in Europe, especially in the Mediterranean area (Tropeano, 1983; Kosmas et al.,  
31 1997; Cerdà and Doerr, 2007; García-Ruiz, 2010; García-Ruiz et al., 2015). Analysis of data  
32 collected throughout Europe showed that in the Mediterranean region runoff higher than 9% of  
33 annual precipitation (Maetens et al., 2012) and the highest erosion rates ( $17.4 \text{ Mg ha}^{-1} \text{ year}^{-1}$ ) are  
34 related to vineyard land use (Cerdan et al., 2010).

35 Some typical features of the vine-growing system, such as location on hillslopes and disposition of  
36 rows along the slope, make runoff and erosion stronger (Corti et al., 2011). Furthermore, some  
37 practices usually adopted in vineyards' installation (land levelling works and deep tillage) and  
38 vineyards' management (maintenance of bare soil by mechanical or chemical weeding, intense  
39 tractor traffic along fixed paths) are favoring runoff, erosion and further threats as compaction,  
40 nutrient losses and reduction of soil water holding capacity (Tropeano, 1984; Ramos and Martínez-  
41 Casasnovas, 2004; Ferrero et al., 2005; Ramos and Martínez-Casasnovas, 2007; Arnáez et al.,  
42 2007). The effects of the inter-rows soil management on runoff and soil erosion in vineyards of  
43 southern Europe was evaluated in several studies under natural rainfall, at different spatial scales  
44 (from plot to catchment) and from event to multi-year temporal scales (Tropeano, 1983; Kosmas et  
45 al., 1997; Arnaez et al., 2007; Brenot et al., 2008; Casalí et a., 2008; Raclot et al., 2009; Ruiz-  
46 Colmenero et al., 2011; Novara et al., 2011; Corti et al., 2011; Biddoccu et al., 2016). The use of  
47 grass cover in the inter-rows is one of the most common and effective soil management practices  
48 adopted in order to reduce runoff and soil erosion in vineyards (Blavet et al., 2009; Novara et al.,  
49 2011; Ruiz-Colmenero et al., 2011; Prosdocimi et al., 2016) and other land uses which are  
50 especially subjected to erosion as olive groves (Gómez et al., 2009). Under the indication of the

51 CAP agro-environmental requirements, some Rural Development Programmes (i.e., Regione  
52 Piemonte, NW-Italy) introduced during the period 2007-2013 specific subsidies to encourage the  
53 adoption of grass cover in vineyards and orchards in order to protect soil from degradation.  
54 However, tillage is still used in vineyards growing on low-permeability soils as a practice to remove  
55 grass in summer and improve water infiltration, particularly during autumn and winter time. In fact,  
56 growers are often worried that competition for soil resources, namely water and nutrients, between  
57 the grass cover and grapevines could affect grape yield and quality.

58 Most studies on runoff and erosion in vineyards consider topographic features, soil properties,  
59 rainfall characteristics, and soil management techniques in relation to the hydrological and erosive  
60 response of the vineyard at yearly or multi-annual scales (Prosdocimi et al., 2016). Nevertheless,  
61 annual runoff and soil losses could be strongly conditioned by few rainfall events (Gómez et al.,  
62 2014; González-Hidalgo et al., 2009). The adopted soil management influences strongly the  
63 temporal and spatial variations of the soil surface characteristics (soil cover, topsoil structure and  
64 soil crusting) and soil hydrological characteristics, which drive the partition of rainfall between  
65 runoff and infiltration at the field-scale (Leonard and Andrieux, 1998; Pare et al., 2011). There is  
66 still a gap in knowledge about the effect of the temporal variations of topsoil conditions on the  
67 triggering of runoff and soil water erosion throughout the year. A better understanding of the field  
68 response to rainfall events, taking into account the variability of the soil conditions during the year,  
69 could be useful for water balance and erosion modelling purposes (Celette et al., 2010) and to  
70 support soil management decisions in vineyards, in order to reduce runoff and erosion.

71 This study presents the results of a 2-year experiment monitoring topsoil hydrological properties  
72 and recording runoff and soil erosion in two vineyard field-scale plots with different inter-row soil  
73 management, conventional tillage and grass cover, respectively. The objectives were: (i) to evaluate  
74 the effects of soil management, at different temporal scales, namely at yearly, seasonal and single  
75 event ones; (ii) to identify in each event the prevalent runoff mechanism (either infiltration or



76 saturation excess) in relation to soil management, soil hydraulic conductivity and bulk density, soil  
77 moisture and precipitation characteristics.

## 78 **2. Materials and Methods**

### 79 **2.1 Study site**

80 The study was carried out within the “Tenuta Cannona Experimental Vine and Wine Centre of  
81 Regione Piemonte” (44°40’ N, 8°37’ E, 296 m asl), which is located in the Alto Monferrato hilly  
82 area of Piemonte, North-West Italy . The climate is sublitoranean, (average annual precipitation of  
83 965 mm at the Ovada station, in the period 1951-1990), mainly concentrated in October, November  
84 and March. The driest month was July. The mean annual temperature measured at Alessandria  
85 during the same period of observation was 12.6°C (Biancotti et al. 1998). At the study site, the  
86 average annual precipitation in the period 2000-2014 was 905 mm and the mean annual air  
87 temperature was 14.5°C. The Cannona vineyards lie on Pleistocenic fluvial terraces in the Tertiary  
88 Piedmont Basin, including highly altered gravel, sand and silty-clay deposits, with red alteration  
89 products. The soils derived from reworked Pleistocene alluvium, and they have a clay to clay-loam  
90 texture.

91 The experiment was conducted in two vineyard plots, which are part of a larger vineyard, lying on a  
92 hillslope with SE aspect and average 15% slope. Each plot is 1221 m<sup>2</sup> (74 m long and 16.5 m wide)  
93 and includes 7 vine rows aligned along the slope, where the vines are spaced 1.0 m along the row  
94 and 2.75 m between the rows. The soil has been managed with different techniques since 2000. The  
95 first plot has been managed with conventional tillage (CT, cultivated with chisel to a depth of about  
96 0.25 m), while in the second plot grass cover has been adopted (GC, with spontaneous grass  
97 controlled with mulcher during the year). The mulcher mows and chips the grass, and residues are  
98 left on the soil surface. Soil tillage (in CT) and grass mulching (in GC) were usually carried out  
99 twice a year, in spring and autumn. In autumn 2011, the inter-rows of the GC plot were tilled and a  
100 grass mixture was sown, to renew the grass cover. The grass mixture was composed of: *Lolium*

101 *perenne* 20%, *Festuca rubra* 60%, *Poa nemoralis* 15%, *Poa trivialis* 5%. Weeds under the rows of  
102 the two plots were controlled with Glyphosate in spring, 0.6 m across the vine row. Most of the  
103 farming operations in the vineyard were carried out using tracked or tyred tractors, with  
104 intensification from spring to the grape harvest time. During the period of the present study, soil  
105 tillage (in CT) and grass mulching (in GC) were carried out five times (on: 24/10/2012, 05/06/2013,  
106 11/11/2013, 16/05/2014, 24/10/2014). The soil is classified as *Typic Ustorthents, fine-loamy, mixed,*  
107 *calcareous, mesic* (Soil Survey Staff, 2010) or *Dystric Cambisols* (FAO/ISRIC/ISSS, 1998). Soil  
108 textural composition obtained from soil samples taken in 2014, at 0-10 cm depth, indicated a silty  
109 clay loam soil in the GC plot, with 15% sand, 53% silt and 32% clay content; and a silt loam soil in  
110 the CT plot, with 28% sand, 54% silt and 18% clay content.

## 111 **2.2 Measurements**

112 The experiment was conducted from October, 2012 to November, 2014. A monitoring system  
113 provided continuous measurements of rainfall, runoff and topsoil water content for the two  
114 experimental plots. Runoff samples were also collected to obtain sediment yield for erosive events.  
115 Periodic measurements were carried out to obtain values of saturated hydraulic conductivity ( $K_{fs}$ ),  
116 bulk density (BD) and initial soil water content ( $SWC_i$ ) in the two plots, in order to detect the  
117 temporal variability of the field-saturated soil hydraulic conductivity at the surface of the vineyard  
118 inter-rows, with different conditions depending on soil management. Measurements were carried  
119 out both in the no-track (indicated as NT) and in the track position (indicated as T), which is the  
120 portion of soil affected by the passage of tractor wheels or tracks.

### 121 ***Rainfall, runoff, erosion and soil water content***

122 Rainfall measurements were obtained from an automatic rainfall gauge, with 0.2 mm resolution, at  
123 about 200 m from the plots. Rainfall data were recorded at 10-min intervals since August, 2013,  
124 whereas only hourly measurements were available for the previous period. Since June, 2014, 1-min

125 rainfall intensity data have been obtained from an optical disdrometer installed near the plots (Laser  
126 Precipitation Monitor, Thies-Clima, Germany).

127 Each plot was hydraulically bounded: a channel at the top of the plots collected upstream water.  
128 Runoff and sediments were collected by a channel, connected to a sedimentation trap and then to a  
129 tipping bucket device to measure the discharge of runoff from each plot. A portion of the runoff-  
130 sediment mixture was sampled for each tip. The tipping bucket devices were calibrated to measure  
131 runoff with 0.1 mm resolution. In addition, hourly measurements of the runoff volumes were  
132 obtained from electro-magnetic counters. After each erosive event, a 1.5 L sample of runoff-  
133 sediment mixture was collected. Sediments deposited along drains and in the sedimentation traps  
134 were also collected and dry-weighed. To obtain the sediment yield from each erosive event,  
135 sediment concentration was multiplied by the runoff volume and added to the weight of deposited  
136 sediments. Four soil moisture 5 TM sensors (Decagon Devices) were gravimetrically calibrated and  
137 installed at 10 cm depth in each plot in NT and T positions. Soil water content measurements were  
138 recorded every 60 minutes.

### 139 *Infiltration tests*

140 Several series of infiltration tests were carried out in the 2-year period of observations, using the  
141 simplified falling head technique (SFH), proposed by Bagarello et al. (2004). Eight series of tests  
142 were done in the CT plot and seven series in the GC plot. The tests were conducted on the same  
143 date in the two plots, except from November to December 2012, when they were carried out with a  
144 delay of three weeks in GC. At each measurement date, four to eight SFH experiments were  
145 performed, with 2-4 measurements carried out in the no-track position of the inter-row and 2-4 in  
146 the track position. To assure one-dimensional flow, a second ring was inserted concentric to the  
147 inner one. The two PVC cylinders had a height of 0.30 m, and inner diameters of 0.305 m and 0.486  
148 m. They were inserted in the soil to a minimum depth of 0.06 m. The applied volumes of water  
149 were 7.0 L in the inner ring and 10.8 L in the bigger cylinder. We kept a minimum height of 0.06 m

150 of water on the sloping soil surface. Bodhinayake et al. (2004) have demonstrated that slope does  
151 not affect the measurements significantly. Before the execution of each test an undisturbed soil core  
152 ( $V= 100 \text{ cm}^3$ ) was collected next to the investigated area at depth of 0 to 0.07 m, to determine the  
153 soil bulk density. For the same purpose a sample was collected after the water infiltration inside the  
154 inner ring. Initial and saturated volumetric water contents ( $\text{SWC}_i$  and  $\text{SWC}_s$ ) were also obtained  
155 from the collected soil samples.

156 Each BD,  $\text{SWC}_i$ ,  $K_{fs}$ , dataset was summarized by calculating the mean and the associated  
157 coefficient of variation (CV), in order to compare the data obtained by the infiltration experiments.  
158 The statistical frequency distributions of the data were assumed to be normal for the initial soil  
159 water content and the soil bulk density and log-normal for the field-saturated hydraulic conductivity  
160 (Warrick, 1998). Differences between positions (NT vs T) in the same plot, and differences between  
161 the two plots , were evaluated by using *t*-test at 0.05 probability level.

### 162 **2.3 Rainfall-runoff events analysis**

163 Rainfall events occurring after August 2013 for which precipitation was recorded at 10-min  
164 intervals, were analysed in order to evaluate the relationships among rainfall and soil hydrological  
165 characteristics and the runoff and erosion processes. For each event, rainfall amount and duration,  
166 maximum rainfall intensity at different time intervals (10, 30, and 60 minutes), and cumulative  
167 precipitation (during the previous 7, 15, 30, and 45 days) were obtained. Rainfall energy (Brown  
168 and Foster, 1987) and rainfall event erosivity (Renard et al., 1997) were computed, by means of  
169 RIST (Rainfall Intensity Summarization Tool) (ARS-USDA, 2015). Rainfall events were  
170 considered as significant when cumulative rainfall was larger than 12.7 mm, according to the  
171 RUSLE procedure. Only one smaller rainfall event (on 14/08/2013) was analysed, because of its  
172 high intensity ( $21.6 \text{ mm h}^{-1}$  in 10 min). A total of 29 rainfall events were selected, each one with its  
173 own measured values of the following soil properties:  $K_{fs}$ ,  $\text{SWC}_s$ , BD, for T and NT positions.

174 Firstly, Principal Component Analysis (PCA) was performed. Afterwards, stepwise multiple linear  
175 regression was applied. Finally, each of the significant rainfall events was analyzed in order to  
176 identify the surface runoff occurrence and its type (Hortonian or saturation of the soil surface  
177 horizon).

178 For this last purpose, the method of Dehotin et al. (2015) was applied, by comparing time series of  
179 rainfall intensities with  $K_{fs}$  values, and topsoil SWC with  $SWC_s$  values, respectively in order to  
180 detect either Hortonian or saturation excess runoff type. The measured values of  $K_{fs}$  and  $SWC_s$  in  
181 the CT and GC plots in the period of occurrence of each rainfall event were used as reference  
182 values. The lower mean values between  $K_{fs}(T)$  and  $K_{fs}(NT)$ , and between  $SWC_s(T)$  and  $SWC_s(NT)$ ,  
183 respectively, were chosen as reference  $K_{fs}$  and  $SWC_s$  for the event. Hortonian runoff was detected if  
184 rainfall intensity values were higher than the  $K_{fs}$  reference value. The runoff occurrence due to  
185 saturation of the soil surface horizon was detected by comparing soil water content time series with  
186 the  $SWC_s$  reference for each event. It was assumed that if the soil water content time series reached  
187 the value of  $SWC_s$  and remained almost constant, the first horizon was saturated and additional  
188 rainfall was generating surface runoff.

## 189 **Results**

### 190 **3.1. Soil hydrological properties**

191 Table 1 summarizes the infiltration tests, which were conducted with initial soil water content  
192 ranging between 0.158 and 0.357  $\text{cm}^3\text{cm}^{-3}$  in CT, and between 0.191 and 0.405  $\text{cm}^3\text{cm}^{-3}$  in GC. On  
193 most dates,  $CV(SWC_i)$  was lower than 10% in CT. Higher variations were obtained in GC,  
194 especially in the NT position. Soil water content was usually higher in T than in NT in the two  
195 plots, with significant differences only in October, 2012, in CT and in December, 2012, and in July,  
196 2013, in GC. For measurements carried out on the same date, soil water content was always higher  
197 in GC than in CT.

198 At the time of the execution of the infiltration tests, the bulk density varied between 1.19 and 1.55 g  
199  $\text{cm}^{-3}$  in CT and between 1.24 and 1.47  $\text{g cm}^{-3}$  in GC. The coefficient of variation of bulk density  
200 was generally  $< 10\%$ , with some exceptions in GC. In the CT plot, bulk density differences between  
201 positions were significant in most of the sampling dates. Unexpectedly, in July 2013, about a month  
202 after tillage, bulk density was higher than before tillage (May, 2013), both in T and in NT position.  
203 Further increase in bulk density was recorded in the T position during the following months, up to  
204  $1.55 \text{ g cm}^{-3}$  (October, 2013). A decrease of bulk density was usually observed between  
205 measurements done in autumn and in the following spring, for both treatments.

206 The  $K_{fs}$  was significantly lower in GC(T) than in CT(T) comparing mean values which were  
207 obtained in November and December, 2012, some days after tillage. For the remaining dates, the  
208 lowest hydraulic conductivity was always measured in CT(T), and the difference was statistically  
209 significant in May 2014 and October 2014. In NT,  $K_{fs}$  was higher in CT than in GC in most of the  
210 dates, being the difference statistically significant in autumn 2012, October 2013 and May 2014.  
211 Higher field-saturated hydraulic conductivity values in the less compacted portion of the GC inter-  
212 row could also relate to different texture of the topsoil.

### 213 **3.2 Runoff and soil erosion seasonal distribution**

214 Table 2 presents a summary of the annual precipitation, runoff and sediment yield during the years  
215 2013 and 2014. During 2013 the annual precipitation was 971 mm, higher than the mean of the  
216 period 2000-2013 (849 mm) (Biddoccu et al., 2016). The rainiest seasons were spring and then  
217 winter, when more than 76% of the cumulative precipitation fell, whereas autumn and especially  
218 summer were drier than usual. The highest runoff coefficients were measured in spring (in CT) and  
219 winter (in GC). In CT sediment yield was much greater in winter than in other seasons, due to a  
220 single event (19-28/12/2013). Precipitation measured in 2014 was more than 40% greater than the  
221 above cited average. The 40% and 35% of annual precipitation were recorded in autumn and in

222 winter, respectively. Highest runoff volumes and runoff coefficients were measured in winter in  
223 both plots.

### 224 **3.3 Influence of rainfall and soil properties on runoff and sediment yield**

225 Table 3 summarises the results of the PCA for the rainfall events and soil properties. More than  
226 80% of the variance among events can be explained by four principal components for CT (83%) and  
227 for GC (86%). The first principal component for CT represents 29% of the variance of the system  
228 and is a good indicator (loadings > 0.90) of rainfall and runoff depth, and rainfall duration of the  
229 event. The intensity of the rainfall is highly correlated with the second principal component (26% of  
230 the variance). Sediment yield is also moderately correlated with this component. The antecedent  
231 precipitation (during the previous 7, 15 and 30 days) and the field-saturated hydraulic conductivity  
232 were the variables best correlated with the third and fourth component, respectively. The first  
233 principal component for the GC events (30% of the variance) is a good indicator of the rainfall and  
234 runoff depths, of rainfall duration and sediment yield. Initial soil water content and antecedent  
235 precipitation in the previous 30 and 15 days are moderately well-represented in the second principal  
236 component (21 % of the variance of the system). The PC3-GC component also represents 21% of  
237 the variance and it is well correlated with maximum rainfall intensity. The PC4-GC is highly  
238 correlated with field-saturated hydraulic conductivity and bulk density.

239 Figure 1 shows the rainfall events represented as individuals on the principal component plans, and  
240 classified by season. Fig.1a represents the events measured in the CT plot in the PC1\_CT-PC2\_CT  
241 plan. In the first quadrant larger rainfall events that produced highest runoff and sediment yield in  
242 CT are represented. They occurred in autumn and winter, when most precipitations greater than 100  
243 mm and long duration (>60 hours) produced significant runoff and erosion, up to 4.9 Mg ha<sup>-1</sup>. The  
244 highest runoff coefficients were recorded with rain causing snowmelt and also relevant erosion  
245 (29/1-13/2/2013 and 26/2-4/3/2013). Most of the high intensity and potentially erosive events  
246 (positive values of PC2\_CT) occurred in summer and early autumn. In that period, runoff occurred

247 when high intensity rainfall ( $I_{max10} > 30 \text{ mm h}^{-1}$ ) was preceded by rainfall in the previous days. The  
248 storm event of 7-8/7/2014 occurred three days after another storm, which did not produce  
249 significant runoff. In the second event, 36 mm of rain fell with the highest 10-min intensity ( $59.15$   
250  $\text{mm h}^{-1}$ ) and produced 9.5 mm of runoff. This resulted in the highest erosion recorded during the  
251 period of observation ( $5.6 \text{ Mg ha}^{-1}$ ). Figure 1b shows events recorded in the GC plot in the  
252 PC1\_GC-PC3\_GC plan. Events represented in the first and fourth quadrant ( $PC1\_GC > 0$ ) occurred  
253 in autumn and winter. Among them, events with  $P > 90 \text{ mm}$  produced high runoff ( $RC > 20\%$ ) and  
254 erosion. Summer and spring events produced negligible runoff and erosion, even with high rainfall  
255 intensities.

256 Table 4 presents the summary of the multiple linear regression models for runoff and sediment yield  
257 variables. In predicting runoff, there is significant correlation with rainfall depth, with a higher  
258 determination coefficient for CT. The runoff model for GC included firstly the rainfall duration.  
259 The variables which were included in the following steps in the runoff model for CT were rainfall  
260 erosivity, maximum hourly intensity and 7-day antecedent precipitation. In predicting sediment  
261 yield there was a significant correlation with erosivity and rainfall depth. The second variable was  
262 the 7-day antecedent precipitation (lower determination coefficient for CT). Figure 2 shows the  
263 accuracy of the predictions with the best multiple linear regression models. The prediction models  
264 resulted in an overestimation of runoff and sediment yield. For the CT treatment this was  
265 particularly evident in autumn and winter events without snowfall.

### 266 **3.4 Analysis of single events to identify the runoff occurrence and mechanism**

267 Runoff was considered as *significant* when its depth was greater than 1 mm or greater than 2% of  
268 the rainfall depth: 14 and 9 rainfall events produced significant runoff in CT and GC, respectively.  
269 Some cases of surface runoff are shown in Figs.3 and 4. The orange lines represent the reference  
270 value of  $K_{fs}$  measured in the GC plot (discontinuous) and in the CT plot (continuous). In some cases  
271  $K_{fs}$  reference values could not be represented in the graph, because of their greater order of



272 magnitude, with respect to rainfall intensity. The blue lines represent the values of saturated soil  
273 water content for CT and GC. The grey band indicates the uncertainty range of sensors (3%). Green  
274 symbols indicate hourly mean values of soil water content measured by the sensors in the plots.

275 Fig. 3a represents a typical winter rainfall event which caused high runoff volumes both from CT  
276 and from GC. The rainfall event accounted for 216.2 mm of rainfall, with maximum 10-min rainfall  
277 intensity of 16.8 mm h<sup>-1</sup>. Measured runoff coefficients and sediment yields accounted for 42% of  
278 rainfall depth and 4.9 Mg ha<sup>-1</sup> in the CT plot, and 20% and 0.49 Mg ha<sup>-1</sup> in the GC plot. Fig. 3b  
279 shows a spring event for which light runoff was measured, that caused little soil erosion. The  
280 rainfall intensity did not overcome K<sub>fs</sub> in the plots, but soil water content increased to reach  
281 saturation of the soil surface and to generate light saturation-excess runoff in CT. The summer  
282 rainfall event in Fig. 3c accounted for 35.8 mm of rainfall. Although the rainfall intensity was the  
283 highest (59 mm h<sup>-1</sup>), it did not overcome the K<sub>fs</sub> minimum value in GC (106 mm h<sup>-1</sup>). The fast  
284 increase of soil water content made the soil saturated in GC for most of the event duration, both in T  
285 than in NT, and little runoff (0.6 mm) was thus originated in this plot. An increase of the topsoil  
286 water content was also measured in CT, but saturation of the soil surface was not reached. Rainfall  
287 intensity overcame K<sub>fs</sub> in CT, thus runoff caused by infiltration excess occurred (9.5 mm) in this  
288 plot. Sediment yield was very high in the tilled plot (5.6 Mg ha<sup>-1</sup>) and negligible in the grassed  
289 vineyard.

290 Fig. 3d shows a rainfall occurring in autumn, before the execution of tillage. After the 10-minute  
291 rainfall intensity exceeded the K<sub>fs</sub> measured in CT, 7.2 mm of runoff were recorded in this plot.  
292 Very low runoff (only 0.4 mm after the whole rainfall event) was measured in GC. For the same  
293 rainfall event 1-min rainfall intensity was also obtained from disdrometer records (Fig. 4a). The  
294 maximum 1-min rainfall intensity was near the 140 mm h<sup>-1</sup> peak measured by the pluviometer of  
295 34.8 mm h<sup>-1</sup> (over 10-min interval). Erosion was detected in the two plots, and in CT it was nearly  
296 40 times higher than in GC. Fig. 4b presents the 1-min rainfall intensity recorded during another

297 event. The rainfall intensity peak did not exceed the  $K_{fs}$ : The soil in CT was tilled 20 days before  
298 the rainfall event, so its conductivity was assumed to be greater than  $2800 \text{ mm h}^{-1}$  and the mean  $K_{fs}$   
299 measured in GC ranged between  $395 \text{ mm h}^{-1}$  and  $967 \text{ mm h}^{-1}$ . However, soil water content was  
300 close to the saturation level, due to the 236 mm of rainfall in the previous two weeks. Runoff  
301 appeared in the first hours of the event, when 1-min rainfall intensity was higher than  $10 \text{ mm h}^{-1}$   
302 and soil was saturated in CT. Thus, runoff was due to saturation of the soil surface in CT. Within a  
303 few hours high rainfall intensity induced saturation of the upper horizon also in GC. When soil  
304 saturation was reached, both in CT and in CG, runoff depth increased in consequence of higher  
305 rainfall intensities. Sediment yield measured in the two plot was nearly  $1.4 \text{ Mg ha}^{-1}$  and  $0.5 \text{ Mg ha}^{-1}$   
306 in CT and GC, respectively.

307 The rainfall depth of autumn and winter events which caused runoff due to saturation in CT ranged  
308 between 36 and 216 mm and the 10-min maximum rainfall intensity varied between 5 to  $17 \text{ mm h}^{-1}$ .  
309 The soil was wet, with soil water content between  $0.267$  and  $0.382 \text{ cm}^3\text{cm}^{-3}$  and it was characterized  
310 by  $K_{fs}$  greater than  $1000 \text{ mm h}^{-1}$  and bulk density of about  $1.41 \text{ g cm}^{-3}$ . The mean runoff coefficient  
311 which was observed in the CT plot for saturation runoff events occurring in autumn and winter was  
312 79%. A light saturation-excess runoff was also detected in spring, with a very low rainfall depth  
313 ( $P=19.60 \text{ mm}$ ,  $I_{\max 10} = 4.8 \text{ mm h}^{-1}$ ) and field-saturated hydraulic conductivity (minimum  $K_{fs} =$   
314  $18.5 \text{ mm h}^{-1}$ ), and wet soil ( $SWC_{\max} = 0.373 \text{ cm}^3\text{cm}^{-3}$ ). Rainfall events that caused Hortonian runoff  
315 in CT were characterized by 10-min maximum rainfall intensity ranging between 37 and  $59 \text{ mm h}^{-1}$   
316 and rainfall depth from 14 to 36 mm. Although the  $K_{fs}$  values which were measured in autumn and  
317 winter in GC were the lowest (ranging between  $41 \text{ mm h}^{-1}$  and  $85 \text{ mm h}^{-1}$ ), rainfall intensities in this  
318 period (maximum 10-min rainfall intensity ranging between 4 and  $35 \text{ mm h}^{-1}$ ) did not cause  
319 Hortonian runoff.

320 Runoff occurred for 50% of the observations in CT and 20% in GC (Fig. 5a). Relationships between  
321 surface runoff type and the season of rainfall event occurrence was firstly examined, after

322 identification of runoff occurrence on single events (Fig. 5b). Only saturation excess runoff was  
323 found during winter, both in CT and in GC. In spring runoff was detected only in CT, for 20% of  
324 the events. In autumn runoff was detected during 40% and 20% of the events, for CT and GC,  
325 respectively. Half of the runoff events which occurred in autumn in CT were due to Hortonian  
326 runoff. In summer less than 20% of the rainfall events caused saturation excess runoff in GC,  
327 whereas 50 % of the rainfall events produced Hortonian runoff in CT. Relationships between  
328 surface runoff frequency and some rainfall characteristics were examined in the following. Fig. 5c  
329 shows that surface runoff by infiltration excess was detected only in CT for 10-min maximum  
330 rainfall intensity higher than  $20 \text{ mm h}^{-1}$  and it occurred for 60% of rainfall events. In relation to the  
331 rainfall depth (Fig. 5d), rainfall events whose depth was lower than 20 mm, produced some  
332 Hortonian and saturation-excess runoff in CT, but no runoff in GC. Most of the Hortonian runoff  
333 events were induced in CT for rainfall events with a depth ranging from 20 to 40 mm. Every  
334 analyzed rainfall event greater than 40 mm produced surface runoff due to saturation excess in both  
335 plots. Finally, relationships are shown between surface runoff and soil water content at the rainfall  
336 occurrence (Fig. 5e) and field-saturated hydraulic conductivity (Fig. 5f). When soil water content  
337 was lower than  $0.250 \text{ cm}^3 \text{ cm}^{-3}$ , no runoff was measured and detected in GC, and only Hortonian  
338 runoff appeared in CT. Both in CT and in GC, the frequency of saturation excess runoff increased  
339 as initial  $\text{SWC}_i$  was higher. In CT some cases of Hortonian runoff were detected with initial  $\text{SWC}_i$   
340 greater than  $0.250 \text{ cm}^3 \text{ cm}^{-3}$ . In relation to the field-saturated hydraulic conductivity of the soil, the  
341 frequency of events that caused Hortonian runoff was about 30% for  $K_{fs} < 100 \text{ mm h}^{-1}$  in CT,  
342 whereas more than 70% of the rainfall events produced runoff due to saturation excess when  $K_{fs}$   
343  $> 1000$ . On the contrary, most of the runoff events occurred in GC soil when  $K_{fs}$  was lower than 100  
344  $\text{mm h}^{-1}$ .

### 345 **3. Discussions**

#### 346 **4.1. Temporal and management effects on soil hydrological properties**

347 With a few exception, soil water content was higher in the T position than in NT, so soil water  
348 saturation was reached sooner in the T position. The topsoil water content was always higher in GC  
349 than in CT, in both T and NT positions.

350 In both treatments, from late autumn and winter to spring, a decrease of bulk density was observed.  
351 Differences between T and NT were significant in CT, except for one month after autumn tillage.  
352 Bagarello et al. (2014) measured  $K_{fs}$  of 838 mm h<sup>-1</sup> and 7424 mm h<sup>-1</sup>, in the clay soil of a Sicilian  
353 vineyard. In a sandy loam soil, previously tilled but then undisturbed over the 2 years of  
354 observation, Bagarello & Sgroi (2007) obtained mean values ranging from 20 mm h<sup>-1</sup> to 952 mm h<sup>-1</sup>,  
355 a range that was very similar to that the one obtained in the GC plot in the present study. The  
356 coefficients of variation obtained in this study were also comparable to those in Bagarello & Sgroi  
357 (2007).

358 During most of the sampling dates the mean values of  $K_{fs}$  in NT position were higher in CT than in  
359 GC. On the contrary, with the only exception of measurements carried out after the autumn tillage,  
360 the lowest mean values in the T position were observed in CT. In the T portion of the inter-row the  
361 increase in water infiltration with respect to GC was evident only within few weeks after the  
362 autumn tillage, whereas tillage was effective in increasing the hydraulic conductivity in the central  
363 portion of the inter-row for a longer period. From spring to autumn, hydraulic conductivity tended  
364 to be higher in CT than in GC in the central part of the inter-row, but it was lower in the T portion ,  
365 showing mean  $K_{fs}$  values lower than 75 mm h<sup>-1</sup>.

366 Both in CT and in GC, during summer and autumn, bulk density showed an increasing trend, and  
367 hydraulic conductivity a decreasing one, that was particularly evident in the T position. Such  
368 tendency was likely the effect of compaction, due to rainfall and especially to intense tractors traffic  
369 during farming and harvesting operations which were carried out in summer and early autumn.  
370 After harvest, mean hydraulic conductivity lower than 100 mm h<sup>-1</sup> was observed in the GC plot and  
371 the lowest  $K_{fs}$  value (40.5 mm h<sup>-1</sup>) was obtained in winter in the T position. The topsoil showed

372 higher compaction and lower hydraulic conductivity after the productive season, especially after  
373 grape harvest. Indeed, worst conditions for water infiltration were observed during autumn (before  
374 tillage in CT) and also in winter for GC. In 2014 the CT topsoil showed higher sand content than in  
375 GC, whereas the clay content was the highest in GC. Since soil erosion by overland flow is a  
376 selective process (Alberts et al., 1980), the more intense erosion which was observed in CT rather  
377 than in GC in the period 2000-2013 (Biddoccu et al., 2016) may have caused the loss of the finest  
378 particles of soil in CT. Differences in field-saturated hydraulic conductivity, namely the higher  
379 values which were observed in the less compacted portion of the inter-row of the GC plot with  
380 respect to the CT plot, could also be related to these differences in the texture of the surface soil.

#### 381 **4.2. Impact of soil management, soil properties and rainfall on runoff**

382 The seasonal and annual runoff amount was usually 2 to 3.6 times higher in CT than in GC. During  
383 summer 2013, which was relatively dry, no runoff was measured, whereas in summer 2014 the CT  
384 runoff was 38% higher than in GC. In 2013 the highest runoff was observed during spring, the most  
385 rainy season, both in CT and in GC. The highest differences in runoff between CT and GC occurred  
386 in summer and spring, when grass cover was higher.

387 In 2014, the highest runoff volumes and runoff coefficients were observed in winter, especially  
388 when snowfall was followed by rainfall. The seasonal runoff coefficient in CT was 83%, whereas it  
389 was 28% in GC. In CT, the winter season showed the highest runoff coefficient also during the  
390 previous decade (Biddoccu et al., 2014; Biddoccu et al., 2016).

391 Winter precipitation events had also the greatest values on the PC1 axis in Fig.1. Despite the  
392 autumn tillage, runoff was much more abundant in CT than in GC, if one or more rainfall events  
393 had already occurred after the execution of tillage. In CT the runoff response to autumn and winter  
394 relevant rainfall events was confirmed by the single event analysis, carried out to identify the type  
395 of runoff that was originated. Among the events for which runoff was identified, all the winter  
396 precipitation events and 20% of the autumn events caused runoff due to topsoil saturation. In both

397 CT and GC the highest runoff coefficients were observed during events when precipitation included  
398 snowfall. Saturation excess runoff was mainly observed after tillage, in late autumn, despite  $K_{fs}$   
399 showing the highest mean values. The water infiltration could be limited at greater depth, because  
400 of the subsoil compaction. In tilled vineyards van Dijk & van Asch (2002) measured in the subsoil  
401 higher bulk density and penetration resistance than in the topsoil, due to the effect of compaction of  
402 wheel load in tilled vineyards. In autumn and winter the grass cover was less effective in reducing  
403 runoff than in other seasons, however runoff measured in GC was at least more than 50% lower  
404 than in CT.

405 The multiple linear model showed the variable response of the CT plot to rainfall characteristics, in  
406 relation to the main mechanism that generated runoff. Runoff was mainly correlated with  
407 precipitation amount, and also moderately correlated with EI30, maximum hourly rainfall intensity  
408 and 7-day antecedent precipitation. In CT, runoff was generated by infiltration excess during  
409 rainfall events characterized by short duration and low depth, and relevant rainfall intensities and  
410 erosivity, that typically occurred in summer and early autumn (before the execution of tillage),  
411 especially when some precipitation was recorded in the previous 7 days.

412 On the contrary, only duration and depth of the precipitation event were correlated with runoff in  
413 GC, which occurred mainly during large events, because of the saturation-excess effect. Indeed,  
414 infiltration-excess runoff was not identified by the rainfall-runoff event analysis in GC, even in  
415 summer. Despite the low hydraulic conductivity, runoff was caused by saturation of topsoil, as  
416 shown by the fast increase of soil water content up to the saturation level.

417 In summer, the very low hydraulic conductivity of the topsoil in the CT plot was likely due also to  
418 the presence of a structural crust, which was observed after first rainfall events following the late-  
419 spring tillage. As Pare et al. (2011) reported in tilled vineyards, cumulative rainfall and kinetic  
420 energy are the main predicting factors of soil reconsolidation, especially from the fresh tillage to

421 crusting. During summer the gradual increase of crusting and compaction due to the tractors traffic  
422 made the runoff larger in CT than in GC.  
423 The analysis of runoff occurrence of each single event was carried out by comparing soil water  
424 content time series with the saturated values obtained in T and NT positions. Similarly the rainfall  
425 intensity time series were compared with  $K_{fs}$ . During most of the events, the value which was  
426 overcome determining the runoff occurrence, was the value of saturated water content (in case of  
427 saturation-excess runoff) or  $K_{fs}$  (in case of Hortonian runoff), in the T position. This effect was  
428 especially evident in summer, when infiltration-excess runoff occurred in CT and not in GC, in  
429 consequence of the lower  $K_{fs}$  in the T position. The winter rainfall-runoff events represented the  
430 most frequent exception to this, because the differences between values of  $K_{fs}$  and  $SWC_s$  were not  
431 significant between the two positions.

#### 432 **4.3. Impact of soil management and rainfall characteristics on sediment yield**

433 Sediment yields of 5.3 and 9.3 Mg ha<sup>-1</sup> were measured during the two years, respectively, in the CT  
434 plot. The average value was close to 7.4 Mg ha<sup>-1</sup> year<sup>-1</sup>, of the period 2000-2012 for the same plot  
435 (Biddoccu et al., 2016). Tropeano (1984) reported annual soil loss of 47.4 Mg ha<sup>-1</sup> in a tilled  
436 vineyard in Piedmont and annual sediment yields of 31.4 and 88.71 Mg ha<sup>-1</sup> were measured in 2  
437 years of observation in a tilled vineyard in Sicily (Novara et al., 2011). In 2013 erosion was higher  
438 in CT than in GC in each season. During the most erosive winter event, erosion in CT was 10 times  
439 greater than in GC. Also in 2014 seasonal erosion was greater in CT than in GC. In GC annual soil  
440 erosion was 1.5 Mg ha<sup>-1</sup> in 2013, and 1.0 Mg ha<sup>-1</sup>, in 2014, lower than the mean observed in 2000-  
441 2012, of 1.8 Mg ha<sup>-1</sup> year<sup>-1</sup>. Annual sediment yield in GC was 72 and 89% less than in CT. The  
442 difference between CT and GC was greater than in other studies. Novara et al. (2011) observed that  
443 the use of different cover crops in the inter-row reduced soil losses by 56%. Ruiz-Colmenero et al.  
444 (2011), in a 2-year study at plot scale, observed that vineyards with a cover crop lost between 50%  
445 and 75% less soil than with tilled soil.

446 Differences among seasonal amounts of sediment yields were due to the variability of rainfall  
447 during the years. Seasonal distribution of rainfall, runoff and soil erosion in 2014 was similar to the  
448 period 2000-2013 (Biddoccu et al., 2016). Nevertheless in both years just a single event per year  
449 had a great influence in determining the annual sediment yield. In 2013 the rainfall event which  
450 occurred in the period 19-28/12/2013 was the largest (216.2 mm). It was the most erosive event  
451 observed in GC during the study, and the second most erosive event measured in CT, where it  
452 caused 93% of the annual erosion. In 2014, the highest erosion occurred during the summer storm  
453 on 7-8/7/2014, when the rainfall of highest intensity (59 mm h<sup>-1</sup> over 10 minutes) caused more than  
454 5.6 Mg ha<sup>-1</sup> of sediment yield in CT, whereas in GC only 9 kg ha<sup>-1</sup>. Those two extreme events, with  
455 the highest precipitation depth and 10-min maximum intensity, respectively, caused the highest  
456 erosion. These results confirmed the observation of Gómez et al. (2014) and González-Hidalgo et  
457 al. (2009). The multiple linear model predicting sediment yield in CT showed the highest  
458 correlation with erosivity, which depends on the 30-min rainfall intensity and on the energy of  
459 precipitation, and then with the 7-day antecedent precipitation. Hortonian runoff was mainly  
460 observed in summer and early autumn, for rainfall events with 10-min maximum intensity greater  
461 than 20 mm h<sup>-1</sup>, and with significant the 7-day antecedent rainfall. In those cases, although runoff  
462 coefficients were limited (average value of 7%), sediment yield ranged between 289 kg ha<sup>-1</sup> and  
463 5658 kg ha<sup>-1</sup>. Apart from the first precipitation event after tillage, most of the events in autumn and  
464 winter generated saturation-excess runoff in CT, whose rates were greater than 42%, and caused  
465 sediment yield up to 4.9 Mg ha<sup>-1</sup>.

466 The high correlation between sediment yield and erosivity for CT was related to the absence of soil  
467 protection and low hydraulic conductivity in summer and early autumn, when most intense and  
468 erosive rainfall was observed. Nevertheless, high erosivity is also related to the rainfall energy, that  
469 is high for large precipitation events in autumn and winter. Similarly to this study, Raclot et al.  
470 (2009), in tilled vineyards, at event temporal scale and at field spatial scale, found significant



471 correlation between total suspended sediment and rainfall depth and between erosion and maximum  
472 rainfall intensity over a 5-min period.

473 In GC in summer and early-autumn negligible erosion was observed. The protective role of grass  
474 was little effective when saturation-excess runoff was generated by large precipitation events. In  
475 autumn and winter, when grass is sparse and soil is more compacted and less conductive than in  
476 summer (runoff coefficients and erosion up to 53% and 1181 kg ha<sup>-1</sup>). The multiple linear model  
477 showed sediment yield mainly correlated with precipitation depth, and, secondly, to rainfall in the  
478 previous 7 days. The sediment yield during single events, in any case, was lower than in CT. The  
479 results obtained in the GC plot were similar to those of Gómez et al. (2014) in a grassed olive  
480 orchard, with higher correlation between sediment yield and rainfall depth than with rainfall  
481 erosivity and short term intensity. They also observed the largest erosive events in GC in late  
482 autumn and winter, when grass cover is scarce and soil compacted after the productive season.

#### 483 **4. Conclusions**

484 The soil management effects on soil hydraulic conductivity, bulk density, soil moisture, runoff and  
485 water erosion were evaluated in a two-year period in two vineyard field-scale plots, where inter-  
486 rows were managed with conventional tillage and grass cover, respectively. Eighty infiltration tests  
487 were carried out and a dataset of 29 rainfall-runoff events was collected, covering a wide range of  
488 topsoil characteristics.

489 The results highlighted how the tillage increased field-saturated hydraulic conductivity only for a  
490 short period. It tended to be higher in CT than in GC in the central part of the inter-row, but in track  
491 it quickly decreased to such low values that Hortonian runoff was produced during intense summer  
492 storms. Sediment yield in the tilled plot was up to 9 times higher than in the grass cover plot. Due to  
493 high compaction after grape harvest, the worst condition for infiltration were found before autumn  
494 tillage in CT and in autumn and winter for GC. The main runoff events were related to the  
495 saturation-excess mechanism, which was the only one observed in the GC plot and that was

496 frequently observed in the tilled one, particularly in late autumn and in long-duration winter  
497 precipitation events. Despite the autumn tillage, in CT the winter 2014 runoff reached 83% of the  
498 precipitation amount and was nearly 4 times greater than in the grassed plot. The largest runoff  
499 occurred in case of snowfall events followed by rainfall. Very high sediment yield in the tilled plot  
500 was mainly related to rainfall intensity (during summer storm) and rainfall depth (in autumn and  
501 winter). The annual reduction of runoff in the grassed plot was 63% in comparison with the tilled  
502 plot. In autumn and winter the grass cover was less effective in reducing runoff than in summer.  
503 Erosion was relevant in winter when large saturation-excess runoff was generated by long-lasting  
504 rainfall and snowfall. However, the grass cover was effective in reducing annual soil losses (up to  
505 90%) and especially during most erosive events that occurred in summer and early-autumn.

#### 506 **Acknowledgements**

507 This research was funded by the “Centro Sperimentale Vitivinicolo Regionale Tenuta Cannona”  
508 and the Regione Piemonte - Office for Agricultural Development and Office for Agricultural  
509 Enterprises (research project “Tutela del suolo e delle acque superficiali” 2012-2014). The study  
510 was partially carried out within the project CIRCE (Cooperative Internet-of-Data Rural-alpine  
511 Community Environment), funded by EU FESR, by Italian Government and Regione Piemonte  
512 within the programme Regione Piemonte POR/FESR 2007–2013. This work was partially funded  
513 by the project PRIN 2010-2011 2010JHF437\_004. We are grateful to: the staff of the “CSV Tenuta  
514 Cannona”, which collaborated managing the vineyards and in sample collections; Mr. Giorgio  
515 Capello who collaborated in the fieldwork; Dr. Marco Milan and the DISAFA (University of  
516 Torino), for assistance in analysis of runoff samples.

517

#### 518 **References**

519 Alberts, E.E., Moldenhauer, W.C. & Foster, G.R., 1980. Soil Aggregates and Primary Particles  
520 Transported in Rill and Interrill Flow. *Soil Sci. Soc. Am. J.* 44 (3), 590-595.  
521 Doi:10.2136/sssaj1980.03615995004400030032x

522 Arnaez, J., Lasanta, T., Ruiz-Flano, P., Ortigosa, L., 2007. Factors affecting runoff and erosion  
523 under simulated rainfall in Mediterranean vineyards. *Soil and Tillage Research* 93 (2), 324–334.

524 ARS-USDA, 2015. RIST - Rainfall Intensity Summarization Tool.  
525 <http://www.ars.usda.gov/Research/docs.htm?docid=3251> Last accessed on 03.11.2015

526 Bagarello, V., Iovino, M., Elrick, D., 2004. A Simplified Falling-Head Technique for Rapid  
527 Determination of Field-Saturated Hydraulic Conductivity. *Soil Sci. Soc. Am. J.*, 68, 66-73.

528 Bagarello, V., Sgroi, A., 2007. Using the simplified falling head technique to detect temporal  
529 changes in field-saturated hydraulic conductivity at the surface of a sandy loam soil. *Soil Till. Res.*  
530 94, 283-294. doi:10.1016/j.still.2006.08.001

531 Bagarello, V., Baiamonte, G., Castellini, M., Di Prima, S., Iovino M., 2014. A comparison between  
532 the single ring pressure infiltrometer and simplified falling head techniques. *Hydrol. Process.* 28,  
533 4843–4853. doi: 10.1002/hyp.9980

534 Biancotti, A., Bellardone, G., Bovo, S., Cagnazzi, B., Giacomelli, L., Marchisio, C., 1998.  
535 Distribuzione Regionale di Piogge e Temperature. *Collana Studi Climatologici del Piemonte, Vol.1.*  
536 Regione Piemonte. Torino.

537 Biddoccu, M., Opsi, F., Cavallo, E., 2014. Relationships between runoff and soil losses with rainfall  
538 characteristics and a long-term soil management practices in a hilly vineyard (Piedmont, NW Italy).  
539 *Soil Sci. Plant Nutr.* 60(1), 92-99. <http://dx.doi.org/10.1080/00380768.2013.862488>

540 Biddoccu, M., Ferraris, S., Opsi, F., Cavallo, E. 2016. Long-term monitoring of soil management  
541 effects on runoff and soil erosion in sloping vineyards in Alto Monferrato (North-West Italy). *Soil*  
542 *Till. Res.* 155, 176-189, DOI: 10.1016/j.still.2015.07.005

543 Blavet, D., De Noni, G., Le Bissonnais, Y., Leonard, M., Maillo, L., Laurent, J.Y., Asseline, J.,  
544 Leprun, J.C., Arshad, M.A., Roose, E., 2009. Effect of land use and management on the early stages  
545 of soil water erosion in French Mediterranean vineyards. *Soil Till. Res.* 106, 124-136.

546 Bodhinayake, W., Si, B.C., Noborio, K., 2004. Determination of hydraulic properties in sloping  
547 landscapes from tension and double-ring infiltrometers. *Vadose Zone Journal* 3, 964-970.

548 Brenot, J., Quiquerez, A., Petit, C., Garcia, J.-P., 2008. Erosion rates and sediment budgets in  
549 vineyards at 1-m resolution based on stock unearthing (Burgundy, France). *Geomorphology* 100,  
550 345-55.

551 Brown, L.C., Foster, G.R., 1987. Storm erosivity using idealized intensity distributions.  
552 *Transactions of ASAE* 30 (2), 379-386.

553 Casalí, J., Gastesi, R., Álvarez-Mozos, J., De Santisteban, L.M., Lersundi, J.D.V.d., Giménez, R. ,  
554 Larrañaga, A., Goñi, M., Agirre, U., Campo, M.A., López, J.J., Donézar, M., 2008. Runoff, erosion,  
555 and water quality of agricultural watersheds in central Navarre (Spain), *Agr. Water Manage.* 95,  
556 1111-1128.

557 Celette, F., Ripoche, A., Gary, C., 2010. WaLIS—A simple model to simulate water partitioning in  
558 a crop association: The example of an intercropped vineyard. *Agr. Water Manage.* 97, 1749-1759.

559 Cerdà, A., Doerr, S.H., 2007. Soil wettability, runoff and erodibility of major dry-Mediterranean  
560 land use types on calcareous soils. *Hydrol. Process.* 21, 2325-2336.

561 Cerdan, O., Govers, G., Le Bissonnais, Y., Van Oost, K., Poesen, J., Saby, N., Gobin, A., Vacca,  
562 A., Quinton, J., Auerwald, K., Klik, A., Kwaad, F.J.P.M., Raclot, D., Ionita, I., Rejman, J.,  
563 Rousseva, S., Muxart, T., Roxo, M.J. & Dostal, T., 2010. Rates and spatial variations of soil erosion  
564 in Europe: A study based on erosion plot data. *Geomorphology* 122, 167-177.

565 Corti, G., Cavallo, E., Cocco, S., Biddoccu, M., Brecciaroli, G., Agnelli, A., 2011. Evaluation of  
566 erosion intensity and some of its consequences in vineyards from two hilly environments under a  
567 Mediterranean type of climate, Italy. In: Godone, D. & Stanchi, S. (Eds.). *Soil Erosion in*  
568 *Agriculture*. Intech Open Access Publisher Eds., pp. 113–160.

569 Dehotin, J., Breil, P., Braud, I., de Lavenne, A., Lagouy, M., Sarrazin, B., 2015. Detecting surface  
570 runoff location in a small catchment using distributed and simple observation method. *J. Hydrol.*  
571 *525*, 113-129.

572 FAO/ISRIC/ISSS, 1998. *World Reference Base for Soil Resources*. World Soil Resources Report,  
573 No 84. FAO, Rome

574 Ferrero, A., Usowicz, B., Lipiec, J., 2005. Effects of tractor traffic on spatial variability of soil  
575 strength and water content in grass covered and cultivated sloping vineyard. *Soil Till. Res.* 84, 127–  
576 138.

577 García-Ruiz, J.M., 2010. The effects of land uses on soil erosion in Spain: a review. *Catena* 81, 1–  
578 11.

579 García-Ruiz, J.M., Beguería, S., Nadal-Romero, E., Gonzalez-Hidalgo, J.C., Lana-Renault, N.,  
580 Sansjuan, Y., 2015. A meta-analysis of soil erosion rates across the world. *Geomorphology* 239,  
581 160–173.

582 Gómez, J.A., Sobrinho, T.A., Giráldez., J.V., Fereres, E., 2009. Soil management effects on runoff,  
583 erosion and soil properties in an olive grove of Southern Spain. *Soil Till. Res.* 102, 5-13.  
584 doi:[10.1016/j.still.2008.05.005](https://doi.org/10.1016/j.still.2008.05.005)

585 Gómez, J.A., Vanwallenghem, T., De Hoces, A., Taguas, E.V., 2014. Hydrological and erosive  
586 response of a small catchment under olive cultivation in a vertic soil during a five-year period:  
587 Implications for sustainability. *Agr. Ecosyst. Environ.* 188, 229-244.

588 González-Hidalgo, J.C., de Luis, M., Batalla, R.J., 2009. Effects of the largest daily events on total  
589 soil erosion by rainwater. An analysis of the USLE database. *Earth Surface Processes and*  
590 *Landforms* 34 (15), 2070–2077.

591 Kosmas, C., Danalatos, N., Cammeraat, L.H. et al., 1997. The effect of land use on runoff and soil  
592 erosion rates under Mediterranean conditions. *Catena*, 29, 45–59.

593 Leonard, J., Andrieux, P., 1998. Infiltration characteristics of soils in Mediterranean vineyards in  
594 Southern France. *Catena* 32, 209–223.

595 Maetens, W., Vamaercke, M., Poesen, J., Jankauskas, B., Jankauskiene, G., Ionita, I., 2012. Effect  
596 of land use on annual runoff and soil loss in Europe and the Mediterranean: A meta-analysis of plot  
597 data. *Prog. Phys. Geog.* 36 (5), 599-653.

598 Novara, A., Gristina, L., Saladino, S.S., Santoro, A., Cerdà, A. 2011. Soil erosion assessment on  
599 tillage and alternative soil managements in a Sicilian Vineyard. *Soil Till. Res.* 117, 140-147.

600 Pare, N., Andrieux, P., Louchart, X., Biarnes, A., Voltz, M., 2011. Predicting the spatio-temporal  
601 dynamic of soil surface characteristics after tillage. *Soil Till. Res.* 114, 135-145.  
602 doi:[10.1016/j.still.2011.04.003](https://doi.org/10.1016/j.still.2011.04.003)

603 Prosdocimi, M., Cerdà, A., Tarolli, P., 2016. Soil water erosion on Mediterranean vineyards: A  
604 review. *Catena* 141 (2016) 1–21.

605 Raclot, D., Le Bissonais, Y., Louchart, Y., Andrieux, P., Moussa, R., Voltz, M., 2009. Soil tillage  
606 and scale effects on erosion from fields to catchment in a Mediterranean vineyard area. *Agr.*  
607 *Ecosyst. Environ.*, *Catena* 66, 198-210.

608 Ramos, M.C., Martínez-Casasnovas, J.A., 2004. Nutrient losses from a vineyard soil in  
609 Northeastern Spain caused by an extraordinary rainfall event. *Catena* 55,79–90. doi:[10.1016/S0341-](https://doi.org/10.1016/S0341-8162(03)00074-2)  
610 [8162\(03\)00074-2](https://doi.org/10.1016/S0341-8162(03)00074-2)

611 Ramos, M.C., Martínez-Casasnovas, J.A., 2007. Soil loss and soil water content affected by land  
612 leveling in Penedès vineyards. *Catena* 71, 210-217. doi:10.1016/j.catena.2007.03.001

613 Renard, K.G., Foster, G.R., Weesies, G.A., McCool, D.K., Yoder, D.C. 1997. Predicting soil erosion  
614 by water: a guide to conservation planning with the revised universal soil loss equation (RUSLE).  
615 US Department of Agriculture Agricultural Handbook No. 703. USDA Washington DC.

616 Ruiz-Colmenero, M., Bienes, R., Marques, M.J., 2011. Soil and water conservation dilemmas  
617 associated with the use of green cover in steep vineyards. *Soil Till. Res.* 117, 211–223.

618 Soil Survey Staff, 2010. *Keys to Soil Taxonomy*. 11th ed. Washington, DC: USDA-Natural  
619 Resources Conservation Service.

620 Tropeano, D., 1983. Soil-erosion on vineyards in the tertiary piedmontese basin (NorthWestern  
621 Italy) studies on experimental areas. *Catena (Suppl. 4)*, 115–127.

622 Tropeano, D., 1984. Rate of soil erosion processes on vineyards in Central Piedmont (NW Italy).  
623 *Earth Surf. Proc. Land.* 9, 253-266.

624 Van Dijk, S.J.E., van Asch Th.V.J., 2002. Compaction of loamy soils due to tractor traffic in  
625 vineyards and orchards and its effects on infiltration in Southern France. *Soil Till. Res.* 63, 141-153.

626 Warrick, A.W., 1998. Spatial variability. In: Hillel, D. (Ed.), *Environmental Soil Physics*. Academic  
627 Press, San Diego, CA, USA, pp.665–675.

Table 1 – Mean values and coefficient of variation (CV, %) of the bulk density (BD), initial soil water content (SWC<sub>i</sub>) and field-saturated hydraulic conductivity (K<sub>fs</sub>) measured with the SFH techniques on each sampling date in the CT and GC treatments in track (T) and no-track (NT) positions. Geometric mean was used for K<sub>fs</sub>. Bold values are different between positions according to t-test at p=0.05 level. Different letters indicate significant differences between treatments according to t-test at p=0.05 level.

CT		From Tillage Days		P (mm)		BD (g cm <sup>-3</sup> )		SWC <sub>i</sub> (cm <sup>3</sup> cm <sup>-3</sup> )		K <sub>fs</sub> (mm h <sup>-1</sup> )		GC		From Tillage Days		R (mm)		BD (g cm <sup>-3</sup> )		SWC <sub>i</sub> (cm <sup>3</sup> cm <sup>-3</sup> )		K <sub>fs</sub> (mm h <sup>-1</sup> )	
		NT	T	NT	T	NT	T	NT	T	NT	T			NT	T	NT	T	NT	T	NT	T	NT	T
Oct-12	Mean	158	141.	<b>1.3</b>	<b>0</b>	<b>1.50</b>	<b>1</b>	<b>0</b>	<b>71.0</b>	<b>31.3</b>													
	CV (%)		2	6.2	7.3	6.1	4.3	29.9	53.0														
Nov-12	Mean	21	139.	1.3	1.41	0.35	0.35	2886.2	3747.3			Dec-12	Mean	42	259.	1.35	1.41	<b>0.33</b>	<b>0.40</b>	251.	40.5		
	CV (%)		2	8.2	5.5	4.5	4.4	39.1	102.1				CV (%)			2.98	9.79	6.3	2.1	186.	154.		
May-13	Mean	202	698.	<b>1.1</b>	<b>1.38</b>	0.26	0.31	<b>770.1</b>	<b>63.6 a</b>			May-13	Mean	202	698.	1.18	1.29	0.27	0.31	427.	153.		
	CV (%)		6	2.1	7.2	4.5	17.4	34.8	8.6				CV (%)			5.82	11.4	9.8	30.4	45.9	93.6		
Jul-13	Mean	3	16.2	1.3	1.41	0.16	0.18	486.87	44.69			Jul-13	Mean	37	16.2	1.31	1.48	<b>0.09</b>	<b>0.18</b>	105.	156.		
	CV (%)		4	7.0	3.0	17.4	31.6	25.4	102.0				CV (%)			8.22	6.22	19.8	9.3	55.5	152.		
Sep-13	Mean	93	88.2	<b>1.2</b>	<b>1.45</b>	0.15	0.17	<b>332.6</b>	<b>10.0 a</b>			Sep-13	Mean	92	88.2	1.27	1.42	0.19	0.22	93.7	85.2		
	CV (%)			1.1	2.0	4.9	17.1	49.5	11.0				CV (%)			0.74	9.71	20.4	11.6	95.1	120.		
Oct-13	Mean	133	137.	<b>1.3</b>	<b>1.55</b>	0.27	0.29	<b>1456.3</b>	<b>74.4 a</b>			Oct-13	Mean	134	137.	1.34	1.42	0.29	0.33	<b>591.</b>	<b>99.0</b>		
	CV (%)		4	1.2	2.7	11.7	5.2	17.2	12.2				CV (%)			5.35	2.97	5.7	0.0	27.8	25.2		
May-14	Mean	184	740.	<b>1.2</b>	<b>1.51</b>	0.24	0.23	<b>87.1 a</b>	<b>18.5 a</b>			May-14	Mean	183	740.	1.24	1.37	0.25	0.26	525.	423.		
	CV (%)		6	5.8	2.8	8.8	26.1	31.9	45.7				CV (%)			12.8	7.60	17.8	4.5	126.	37.2		
Oct-14	Mean	156	246.	1.2	1.47	0.26	0.32	<b>1343.5</b>	<b>20.5 a</b>			Oct-14	Mean	157	246.	1.24	1.37	0.31	0.32	967.	394.		
	CV (%)		2	6.2	0.5	14.3	0.4	63.1	75.6				CV (%)			6.42	4.76	13.8	20.1	91.3	63.1		

Table 2 – Seasonal and annual records from the experimental vineyard plots (conventional tillage, CT; grass cover, GC) in 2013 and 2014: Precipitation (including snowfall in winter), runoff (RO), runoff coefficient (RC), sediment yield (SY).

	Precipitation (mm)		CT				GC							
	2013	2014	RO (mm)		RC (%)		SY (kg ha <sup>-1</sup> )		RO (mm)		RC (%)		SY (kg ha <sup>-1</sup> )	
Winter	323.2	433.4	108.2	358.4	33	83	4982	1806	51.0	123.4	16	28	498	440
Spring	421.0	125.0	214.1	2.0	51	2	222	2	58.1	0.8	14	1	996	1
Summer	66.8	171.8	0.0	21.4	0	12	0	5657	0.0	1.6	0	1	0	9
Autumn	159.8	480.0	3.1	196.7	2	41	107	1911	1.4	86.0	1	18	6	593
Total	970.8	1210.2	325.5	578.5	34	48	5311	9377	110.5	211.9	11	18	1501	1043



Table 3 – Results of factor analysis of the rainfall events variables and soil variables measured in the two plots, for the extraction of principal components. Values in *italic* and **bold** indicated the moderately high (>0.70) and high (>0.90) loadings. (CT = conventional tilled, GC = grass cover, I max X min = maximum intensity in X min, EI30 = erosivity, RO = runoff, RC = runoff coefficient, SY = sediment yield, Ant. Prec. Y days = antecedent precipitation in previous Y days, SWC = soil water content,  $K_{fs}$  = field-saturated hydraulic conductivity, BD = bulk density).

CT	PC1-CT	PC2-CT	PC3-CT	PC4-CT	GC	PC1-GC	PC2-GC	PC3-GC	PC4-GC
Precipitation	<b>0.971</b>	0.096	-0.018	-0.001	Precipitation	<b>0.966</b>	0.117	0.058	-0.103
Precip. duration	<b>0.925</b>	-0.201	0.035	-0.112	Precip. duration	<b>0.904</b>	0.161	-0.206	0.106
I max 10 min	-0.235	<b>0.870</b>	-0.208	0.021	I max 10 min	-0.173	-0.207	<b>0.914</b>	0.005
I max 30 min	-0.220	<b>0.909</b>	-0.194	0.058	I max 30 min	-0.112	-0.190	<b>0.960</b>	-0.039
I max 60 min	-0.089	<b>0.950</b>	-0.141	0.113	I max 60 min	0.039	-0.112	<b>0.957</b>	-0.082
EI30	0.275	<b>0.915</b>	-0.109	0.015	EI30	0.677	-0.069	0.641	-0.144
RO CT	<b>0.964</b>	0.011	0.037	-0.045	RO GC	<b>0.930</b>	0.166	-0.090	0.150
RC CT	<i>0.899</i>	0.066	0.287	-0.033	RC GC	<i>0.798</i>	0.383	-0.156	0.234
SY_CT	0.111	<i>0.787</i>	0.251	-0.077	SY_GC	<b>0.915</b>	-0.116	0.034	0.091
Ant. Prec. 7 days	-0.148	0.002	<i>0.780</i>	0.003	Ant. Prec. 7 days	-0.001	<i>0.708</i>	0.023	-0.386
Ant. Prec. 15 days	0.159	-0.223	<i>0.703</i>	0.002	Ant. Prec. 15 days	-0.041	<b>0.824</b>	-0.191	-0.062
Ant. Prec.30 days	0.460	-0.220	<i>0.719</i>	-0.218	Ant. Prec.30 days	0.124	<b>0.826</b>	-0.247	0.302
Ant. Prec. 45 days	<i>0.709</i>	-0.250	0.521	-0.209	Ant. Prec. 45 days	0.324	<i>0.701</i>	-0.322	0.381
SWC CT	0.141	0.130	0.605	0.448	SWC GC	0.226	<b>0.830</b>	-0.070	0.017
$K_{fs}$ CT	-0.102	-0.014	0.031	<b>0.964</b>	$K_{fs}$ GC	-0.013	0.089	0.019	<b>-0.921</b>
BD CT	0.368	-0.086	0.504	-0.513	BD GC	0.179	0.132	-0.084	<b>0.906</b>
Eigenvalues	4.661	4.181	2.773	1.522	Eigenvalues	4.797	3.389	3.377	2.195
Accumulated variance	29.134	26.130	17.330	9.514	Accumulated variance	29.982	21.183	21.106	13.716

Table 4 – Summary of the stepwise multiple linear regression model for runoff and sediment yield in the two plots ( $R^2_{adj}$  = adjusted coefficient of determination, EI30 = erosivity, I max 60 min = maximum intensity in 60 min, Ant. Prec.7 days = antecedent rainfall in previous 7 days).

Runoff Conventional Tillage						Runoff Grass Cover					
Variable	Step	Value	$R^2_{adj}$	Increase in $R^2_{adj}$	Sign.		Step	Value	$R^2_{adj}$	Increase in $R^2_{adj}$	Sign.
Intercept		-40.939				Intercept		-6.798			
Precipitation	1	1.366	0.899	0.899		Precipitation duration	1	0.230	0.769	0.769	0.000
EI30	2	-0.387	0.918	0.019	0.000	Precipitation	2	0.129	0.810	0.041	0.000
I max 60 min	3	3.488	0.947	0.029	0.000						
Ant. Prec.7 days	4	0.337	0.957	0.010	0.000						
Sediment Yield Conventional Tillage						Sediment Yield Grass Cover					
Variable	Step	Value	$R^2_{adj}$	Increase in $R^2_{adj}$	Sign.		Step	Value	$R^2_{adj}$	Increase in $R^2_{adj}$	Sign.
Intercept		-855.559				Intercept		-33.434			
EI30	1	13.696	0.507	0.507	0.000	Precipitation	1	1.882	0.753	0.753	0.000
Ant. Prec. 7 days	2	32.758	0.585	0.078	0.000	Ant. Prec.7 days	2	-1.335	0.819	0.066	0.000

## Figure captions

Fig.1 – Representation of events as individuals on the principal component plan, classified by season. (a) Rainfall events associated with runoff-erosion and soil characteristics measured in the CT plot represented in the PC1\_CT-PC2\_CT plan and (b) Rainfall events associated with runoff-erosion and soil characteristics measured in the GC plot represented in the PC1\_GC-PC3\_GC plan. Different symbols indicate season of rainfall occurrence.

Fig.2 – Comparison of values observed and predicted by the multiple linear regression models for (a) runoff in the CT plot, (b) sediment yield in the CT plot, (c) runoff in the GC plot, (b) sediment yield in the GC plot.



Fig.3 – Examples of runoff detection graphs with pluviometer data (10 minutes step) for some rainfall events. R= rain intensity at 10 min step, SWC = measured volumetric soil water content (1 hour step), SWCs = reference value of saturated water content,  $K_{fs}$  = reference value of field-saturated hydraulic conductivity, RO cum = cumulated measured runoff, P = total event precipitation, I10 = maximum rain intensity at 10 min step, P7 = antecedent precipitation at 7-day step, RO= measured runoff, SY = measured sediment yield.

Fig.4 – Examples of runoff detection graphs with disdrometer data (1 minutes step) for some rainfall events. TotalPIntensity\_1min= rain intensity at 1 min step, SWC = measured volumetric soil water content (1 hour step) in the plot, SWCs = reference value of saturated water content,  $K_{fs}$  = reference value of field-saturated hydraulic conductivity, RO cum = cumulated measured runoff, P = total event precipitation, I1 = maximum rain intensity at 1 min step, P7 = antecedent precipitation at 7-day step, RO = measured runoff plot, SY= measured sediment yield, T = track position, NT = no track position.

Fig.5 – Influence of soil management (a), season (b), rainfall maximum intensity (c), rainfall depth (d), initial soil water content (e) and field-saturated hydraulic conductivity (f) on the surface runoff occurrence and type of runoff. Each bar represents the totality of analyzed events for each category, and among those events they indicate the fractions of: (i) events without runoff occurrence (No runoff, blue bars), (ii) events for which hortonian runoff was detected (red bars), (iii) events for which saturation-excess runoff was identified (green bars).

Figure1

[Click here to download high resolution image](#)



◆ Summer ■ Winter ▲ Spring × Autumn

Figure2

[Click here to download high resolution image](#)








Figure3a

[Click here to download high resolution image](#)



a

Figure3b

[Click here to download high resolution image](#)







Figure3c

[Click here to download high resolution image](#)



**C**



Figure3d

[Click here to download high resolution image](#)




Figure4a

[Click here to download high resolution image](#)




Figure4b

[Click here to download high resolution image](#)




Figure 5

[Click here to download high resolution image](#)






Figure1\_gray

[Click here to download high resolution image](#)

◆ Summer ■ Winter ▲ Spring × Autumn









Figure3a\_gray  
[Click here to download high resolution image](#)




Figure3b\_gray  
[Click here to download high resolution image](#)








Figure3c\_gray  
[Click here to download high resolution image](#)



**C**

Figure3d\_gray  
[Click here to download high resolution image](#)



d

Figure4a\_gray  
[Click here to download high resolution image](#)






Figure4b\_gray  
[Click here to download high resolution image](#)



b

Figure5\_gray

[Click here to download high resolution image](#)



**Supplementary material for on-line publication only**

**[Click here to download Supplementary material for on-line publication only: Supplementary\\_file\\_for\\_download\\_19032016.doc](#)**