



Rickettsia raoultii and *Rickettsia sibirica* in ticks from the long-tailed ground squirrel near the China–Kazakhstan border

Shuo Zhao¹ · Meihua Yang² · Mengmeng Jiang³ · Bin Yan¹ · Shanshan Zhao¹ ·
Wumei Yuan¹ · Baoju Wang⁴ · Sándor Hornok⁵ · Yuanzhi Wang¹

Received: 28 October 2018 / Accepted: 20 February 2019 / Published online: 25 February 2019
© The Author(s) 2019

Abstract

Spotted fever group (SFG) rickettsiae cause infection in humans, domestic animals and wildlife. To date, no rickettsial agents have been reported in hard ticks from the long-tailed ground squirrel (*Spermophilus undulatus*). A total of 50 adult ticks and 48 nymphs were collected from *S. undulatus* in the border region of northwestern China. Tick species (identified according to morphological and molecular characteristics) included *Dermacentor nuttalli*, *Dermacentor silvarum* and *Ixodes kaiseri*. Based on the cytochrome *c* oxidase subunit I (*COI*) haplotype analysis, *I. kaiseri* from *S. undulatus* belongs to an ancestral. In addition, all tick samples were analyzed for the presence of rickettsiae by PCR amplification and sequencing of six genetic markers. *Rickettsia raoultii* and *Rickettsia sibirica* subsp. *sibirica* were shown to occur in adults and nymphs of *D. nuttalli* and *D. silvarum*. *Rickettsia sibirica* subsp. *sibirica* was also detected in an *I. kaiseri* adult. *Dermacentor silvarum* and *I. kaiseri* were found for the first time on *S. undulatus*. *Rickettsia raoultii* and *R. sibirica* subsp. *sibirica* were detected in two *Dermacentor* and one *Ixodes* species, respectively, suggesting that these rickettsiae circulate in the region of the China-Kazakhstan border by hard ticks infesting *S. undulatus*.

Keyword *Spermophilus undulatus* · Ticks · *Rickettsia raoultii* · *Rickettsia sibirica* · Northwestern China

Shuo Zhao, Meihua Yang, and Mengmeng Jiang contributed equally to this work.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10493-019-00349-5>) contains supplementary material, which is available to authorized users.

✉ Yuanzhi Wang
wangyuanzhi621@126.com

Extended author information available on the last page of the article

Introduction

The long-tailed ground squirrel (*Spermophilus undulatus*) has been listed as globally vulnerable by the International Union for Conservation of Nature (IUCN) since 2008 (Ramolara et al. 2014). This species is mainly distributed in Kazakhstan, Mongolia, Russia and northern China including Heilongjiang, Inner Mongolia and Xinjiang Uygur Autonomous Region (XUAR) (ZipcodeDev Team 2018). According to previous reports, *S. undulatus* and its ectoparasites are reservoirs of *Yersinia pestis*, *Francisella tularensis* and tick-borne encephalitis virus (Zhao et al. 2017; Wang and Yang 1983; Demina et al. 2017). However, only few tick species (i.e., *Dermacentor marginatus*, *Dermacentor nuttalli*, *Rhipicephalus schulzei* and *Ixodes persulcatus* (Wang and Yang 1983; Demina et al. 2017)) are known to infest *S. undulatus*.

Members of the tick subgenus *Pholeoixodes* prefer burrowing mammals as hosts (Hornok et al. 2017), thus can be expected to occur on *S. undulatus*. Nevertheless, no previous reports confirmed this. In a study comparing three species of this subgenus (*Ixodes kaiseri*, *Ixodes hexagonus* and *Ixodes canisuga*) the 16S rDNA phylogenetic tree reflected that *I. kaiseri* is divided into at least nine COI haplotypes in Europe (Hornok et al. 2017). However, at that time no specimens of *I. kaiseri* were available from Asia for comparison.

Spotted fever group (SFG) rickettsiae cause infection in humans, domestic animals and wildlife (Maina et al. 2014). At least nine *Rickettsia* spp. of the SFG had been detected in XUAR (LopezVelez et al. 2015; Guo 2017). According to previous studies, *Rickettsia sibirica* subsp. *sibirica*, responsible for Siberian tick typhus, is widely distributed in China, Mongolia, Kazakhstan and Russia, and was also molecularly detected in Spain (Guo 2017; Parola et al. 2013; Byambaa et al. 2008; Palomar et al. 2012). *Rickettsia raoultii*, the causative agent of tick-borne lymphadenopathy (TIBOLA) or *Dermacentor*-borne necrosis erythema lymphadenopathy (DEBONEL) (Mediannikov et al. 2008), is prevalent in Mongolia, Europe and the Russian Federation (Oteo and Portillo 2012; Boldbaatar et al. 2017). These examples justify the importance of studying rickettsiae with broad distribution range in Eurasia. Therefore, the aim of this study was to identify ticks infesting *S. undulatus*, and to molecularly identify *Rickettsia* species in these ticks.

Materials and methods

Tick sampling and identification

A total of 36 *S. undulates* were captured in Jinghe County (2023 m above sea level; 44°35'59N, 82°53'28E), near the wetlands around Ebinur Lake, in the Northwest region of XUAR in July 2017. For this purpose, Sherman traps (H.B. Sherman Traps, Tallahassee, FL, USA) were used, which were placed at the entrances of occupied burrows (Torres-Perez et al. 2010). Each survey site included 150 traps that were checked twice a day. Each trap was removed before nightfall and replaced on the survey site the following day.

Ticks were sampled from the entire body of each *S. undulatus*, and were first identified with stereomicroscope (LEICA M165 C) according to standard morphological keys (Hornok et al. 2017). Then the genomic DNA was extracted from all ticks by using the 96 flux automatic nucleic acid extraction instrument with a matching commercial kit (Cell & Tissue Kit, Biotek, Beijing, China) according to our previous report (Liu et al. 2018).

To confirm the morphological identification of tick species, two mitochondrial markers, the *16S rDNA* and the cytochrome *c* oxidase subunit I (*COI*) genes (Hornok et al. 2017), were analyzed according to five representative ticks for each tick species. The phylogenetic relationships among the representative tick specimens were inferred using MEGA 6.0 software. Twelve nucleotide sequences from our study have been deposited in the GenBank database (*16S rDNA*: MG656445, MH324406–MH324409; and *COI*: MH079424, MH279561).

Detection of rickettsial agents and sequence analyses

Six genetic markers, including 17-kilodalton antigen (*17-kDa*), surface cell antigen 4 (*sca4*), citrate synthase (*gltA*), cell surface antigen 1 (*sca1*), and outer membrane proteins A and B (*ompA* and *ompB*), were amplified from each sample to investigate the presence of SFG rickettsiae (Anstead and Chilton 2013a, b). The primers and PCR cycling conditions in this study are shown in Supplement Appendix Table 1. Each PCR assay included a negative control (distilled water instead of tick DNA template) and a positive control (with DNA from *R. raoultii* obtained from wetlands of Ebinur Lake in XUAR). Purification and sequencing of the PCR products were done as described above (Anstead and Chilton 2013a, b). Phylogenetic trees were constructed using the maximum-likelihood and neighbor-joining methods in MEGA 6.0 software (Guo et al. 2015).

Results

A total of 98 ticks (50 adult ticks and 48 nymphs) were collected from captured *S. undulatus*. According to morphological characters and the *16S rDNA* phylogenetic tree, ticks in this study were identified as *D. nuttalli*, *D. silvarum* and *I. kaiseri* (shown in Fig. 1). Based on analysis of *COI* haplotype, 1–5 nucleotide differences were found in comparison with *I. kaiseri* from Europe. The phylogenetic analysis indicated that *I. kaiseri* from *S. undulatus* was in an ancestral position to nine European haplotypes (“L to T”) (Hornok et al. 2017) (shown in Fig. 2).

Out of 50 adult ticks (26 *D. nuttalli*, 21 *D. silvarum* and 3 *I. kaiseri*) and 48 nymphs (27 *D. nuttalli*, 21 *D. silvarum*), 25 adults and 16 nymphs were positive for six *Rickettsia* genetic markers (*17-kDa*, *gltA*, *ompA*, *sca1*, *sca4* and *ompB*). Among them, *R. sibirica* subsp. *sibirica* was found in nineteen *D. nuttalli* (8 nymphs and 11 adult ticks), 13 *D. silvarum* (6 nymphs and 7 adult ticks) and an adult *I. kaiseri*. In addition, *R. raoultii* was found in five *D. nuttalli* (one nymph and four adult ticks), three *D. silvarum* (one nymph and two adult ticks) (shown in Fig. 3; Table 1). As indicated by five genetic markers (*17-kDa*, *gltA*, *sca1*, *sca4* and *ompB*), *R. sibirica* subsp. *sibirica* in this study had sequence similarities in the range of 98.7–100% when compared to *R. sibirica* from *D. nuttalli* infesting sheep in Jimunai County, XUAR (Guo 2017). In addition, *R. raoultii* had 99.1–100% similarity compared to *R. raoultii* strain Khabarovsk (CP010969). The detailed similarities and divergences of the sequences from this study are shown in the Supplement Appendix Table 2.




Fig. 1 The phylogenetic tree inferred from the *16S rDNA* sequences of representative tick specimens. The evolutionary history was inferred using the maximum-likelihood method. The new sequences provided by the present study are indicated by a black triangle (containing the accession number). The phylogenetic analyses were conducted using MEGA 6.0 software

Discussion

Spermophilus undulatus is a burrowing mammalian species, inhabiting mountain areas 1600–3000 m above sea level (a.s.l.), with habitats along wetter front hills, forest margins and river valleys (Wang and Yang 1983). Previously, *D. marginatus*, *D. nuttalli*, *Rh. schulzei* and *Ixodes persulcatus* were sampled from *S. undulatus* (Wang and Yang 1983).

Dermacentor silvarum, a three-host tick species, is widely distributed in North China, Russia and Mongolia (Deng and Jiang 1991; Kulik and Vinokurova 1983). The adult ticks parasitize cattle, horses, sheep, goats, hares and hedgehogs, and the larvae and nymphs mainly infest voles, squirrels and birds (Wang and Yang 1983). On the other hand, *I. kaiseri* occurs in the western Palearctic, including Germany, Hungary, southern Moldavia, Ukraine, Crimea, Romania, Egypt, Israel, Azerbaijan and Georgia, where it typically infests badgers, foxes, steppe polecats, raccoon dogs, common hedgehogs and domestic dogs (ZipcodeDev Team 2018; Hornok et al. 2017). In the present study, adult ticks (*D. nuttalli*, *D. silvarum* and *I. kaiseri*) and nymphs (*D. nuttalli* and *D. silvarum*) were found on *S. undulatus* in northwestern China. Our results extend the host range of *D. silvarum* and *I. kaiseri*. The ancestral phylogenetic position of its *COI* haplotype from China supports that the genetic diversity of *I. kaiseri* might be related to various geographic locations (Hornok et al. 2017).

To date, *R. sibirica*, including *R. sibirica* subsp. *sibirica* and *R. sibirica* subsp. *mongolotimonae*, have been reported from *Hyalomma anatolicum*, *Hyalomma truncatum*, *Hy.*




Fig. 2 Phylogenetic tree based on the *COI* gene, including sequences obtained in this study and representative sequences of other *Ixodes* spp.

asiaticum, *Rh. pusillus*, *Dermacentor sinicus*, *D. nuttalli*, *D. marginatus*, *D. reticulatus*, *D. silvarum*, *Haemaphysalis yeni*, *Haemaphysalis concinna* and *I. persulcatus* (Parola et al. 2013). In XUAR, *R. sibirica* subsp. *sibirica* has already been isolated from *D. nuttalli* in Jinghe County in 1974 (Kong et al. 1982). However, our finding is the first molecular evidence on the presence of *R. sibirica* subsp. *sibirica* in *I. kaiseri* from *S. undulatus*.

Another *Rickettsia* species, *R. raoultii*, is highly prevalent in XUAR and its neighboring countries. Among the others, it was detected in *Haemaphysalis erinacei* from marbled polecats in the Ebinur Lake wetlands (189 m above sea level; 82°48'51E 45°04'22N) in northwest China in 2014 (Anstead and Chilton 2013a, b). Jinghe County, neighboring Ebinur Lake wetlands, has a similar geographic habitat. In the present study, *R. raoultii* was detected in *D. nuttalli* and *D. silvarum* from *S. undulatus*. These findings suggest that *D. nuttalli*, *D. silvarum* and *I. kaiseri* parasitizing *S. undulatus* may serve as reservoirs and carriers for *R. raoultii* and *R. sibirica*.

Jinghe County, with the largest density of *S. undulatus* in China (10–50/hectare), was listed as a main *Marmot baibacina*–*S. undulatus* plague focus since 1967 (Wang and Yang 1983). *Yersinia pestis* was isolated from 3.22% (66/2051) of *S. undulatus* (Zhang and Sheng 1991). In our current study, the rickettsial *17-kDa* gene was detected from two




Fig. 3 Phylogenetic tree of the 17-kDa-ompA-gltA-sca4-sca1-ompB concatenated sequence of rickettsial agents in adult ticks and nymphs

Table 1 The prevalence of *Rickettsia sibirica* and *R. raoultii* in *Dermacentor nuttalli*, *D. silvarum* and *Ixodes kaiseri* from *Spermophilus undulatus*

Tick species	<i>Rickettsia</i> species	Prevalence (%) in nymphs	Prevalence (%) in adults
<i>D. nuttalli</i>	<i>R. sibirica</i>	16.7 (8/48)	22.0 (11/50)
	<i>R. raoultii</i>	2.1 (1/48)	8.0 (4/50)
<i>D. silvarum</i>	<i>R. sibirica</i>	12.5 (6/48)	14.0 (7/50)
	<i>R. raoultii</i>	2.1 (1/48)	4.0 (2/50)
<i>I. kaiseri</i>	<i>R. sibirica</i>	0	6.0 (3/50)
	<i>R. raoultii</i>	0	0

naturally killed *S. undulatus* (data not shown). These findings indicate that several vector-borne pathogens co-circulate in Jinghe County, and imply potential risks associated with

populations of *S. undulatus*. In the future, it is necessary to confirm whether *D. nuttalli*, *D. silvarum* and *I. kaiseri* infesting *S. undulatus* could serve as vectors, transmitting rickettsial agents to *S. undulatus*.

Conclusions

Dermacentor nuttalli, *D. silvarum* and *I. kaiseri* were sampled from *S. undulatus* near the China-Kazakhstan border, and *I. kaiseri* in this study had an ancestral *COI* haplotype when compared to conspecific European sequences. *Rickettsia raoultii* was detected in the adult and nymph stages of *D. nuttalli* and *D. silvarum*, and *R. sibirica* also in one *I. kaiseri* adult. These findings extend our knowledge on the range of tick species infesting *S. undulatus* and harboring rickettsial agents.

Acknowledgements The authors are grateful to Dr. Jinliang Sheng (College of Animal Science and Technology) for capturing *Spermophilus undulatus*.

Author contributions SZ, MY, MJ and YW conceived and designed the study. BY, SZ, WY, BW and WH processed the samples and performed molecular and phylogenetic analyses. SZ and SH contributed to writing the manuscript. All authors read and approved the final version of the manuscript.

Funding This research was supported in part by the National Key Research and Development Program of China (2018ZX10101002-002-007 and 2017ZX10304402-002-005), National Natural Science Foundation of China (81560338 and U1503283), and International Scientific and Technological Cooperation in Bingtuan (2016AH001).

Data Availability The datasets supporting the conclusions of this article are available in GenBank (National Center for Biotechnology Information) [unique persistent identifier and hyperlink to datasets in <http://www.ncbi.nlm.nih.gov/genbank/>].

Compliance with ethical standards

Conflict of interest The authors declare that they have no competing interests.

Ethics approval and informed consent This study was approved by the Animal Ethics Committee of Shihezi University (Approval No. AECSU2017-22).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Anstead CA, Chilton NB (2013a) A novel *Rickettsia* species detected in Vole Ticks (*Ixodes angustus*) from Western Canada. *Appl Environ Microbiol* 79:7583–7589
- Anstead CA, Chilton NB (2013b) Detection of a novel Rickettsia, (Alphaproteobacteria: Rickettsiales) in rotund ticks (*Ixodes kingi*) from Saskatchewan, Canada. *Ticks Tick Borne Dis* 4:202–206
- Boldbaatar B, Jiang RR, von Fricken ME, Lkhagvatseren S, Nymadawa P, Baigalmaa B, Wang YW, Anderson BD, Jiang JF, Gray GC (2017) Distribution and molecular characteristics of rickettsiae found in ticks across Central Mongolia. *Parasit Vectors* 10(1):61
- Byambaa B (2008) Nature-focal rickettsioses in Mongolia two decades of Russian–Mongolian scientific collaboration. *Vestn Ross Akad Med Nauk* 7:44–45

- Demina TV, Tkachev SE, Kozlova IV, Doroshchenko EK, Lisak OV, Suntsova OV, Verkhovina MM, Dzhi-oev YP, Paramonov AI, Tikunov AY, Tikunova NV, Zlobin VI, Ruzek D (2017) Comparative analysis of complete genome sequences of European subtype tick-borne encephalitis virus strains isolated from *Ixodes persulcatus*, ticks, long-tailed ground squirrel (*Spermophilus undulatus*), and human blood in the Asian part of Russia. *Ticks Tick Borne Dis* 8:547–553
- Deng GF, Jiang ZJ (1991) Economic insect fauna of China, Fasc 39, Acari: Ixodidae, 1st edn. Science Press, Beijing
- Guo LP (2017) Distribution and molecular characteristics of Tick species and tick-borne Rickettsiae, Xinjiang border areas. Shihezi University, Shihezi
- Guo LP, Mu LM, Xu J, Jiang SH, Wang AD, Chen CF, Guo G, Zhang WJ, Wang YZ (2015) Rickettsia raoultii in Haemaphysalis erinacei from marbled polecats, China–Kazakhstan border. *Parasit Vectors* 8:461
- Hornok S, Sándor AD, Beck R, Farkas R, Beati L, Kontschán J, Takács N, Földvári G, Silaghi C, Meyer-Kayser E, Hodžić A, Tomanović S, Abdullah S, Wall R, Estrada-Peña A, Duscher GG, Plantard O (2017) Contributions to the phylogeny of *Ixodes* (*Pholeoixodes*) *canisuga*, *I. (Ph.) kaiseri*, *I. (Ph.) hexagonus*, and a simple pictorial key for the identification of their females. *Parasit Vectors* 10:545
- Kong SM, Cao GY, Zhang YF, Chen GZ (1982) Isolation and identification of Rickettsia from tick-borne spotted fever group in Jinghe County of Xinjiang. *Mic Bull* 9:13–15
- Kulik IL, Vinokurova NS (1983) Range of the tick *Dermacentor silvarum* in the USSR. *Med Parazitol (Mosk)* 52:23–28
- Liu XF, Yan B, Wang Q, Jiang MM, Tu CC, Chen CF, Hornok S, Wang Y (2018) Babesia vesperuginis in Common Pipistrelle (*Pipistrellus pipistrellus*) and the Bat Soft Tick Argas vespertilionis in the People's Republic of China. *J Wildl Dis* 54:419–421
- LopezVelez R, Palomar AM, Oteo JA, Norman FF, Pérez-Molina JA, Portillo A (2015) Novel *Candidatus rickettsia* species detected in nostril tick from human, Gabon, 2014. *Emerg Infect Dis* 21:325–327
- Maina AN, Jiang J, Omulo SA, Cutler SJ, Ade F, Ogola E, Feikin DR, Njenga MK, Cleaveland S, Mpoke S, Ng'ang'a Z, Breiman RF, Knobel DL, Richards AL (2014) High prevalence of *Rickettsia africae* variants in *Amblyomma variegatum* ticks from domestic mammals in rural western Kenya: implications for human health. *Vector Borne Zoonotic Dis* 14:693–702
- Mediannikov O, Matsumoto K, Samoylenko I, Drancourt M, Roux V, Rydkina E, Davoust B, Tarasevich I, Brouqui P, Fournier PE (2008) *Rickettsia raoultii* sp. nov. a spotted fever group rickettsia associated with Dermacentor ticks in Europe and Russia. *Int J Syst Evol Microbiol* 58:1635–1639
- Oteo JA, Portillo A (2012) Tick-borne rickettsioses in Europe. *Ticks Tick Borne Dis* 3:271–278
- Palomar AM, Santibañez P, Mazuelas D, Roncero L, Santibañez S, Portillo A, Oteo JA (2012) Role of birds in dispersal of etiologic agents of tick-borne zoonoses, Spain, 2009. *Emerg Infect Dis* 18:1188–1191
- Parola P, Paddock CD, Socolovschi C, Labruna MB, Mediannikov O, Kernif T, Abdad MY, Stenos J, Bitam I, Fournier PE, Raoult D (2013) Update on tick-borne rickettsioses around the world: a geographic approach. *Clin Microbiol Rev* 26:657–702
- Ramoslara N, Koprowski JL, Kryštufek B, Hoffmann IE (2014) *Spermophilus citellus* (Rodentia: Sciuridae). *Mamm Species* 913:71–87
- Torres-Perez F, Wilson L, Collinge SK, Harmon H, Ray C, Medina RA, Hjelle B (2010) Sin Nombre virus infection in field workers, Colorado, USA. *Emerg Infect Dis* 16:308–310
- Wang SB, Yang GY (1983) Rodent fauna of Xinjiang. Iedn Xinjiang People's publishing house, Urumqi
- Zhang HQ, Sheng GJ (1991) Xinjiang long tail mouse The plague overview about *Spermophilus undulatus* in Xinjiang Uygur Autonomous Region. *Endem Dis Bull* 5:114–117
- Zhao SS, Pulati Y, Yin XP, Li W, Wang BJ, Yang K, Chen CF, Wang YZ (2017) Wildlife plague surveillance near the China-Kazakhstan border: 2012–2015. *Transbound Emerg Dis* 64(6):48–51
- Zipcodezoo. ZipcodeDev Team (2018). <http://zipcodezoo.com/>

Affiliations

Shuo Zhao¹ · Meihua Yang² · Mengmeng Jiang³ · Bin Yan¹ · Shanshan Zhao¹ ·
Wumei Yuan¹ · Baoju Wang⁴ · Sándor Hornok⁵ · Yuanzhi Wang¹

Shuo Zhao
954855721@qq.com

Meihua Yang
1328251980@qq.com

Mengmeng Jiang
577674273@qq.com

Bin Yan
1262321593@qq.com

Shanshan Zhao
1393819696@qq.com

Wumei Yuan
348781741@qq.com

Baoju Wang
bjwang73@163.com

Sándor Hornok
hornok.sandor@univet.hu

¹ Department of Basic Medicine, School of Medicine, Shihezi University, Shihezi, Xinjiang, China

² Department of Forestry, College of Agriculture, Shihezi University, Shihezi, Xinjiang, China

³ Department of Veterinary Medicine, College of Animal and Science, Shihezi University, Shihezi, Xinjiang, China

⁴ Department of Infectious Diseases, Union Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

⁵ Department of Parasitology and Zoology, University of Veterinary Medicine, Budapest, Hungary