

EDITORIAL

WEIGHT LOSS IS A MAJOR CAUSE OF FRAILTY

B. FOUGÈRE^{1,2,3}, J.E. MORLEY³

1. Gérontopôle, Centre Hospitalier Universitaire de Toulouse, Toulouse, France; 2. Inserm UMR 1027, Université de Toulouse III Paul Sabatier, Toulouse, France; 3. Division of Geriatric Medicine, Saint Louis University School of Medicine, St. Louis, Missouri, USA. Corresponding author: John E. Morley, MB, BCh, Division of Geriatric Medicine, Saint Louis University School of Medicine, 1402 S. Grand Blvd., M238, St. Louis, MO 63104, Email: john.morley@health.slu.edu

The concept of frailty as a centrally important geriatric syndrome is now well established (1-4). Frailty is defined as a syndrome where an older person is vulnerable to becoming functionally impaired or dying when exposed to a stressor (5). A popular approach to identifying frailty is the physical frailty phenotype as first described by Fried et al. (6). This was developed out of data from the Cardiovascular Health Study and defined frailty as having 3 or more of: (i) unintentional weight loss (10 lbs or 4.5 kg in past year); (ii) self-reported exhaustion, (iii) weakness (grip strength); (iv) slow walking speed and (v) low physical activity. To accommodate the busy schedule of clinical practitioners two rapid assessments for physical frailty have been developed, namely the Gérontopôle Frailty Screening Tool (GFST) (7, 8) and the FRAIL (9-11).

While malnutrition (weight loss) is only one component of the physical phenotype scale, it is also a cause of the other 4 components (12-16). Depression is a common cause of anorexia/weight loss and fatigue in older persons, providing an important connection between these symptoms (17-20). Depression is a major cause of frailty in older persons (21). Loss of muscle mass is associated with fatigue (22).


Weight loss leads not only to loss of fat but also loss of muscle lean mass and bone (23). Obviously, muscle lean mass loss is a major cause of sarcopenia which includes loss of walking speed and/or grip strength (24-28). There is a major overlap between sarcopenia and frailty (29-34). Nutritional supplementation can enhance muscle mass and function in persons with sarcopenia (35, 36) and in frail persons (37-39).

A systematic review has shown that malnutrition as measured by the Mini Nutritional Assessment (MNA) is strongly associated with frailty (40). However, it should be recognized that all frail persons are not malnourished (41-44). When frailty and malnutrition co-exist they are highly predictive of mortality in functionally impaired persons (45). The FRAIL-NH which was developed for functionally impaired persons in the nursing home has been shown to be highly predictive of further functional impairment and mortality (46-48). Like the FRAIL phenotype, it is strongly dependent on weight loss and problems with eating.

There is evidence that diet quality in later life may influence the development of frailty (49-52). Adherence to the Mediterranean diet appears to be particularly protective against developing frailty (53-56). This effect appears to be related predominantly to a higher intake in polyphenols (57).

Overall it would appear that the anorexia of aging (58-60) and associated weight loss represents an important role in the pathophysiology of frailty (Figure 1). For this reason, we recommend early screening for anorexia utilizing the Simplified Nutrition Assessment Questionnaire (SNAQ) and/or the Mini Nutritional Assessment (MNA) (61-65). Simple approaches such as the St. Louis University Rapid Geriatric Assessment or the Gérontopôle Frailty Screening Tool (GFST) need to be incorporated into all primary care practices (66-70).

Figure 1
The Interrelationship Between Weight Loss and Frailty



References

1. Dent E, Lien C, Lim WS, et al. The Asia-Pacific clinical practice guidelines for the management of frailty. *J Am Med Dir Assoc* 2017;18:564-575.
2. Morley JE, Vellas B, van Kan GA, Anker SD, et al. Frailty consensus: A call to action. *J Am Med Dir Assoc* 2013;14:392-397.
3. Sinclair A, Vellas B. (Editorial: Frailty – an emerging high impact complication of diabetes requiring specific management approaches. *J Nutr Health Aging* 2017;21:193-194.
4. Cesari M, Prince M, Thiyagarajan JA, De Carvalho IA, et al. Frailty: An emerging public health priority. *J Am Med Dir Assoc* 2016;17:188-192.
5. Morley JE, von Haehling S, Anker SD, Vellas B. From sarcopenia to frailty: A road less traveled. *J Cachexia Sarcopenia Muscle* 2014;5:5-8.
6. Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: Evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56:M146-156.
7. Vellas B. Implementing frailty screening, assessment, and sustained intervention: The experience of the Gérontopôle. *J Nutr Health Aging* 2015;19:673-680.
8. Hoogendijk EO, van Kan GA, Guyonnet S, Vellas B, Cesari M. Components of the frailty phenotype in relation to the frailty index: Results from the Toulouse Frailty platform. *J Am Med Dir Assoc* 2015;16:855-859.
9. Morley JE, Malmstrom TK, Miller DK. A simple frailty questionnaire (FRAIL) predicts outcomes in middle aged African Americans. *J Nutr Health Aging* 2012;16:601-608.

WEIGHT LOSS IS A MAJOR CAUSE OF FRAILTY

10. Gardiner PA, Mishra GD, Dobson AJ. Validity and responsiveness of the FRAIL scale in a longitudinal cohort study of older Australian women. *J Am Med Dir Assoc* 2015;16:781-783.
11. Li Y, Zou Y, Wang S, et al. A pilot study of the FRAIL scale on predicting outcomes in Chinese elderly people with type 2 diabetes. *J Am Med Dir Assoc* 2015;16:714.e7-714.e12.
12. Alesi ER, del Fabbrio E. Opportunities for targeting the fatigue-anorexia-cachexia symptom cluster. *Cancer J* 2014;20:325-329.
13. Soenen S, Chapman IM. Body weight, anorexia, and undernutrition in older people. *J Am Med Dir Assoc* 2013;14:642-648.
14. Nguyen TN, Cumming RG, Hilmer SN. A review of frailty in developing countries. *J Nutr Health Aging* 2015;19:941-946.
15. Morley JE. Anorexia of ageing: A key component in the pathogenesis of both sarcopenia and cachexia. *J Cachexia Sarcopenia Muscle* 2017;8:523-526.
16. Anker SD, Morley JE. Cachexia: A nutritional syndrome? *J Cachexia Sarcopenia Muscle* 2015;6:269-271.
17. Thakur M, Blazer DG. Depression in long-term care. *J Am Med Dir Assoc* 2008;9:82-87.
18. Tamura BK, Bell CL, Masaki KH, Amella EJ. Factors associated with weight loss, low BMI, and malnutrition among nursing home patients: A systematic review of the literature. *J Am Med Dir Assoc* 2013;14:649-655.
19. Luo H, Tang JY, Wong GH, et al. The effect of depressive symptoms and antidepressant use on subsequent physical decline and number of hospitalizations in nursing home residents: A 9-year longitudinal study. *J Am Med Dir Assoc* 2015;16:1048-1054.
20. Almeida OP, Hankey GJ, Yeap BB, et al. Depression among nonfrail old men is associated with reduced physical function and functional capacity after 9 years follow-up: The Health in Men Cohort study. *J Am Med Dir Assoc* 2017;18:65-69.
21. Almeida OP, Hankey GJ, Yeap BB, et al. Depression, frailty, and all-cause mortality: A cohort study of men older than 75 years. *J Am Med Dir Assoc* 2015;16:296-300.
22. Anker SD, Sharma R. The syndrome of cardiac cachexia. *Int J Cardiol* 2002;85:51-66.
23. Huo YR, Suriyaarachchi P, Gomez F et al. Comprehensive nutritional status in sarcopenic osteoporotic older fallers. *J Nutr Health Aging* 2015;19:474-480.
24. Masanes F, Rojano I, Luque X, Salva A, et al. Cut-off points for muscle mass – not grip strength or gait speed – determine variations in sarcopenia prevalence. *J Nutr Health Aging* 2017;21:825-829.
25. Balogun S, Winzenberg T, Wills K, et al. Prospective associations of low muscle mass and function with 10-year falls risk, incident fracture and mortality in community-dwelling older adults. *J Nutr Health Aging* 2017;21:843-848.
26. Morley JE. Sarcopenia in the elderly. *Fam Pract* 2012;29(Suppl 1):i44-i48.
27. Chen LK, Lee WJ, Peng LN, et al. Recent advances in sarcopenia research in Asia: 2016 Update from the Asian Working Group for Sarcopenia. *J Am Med Dir Assoc* 2016;17:767.e1-7.
28. Shimada H, Makizako H, Doi T, Tsutsumimoto K, Suzuki T. Incidence of disability in frail older persons with or without slow walking speed. *J Am Med Dir Assoc* 2015;16:690-696.
29. Molina-Garrido MJ, Guillen-Ponce C, Fernandez-Felix BM. Relationship between sarcopenia and frailty in a Spanish cancer in the elderly unit: The ONCOSARCO project. *J Am Med Dir Assoc* 2016;17:760-761.
30. Liccini A, Malmstrom TK. Frailty and sarcopenia as predictors of adverse health outcomes in persons with diabetes mellitus. *J Am Med Dir Assoc* 2016;17:846-851.
31. Woo J, Yu R, Wong M, Yeung F, Wong M, Lum C. Frailty screening in the community using the FRAIL scale. *J Am Med Dir Assoc* 2015;16:412-419.
32. Reijnierse EM, Trappenburg MC, Blauw GJ, et al. Common ground? The concordance of sarcopenia and frailty definitions. *J Am Med Dir Assoc* 2016;17:371.e7-12.
33. Yoshimura Y, Uchida K, Jeong S, Yamaga M. Effects of nutritional supplements on muscle mass and activities of daily living in elderly rehabilitation patients with decreased muscle mass: A randomized controlled trial. *J Nutr Health Aging* 2016;20:185-191.
34. Fougère B, Vellas B, Abellan van Kan G, Cesari M. Identification of biological markers for better characterization of older subjects with physical frailty and sarcopenia. *Translational Neuroscience* 2015;6:103-10.
35. Bauer JM, Verlaan S, Bautmans I, et al. Effects of a vitamin D and leucine-enriched whey protein nutritional supplement on measures of sarcopenia in older adults, the PROVIDE study: A randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc* 2015;16:740-747.
36. Brown JC, Harhay MO, Harhay MN. Appendicular lean mass and mortality among prefrail and frail older adults. *J Nutr Health Aging* 2017;21:342-345.
37. Ng TP, Feng L, Nyunt MS, et al. Nutritional, physical, cognitive, and combination interventions and frailty reversal among older adults: A randomized controlled trial. *Am J Med* 2015;128:1225-1236.e1.
38. Tarazona-Santabalbina FJ, Gomez-Cabrera MC, Perez-Ros P, et al. A multicomponent exercise intervention that reverses frailty and improves cognition, emotion, and social networking in the community-dwelling frail elderly: A randomized clinical trial. *J Am Med Dir Assoc* 2016;17:426-433.
39. Abizanda P, Lopez MD, Garcia VP, et al. Effects of an oral nutritional supplementation plus physical exercise intervention on the physical function, nutritional status, and quality of life in frail institutionalized older adults: The ACTIVNES study. *J Am Med Dir Assoc* 2015;16:439.e9-439.e16.
40. Lorenzo-Lopez L, Maseda A, de labra C, et al. Nutritional determinants of frailty in older adults: A systematic review. *BMC Geriatr* 2017;17:108.
41. Lilamand M, Kelaidit E, Cesari M; Toulouse Frailty Platform Team. Validation of the Mini Nutritional Assessment-Short Form in a population of frail elders without disability. Analysis of the Toulouse frailty platform population in 2013. *J Nutr Health Aging* 2015;19:570-574.
42. Wei K, Nyung MSZ, Gao Q, Wee SL, Ng TP. Frailty and malnutrition: Related and distinct syndrome prevalence and association among community-dwelling older adults: Singapore Longitudinal Ageing studies. *J Am Med Dir Assoc* 2017;18:____ doi: 10.1016/j.jamda.2017.06.017. (In press).
43. Bollwein J, Volkert D, Kiekmann R, et al. Nutritional status according to the mini nutritional assessment (MNA®) and frailty in community dwelling older persons: A close relationship. *J Nutr Health Aging* 2013;17:351-356.
44. Dent E, Lien C, Lim WS, et al. The Asia-Pacific clinical practice guidelines for the management of frailty. *J Am Med Dir Assoc* 2017;18:564-575.
45. Kamo T, Takayama K, Ishii H, Suzuki K, Eguchi K, Nishida Y. Coexisting severe frailty and malnutrition predict mortality among the oldest old in nursing homes: A 1-year prospective study. *Arch Gerontol Geriatr* 2017;70:99-104.
46. Kaehr E, Visvanathan R, Malmstrom TK, Morley JE. Frailty in nursing homes: The FRAIL-NH scale. *J Am Med Dir Assoc* 2015;16:87-89.
47. Kaehr EW, Pape LC, Malmstrom TK, Morley JE. FRAIL-NH predicts outcomes in long term care. *J Nutr Health Aging* 2016;20:192-198.
48. Luo H, Lum TY, Wong GH, Kwan JS, Tang JY, Chi I. Predicting adverse health outcomes in nursing homes: A 9-year longitudinal study and development of the FRAIL-Minimum Data Set (MDS Quick Screening Tool). *J Am Med Dir Assoc* 2015;16:1042-1047.
49. Bonnefoy M, Berrut G, Lesourd B, et al. Frailty and nutrition: Searching for evidence. *J Nutr Health Aging* 2015;19:250-257.
50. Bollwein J, Volkert D, Kiekmann R, Kaiser MJ, et al. Nutritional status according to the mini nutritional assessment (MNA®) and frailty in community dwelling older persons: A close relationship. *J Nutr Health Aging* 2013;17:351-356.
51. Pilleron S, Ajana S, Jutand MA, et al. Dietary patterns and 12-year risk of frailty: Results from the Three-City Bordeaux study. *J Am Med Dir Assoc* 2017;18:169-175.
52. Chan R, Leung J, Woo J. Dietary patterns and risk of frailty in Chinese community-dwelling older people in Hong Kong: A prospective cohort study. *Nutrients* 2015;7:7070-7084.
53. León-Muñoz, Guallar-Castillón P, López-García E, Rodríguez-Artalejo F. Mediterranean diet and risk of frailty in community-dwelling older adults. *J Am Med Dir Assoc* 2014;15:899-903.
54. Talegawkar SA, Bandinelli S, Bandeen-Roche K, et al. A higher adherence to a Mediterranean-style diet is inversely associated with the development of frailty in community-dwelling elderly men and women. *J Nutr* 2012;142:2161-2166.
55. Rahi B, Ajana S, Tabue-Teguio M, et al. High adherence to a Mediterranean diet and lower risk of frailty among French older adult community-dwellers: Results from the Three-City-Bordeaux study. *Clin Nutr* May 2017;31. Doi: 10.1016/j.clnu.2017.05.020 [Epub ahead of print].
56. Fougère B, Mazzuco S, Spagnolo P, Guyonnet S, Vellas B, Cesari M, Gallucci M. Association between the Mediterranean-Style Dietary Pattern Score and Physical Performance: results from TRELONG study. *JNH* 2016;20:415-9.
57. Urpi-Sarda M, Andres-Lacueva C, Rabassa M, et al. The relationship between urinary total polyphenols and the frailty phenotype in a community-dwelling older population: The INCHIANTI study. *J Gerontol A Biol Sci Med Sci* 2015;70:1141-1147.
58. Mir F, Zafar F, Morley JE. Anorexia of aging: Can we decrease protein energy undernutrition in the nursing home? *J Am Med Dir Assoc* 2013;14:77-79.
59. Morley JE. Pathophysiology of the anorexia of aging. *Curr Opin Clin Nutr Metab Care* 2013;16:27-32.
60. Morley JE. Anorexia of aging: A true geriatric syndrome. *J Nutr Health Aging* 2012;16:422-425.
61. Tokudome Y, Okumura K, Kumagai Y, et al. Development of the Japanese version of the Council on Nutrition Appetite Questionnaire and its simplified versions, and evaluation of their reliability, validity, and reproducibility. *J Epidemiol* Feb 2. 2017;Doi: 10.1016/j.je.2016.11.002 [Epub ahead of print].
62. Pilgrim AL, Baylis D, Jameson KA, et al. Measuring appetite with the Simplified Nutritional Appetite Questionnaire identifies hospitalised older people at risk of worse health outcomes. *J Nutr Health Aging* 2016;20:3-7.
63. Schrader E, Grosch E, Bertsch T, Sieber CC, Volkert D. Nutritional and functional status in geriatric day hospital patients – MNA sort form versus full MNA. *J Nutr Health Aging* 2016;20:918-926.
64. Van Wissen J, van Stijn MF, Doodeman HJ, Houdijk AP. Mini nutritional assessment and mortality after hip fracture surgery in the elderly. *J Nutr Health Aging* 2016;20:964-968.
65. Kellest J, Kyle G, Ispopoulos C, Naunton M. Nutrition screening practices amongst Australian residential aged care facilities. *J Nutr Health Aging* 2016;20:1040-1044.
66. Morley JE. Rapid Geriatric Assessment: Secondary prevention to stop age-associated

THE JOURNAL OF NUTRITION, HEALTH & AGING©

- disability. *Clin Geriatr Med* 2017;33:431-440.
67. Morley JE, Arai H, Cao L, Dong B, Merchant RA, et al. Integrated care: Enhancing the role of the primary health care professional in preventing functional decline: A systematic review. *J Am Med Dir Assoc* 2017;18:489-494.
68. Morley JE, Little MO, Berg-Weger M. Rapid geriatric assessment: A tool for primary care physicians. *J Am Med Dir Assoc* 2017;18:195-199.
69. Tavassoli N, Guyonnet S, Abelland van Kan G, et al. Description of 1,108 older patients referred by their physician to the "Geriatric Frailty Clinic (G.F.C.) for assessment of frailty and prevention of disability" at the Gerontopole. *J Nutr Health Aging* 2014;18:457-464.
70. Vellas B, Balardy L, Gillette-Guyonnet S, et al. Looking for frailty in community-dwelling older persons: The Gerontopole Frailty Screening tool (FGST). *J Nutr Health Aging* 2013;17:629-631.