

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
21 December 2017 (21.12.2017)



(10) International Publication Number
WO 2017/216685 A1

(51) International Patent Classification:

C07D 491/14 (2006.01) A61P 31/12 (2006.01)
A61K 31/4741 (2006.01)

(21) International Application Number:

PCT/IB2017/053365

(22) International Filing Date:

07 June 2017 (07.06.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/351,037 16 June 2016 (16.06.2016) US
62/479,052 30 March 2017 (30.03.2017) US

(71) Applicant: **NOVARTIS AG** [CH/CH]; Lichtstrasse 35,
4056 Basel (CH).

(72) Inventors: **FU, Jiping**; Novartis Institutes for BioMedical
Research Inc., 5300 Chiron Way, Emeryville, CA 94608
(US). **JIN, Xianming**; Novartis Institutes for BioMedical
Research Inc., 5300 Chiron Way, Emeryville, CA 94608
(US). **LEE, Patrick**; Novartis Institutes for BioMedical
Research Inc., 5300 Chiron Way, Emeryville, CA 94608
(US). **LU, Peichao**; Novartis Institutes for BioMedical Re-
search Inc., 5300 Chiron Way, Emeryville, CA 94608 (US).
YOUNG, Joseph Michael; Novartis Institutes for Bio-
Medical Research Inc., 5300 Chiron Way, Emeryville, CA
94608 (US).

(74) Agent: **NOVARTIS AG**; Lichtstrasse 35, 4056 Basel
(CH).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR,
KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))

(54) Title: PENTACYCLIC PYRIDONE COMPOUNDS AS ANTIVIRALS



(57) Abstract: The invention provides compounds of Formula (I), wherein R⁴ and R⁵ are joined together to form a ring as described herein, along with pharmaceutically acceptable salts, pharmaceutical compositions containing such compounds, as well as methods to use these compounds, salts and compositions for treating viral infections, particularly infections caused by hepatitis B virus, and for reducing the occurrence of serious conditions associated with HBV.



WO 2017/216685 A1

PENTACYCLIC PYRIDONE COMPOUNDS AS ANTIVIRALS

FIELD OF THE INVENTION

The present invention relates to novel pentacyclic pyridone compounds that are inhibitors of hepatitis virus replication, and are thus useful to treat viral infections, and particularly hepatitis B virus (HBV). The invention provides novel pentacyclic pyridone compounds as disclosed herein, pharmaceutical compositions containing such compounds, and methods of using these compounds and compositions in the treatment and prevention of HBV infections.

BACKGROUND

Globally, over 400 million people are chronically infected with hepatitis B virus (HBV), and more than 12 million reside in the United States alone. Of those chronically infected patients, up to 40 percent will eventually develop complications of liver failure from cirrhosis or development of hepatocellular carcinoma (HCC). HBV belongs to the family of Hepadnaviridae, a group of small hepatotropic DNA viruses that replicate through the reverse transcription of an RNA intermediate. The 3.2-kb HBV genome in viral particles is in a circular, partially double-stranded DNA conformation (relaxed circular DNA or rcDNA). The HBV genome consists of four overlapping open reading frames (ORFs), which encode for the core, polymerase (Pol), envelope, and X proteins. rcDNA is transcriptionally inert and must be converted into covalently closed circular DNA (cccDNA) in the nucleus of infected cells before viral RNAs can be transcribed. cccDNA is the only template for HBV transcription and, because HBV RNA templates genomic reverse transcription, its persistence is required for persistent infection.

The envelope of HBV comprises a mixture of surface antigen proteins (HBsAg). The HBsAg coat is a mixture of three overlapping proteins: all three share a common region, which corresponds to the smallest of the three proteins (SHBsAg). The mixture consists mostly of SHBsAg, but also includes Medium HBsAg, which comprises SHBsAg plus an additional polypeptide segment, and Large HBsAg, which comprises M HBsAg plus another added polypeptide segment. In addition to forming the infectious virion particle, the S, M and L HBsAg proteins also assemble into a subviral particle known as the 22-nm particle, which is not infectious but contains the same proteins that envelope the infectious virus particles. Indeed, these subviral, non-infectious particles have been used as a vaccine, since they contain the same antigenic surface proteins as the infectious HBV virion, and thus elicit antibodies that recognize the infectious agent. Interestingly, these subviral particles greatly outnumber infectious virions, and are believed to protect the infectious virions from the immune system of the infected host. By sheer numbers, they may act as

decoys, distracting immune responses from the infectious virus particles, but in addition they are reported to suppress the function of immune cells (monocytes, dendritic cells and natural killer cells) and may thus impair the immune response to HBV. Because these subviral particles protect infectious HBV from the host immune system, reducing the level of subviral particles has been recognized as a viable therapeutic approach. See, e.g., WO2015/113990.

One of the key diagnostic symptoms of chronic HBV is the high serum levels of the hepatitis B surface antigen (HBsAg). Clinical data in recent years suggest that sustained virologic response is often associated with on-treatment HBsAg decline during the early phase of the treatment as early as week 8, while sustained exposure to HBsAg and other viral antigens may lead to HBV-specific immune-tolerance. Chronic HB patients who experienced larger and faster decreases in serum HBsAg levels achieved significantly higher rate (~40%) of sustained virologic response as defined by sustained viral control post treatment.


Current treatment options for HBV include interferon therapies and nucleoside/nucleotide inhibitors of the viral DNA polymerase, such as entecavir and tenofovir. These focus on reduction in the level of viremia and toleration of hepatic dysfunction, and may have adverse side-effects and also select for drug-resistant virus variants during long term therapy. More importantly, these therapies cannot eradicate the intrahepatic HBV cccDNA pool in chronic hepatitis B patients or limit the transcription of HBsAg from the pre-existing cccDNA, nor do they affect the secretion of synthesized HBsAg into patients' blood to counteract the host innate immune response. As a result, these HBV treatments are in most cases life-long therapies, and discontinuation often leads to virological relapse.

Accordingly, there remains a need for more effective treatments for HBV, especially for treating chronic HBV infections (cHBV). The invention provides compounds that are believed to operate by suppression of the secretion of the 22 nm subviral particles containing HBsAg. These compounds are useful to treat HBV infections and to reduce the incidence of serious liver disorders caused by HBV infections.

SUMMARY

The present invention provides novel compounds that inhibit secretion of HBsAg from cells infected with hepatitis B virus and thereby reduce viral load and viral replication in patients having chronic HBV infection. Thus the compounds of the invention are suitable for treatment of patients with HBV, including cHBV.

In one aspect, the invention provides compounds of Formula (I):



(I) wherein:

R^1 is H, halo, or C_1 - C_3 alkyl;

R^2 is H, halo, CN, C_1 - C_3 alkyl or C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy;

R^3 is H, OH, halo, CN, C_1 - C_3 alkyl, C_3 - C_6 cycloalkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, or C_1 - C_3 haloalkoxy;

R^4 and R^5 taken together form a 5 or 6 membered ring that is fused to the phenyl ring to which R^4 and R^5 are attached, wherein the 5 or 6 membered ring contains one or two oxygen atoms as ring members and is optionally substituted by up to four groups independently selected from $-CN$, $-OR$, $-SR$, $-SO_2R$, and C_1 - C_4 alkyl optionally substituted with one or two groups selected from halo, $-OR$, $-SR$, $-SO_2R$, C_1 - C_3 haloalkoxy, CN, and $-N(R)_2$;

R^6 is H, halo, C_1 - C_3 alkoxy, or C_1 - C_6 alkyl, or is taken together with R^9 to form a ring;

R^7 is H, halo, C_1 - C_3 alkoxy, or C_1 - C_6 alkyl, or is taken together with R^9 to form a ring;

R^8 is H or C_1 - C_6 alkyl, or is taken together with R^9 to form a ring;

R^9 taken together with exactly one group selected from R^6 , R^7 and R^8 forms a 3-7 membered cycloalkyl ring or a 3-7 membered heterocyclic ring containing N, O or S as a ring member; wherein the cycloalkyl or heterocyclic ring is optionally substituted with up to three groups selected from R, $-OR$, $-NR_2$, halo, CN, COOR, $CONR_2$, and oxo;

W is $-COOR^*$, $-C(O)NH-SO_2R$, $-C(O)NH-SO_2NR_2$, 5-tetrazolyl, or 1,2,4-oxadiazol-3-yl-5(4H)-one;

R^* is H or C_1 - C_6 alkyl that is optionally substituted with one or two groups selected from halo, $-OR$, oxo, CN, $-NR_2$, COOR, and $CONR_2$;

R is independently selected at each occurrence from H and C₁-C₄ alkyl optionally substituted with one to three groups selected from halo, -OH, C₁-C₃ alkoxy, oxo, CN, -NH₂, -NH(C₁-C₃ alkyl), -N(C₁-C₃ alkyl)₂, and cyclopropyl;

and two R groups directly attached to the same atom, which may be C or N, can optionally be taken together to form a 3-6 membered ring that can optionally contain an added heteroatom selected from N, O and S as a ring member, and can be substituted by up to two groups selected from -OH, oxo, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

or a pharmaceutically acceptable salt thereof.

The invention also provides pharmaceutical compositions containing the novel compounds as well as methods to use the compounds and compositions to inhibit hepatitis B virus replication, and to treat disease conditions associated with or caused by HBV. Further objects of this invention are described in the following description and the examples.

DETAILED DESCRIPTION

For purposes of interpreting this specification, the following definitions will apply, and whenever appropriate, terms used in the singular will also include the plural.

Terms used in the specification have the following meanings unless the context clearly indicates otherwise:

As used herein, the term "subject" refers to an animal. In certain aspects, the animal is a mammal. A subject also refers to for example, primates (e.g., humans), cows, sheep, goats, horses, dogs, cats, rabbits, rats, mice, fish, birds and the like. In certain embodiments, the subject is a human. A "patient" as used herein refers to a human subject.

As used herein, the term "inhibition" or "inhibiting" refers to the reduction or suppression of a given condition, symptom, or disorder, or disease, or a significant decrease in the baseline activity of a biological activity or process.

As used herein, the term "treating" or "treatment" of any disease or disorder refers in one embodiment, to ameliorating the disease or disorder (i.e., slowing or arresting or reducing the development of the disease or at least one of the clinical symptoms thereof). In another embodiment "treating" or "treatment" refers to alleviating or ameliorating at least one physical parameter including those which may not be discernible by the patient. In yet another embodiment, "treating" or "treatment" refers to modulating the disease or disorder, either physically, (e.g., stabilization of a discernible symptom), physiologically, (e.g., stabilization of a physical parameter), or both. In yet another embodiment, "treating" or

"treatment" refers to preventing or delaying the onset or development or progression of the disease or disorder.

As used herein, the term "a," "an," "the" and similar terms used in the context of the present invention (especially in the context of the claims) are to be construed to cover both the singular and plural unless otherwise indicated herein or clearly contradicted by the context.

All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g. "such as") provided herein is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention otherwise claimed.

"Optionally substituted" means the group referred to can be substituted at one or more positions by any one or any combination of the radicals listed thereafter. The number, placement and selection of substituents is understood to encompass only those substitutions that a skilled chemist would expect to be reasonably stable; thus 'oxo' would not be a substituent on an aryl or heteroaryl ring, for example, and a single carbon atom would not have three hydroxy or amino substituents. Unless otherwise specified, optional substituents are typically up to four groups selected from halo, oxo, CN, amino, hydroxy, -C₁₋₃ alkyl, -OR*, -NR*₂, -SR*, -SO₂R*, -COOR*, and -CONR*₂, where each R* is independently H or C₁₋₃ alkyl.

"Aryl" as used herein refers to a phenyl or naphthyl group unless otherwise specified. Aryl groups unless otherwise specified may be optionally substituted with up to four groups selected from halo, CN, amino, hydroxy, C₁₋₃ alkyl, -OR*, -NR*₂, -SR*, -SO₂R*, -COOR*, and -CONR*₂, where each R* is independently H or C₁₋₃ alkyl.

"Halo" or "halogen", as used herein, may be fluorine, chlorine, bromine or iodine.

"C₁₋₆ alkyl" or "C₁-C₆ alkyl", as used herein, denotes straight chain or branched alkyl having 1-6 carbon atoms. If a different number of carbon atoms is specified, such as C₄ or C₃, then the definition is to be amended accordingly, such as "C₁₋₄ alkyl" will represent methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and tert-butyl.

"C₁₋₆ alkylene" or "C₁-C₆ alkylene", as used herein, denotes straight chain or branched alkyl having 1-6 carbon atoms and two open valences for connection to two other groups. If a different number of carbon atoms is specified, such as C₄ or C₃, then the definition is to be amended accordingly, such as "C₁₋₄ alkylene" will represent methylene (-CH₂-), ethylene (-CH₂CH₂-), straight chain or branched propylene (-CH₂CH₂CH₂- or -CH₂-CHMe-CH₂-), and the like.

"C₁₋₆ alkoxy", as used herein, denotes straight chain or branched alkoxy (-O-Alkyl) having 1-6 carbon atoms. If a different number of carbon atoms is specified, such as C₄ or

C₃, then the definition is to be amended accordingly, such as "C₁₋₄ alkoxy" will represent methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy and tert-butoxy.

"C₁₋₄ Haloalkyl" or "C₁-C₄ haloalkyl" as used herein, denotes straight chain or branched alkyl having 1-4 carbon atoms wherein at least one hydrogen has been replaced with a halogen. The number of halogen replacements can be from one up to the number of hydrogen atoms on the unsubstituted alkyl group. If a different number of carbon atoms is specified, such as C₆ or C₃, then the definition is to be amended accordingly. Thus "C₁₋₄ haloalkyl" will represent methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and tert-butyl that have at least one hydrogen substituted with halogen, such as where the halogen is fluorine: CF₃CF₂-, (CF₃)₂CH-, CH₃-CF₂-, CF₃CF₂-, CF₃, CF₂H-, CF₃CF₂CH(CF₃)- or CF₃CF₂CF₂CF₂-.

"C₃₋₈ cycloalkyl" as used herein refers to a saturated monocyclic hydrocarbon ring of 3 to 8 carbon atoms. Examples of such groups include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. If a different number of carbon atoms is specified, such as C₃-C₆, then the definition is to be amended accordingly.

"4- to 8-Membered heterocyclyl", "5- to 6- membered heterocyclyl", "3- to 10-membered heterocyclyl", "3- to 14-membered heterocyclyl", "4- to 14-membered heterocyclyl" and "5- to 14-membered heterocyclyl", refers, respectively, to 4- to 8-membered, 5- to 6-membered, 3- to 10-membered, 3- to 14-membered, 4- to 14-membered and 5- to 14-membered heterocyclic rings; unless otherwise specified, such rings contain 1 to 7, 1 to 5, or 1 to 3 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur as ring members, and the rings may be saturated, or partially saturated but not aromatic. The heterocyclic group can be attached to another group at a nitrogen or a carbon atom. The term "heterocyclyl" includes single ring groups, fused ring groups and bridged groups. Examples of such heterocyclyl include, but are not limited to pyrrolidine, piperidine, piperazine, pyrrolidinone, morpholine, tetrahydrofuran, tetrahydrothiophene, tetrahydrothiopyran, tetrahydropyran, 1,4-dioxane, 1,4-oxathiane, 8-aza-bicyclo[3.2.1]octane, 3,8-diazabicyclo[3.2.1]octane, 3-Oxa-8-aza-bicyclo[3.2.1]octane, 8-Oxa-3-aza-bicyclo[3.2.1]octane, 2-Oxa-5-aza-bicyclo[2.2.1]heptane, 2,5-Diaza-bicyclo[2.2.1]heptane, azetidine, ethylenedioxo, oxetane or thiazole. In certain embodiments, if not otherwise specified, heterocyclic groups have 1-2 heteroatoms selected from N, O and S as ring members, and 4-7 ring atoms, and are optionally substituted with up to four groups selected from halo, oxo, CN, amino, hydroxy, C₁₋₃ alkyl, -OR*, -NR*₂, -SR*, -SO₂R*, -COOR*, and -CONR*₂, where each R* is independently H or C₁₋₃ alkyl. In particular, heterocyclic groups containing a sulfur atom are optionally substituted with one or two oxo groups on the sulfur.

"Heteroaryl" is a completely unsaturated (aromatic) ring. The term "heteroaryl" refers to a 5-14 membered monocyclic- or bicyclic- or tricyclic-aromatic ring system, having


1 to 8 heteroatoms selected from N, O or S. Typically, the heteroaryl is a 5-10 membered ring or ring system (e.g., 5-7 membered monocyclic group or an 8-10 membered bicyclic group), often a 5-6 membered ring containing up to four heteroatoms selected from N, O and S, though often a heteroaryl ring contains no more than one divalent O or S in the ring. Typical heteroaryl groups include furan, isothiazole, thiadiazole, oxadiazole, indazole, indole, quinoline, 2- or 3-thienyl, 2- or 3-furyl, 2- or 3-pyrrolyl, 2-, 4-, or 5-imidazolyl, 3-, 4-, or 5-pyrazolyl, 2-, 4-, or 5-thiazolyl, 3-, 4-, or 5-isothiazolyl, 2-, 4-, or 5-oxazolyl, 3-, 4-, or 5-isoxazolyl, 3- or 5-(1,2,4-triazolyl), 4- or 5-(1,2,3-triazolyl), tetrazolyl, triazine, pyrimidine, 2-, 3-, or 4-pyridyl, 3- or 4-pyridazinyl, 3-, 4-, or 5-pyrazinyl, 2-pyrazinyl, and 2-, 4-, or 5-pyrimidinyl. Heteroaryl groups are optionally substituted with up to four groups selected from halo, CN, amino, hydroxy, C₁₋₃ alkyl, -OR*, -NR*₂, -SR*, -SO₂R*, -COOR*, and -CONR*₂, where each R* is independently H or C₁₋₃ alkyl.

The term "hydroxy" or "hydroxyl" refers to the group -OH.

Various embodiments of the invention are described herein. It will be recognized that features specified in each embodiment may be combined with other specified features to provide further embodiments.

The following enumerated embodiments are representative of the invention:

1. A compound of formula (I):



(I) wherein:

R¹ is H, halo, or C₁-C₃ alkyl;

R² is H, halo, CN, C₁-C₃ alkyl or C₁-C₃ haloalkyl, C₁-C₃ alkoxy;

R³ is H, OH, halo, CN, C₁-C₃ alkyl, C₃-C₆ cycloalkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, or C₁-C₃ haloalkoxy;

R⁴ and R⁵ taken together form a 5 or 6 membered ring that is fused to the phenyl ring to which R⁴ and R⁵ are attached, wherein the 5 or 6 membered ring contains one or two oxygen atoms as ring members and is optionally substituted by up to four groups independently selected from -CN, -OR, -SR, -SO₂R, and C₁-C₄ alkyl optionally substituted with one or two groups selected from halo, -OR, -SR, -SO₂R, C₁-C₃ haloalkoxy, CN, and -N(R)₂;

R⁶ is H, halo, C₁-C₃ alkoxy, or C₁-C₆ alkyl, or is taken together with R⁹ to form a ring;

R⁷ is H, halo, C₁-C₃ alkoxy, or C₁-C₆ alkyl, or is taken together with R⁹ to form a ring;

R⁸ is H or C₁-C₆ alkyl, or is taken together with R⁹ to form a ring;

R⁹ taken together with exactly one group selected from R⁶, R⁷ and R⁸ forms a 3-7 membered cycloalkyl ring or a 3-7 membered heterocyclic ring containing N, O or S as a ring member; wherein the cycloalkyl or heterocyclic ring is optionally substituted with up to three groups selected from R, -OR, -NR₂, halo, CN, COOR, CONR₂, and oxo;

W is -COOR*, -C(O)NH-SO₂R, -C(O)NH-SO₂NR₂, 5-tetrazolyl, or 1,2,4-oxadiazol-3-yl-5(4H)-one;

R* is H or C₁-C₆ alkyl that is optionally substituted with one or two groups selected from halo, -OR, oxo, CN, -NR₂, COOR, and CONR₂;

R is independently selected at each occurrence from H and C₁-C₄ alkyl optionally substituted with one to three groups selected from halo, -OH, C₁-C₃ alkoxy, oxo, CN, -NH₂, -NH(C₁-C₃ alkyl), -N(C₁-C₃ alkyl)₂, and cyclopropyl;

and two R groups directly attached to the same atom, which may be C or N, can optionally be taken together to form a 3-6 membered ring that can optionally contain an added heteroatom selected from N, O and S as a ring member, and can be substituted by up to two groups selected from -OH, oxo, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

or a pharmaceutically acceptable salt thereof.

In some examples of this embodiment, R⁸ is H and R⁹ forms a ring with R⁶. Optionally, R⁷ is also H. Typically, this is a 5-6 membered carbocyclic ring; in some embodiments, this ring is cis-fused onto the ring in Formula I to which it is attached.


2. The compound of embodiment 1, or a pharmaceutically acceptable salt thereof, wherein R¹ is H.

3. The compound according to any one of the preceding embodiments, or a pharmaceutically acceptable salt thereof, wherein R² is H or halo. In some of these embodiments, R² is H.

4. A compound according to any one of embodiments 1 to 3, or a pharmaceutically acceptable salt thereof, wherein R³ is C₁-C₃ alkoxy or halo. In some of these embodiments, R² is methoxy.

5. The compound of any of embodiments 1-4, wherein W is $-\text{COOR}^*$; or a pharmaceutically acceptable salt thereof. In some of these embodiments, R^* is H.

6. The compound of any of embodiments 1-5, which is of the formula (II):



wherein each R^{10} is independently selected from H, R^{11} , $-\text{OR}$, $-\text{SR}$, $-\text{SO}_2\text{R}$, and $-\text{N}(\text{R})_2$;

R^{11} is C_1 - C_4 alkyl, optionally substituted with one or two groups selected from CN, $-\text{OR}$, C_1 - C_3 haloalkoxy, $-\text{NR}_2$, and $-\text{SO}_2\text{R}$;


n is 1 or 2;

Z is O or CR_2 ;

or a pharmaceutically acceptable salt thereof.

In some of these embodiments, Z is O or CH_2 . In some of these embodiments, one or two of the R^{10} groups that are present represent H, and typically at least one R^{10} group represents R^{11} or $-\text{OR}$.

7. The compound of embodiment 6, which is of the formula (III), or a pharmaceutically acceptable salt thereof:



wherein R^{10} is H, R^{11} , $-OR$, $-SR$, $-SO_2R$, or $-N(R)_2$;

R^{11} is C_1 - C_4 alkyl optionally substituted with $-OH$ or OMe ;

R^{12} and R^{13} are each selected from H and C_1 - C_3 alkyl,


or R^{12} and R^{13} taken together form a cyclopropyl or cyclobutyl ring;

and Z is O or CH_2 .

In some of these embodiments, R^{12} and R^{13} are both methyl. In some of these embodiments, Z is CH_2 and n is 0. In other of these embodiments, n is 1 and Z is CH_2 or O.


8. A compound according to any of embodiments 6-7, or a pharmaceutically acceptable salt thereof, wherein R^{10} is $-CH_2OMe$, $-(CH_2)_2-OMe$, or $-OR$. In some of these embodiments, $-OR$ represents $-OMe$, $-O-CH_2OMe$, or $-O-(CH_2)_2-OMe$.

9. A compound according to any of embodiments 1-8, or a pharmaceutically acceptable salt thereof, which is of the formula:




wherein R^{10} is $-CH_2OMe$, $-(CH_2)_2-OMe$, or $-OR$.

In some of these embodiments, the compound is of this formula:




10. A compound according to embodiment 9, which is of the formula:



wherein R^{10} is $-\text{CH}_2\text{OMe}$, $-(\text{CH}_2)_2\text{-OMe}$, or $-\text{OR}$;

or a pharmaceutically acceptable salt thereof.


In certain embodiments, the compound is of this formula:



11. A compound according to any of embodiments 6-10, or a pharmaceutically acceptable salt thereof, wherein one R^{10} represents $-\text{OR}'$, wherein R' is $\text{C}_1\text{-C}_3$ alkyl substituted with up to two groups selected from OH and OMe.

12. A compound according to any of embodiments 6-10, or a pharmaceutically acceptable salt thereof, wherein one R^{10} is selected from $-\text{CH}_2\text{OMe}$, $-\text{CH}_2\text{CH}_2\text{OMe}$, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{OMe}$.

13. A compound according to any one of embodiments 1-9, which is of the formula:




wherein R¹⁰ is -CH₂OMe, -(CH₂)₂-OMe, or -OR;


or a pharmaceutically acceptable salt thereof.

14. The compound of embodiment 13, wherein R¹⁰ is selected from -OMe, -CH₂OMe, -CH₂CH₂OMe, and -CH₂CH₂CH₂OMe;

or a pharmaceutically acceptable salt thereof.

15. A compound selected from:






wherein R¹ is either H or F;

or a pharmaceutically acceptable salt thereof.

16. The compound of embodiment 1, which is:



or a pharmaceutically acceptable salt thereof.

16a. The compound of embodiment 1 or any of embodiments 3-15, wherein R¹ is F.

17. A pharmaceutical composition, comprising a compound of any of the preceding embodiments admixed with at least one pharmaceutically acceptable carrier.

18. A method to treat a subject having a hepatitis B infection, which comprises administering to the subject a compound of any of embodiments 1-16 or a pharmaceutical composition of embodiment 17.

19. The method of embodiment 18, wherein the compound of any one of embodiments 1-16 or the pharmaceutical composition of embodiment 17 is used in combination with an additional therapeutic agent selected from an interferon or peginterferon, an HBV polymerase inhibitor, a viral entry inhibitor, a viral maturation inhibitor, a capsid assembly inhibitor, an HBV core modulator, a reverse transcriptase inhibitor, a TLR-agonist, or an immunomodulator.

20. A method to inhibit replication of hepatitis B virus, which comprises contacting the hepatitis B virus, either in vitro or in vivo, with a compound according to any one of embodiments 1-16.

21. A pharmaceutical combination, comprising a compound of any of embodiments 1-16 and at least one additional therapeutic agent.

Another embodiment of the invention provides a compound as described above, or a pharmaceutically acceptable salt thereof, for use as a medicament. In one aspect, the medicament is for treatment of a subject having an HBV infection. In a particular embodiment, the subject is a human diagnosed with chronic HBV.

Also within the scope of this invention is the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament; in some embodiments, this medicament is for the treatment or prevention of a viral disease and/or infection in a human being, particularly where the virus involved is HBV.

Included within the scope of this invention is a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient. Optionally, the composition comprises at least two pharmaceutically acceptable carriers and/or excipients.

According to a further aspect of this embodiment the pharmaceutical composition according to this invention further comprises a therapeutically effective amount of at least one other antiviral agent.

The invention also provides the use of a pharmaceutical composition as described hereinabove for the treatment of a HBV infection in a human being having or at risk of having the infection.

The invention also provides the use of a pharmaceutical composition as described hereinabove for the treatment of HBV infection in a human being having or at risk of having the disease.

Another aspect of the invention involves a method of treating or preventing a hepatitis B viral disease and/or infection in a human being by administering to the human being an antivirally effective amount of a compound of the invention, a pharmaceutically

acceptable salt thereof, or a composition as described above, alone or in combination with at least one other antiviral agent, administered together or separately.

An additional aspect of this invention refers to an article of manufacture comprising a composition of the invention that is effective to treat a hepatitis B viral disease and/or infection; and packaging material comprising a label which indicates that the composition can be used to treat disease and/or infection by a hepatitis B virus; wherein the composition comprises a compound of formula (I) according to this invention or a pharmaceutically acceptable salt thereof.

Still another aspect of this invention relates to a method of inhibiting the replication of HBV, comprising exposing the virus to an effective amount of the compound of formula (I), or a salt thereof, under conditions where replication of the virus is inhibited. This method can be practiced in vitro or in vivo.

Further included in the scope of the invention is the use of a compound of formula (I), or a salt thereof, to inhibit the replication of HBV.

In all of the embodiment referring to a compound of Formula (I), the compound of Formula (I) can be a compound according to any of embodiments 1-16 described above.

In some embodiments, the compound of Formula (I) is co-administered with or used in combination with at least one additional therapeutic agent selected from: an interferon or peginterferon, an HBV polymerase inhibitor, a viral entry inhibitor, a viral maturation inhibitor, a capsid assembly inhibitor, an HBV core modulator, a reverse transcriptase inhibitor, a TLR-agonist, or an immunomodulator. Optionally, the compound of Formula (I) may be prepared for simultaneous or sequential use in combination with an additional therapeutic agent; or the compound of Formula (I) may be combined into a pharmaceutical combination comprising a compound of Formula (I) and at least one additional therapeutic agent. Some particular therapeutic agents that may be used in combination with the compounds of the invention include immunomodulators described herein, interferon alfa 2a, interferon alfa-2b, pegylated interferon alfa-2a, pegylated interferon alfa-2b, TLR-7 and TLR-9 agonists, entecavir, tenofovir, cidofovir, telbivudine, didanosine, zalcitabine, stavudine, lamivudine, abacavir, emtricitabine, apricitabine, atevirapine, ribavirin, acyclovir, famciclovir, valacyclovir, ganciclovir, adefovir, efavirenz, nevirapine, delavirdine, and etravirine. Suitable core modulators are disclosed in WO2013/096744; suitable HBV capsid inhibitors are described in US2015/0252057.

These additional agents may be combined with the compounds of this invention to create a single pharmaceutical dosage form. Alternatively these additional agents may be separately administered to the patient as part of a multiple dosage form, for example, using a kit. Such additional agents may be administered to the patient prior to, concurrently with, or following the administration of a compound of the invention, or a pharmaceutically acceptable salt thereof. Alternatively, these additional therapeutic agents may be

administered separately from and optionally by different routes of administration and on different dosing schedules from the compound of the invention, provided the compound of the invention and the additional therapeutic agent are used concurrently for treatment of an HBV infection or a disorder caused or complicated by an HBV infection.

The dose range of the compounds of the invention applicable per day is usually from 0.01 to 100 mg/kg of body weight, preferably from 0.1 to 50 mg/kg of body weight. Each dosage unit may conveniently contain from 5% to 95% active compound (w/w). Preferably such preparations contain from 20% to 80% active compound.

The actual pharmaceutically effective amount or therapeutic dosage will of course depend on factors known by those skilled in the art such as age and weight of the patient, route of administration and severity of disease. In any case the combination will be administered at dosages and in a manner which allows a pharmaceutically effective amount to be delivered based upon patient's unique condition.

When the composition of this invention comprises a combination of a compound of the invention and one or more additional therapeutic or prophylactic agent, both the compound and the additional agent should be present at dosage levels of between about 10 to 100%, and more preferably between about 10 and 80% of the dosage normally administered in a monotherapy regimen.

Antiviral agents contemplated for use in such combination therapy include agents (compounds or biologicals) that are effective to inhibit the formation and/or replication of a virus in a human being, including but not limited to agents that interfere with either host or viral mechanisms necessary for the formation and/or replication of a virus in a human being. Such agents can be selected from entecavir, tenofovir, cidofovir, telbivudine, didanosine, zalcitabine, stavudine, lamivudine, abacavir, emtricitabine, apricitabine, atevirapine, ribavirin, acyclovir, famciclovir, valacyclovir, ganciclovir, adefovir, efavirenz, nevirapine, delavirdine, and etravirine, and immunomodulators described herein including interferons and pegylated interferons, TLR-7 agonists, and TLR-9 agonists.


Many compounds of the invention contain one or more chiral centers. These compounds may be made and used as single isomers or as mixtures of isomers. Methods for separating the isomers, including diastereomers and enantiomers, are known in the art, and examples of suitable methods are described herein. In certain embodiments, the compounds of the invention are used as a single substantially pure isomer, meaning at least 90% of a sample of the compound is the specified isomer and less than 10% of the sample is any other isomer or mixture of isomers. Preferably, at least 95% of the sample is a single isomer. Selection of a suitable isomer is within the ordinary level of skill, as one isomer will typically be more active in the *in vivo* or *in vitro* assay described herein for measuring HBV activity, and will be the preferred isomer. Where *in vitro* activity differences between isomers are relatively small, e.g. less than about a factor of 4, a preferred isomer may be

selected based on activity level against viral replication in cell culture, using methods such as those described herein: the isomer having a lower MIC (minimum inhibitory concentration) or EC-50 is preferred.

The compounds of the invention may be synthesized by the general synthetic routes illustrated below, specific examples of which are described in more detail in the Examples. Additional guidance for synthesis of the compounds of Formula (I) and synthetic intermediates useful for these syntheses are disclosed in published PCT applications WO2015/113990 and WO2015/173164.


A versatile intermediate useful to make many compounds of the invention is readily prepared from meta-bromophenols as illustrated in Scheme 1. This provides an ortho-allylphenol, which can be used by known methods to form benzo-dihydrofuran and benzo-dihydropyran compounds. The choice of conditions for this cyclization can provide further opportunities to functionalize the newly-formed ring. The Examples that follow illustrate such methods, and persons of skill in the art will recognize modifications and variations that can be used to make many other compounds within the scope of the invention.

Scheme 1. General method to synthesize key intermediates



Using this intermediate and cyclic ketones, a variety of compounds of the invention can readily be prepared by methods such as those in the Examples and illustrated generally in Scheme 2. In step 1, a palladium-catalyzed coupling is used to attach the cyclic ketone, and the carbonyl is then elaborated to a formamide moiety that can be used (step 4) to form dihydroisoquinoline intermediates. These are readily annulated (step 6) and dehydrated (step 7) to provide compounds of Formula (I).

Scheme 2. General method to synthesize final compounds



The term "an optical isomer" or "a stereoisomer" refers to any of the various stereoisomeric configurations which may exist for a given compound of the present invention and includes geometric isomers. It is understood that a substituent may be attached at a chiral center of a carbon atom. The term "chiral" refers to molecules which have the property of non-superimposability on their mirror image partner, while the term "achiral" refers to molecules which are superimposable on their mirror image partner. Therefore, the invention includes enantiomers, diastereomers or racemates of the compound. "Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other. A 1:1 mixture of a pair of enantiomers is a "racemic" mixture. The term is used to designate a racemic mixture where appropriate. "Diastereoisomers" are stereoisomers that have at least two asymmetric atoms, but which are not mirror-images of each other. The absolute stereochemistry is specified according to the Cahn-Ingold-Prelog R-S system. When a compound is a pure enantiomer the stereochemistry at each chiral carbon may be specified by either R or S. Resolved compounds whose absolute configuration is unknown can be designated (+) or (-) depending on the direction (dextro- or levorotatory) which they rotate plane polarized light at the wavelength of the sodium D line. Certain compounds described herein contain one or more asymmetric centers or axes and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (R)- or (S)-.

Depending on the choice of the starting materials and procedures, the compounds can be present in the form of one of the possible isomers or as mixtures thereof, for example as pure optical isomers, or as isomer mixtures, such as racemates and diastereoisomer mixtures, depending on the number of asymmetric carbon atoms. The present invention is meant to include all such possible stereoisomers, including racemic mixtures, diastereomeric mixtures and optically pure forms. Optically active (R)- and (S)-isomers may be prepared using chiral synthons or chiral reagents, or resolved using

conventional techniques. If the compound contains a double bond, the substituent may be E or Z configuration. If the compound contains a disubstituted cycloalkyl, the cycloalkyl substituent may have a cis- or trans-configuration. All tautomeric forms are also intended to be included.

Any resulting mixtures of isomers can be separated on the basis of the physicochemical differences of the constituents, into the pure or substantially pure geometric or optical isomers or diastereomers, for example, by chromatography and/or fractional crystallization.

Any resulting racemates of final products or intermediates can be resolved into the optical antipodes by known methods, e.g., by separation of the diastereomeric salts thereof, obtained with an optically active acid or base, and liberating the optically active acidic or basic compound. In particular, a basic moiety may thus be employed to resolve the compounds of the present invention into their optical antipodes, e.g., by fractional crystallization of a salt formed with an optically active acid, e.g., tartaric acid, dibenzoyl tartaric acid, diacetyl tartaric acid, di-O,O'-p-toluoyl tartaric acid, mandelic acid, malic acid or camphor-10-sulfonic acid. Racemic products can also be resolved by chiral chromatography, e.g., high pressure liquid chromatography (HPLC) using a chiral adsorbent.

Furthermore, the compounds of the present invention, including their salts, can also be obtained in the form of their hydrates, or include other solvents used for their crystallization. The compounds of the present invention may inherently or by design form solvates with pharmaceutically acceptable solvents (including water); therefore, it is intended that the invention embrace both solvated and unsolvated forms. The term "solvate" refers to a molecular complex of a compound of the present invention (including pharmaceutically acceptable salts thereof) with one or more solvent molecules. Such solvent molecules are those commonly used in the pharmaceutical art, which are known to be innocuous to the recipient, e.g., water, ethanol, and the like. The term "hydrate" refers to the complex where the solvent molecule is water.

The compounds of the present invention, including salts, hydrates and solvates thereof, may inherently or by design form polymorphs.

As used herein, the terms "salt" or "salts" refers to an acid addition or base addition salt of a compound of the present invention. "Salts" include in particular "pharmaceutically acceptable salts". The term "pharmaceutically acceptable salts" refers to salts that retain the biological effectiveness and properties of the compounds of this invention and, which typically are not biologically or otherwise undesirable. In many cases, the compounds of the present invention are capable of forming acid and/or base salts by virtue of the presence of amino and/or carboxyl groups or groups similar thereto.

Pharmaceutically acceptable acid addition salts can be formed with inorganic acids

and organic acids, e.g., acetate, aspartate, benzoate, besylate, bromide/hydrobromide, bicarbonate/carbonate, bisulfate/sulfate, camphorsulfonate, chloride/hydrochloride, chlorotheophyllonate, citrate, ethandisulfonate, fumarate, gluceptate, gluconate, glucuronate, hippurate, hydroiodide/iodide, isethionate, lactate, lactobionate, laurylsulfate, malate, maleate, malonate, mandelate, mesylate, methylsulphate, naphthoate, napsylate, nicotinate, nitrate, octadecanoate, oleate, oxalate, palmitate, pamoate, phosphate/hydrogen phosphate/dihydrogen phosphate, polygalacturonate, propionate, stearate, succinate, sulfosalicylate, tartrate, tosylate and trifluoroacetate salts.

Inorganic acids from which salts can be derived include, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like.

Organic acids from which salts can be derived include, for example, acetic acid, propionic acid, glycolic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, toluenesulfonic acid, sulfosalicylic acid, and the like. Pharmaceutically acceptable base addition salts can be formed with inorganic and organic bases.

Inorganic bases from which salts can be derived include, for example, ammonium salts and metals from columns I to XII of the periodic table. In certain embodiments, the salts are derived from sodium, potassium, ammonium, calcium, magnesium, iron, silver, zinc, and copper; particularly suitable salts include ammonium, potassium, sodium, calcium and magnesium salts.

Organic bases from which salts can be derived include, for example, primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, basic ion exchange resins, and the like. Certain organic amines include isopropylamine, benzathine, choline, diethanolamine, diethylamine, lysine, meglumine, piperazine and tromethamine.

The pharmaceutically acceptable salts of the present invention can be synthesized from a basic or acidic moiety, by conventional chemical methods. Generally, such salts can be prepared by reacting free acid forms of these compounds with a stoichiometric amount of the appropriate base (such as Na, Ca, Mg, or K hydroxide, carbonate, bicarbonate or the like), or by reacting free base forms of these compounds with a stoichiometric amount of the appropriate acid. Such reactions are typically carried out in water or in an organic solvent, or in a mixture of the two. Generally, use of non-aqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile is desirable, where practicable. Lists of additional suitable salts can be found, e.g., in "Remington's Pharmaceutical Sciences", 20th ed., Mack Publishing Company, Easton, Pa., (1985); and in "Handbook of Pharmaceutical Salts: Properties, Selection, and Use" by Stahl and Wermuth (Wiley-VCH, Weinheim, Germany, 2002).

Any formula given herein is intended to represent unlabeled forms as well as

isotopically labeled forms of the compounds of the present invention having up to three atoms with non-natural isotope distributions, e.g., sites that are enriched in deuterium or ^{13}C or ^{15}N . Isotopically labeled compounds have structures depicted by the formulas given herein except that one or more atoms are replaced by an atom having a selected atomic mass or mass number other than the natural-abundance mass distribution. Examples of isotopes that can be usefully over-incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, and chlorine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{18}F , ^{31}P , ^{32}P , ^{35}S , ^{36}Cl , ^{125}I respectively. The invention includes various isotopically labeled compounds of the present invention, for example those into which radioactive isotopes, such as ^3H and ^{14}C , or those in which non-radioactive isotopes, such as ^2H and ^{13}C are present at levels substantially above normal isotope distribution. Such isotopically labelled compounds are useful in metabolic studies (with ^{14}C , for example), reaction kinetic studies (with, for example ^2H or ^3H), detection or imaging techniques, such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT) including drug or substrate tissue distribution assays, or in radioactive treatment of patients. In particular, an ^{18}F labeled compound of the present invention may be particularly desirable for PET or SPECT studies. Isotopically-labeled compounds of the present invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Examples and Preparations using an appropriate isotopically-labeled reagent in place of the non-labeled reagent typically employed. Labeled samples may be useful with quite low isotope incorporation, such as where a radiolabel is used to detect trace amounts of the compound.

Further, site-specific substitution with heavier isotopes, particularly deuterium (i.e., ^2H or D), may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements or an improvement in therapeutic index. It is understood that deuterium in this context is regarded as a substituent of a compound of the present invention, and typically a sample of a compound having deuterium as a substituent has at least 50% deuterium incorporation at the labeled position(s). The concentration of such a heavier isotope, specifically deuterium, may be defined by the isotopic enrichment factor. The term "isotopic enrichment factor" as used herein means the ratio between the isotopic abundance and the natural abundance of a specified isotope. If a substituent in a compound of this invention is denoted deuterium, such compound has an isotopic enrichment factor for each designated deuterium atom of at least 3500 (52.5% deuterium incorporation at each designated deuterium atom), at least 4000 (60% deuterium incorporation), at least 4500 (67.5% deuterium incorporation), at least 5000 (75% deuterium incorporation), at least 5500 (82.5% deuterium incorporation), at least 6000 (90% deuterium incorporation), at least 6333.3 (95% deuterium incorporation), at least

6466.7 (97% deuterium incorporation), at least 6600 (99% deuterium incorporation), or at least 6633.3 (99.5% deuterium incorporation).

Pharmaceutically acceptable solvates in accordance with the invention include those wherein the solvent of crystallization may be isotopically substituted, e.g. D₂O, d⁶-acetone, d⁶-DMSO.

Compounds of the present invention that contain groups capable of acting as donors and/or acceptors for hydrogen bonds may be capable of forming co-crystals with suitable co-crystal formers. These co-crystals may be prepared from compounds of the present invention by known co-crystal forming procedures. Such procedures include grinding, heating, co-subliming, co-melting, or contacting in solution compounds of the present invention with the co-crystal former under crystallization conditions and isolating co-crystals thereby formed. Suitable co-crystal formers include those described in WO 2004/078163. Hence the invention further provides co-crystals comprising a compound of the present invention.

Methods of Use

All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g. "such as") provided herein is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention otherwise claimed.

The compounds of the invention can be administered by known methods, including oral, parenteral, inhalation, and the like. In certain embodiments, the compound of the invention is administered orally, as a pill, lozenge, troche, capsule, solution, or suspension. In other embodiments, a compound of the invention is administered by injection or infusion. Infusion is typically performed intravenously, often over a period of time between about 15 minutes and 4 hours. In other embodiments, a compound of the invention is administered intranasally or by inhalation; inhalation methods are particularly useful for treatment of respiratory infections. Compounds of the present invention exhibit oral bioavailability, so oral administration is sometimes preferred.

In certain embodiments of the present invention, a compound of the present invention is used in combination with a second antiviral agent, such as those named herein.

By the term "combination", is meant either a fixed combination in one dosage unit form, as separate dosage forms suitable for use together either simultaneously or sequentially, or as a kit of parts for the combined administration where a compound of the present invention and a combination partner may be administered independently at the same time or separately within time intervals that especially allow that the combination partners show a cooperative, e.g., synergistic, effect, or any combination thereof.

The second antiviral agent may be administered in combination with the compounds of the present inventions wherein the second antiviral agent is administered prior to, simultaneously, or after the compound or compounds of the present invention. When simultaneous administration of a compound of the invention with a second agent is desired and the route of administration is the same, then a compound of the invention may be formulated with a second agent into the same dosage form. An example of a dosage form containing a compound of the invention and a second agent is a tablet or a capsule.

In some embodiments, a combination of a compound of the invention and a second antiviral agent may provide synergistic activity. The compound of the invention and second antiviral agent may be administered together, separate but simultaneously, or sequentially.

An "effective amount" of a compound is that amount necessary or sufficient to treat or prevent a viral infection and/or a disease or condition described herein. In an example, an effective amount of a compound of Formula I is an amount sufficient to treat viral infection in a subject. In another example, an effective amount is an amount sufficient to treat HBV in a subject in need of such treatment. The effective amount can vary depending on such factors as the size and weight of the subject, the type of illness, or the particular compound of the invention. For example, the choice of the compound of the invention can affect what constitutes an "effective amount." One of ordinary skill in the art would be able to study the factors contained herein and make the determination regarding the effective amount of the compounds of the invention without undue experimentation.

The regimen of administration can affect what constitutes an effective amount. The compound of the invention can be administered to the subject either prior to or after the onset of a viral infection. Further, several divided dosages, as well as staggered dosages, can be administered daily or sequentially, or the dose can be continuously infused, or can be a bolus injection. Further, the dosages of the compound(s) of the invention can be proportionally increased or decreased as indicated by the exigencies of the therapeutic or prophylactic situation.

Compounds of the invention may be used in the treatment of states, disorders or diseases as described herein, or for the manufacture of pharmaceutical compositions for use in the treatment of these diseases. The invention provides methods of use of compounds of the present invention in the treatment of these diseases or for preparation of pharmaceutical compositions having compounds of the present invention for the treatment of these diseases.

The language "pharmaceutical composition" includes preparations suitable for administration to mammals, *e.g.*, humans. When the compounds of the present invention are administered as pharmaceuticals to mammals, *e.g.*, humans, they can be given *per se* or as a pharmaceutical composition containing, for example, 0.1 to 99.5% (more preferably, 0.5 to 90%) of at least one compound of Formula (I) or any subgenus thereof as active

ingredient in combination with a pharmaceutically acceptable carrier, or optionally two or more pharmaceutically acceptable carriers.

The phrase "pharmaceutically acceptable carrier" is art recognized and includes a pharmaceutically acceptable material, composition or vehicle, suitable for administering compounds of the present invention to mammals. The carriers include liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting the subject agent from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically acceptable carriers include: sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients, such as cocoa butter and suppository waxes; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; phosphate buffer solutions; and other non-toxic compatible substances employed in pharmaceutical formulations. Typically, pharmaceutically acceptable carriers are sterilized and/or substantially pyrogen-free.

Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions.

Examples of pharmaceutically acceptable antioxidants include: water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, α -tocopherol, and the like; and metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

Formulations of the present invention include those suitable for oral, nasal, inhalation, topical, transdermal, buccal, sublingual, rectal, vaginal and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient that can be combined with a carrier material to produce a single dosage form will generally be that amount of the compound that produces a therapeutic effect. Generally,

out of one hundred per cent, this amount will range from about 1 per cent to about ninety-nine percent of active ingredient, preferably from about 5 per cent to about 70 per cent, most preferably from about 10 per cent to about 30 per cent.

Methods of preparing these formulations or compositions include the step of bringing into association a compound of the present invention with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a compound of the present invention with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

Formulations of the invention suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored base, for example, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of a compound of the present invention as an active ingredient. A compound of the present invention may also be administered as a bolus, electuary or paste.

In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), the active ingredient is mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; humectants, such as glycerol; disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; solution retarding agents, such as paraffin; absorption accelerators, such as quaternary ammonium compounds; wetting agents, such as, for example, cetyl alcohol and glycerol monostearate; absorbents, such as kaolin and bentonite clay; lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions that can be dissolved in sterile water, or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. The active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

Liquid dosage forms for oral administration of the compounds of the invention include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluent commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

Formulations of the pharmaceutical compositions of the invention for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing one or more compounds of the invention with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the rectum or vaginal cavity and release the active compound.

Formulations of the present invention which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

Dosage forms for the topical or transdermal administration of a compound of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants that may be required.

The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

Powders and sprays can contain, in addition to a compound of this invention, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

Transdermal patches have the added advantage of providing controlled delivery of a compound of the present invention to the body. Such dosage forms can be made by dissolving or dispersing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate of such flux can be controlled by either providing a rate controlling membrane or dispersing the active compound in a polymer matrix or gel.

Ophthalmic formulations, eye ointments, powders, solutions and the like, are also contemplated as being within the scope of this invention.

Pharmaceutical compositions of this invention suitable for parenteral administration may comprise one or more compounds of the invention in combination with one or more pharmaceutically acceptable carriers such as sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

Examples of suitable aqueous and nonaqueous carriers that may be employed in the pharmaceutical compositions of the invention include water, ethanol, glycol ethers, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions,

and by the use of surfactants.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents that delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally-administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the subject compounds in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions that are compatible with body tissue.

The preparations of the present invention may be given orally, parenterally, topically, or rectally. They are of course given by forms suitable for each administration route. For example, they are administered in tablets or capsule form, by injection, inhalation, eye lotion, ointment, suppository, etc., administration by injection, infusion or inhalation; topical by lotion or ointment; and rectal by suppositories.

The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion. Intravenous infusion is sometimes a preferred method of delivery for compounds of the invention. Infusion may be used to deliver a single daily dose or multiple doses. In some embodiments, a compound of the invention is administered by infusion over an interval between 15 minutes and 4 hours, typically between 0.5 and 3 hours. Such infusion may be used once per day, twice per day or up to three times per day.

The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" as used herein mean the administration of a compound, drug or other material other than directly into the central nervous system, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

These compounds may be administered to humans and other animals for therapy by any suitable route of administration, including orally, nasally, as by, for example, a spray, rectally, intravaginally, parenterally, intracisternally and topically, as by powders, ointments or drops, including buccally and sublingually.

Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

The selected dosage level will depend upon a variety of factors including the activity of the particular compound of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compound employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the compounds of the invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

In general, a suitable daily dose of a compound of the invention will be that amount of the compound that is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. Generally, intravenous and subcutaneous doses of the compounds of this invention for a patient, when used for the indicated effects, will range from about 0.0001 to about 100 mg per kilogram of body weight per day, more preferably from about 0.01 to about 50 mg per kg per day, and still more preferably from about 0.1 to about 20 mg per kg per day. An effective amount is that amount which prevents or treats a viral infection, such as HBV.

Treatment with a compound or composition described herein may be repeated daily for a period sufficient to reduce or substantially eliminate an HBV infection or viral load. For example, treatment may be continued for a week, or two weeks, or 3-4 weeks, or 4-8 weeks, or 8-12 weeks, 2-6 months, or longer, e.g., until viral load or other measure of infection shows a substantial reduction in viral load or viral activity or other signs or symptoms of HBV infection. The skilled treating physician can readily determine a suitable duration of treatment.

If desired, the effective daily dose of the active compound may be administered as a single dose per day, or as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms. Compounds delivered orally or by inhalation, are commonly administered in one to four doses per day. Compounds delivered by injection are typically administered once per day, or once every other day. Compounds delivered by infusion are typically administered in one to three doses per day. When multiple doses are administered within a day, the doses may be administered at intervals of about 4 hours, about 6 hours, about 8 hours or about 12 hours.

While it is possible for a compound of the present invention to be administered alone, it is preferable to administer the compound as a pharmaceutical composition such as those described herein. Thus methods of using the compounds of the invention include administering the compound as a pharmaceutical composition, wherein at least one compound of the invention is admixed with a pharmaceutically acceptable carrier prior to administration.

Use of Compounds of the Invention in combination with immunomodulators

The compounds and compositions described herein can be used or administered in combination with one or more therapeutic agents that act as immunomodulators, e.g., an activator of a costimulatory molecule, or an inhibitor of an immune-inhibitory molecule, or a vaccine. The Programmed Death 1 (PD-1) protein is an inhibitory member of the extended CD28/CTLA4 family of T cell regulators (Okazaki et al. (2002) *Curr Opin Immunol* 14: 391779-82; Bennett et al. (2003) *J. Immunol.* 170:711-8). PD-1 is expressed on activated B cells, T cells, and monocytes. PD-1 is an immune-inhibitory protein that negatively regulates TCR signals (Ishida, Y. et al. (1992) *EMBO J.* 11:3887-3895; Blank, C. et al. (Epub 2006 Dec. 29) *Immunol. Immunother.* 56(5):739-745), and is up-regulated in chronic infections. The interaction between PD-1 and PD-L1 can act as an immune checkpoint, which can lead to, e.g., a decrease in infiltrating lymphocytes, a decrease in T-cell receptor mediated proliferation, and/or immune evasion by cancerous or infected cells (Dong et al. (2003) *J. Mol. Med.* 81:281-7; Blank et al. (2005) *Cancer Immunol. Immunother.* 54:307-314; Konishi et al. (2004) *Clin. Cancer Res.* 10:5094-100). Immune suppression can be

reversed by inhibiting the local interaction of PD-1 with PD-L1 or PD-L2; the effect is additive when the interaction of PD-1 with PD-L2 is blocked as well (Iwai et al. (2002) Proc. Nat'l. Acad. Sci. USA 99:12293-7; Brown et al. (2003) J. Immunol. 170:1257-66).

Immunomodulation can be achieved by binding to either the immune-inhibitory protein (e.g., PD-1) or to binding proteins that modulate the inhibitory protein (e.g., PD-L1, PD-L2).

In one embodiment, the combination therapies of the invention include an immunomodulator that is an inhibitor or antagonist of an inhibitory molecule of an immune checkpoint molecule. In another embodiment the immunomodulator binds to a protein that naturally inhibits the immuno-inhibitory checkpoint molecule. When used in combination with antiviral compounds, these immunomodulators can enhance the antiviral response, and thus enhance efficacy relative to treatment with the antiviral compound alone.

The term "immune checkpoints" refers to a group of molecules on the cell surface of CD4 and CD8 T cells. These molecules can effectively serve as "brakes" to down-modulate or inhibit an adaptive immune response. Immune checkpoint molecules include, but are not limited to, Programmed Death 1 (PD-1), Cytotoxic T-Lymphocyte Antigen 4 (CTLA-4), B7H1, B7H4, OX-40, CD137, CD40, and LAG3, which directly inhibit immune cells.

Immunotherapeutic agents which can act as immune checkpoint inhibitors useful in the methods of the present invention, include, but are not limited to, inhibitors of PD-L1, PD-L2, CTLA4, TIM3, LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4 and/or TGFR beta. Inhibition of an inhibitory molecule can be performed by inhibition at the DNA, RNA or protein level. In some embodiments, an inhibitory nucleic acid (e.g., a dsRNA, siRNA or shRNA), can be used to inhibit expression of an inhibitory molecule. In other embodiments, the inhibitor of an inhibitory signal is a polypeptide, e.g., a soluble ligand, or an antibody or antigen-binding fragment thereof, that binds to the inhibitory molecule.

By "in combination with," it is not intended to imply that the therapy or the therapeutic agents must be administered at the same time and/or formulated for delivery together, although these methods of delivery are within the scope described herein. The immunomodulator can be administered concurrently with, prior to, or subsequent to, one or more compounds of the invention, and optionally one or more additional therapies or therapeutic agents. The therapeutic agents in the combination can be administered in any order. In general, each agent will be administered at a dose and/or on a time schedule determined for that agent. It will further be appreciated that the therapeutic agents utilized in this combination may be administered together in a single composition or administered separately in different compositions. In general, it is expected that each of the therapeutic agents utilized in combination be utilized at levels that do not exceed the levels at which they are utilized individually. In some embodiments, the levels utilized in combination will be lower than those utilized individually.

In certain embodiments, the antiviral compounds described herein are administered

in combination with one or more immunomodulators that are inhibitors of PD-1, PD-L1 and/or PD-L2. Each such inhibitor may be an antibody, an antigen binding fragment thereof, an immunoadhesin, a fusion protein, or an oligopeptide. Examples of such immunomodulators are known in the art.

In some embodiments, the immunomodulator is an anti-PD-1 antibody chosen from MDX-1106, Merck 3475 or CT- 011.

In some embodiments, the immunomodulator is an immunoadhesin (e.g., an immunoadhesin comprising an extracellular or PD-1 binding portion of PD-L1 or PD-L2 fused to a constant region (e.g., an Fc region of an immunoglobulin sequence).

In some embodiments, the immunomodulator is a PD-1 inhibitor such as AMP-224.

In some embodiments, the the immunomodulator is a PD-L1 inhibitor such as anti-PD-L1 antibody.

In some embodiments, the immunomodulator is an anti-PD-L1 binding antagonist chosen from YW243.55.S70, MPDL3280A, MEDI-4736, MSB-0010718C, or MDX-1105. MDX-1105, also known as BMS-936559, is an anti-PD-L1 antibody described in WO2007/005874. Antibody YW243.55.S70 is an anti-PD-L1 described in WO 2010/077634.

In some embodiments, the immunomodulator is nivolumab (CAS Registry Number: 946414-94-4). Alternative names for nivolumab include MDX-1106, MDX-1106-04, ONO-4538, or BMS-936558. Nivolumab is a fully human IgG4 monoclonal antibody which specifically blocks PD-1. Nivolumab (clone 5C4) and other human monoclonal antibodies that specifically bind to PD-1 are disclosed in US 8,008,449, EP2161336 and WO2006/121168.

In some embodiments, the immunomodulator is an anti-PD-1 antibody Pembrolizumab. Pembrolizumab (also referred to as Lambrolizumab, MK-3475, MK03475, SCH-900475 or KEYTRUDA®; Merck) is a humanized IgG4 monoclonal antibody that binds to PD-1. Pembrolizumab and other humanized anti-PD-1 antibodies are disclosed in Hamid, O. et al. (2013) New England Journal of Medicine 369 (2): 134–44, US 8,354,509, WO2009/114335, and WO2013/079174.

In some embodiments, the immunomodulator is Pidilizumab (CT-011; Cure Tech), a humanized IgG1k monoclonal antibody that binds to PD1. Pidilizumab and other humanized anti-PD-1 monoclonal antibodies are disclosed in WO2009/101611.

Other anti-PD1 antibodies useful as immunomodulators for use in the methods disclosed herein include AMP 514 (Amplimmune), and anti-PD1 antibodies disclosed in US 8,609,089, US 2010028330, and/or US 20120114649. In some embodiments, the anti-PD-L1 antibody is MSB0010718C. MSB0010718C (also referred to as A09-246-2; Merck Serono) is a monoclonal antibody that binds to PD-L1.

In some embodiments, the immunomodulator is MDPL3280A (Genentech / Roche), a human Fc optimized IgG1 monoclonal antibody that binds to PD-L1. MDPL3280A and

other human monoclonal antibodies to PD-L1 are disclosed in U.S. Patent No.: 7,943,743 and U.S. Publication No.: 20120039906. Other anti-PD-L1 binding agents useful as immunomodulators for methods of the invention include YW243.55.S70 (see WO2010/077634), MDX-1105 (also referred to as BMS-936559), and anti-PD-L1 binding agents disclosed in WO2007/005874.

In some embodiments, the immunomodulator is AMP-224 (B7-DCIg; Amplimmune; e.g., disclosed in WO2010/027827 and WO2011/066342), is a PD-L2 Fc fusion soluble receptor that blocks the interaction between PD1 and B7-H1.

In some embodiments, the immunomodulator is an anti-LAG-3 antibody such as BMS-986016. BMS-986016 (also referred to as BMS986016) is a monoclonal antibody that binds to LAG-3. BMS-986016 and other humanized anti-LAG-3 antibodies are disclosed in US 2011/0150892, WO2010/019570, and WO2014/008218

In certain embodiments, the combination therapies disclosed herein include a modulator of a costimulatory molecule or an inhibitory molecule, e.g., a co-inhibitory ligand or receptor.

In one embodiment, the costimulatory modulator, e.g., agonist, of a costimulatory molecule is chosen from an agonist (e.g., an agonistic antibody or antigen-binding fragment thereof, or soluble fusion) of OX40, CD2, CD27, CDS, ICAM-1, LFA-1 (CD11a/CD18), ICOS (CD278), 4-1BB (CD137), GITR, CD30, CD40, BAFFR, HVEM, CD7, LIGHT, NKG2C, SLAMF7, NKp80, CD160, B7-H3 or CD83 ligand.

In another embodiment, the combination therapies disclosed herein include an immunomodulator that is a costimulatory molecule, e.g., an agonist associated with a positive signal that includes a costimulatory domain of CD28, CD27, ICOS and/or GITR.

Exemplary GITR agonists include, e.g., GITR fusion proteins and anti-GITR antibodies (e.g., bivalent anti-GITR antibodies), such as, a GITR fusion protein described in U.S. Patent No.: 6,111,090, European Patent No.: 090505B1, U.S. Patent No.: 8,586,023, PCT Publication Nos.: WO 2010/003118 and 2011/090754, or an anti-GITR antibody described, e.g., in U.S. Patent No.: 7,025,962, European Patent No.: 1947183B1, U.S. Patent No.: 7,812,135, U.S. Patent No.: 8,388,967, U.S. Patent No.: 8,591,886, European Patent No.: EP 1866339, PCT Publication No.: WO 2011/028683, PCT Publication No.: WO 2013/039954, PCT Publication No.: WO2005/007190, PCT Publication No.: WO 2007/133822, PCT Publication No.: WO2005/055808, PCT Publication No.: WO 99/40196, PCT Publication No.: WO 2001/03720, PCT Publication No.: WO99/20758, PCT Publication No.: WO2006/083289, PCT Publication No.: WO 2005/115451, U.S. Patent No.: 7,618,632, and PCT Publication No.: WO 2011/051726.

In one embodiment, the immunomodulator used is a soluble ligand (e.g., a CTLA-4-Ig), or an antibody or antibody fragment that binds to PD-L1, PD-L2 or CTLA4. For example, the anti-PD-1 antibody molecule can be administered in combination with an anti-CTLA-4

antibody, e.g., ipilimumab, for example. Exemplary anti-CTLA4 antibodies include Tremelimumab (IgG2 monoclonal antibody available from Pfizer, formerly known as ticilimumab, CP-675,206); and Ipilimumab (CTLA-4 antibody, also known as MDX-010, CAS No. 477202-00-9).

In one embodiment, an anti-PD-1 antibody molecule is administered after treatment with a compound of the invention as described herein.

In another embodiment, an anti-PD-1 or PD-L1 antibody molecule is administered in combination with an anti-LAG-3 antibody or an antigen-binding fragment thereof. In another embodiment, the anti-PD-1 or PD-L1 antibody molecule is administered in combination with an anti-TIM-3 antibody or antigen-binding fragment thereof. In yet other embodiments, the anti-PD-1 or PD-L1 antibody molecule is administered in combination with an anti-LAG-3 antibody and an anti-TIM-3 antibody, or antigen-binding fragments thereof. The combination of antibodies recited herein can be administered separately, e.g., as separate antibodies, or linked, e.g., as a bispecific or trispecific antibody molecule. In one embodiment, a bispecific antibody that includes an anti-PD-1 or PD-L1 antibody molecule and an anti-TIM-3 or anti-LAG-3 antibody, or antigen-binding fragment thereof, is administered. In certain embodiments, the combination of antibodies recited herein is used to treat a cancer, e.g., a cancer as described herein (e.g., a solid tumor). The efficacy of the aforesaid combinations can be tested in animal models known in the art. For example, the animal models to test the synergistic effect of anti-PD-1 and anti-LAG-3 are described, e.g., in Woo et al. (2012) *Cancer Res.* 72(4):917-27).

Exemplary immunomodulators that can be used in the combination therapies include, but are not limited to, e.g., afutuzumab (available from Roche®); pegfilgrastim (Neulasta®); lenalidomide (CC-5013, Revlimid®); thalidomide (Thalomid®), actimid (CC4047); and cytokines, e.g., IL-21 or IRX-2 (mixture of human cytokines including interleukin 1, interleukin 2, and interferon γ , CAS 951209-71-5, available from IRX Therapeutics).

Exemplary doses of such immunomodulators that can be used in combination with the antiviral compounds of the invention include a dose of anti-PD-1 antibody molecule of about 1 to 10 mg/kg, e.g., 3 mg/kg, and a dose of an anti-CTLA-4 antibody, e.g., ipilimumab, of about 3 mg/kg.

Examples of embodiments of the methods of using the antiviral compounds of the invention in combination with an immunomodulator include these, which may be used along with a compound of Formula I or any subgenus or species thereof that is disclosed herein:

- i. A method to treat a viral infection in a subject, comprising administering to the subject a compound of Formula (I) as described herein, and an immunomodulator.
- ii. The method of embodiment i, wherein the immunomodulator is an activator of a

costimulatory molecule or an inhibitor of an immune checkpoint molecule.

iii. The method of either of embodiments i and ii, wherein the activator of the costimulatory molecule is an agonist of one or more of OX40, CD2, CD27, CDS, ICAM-1, LFA-1 (CD11a/CD18), ICOS (CD278), 4-1BB (CD137), GITR, CD30, CD40, BAFFR, HVEM, CD7, LIGHT, NKG2C, SLAMF7, NKp80, CD160, B7-H3 and CD83 ligand.

iv. The method of any of embodiments i-iii above, wherein the inhibitor of the immune checkpoint molecule is chosen from PD-1, PD-L1, PD-L2, CTLA4, TIM3, LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4 and TGFR beta.

v. The method of any of any of embodiments i-iii, wherein the inhibitor of the immune checkpoint molecule is chosen from an inhibitor of PD-1, PD-L1, LAG-3, TIM-3 or CTLA4, or any combination thereof.

vi. The method of any of embodiments i-v, wherein the inhibitor of the immune checkpoint molecule is a soluble ligand or an antibody or antigen-binding fragment thereof, that binds to the immune checkpoint molecule.

vii. The method of any of embodiments i-vi, wherein the antibody or antigen-binding fragment thereof is from an IgG1 or IgG4 (e.g., human IgG1 or IgG4).

viii. The method of any of embodiments i-vii, wherein the antibody or antigen-binding fragment thereof is altered, e.g., mutated, to increase or decrease one or more of: Fc receptor binding, antibody glycosylation, the number of cysteine residues, effector cell function, or complement function.

ix. The method of any of embodiments i-viii, wherein the antibody molecule is a bispecific or multispecific antibody molecule that has a first binding specificity to PD-1 or PD-L1 and a second binding specificity to TIM-3, LAG-3, or PD-L2.

x. The method of any of embodiments i-ix, wherein the immunomodulator is an anti-PD-1 antibody chosen from Nivolumab, Pembrolizumab or Pidilizumab.

xi. The method of any of embodiments i-x, wherein the immunomodulator is an anti-PD-L1 antibody chosen from YW243.55.S70, MPDL3280A, MEDI-4736, MSB-0010718C, or MDX-1105.

xii. The method of any of embodiments i-x, wherein the immunomodulator is an anti-LAG-3 antibody molecule.

xiii. The method of embodiment xii, wherein the anti-LAG-3 antibody molecule is BMS-986016.

xiv. The method of any of embodiments i-x, wherein the immunomodulator is an anti-PD-1 antibody molecule administered by injection (e.g., subcutaneously or intravenously) at a dose of about 1 to 30 mg/kg, e.g., about 5 to 25 mg/kg, about 10 to 20 mg/kg, about 1 to 5 mg/kg, or about 3 mg/kg., e.g., once a week to once every 2, 3, or 4 weeks.

xv. The method of embodiment xiv, wherein the anti-PD-1 antibody molecule is administered at a dose from about 10 to 20 mg/kg every other week.

xvi. The method of embodiment xv, wherein the anti-PD-1 antibody molecule, e.g., nivolumab, is administered intravenously at a dose from about 1 mg/kg to 3 mg/kg, e.g., about 1 mg/kg, 2 mg/kg or 3 mg/kg, every two weeks.

xvii. The method of embodiment xv, wherein the anti-PD-1 antibody molecule, e.g., nivolumab, is administered intravenously at a dose of about 2 mg/kg at 3-week intervals.

The compounds as described herein may be synthesized by the general synthetic routes below, specific examples of which are described in more detail in the Examples.

General Synthetic Procedures

All starting materials, building blocks, reagents, acids, bases, dehydrating agents, solvents, and catalysts utilized to synthesize the compounds of the invention are either commercially available or can be produced by organic synthesis methods known to one of ordinary skill in the art (Houben-Weyl 4th Ed. 1952, Methods of Organic Synthesis, Thieme, Volume 21). General methods for synthesis of compounds of the invention are illustrated by the Examples below, and by methods disclosed in published PCT applications WO2015/113990 and WO2015/173164.

LIST OF ABBREVIATIONS

Ac	acetyl
ACN	Acetonitrile
AcOEt / EtOAc	Ethyl acetate
AcOH	acetic acid
aq	aqueous
Bn	benzyl
Bu	butyl (nBu = n-butyl, tBu = tert-butyl)
CDI	Carbonyldiimidazole
DBU	1,8-Diazabicyclo[5.4.0]-undec-7-ene
Boc ₂ O	di-tert-butyl dicarbonate
DCE	1,2-Dichloroethane
DCM	Dichloromethane
DIAD	Diisopropyl azodicarboxylate
DiBAL-H	Diisobutylaluminum Hydride
DIPEA	N-Ethyl-diisopropylamine
DMA	N,N-dimethylacetamide
DMAP	Dimethylaminopyridine
DMF	N,N'-Dimethylformamide
DMSO	Dimethylsulfoxide

EDCI	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide
EI	Electrospray ionisation
Et ₂ O	Diethylether
Et ₃ N	Triethylamine
Ether	Diethylether
EtOAc	Ethyl acetate
EtOH	Ethanol
FA	Formic acid
FC	Flash Chromatography
h	hour(s)
HCl	Hydrochloric acid
HOBt	1-Hydroxybenzotriazole
HPLC	High Performance Liquid Chromatography
H ₂ O	Water
IPA	isopropanol
L	liter(s)
LC-MS	Liquid Chromatography Mass Spectrometry
LiHMDS	Lithium bis(trimethylsilyl)amide
Me	methyl
MeI	Iodomethane
MeOH	Methanol
mg	milligram
min	minute(s)
mL	milliliter
MS	Mass Spectrometry
Pd/C	palladium on charcoal
PG	protecting group
Ph	phenyl
Ph ₃ P	triphenyl phosphine
Prep	Preparative
Rf	ratio of fronts
RP	reverse phase
Rt	Retention time
rt	Room temperature
SFC	Supercritical Fluid Chromatography
SiO ₂	Silica gel
T3P®	Propylphosphonic acid anhydride
TBAF	Tetrabutylammonium fluoride

TBDMS	t-Butyldimethylsilyl
TEA	Triethylamine
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran
TLC	Thin Layer Chromatography
TsCl	toluene sulfonyl chloride

Within the scope of this text, a readily removable group that is not a constituent of the particular desired end product of the compounds of the present invention is designated a "protecting group," unless the context indicates otherwise. The protection of functional groups by such protecting groups, the protecting groups themselves, and their cleavage reactions are described for example in standard reference works, such as e.g., *Science of Synthesis: Houben-Weyl Methods of Molecular Transformation*. Georg Thieme Verlag, Stuttgart, Germany. 2005. 41627 pp. (URL: <http://www.science-of-synthesis.com> (Electronic Version, 48 Volumes)); J. F. W. McOmie, "Protective Groups in Organic Chemistry", Plenum Press, London and New York 1973, in T. W. Greene and P. G. M. Wuts, "Protective Groups in Organic Synthesis", Third edition, Wiley, New York 1999, in "The Peptides"; Volume 3 (editors: E. Gross and J. Meienhofer), Academic Press, London and New York 1981, in "Methoden der organischen Chemie" (Methods of Organic Chemistry), Houben Weyl, 4th edition, Volume 15/I, Georg Thieme Verlag, Stuttgart 1974, in H.-D. Jakubke and H. Jeschkeit, "Aminosäuren, Peptide, Proteine" (Amino acids, Peptides, Proteins), Verlag Chemie, Weinheim, Deerfield Beach, and Basel 1982, and in Jochen Lehmann, "Chemie der Kohlenhydrate: Monosaccharide und Derivate" (Chemistry of Carbohydrates: Monosaccharides and Derivatives), Georg Thieme Verlag, Stuttgart 1974. A characteristic of protecting groups is that they can be removed readily (i.e., without the occurrence of undesired secondary reactions) for example by solvolysis, reduction, photolysis or alternatively under physiological conditions (e.g., by enzymatic cleavage).

Salts of compounds of the present invention having at least one salt-forming group may be prepared in a manner known per se. For example, salts of compounds of the present invention having acid groups may be formed, for example, by treating the compounds with metal compounds, such as alkali metal salts of suitable organic carboxylic acids, e.g., the sodium salt of 2-ethyl hexanoic acid, with organic alkali metal or alkaline earth metal compounds, such as the corresponding hydroxides, carbonates or hydrogen carbonates, such as sodium or potassium hydroxide, carbonate or hydrogen carbonate, with corresponding calcium compounds or with ammonia or a suitable organic amine, stoichiometric amounts or only a small excess of the salt-forming agent preferably being used. Acid addition salts of compounds of the present invention are obtained in customary manner, e.g., by treating the compounds with an acid or a suitable anion exchange reagent.

Internal salts of compounds of the present invention containing acid and basic salt-forming groups, e.g., a free carboxy group and a free amino group, may be formed, e.g., by the neutralization of salts, such as acid addition salts, to the isoelectric point, e.g., with weak bases, or by treatment with ion exchangers.

Salts can be converted in customary manner into the free compounds; metal and ammonium salts can be converted, for example, by treatment with suitable acids, and acid addition salts, for example, by treatment with a suitable basic agent.

Mixtures of isomers obtainable according to the invention can be separated in a manner known per se into the individual isomers; diastereoisomers can be separated, for example, by partitioning between polyphasic solvent mixtures, recrystallization and/or chromatographic separation, for example over silica gel or by, e.g., medium pressure liquid chromatography over a reversed phase column, and racemates can be separated, for example, by the formation of salts with optically pure salt-forming reagents and separation of the mixture of diastereoisomers so obtainable, for example by means of fractional crystallization, or by chromatography over optically active column materials.

Intermediates and final products can be worked up and/or purified according to standard methods, e.g., using chromatographic methods, distribution methods, (re-) crystallization, and the like.

EXAMPLES

The invention is illustrated by the following examples, which should not be construed as limiting. The assays used to demonstrate the efficacy of compounds of Formula (I) in these assays are generally regarded as predictive of efficacy in subjects.

General Conditions:

Mass spectra were run on LC-MS systems using electrospray ionization. These were WATERS Acquity Single Quad Detector. $[M+H]^+$ refers to mono-isotopic molecular weights. NMR spectra were run on open access Varian 400 or Varian 500 NMR spectrometers. Spectra were measured at 298K and were referenced using the solvent peak. Chemical shifts for 1H NMR are reported in parts per million (ppm).

Mass spectra were run on LC-MS systems with one of the following conditions:

1. Waters Acquity UPLC-H class system equipped with SQD detector.

Column: ACQUITY UPLC HSS C18 (50*2.1) mm, 1.8 μ .

Column temperature: Ambient.

Mobile Phase: A) 0.1% FA + 5mM Ammonium Acetate in Water.

B) 0.1% FA in Acetonitrile.

Gradient: 5-5% solvent B in 0.40 min, 5-35% solvent B in 0.80 min, 35-55% solvent B in 1.2 min,

55-100% solvent B in 2.5 min.

Flow rate: 0.55mL/min.

Compounds were detected by a Waters Photodiode Array Detector.

2. Waters LCMS system equipped with ZQ 2000 detector.

Column: X-BRIDGE C18 (50*4.6) mm, 3.5u.

Column temperature: Ambient.

Mobile Phase: A) 0.1% NH₃ in Water.

B) 0.1% NH₃ in Acetonitrile.

Gradient: 5-95% solvent B in 5.00 min.

Flow rate: 1.0mL/min.

Compounds were detected by a Waters Photodiode Array Detector.

3. Waters ACQUITY UPLC system and equipped with a ZQ 2000 MS system.

Column: Kinetex by Phenomenex, 2.6 um, 2.1 x 50mm


Column temperature: 50 °C

Gradient: 2-88% (or 00-45%, or 65-95%) solvent B over a 1.29 min period


Flow rate: 1.2mL/min.

Compounds were detected by a Waters Photodiode Array Detector.


Example 1: Synthesis of 6-methoxy-8-(methoxymethyl)-12,12-dimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [1]




Step 1: 2-(Allyloxy)-4-bromo-1-methoxybenzene [1.1a]




A round bottom flask was charged with 5-bromo-2-methoxyphenol (15.0 g, 73.9 mmol), allylbromide (11.6 g, 96 mmol), K₂CO₃ (20.4 g, 148 mmol) and acetone (185 mL). The mixture was heated at 60 °C for 3 hours. After it cooled to rt, the mixture was filtered and the filtrate was concentrated to give product (17.9 g, 100% yield). The mixture was continued to the next step with no further purification. ¹H NMR (400 MHz, CDCl₃): 3.84 (s, 3 H), 4.49 - 4.63 (m, 2 H), 5.31 (m, 1 H), 5.41 (m, 1 H), 5.92 - 6.16(m, 1 H), 6.74 (m, 1 H), 6.98 (m, 1 H), 7.01-7.04 (m, 1H).

Step 2. 2-Allyl-3-bromo-6-methoxyphenol [1.1b]

A sealed tube was charged with 2-(allyloxy)-4-bromo-1-methoxybenzene (14.0 g) and DMF (14.4 mL) and then purged with nitrogen. The solution was heated at 200 °C for 3 hours. The solvent was then removed under reduced pressure and the residue was purified by silica gel column chromatography, EtOAc/heptane 0 to 50%, to give product (12 g, 86% yield). ¹H NMR (400 MHz, CDCl₃): 3.57 (dt, *J*=6.09, 1.38 Hz, 2 H), 3.87 (s, 3 H), 4.98 - 5.14 (m, 2 H), 5.80 (s, 1 H), 5.95 (m, 1 H), 6.63 (d, *J*=8.71 Hz, 1 H), 7.06 (d, *J*=8.66 Hz, 1 H).


Step 3. (4-Bromo-7-methoxy-2,3-dihydrobenzofuran-2-yl)methanol [1.1c]

To a solution of 2-allyl-3-bromo-6-methoxyphenol (5.7g, 23.45 mmol) in DCM (47 mL) at 0 °C was added mCPBA (12.14 g, 50%, 35.2 mmol) and the resulting mixture was stirred at rt for 2 hours. The reaction was quenched by adding sat. aq. Na₂S₂O₃ solution, and sat. aq. NaHCO₃ solution and then extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was dissolved in MeOH (50 mL) and K₂CO₃ (9.7 g, 70 mmol) was added. The mixture was then stirred at rt for 2 hours and then filtered. The filtrate was concentrated and the residue was purified by silica gel column chromatography, EtOAc/heptane 0 to 60% to give product (3.9 g, 64 % yield). ¹H NMR (400 MHz, CDCl₃): 3.01 - 3.44 (m, 2 H), 3.70 - 3.81 (m, 1 H), 3.85 (s, 3 H), 3.88 - 4.05 (m, 1 H), 4.88 - 5.10 (m, 1 H), 6.64 (d, *J*=8.61Hz, 1 H), 6.94 (d, *J*=8.66 Hz, 1 H).

Step 4. 4-Bromo-7-methoxy-2-(methoxymethyl)-2,3-dihydrobenzofuran [1.1d]


To a solution of (4-Bromo-7-methoxy-2,3-dihydrobenzofuran-2-yl)methanol (3.9 g, 15 mmol) in DMF (15 mL) at 0 °C was added NaH (1.2 g, 60%, 30 mmol). After stirring at rt for 20 mins, the mixture was placed in an ice water bath and MeI (6.4 g, 45 mmol) was added and the mixture was stirred at rt for 2 hours. The reaction was quenched by adding sat. aq. NH₄Cl solution. The mixture was then extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, EtOAc/heptane 0 to 40% to give product (3.6 g, 88% yield). LCMS (*m/z*): 273.1 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): 3.05 - 3.16 (m, 1 H), 3.20 - 3.34 (m, 1 H), 3.37 - 3.52 (m, 3 H), 3.55 - 3.71 (m, 2 H), 3.83 (s, 3 H), 5.01 (ddt, *J*=9.58, 7.90, 4.90, 4.90 Hz, 1 H), 6.62 (d, *J*=8.66 Hz, 1 H) 6.91 (d, *J*=8.66 Hz, 1 H).

Step 5. 5-(7-methoxy-2-(methoxymethyl)-2,3-dihydrobenzofuran-4-yl)-2,2-dimethylcyclopentan-1-one [1.1e]




A tube was charged with Pd(OAc)₂ (89 mg, 0.4 mmol), dicyclohexyl(2'-methyl-[1,1'-biphenyl]-2-yl)phosphane (0.29 g, 0.79 mmol), NaO-tBu (1.9 g, 20 mmol), 4-Bromo-7-methoxy-2-(methoxymethyl)-2,3-dihydrobenzofuran (3.6 g, 13.2 mmol), 2,2-dimethylcyclopentanone (3.0 g, 26.4 mmol) and toluene (13 mL). The tube was then purged with nitrogen and sealed, and heated at 60 °C for 6 hours. After cooling to rt, the mixture was then diluted with EtOAc and washed with sat. aq NH₄Cl solution, brine, dried over Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, EtOAc/heptane 0 to 50% to give product (2.8 g, 69.8 % yield). LCMS (m/z): 305.5 [M+H]⁺.

Step 6. 5-(7-methoxy-2-(methoxymethyl)-2,3-dihydrobenzofuran-4-yl)-2,2-dimethylcyclopentan-1-amine [1.1f]



To a solution of **1.1e** (2.8 g, 9.2 mmol) in MeOH was added ammonium acetate (7.1 g, 92 mmol) and sodium cyanoborohydride (1.7 g, 28 mmol). The mixture was stirred at 60 °C for 3 days. The reaction solution was concentrated under vacuum and then quenched by adding 10% NaOH aq. solution. The mixture was extracted with DCM. The organic layer was washed with brine, dried over MgSO₄ and concentrated. The crude material was continued to the next step with no further purification. LCMS (m/z): 306.5 [M+H]⁺.


Step 7. N-(2-(7-methoxy-2-(methoxymethyl)-2,3-dihydrobenzofuran-4-yl)cyclopentyl)formamide N-(5-(7-methoxy-2-(methoxymethyl)-2,3-dihydrobenzofuran-4-yl)-2,2-dimethylcyclopentyl)formamide [1.1g]




To a solution of **1.1f** (2.8 g, 9.17 mmol) in DMF (15 mL) at 0 °C was added EDCl (3.5 g, 18.3 mmol) and DIPEA (4.8 ml, 27.5 mmol) followed by dropwise addition of formic acid (1.6 g, 37 mmol). The mixture was then stirred at rt for 3 hours. The reaction mixture was then diluted with EtOAc and washed with 1.0 N HCl aq solution, water, and brine, dried over MgSO₄ and concentrated. The crude material was purified by silica gel column

chromatography, acetone/heptane 0 to 60% to give product (2.1 g, 69% yield). LCMS (m/z): 334.3 [M+H]⁺.

Step 8. 4-methoxy-2-(methoxymethyl)-8,8-dimethyl-1,7a,8,9,10,10a-hexahydro-2H-cyclopenta[c]furo[3,2-f]isoquinoline [1.1h]




To a solution of **1.1g** (2.1g, 6.30 mmol) in CH₃CN (21 mL) was added POCl₃ (1.06 g, 6.9 mmol) and the solution was then stirred at 70 °C for 2 hours. The solvent was removed under reduced pressure and the residue was dissolved in EtOAc, and basified by adding ammonium hydroxide solution until pH=11. The organic layer was washed with brine, dried over MgSO₄ and concentrated. The residue was purified by silica gel column chromatography, EtOAc/heptane 0% to 100% followed by 100% acetone to give product. The more polar isomer is the desired cis isomer **1.1h** (800 mg, 40% yield).




LCMS (m/z): 316.6 [M+H]⁺.

Step 9. Ethyl 6-methoxy-8-(methoxymethyl)-12,12-dimethyl-3-oxo-3,4,4a,8,9,9c,10,11,12,12a-decahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylate [1.1i]



A vial was charged with **1.1h** (800 mg, 2.5 mmol), ethyl (Z)-2-(ethoxymethylene)-3-oxobutanoate (1.4 g, 7.6 mmol) and EtOH (5.1 mL). The solution was then heated at 120 °C for 24 hours. The solvent was removed under reduced pressure and the residue was continued to the next step with no further purification. LCMS (m/z): 456.2 [M+H]⁺.

Step 10. Ethyl 6-methoxy-8-(methoxymethyl)-12,12-dimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylate [1.1j]




A mixture of **1.1i** (1.1 g, 2.4 mmol), p-chloranil (0.71 g, 2.9 mmol) and DME (2.7 mL) was heated at 100 °C for 2 hours. The solvent was then removed under reduced pressure and the residue was purified by silica gel column chromatography, DCM/MeOH 0 to 50% to give product (750 mg, 69%). LCMS (m/z): 454.3 [M+H]⁺.

The material was then separated by chiral SFC, OD column, flow rate 5mL/min, CO₂/MeOH=80/20 to give four isomers.


1.1j-1: t_R 2.7 min; **1.1j-2**: t_R 3.4 min; **1.1j-3**: t_R 5.4 min; **1.1j-4**: t_R 8.5 min.

Step 11. 6-methoxy-8-(methoxymethyl)-12,12-dimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [1]




To a solution of **1.1j-4** (60 mg, 0.137 mmol) in THF (0.5 mL) was added LiOH (1.0 M in water, 0.27 mL, 0.27 mmol). After stirring at rt for 2 hours, the reaction was acidified by adding 1.0 N HCl aq. solution. The mixture was extracted with DCM. The organic layer was concentrated and purified by reverse phase HPLC to give product **1** (47 mg, 80%). LCMS (m/z): 426.2 [M+H]⁺. ¹H NMR (500 MHz, CDCl₃): 0.60 (s, 3 H), 1.2 (s, 3H), 1.51 - 1.76 (m, 2 H) 1.98 - 2.15 (m, 1 H) 2.45-2.48 (m, 1 H) 3.15 -3.18 (m, 1 H) 3.33-3.35 (m, 1 H) 3.49 (s, 3 H) 3.72 - 3.80 (m, 1 H) 3.82 - 3.92 (m, 2 H) 3.96 (s, 3 H) 4.18-4.20 (m, 1H) 4.94 - 5.14 (m, 1 H) 7.23 (s, 1 H) 7.39 (s, 1 H) 8.46 (s, 1 H)

Example 1.2: Synthesis of 4-fluoro-6-methoxy-8-(methoxymethyl)-12,12-dimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid



1.2


Step 1: ethyl (Z)-2-(ethoxymethylene)-4,4-difluoro-3-((trimethylsilyl)oxy)but-3-enoate [1.2a]



1.2a

Under an argon atmosphere, a mixture of Mg (3.69 g, 152 mmol) and TMSCl (19.43 mL, 152 mmol) was treated with ultrasound irradiation for 15 min. To the mixture was added DMF (30 mL), and ethyl (Z)-2-(ethoxymethylene)-4,4,4-trifluoro-3-oxobutanoate (4.56 g, 19mmol) was added dropwise at 50 °C under an argon atmosphere. The reaction mixture was stirred for additional 3 min. at 50 °C. After removal of excess TMSCl in vacuo, the crude mixture was filtered and the filtrate (containing **1.2a** and DMF) was used in the next step without further purification.

Step 2: Ethyl 4-fluoro-6-methoxy-8-(methoxymethyl)-12,12-dimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylate [1.2b]




1.2b

To a mixture of zinc iodide (961 mg, 3.01 mmol) and 4-methoxy-2-(methoxymethyl)-8,8-dimethyl-1,7a,8,9,10,10a-hexahydro-2H-cyclopenta[c]furo[3,2-f]isoquinoline (950 mg, 3.01 mmol) in acetonitrile (10 ml) was added dropwise the crude solution of (Z)-ethyl 2-(ethoxymethylene)-4,4-difluoro-3-((trimethylsilyl)oxy)but-3-enoate (4433 mg, 15.06 mmol) in DMF (20 ml). The reaction mixture was stirred at 50 °C for 18 hrs and then at 100 °C for 4 hours. The mixture was concentrated by high vacuum oil pump with 50 °C water bath for 1 h. To the residue was added DCM (10 mL) and the resultant mixture was stirred for 5 min

then filtered. The solution was directly loaded to silica gel and flushed with acetone/heptane 70% to give product (1.2 g, 84 % yield). LCMS (m/z): 472.5 [M+H]⁺.

Step 3: 4-fluoro-6-methoxy-8-(methoxymethyl)-12,12-dimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [1.2]



NaOH (1.5 ml, 7.5 mmol, 5N solution) was added to a solution of **1.2b** (600 mg, 1.272 mmol) in THF (10mL) and the mixture was stirred for 30 min. The solution was then acidified by adding 4.0 N HCl, diluted with water, and extracted twice with DCM. The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The material was purified by chiral chromatography (AD column, 4.6x100 mm, SFC 5 mL/min, CO₂/IPA=70/30), and returned as three isomers.

1.2-I: tR 2.32 min; single enantiomer, LC-MS (m/z): 444.5 [M+H]⁺.


1.2-II: tR 3.65 min, mixtures of two isomers; LC-MS (m/z): 444.5 [M+H]⁺.

1.2-III: tR 4.53 min; single enantiomer, LC-MS (m/z): 444.5 [M+H]⁺. ¹H NMR (400 MHz, Acetonitrile-*d*₃) δ 8.37 (s, 1H), 7.78 (d, *J* = 2.8 Hz, 1H), 5.15 (dtd, *J* = 9.6, 5.6, 3.7 Hz, 1H), 4.43 (d, *J* = 9.8 Hz, 1H), 3.88 (s, 3H), 3.63 – 3.46 (m, 3H), 3.34 (s, 3H), 3.02 (dd, *J* = 15.8, 5.8 Hz, 1H), 2.43 (dtd, *J* = 12.9, 8.5, 4.1 Hz, 1H), 1.73 – 1.44 (m, 4H), 1.19 (s, 3H), 0.55 (s, 3H).


For the compound **1.2-I** of Example 1.2, EC₅₀ in the HBsAg assay was 12 nM.

For the compound **1.2-II** of Example 1.2, EC₅₀ in the HBsAg assay was 17 nM.


For the compound **1.2-III** of Example 1.2, EC₅₀ in the HBsAg assay was 1.4 nM. The relative stereochemistry of the methoxymethyl group on the dihydrofuran ring of this isomer was not determined, but it is believed to be that the saturated ring has the absolute configuration shown here:



Example 2: Synthesis of (10cR,13aS)-6-methoxy-13,13-dimethyl-3-oxo-8,9,10,10c,11,12,13,13a-octahydro-3H-cyclopenta[c]pyrano[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [2]




Step 1. 3-bromo-2-(3-hydroxypropyl)-6-methoxyphenol [2.1a]




To a solution of **1.1b** (5 g, 20.6 mmol) in THF (68 ml) at 0 °C was added borane-methyl sulfide complex (3.9 ml, 41.1 mmol) and then the solution was stirred at rt for 2 hours. At 0 °C, 5 mL of water was added followed by aq. NaOH solution (5.0 M, 12 mL, 62 mmol) and sodium perborate tetrahydrate (9.5 g, 62 mmol). The mixture was then heated at 50 °C for 1 hour. After cooling to rt, the mixture was extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated. The residue was purified by silica gel column chromatography, EtOAc/heptane 0 to 60% to give product (4.7 g, 88%). LCMS (m/z): 261.0 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): 1.88 (m, 2 H), 2.93 (m, 2 H), 3.59 (m, 2 H), 3.88 (s, 3 H), 6.08 (s, 1 H) 6.63 (m, 1 H) 7.06 (m, 1 H).

Step 2. 5-bromo-8-methoxychromane [2.1b]




To a solution of **2.1a** (4.7g, 18.00 mmol) in DCM at 0 °C was added PPh₃ (5.6 g, 21.6 mmol) followed by diethyl diazene-1,2-dicarboxylate (3.45 g, 3.1 mmol) and the resulting mixture was stirred at rt for 18 hours. The solvent was removed and to the residue was added diethyl ether. The white precipitate was removed by filtration and the filtrate was concentrated. The residue was dissolved in EtOH (5mL) and H₂O₂ (35%, 1.0 mL) was added. After 20 mins, the solution was concentrated and the residue was purified by silica gel column chromatography, EtOAc/heptane 0 to 30% to give product. LCMS (m/z): 244.0 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): 1.88 - 2.18 (m, 2 H), 2.74 (t, J=6.63 Hz, 2 H), 3.84 (s, 3 H), 4.12 - 4.30 (m, 2 H), 6.62 (d, J=8.71 Hz, 1 H), 7.06 (d, J=8.66 Hz, 1 H).

Step 3. Ethyl 6-methoxy-13,13-dimethyl-3-oxo-8,9,10,10c,11,12,13,13a-octahydro-3H-cyclopenta[c]pyrano[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylate [2.1c]



Compound **2.1c** was synthesized from **2.1b** by the method described in the synthesis of **1.1** Step 5-10. LCMS (m/z): 424.2 [M+H]⁺.


Step 4. (10cR,13aS)-6-methoxy-13,13-dimethyl-3-oxo-8,9,10,10c,11,12,13,13a-octahydro-3H-cyclopenta[c]pyrano[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [2]




Compound **2.1c** was separated by chiral SFC OD column, flow rate 5 mL/min, CO₂/MeOH 70/30 to give two enantiomers **2.1c-1** (tR 1.26 min) and **2.1c-2** (tR 3.13 min).

To a solution of **2.1c-2** (14 mg, 0.034 mmol) in THF (0.17 mL) was added LiOH (1.0 M in water, 0.068, 0.068 mmol). After stirring at rt for 2 hours, the reaction was acidified by adding 1.0 N HCl aq. solution. The mixture was extracted with DCM. The organic layer was concentrated and purified by reverse phase HPLC to give product (9 mg, 66%). LCMS (m/z): 396.2 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): 0.43 (s, 3 H), 1.08 (s, 3 H), 1.41 - 1.57 (m, 2 H), 1.65 - 1.79 (m, 2 H), 2.03-2.07 (m, 1 H), 2.51 - 2.66 (m, 2 H), 2.69 - 2.81 (m, 1 H), 3.77 - 3.90 (m, 1 H), 3.83 (s, 3H), 4.01-4.06 (m, 1 H) 4.37-4.38 (m, 1 H) 4.65-4.68 (m, 1 H) 7.40 (s, 1 H) 7.50 (s, 1 H) 8.52 (s, 1 H).

Example 3. Synthesis of 6-methoxy-12,12-dimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [3]




Step 1. (2-allyl-3-bromo-6-methoxyphenoxy)(tert-butyl)dimethylsilane [3a]



To a solution of 2-allyl-3-bromo-6-methoxyphenol (12.0 g, 49 mmol) and TEA (7.4 g, 74 mmol) at 0 °C was added TBSOTf (17.0 g, 64 mmol) and the resulting solution was


stirred at room temperature for 2 hours. Reaction the mixture was then diluted by DCM and washed with water. The phases were separated and the organic layer was dried over Na_2SO_4 and concentrated. The residue was purified by column chromatography, EtOAc/hexane 20-60%, to give product (16.0 g, 92% yield). ^1H NMR (400 MHz, CDCl_3): 7.28-7.137 (d, 1H), 6.60-6.72 (m, 1H), 6.035-6.098 (m, 1H), 5.034-5.133 (d, 2H), 3.894 (s, 3H), 0.98 (12H, s), 0.204 (6H, s)

Step 2. 2-(6-bromo-2-((tert-butyldimethylsilyl)oxy)-3-methoxyphenyl)acetaldehyde [3b]




To a mixture of (2-allyl-3-bromo-6-methoxyphenoxy)(tert-butyl)dimethylsilane (16.0 g, 44 mmol) and 1,4-Dioxane/ H_2O (1/1) (230mL) at 0 °C was added 2,6-Lutidine (9.6 g, 89 mmol) and OsO_4 (0.22 g, 0.89 mmol). The mixture was stirred for 30 min, after which NaOCl_4 (38.3 g, 179 mmol) was added and the mixture was stirred at 0 °C for 2 hours. The reaction mixture was filtered and filtrate was wash with brine and diluted aq. HCl solution. The organic layer was dried over Na_2SO_4 and concentrated. The crude material was directly taken to the next step without further purification (15.0 g, 93% yield). ^1H NMR (400 MHz, CDCl_3): 9.73 (s, 1H), 7.190-7.282 (d, 1H), 6.62-6.74 (m, 1H) 3.95 (s, 1H), 3.810 (s, 3H), 0.98 (s, 9H), 0.204 (s, 6H)


Step 3. 2-(6-bromo-2-((tert-butyldimethylsilyl)oxy)-3-methoxyphenyl)ethan-1-ol [3c]




NaBH_4 (3.17 g, 83.5 mmol) was added to a solution of 2-(6-bromo-2-((tert-butyldimethylsilyl)oxy)-3-methoxyphenyl)acetaldehyde (15.0 g, 41 mmol) in MeOH (150.0 mL) and the resulting mixture was warmed to rt and stirred for 30 mins. The mixture was then concentrated and to the crude material was added water and EtOAc. NaOH aq. solution (20%) was added and the mixture was stirred for 20 mins. The reaction mixture was extracted with EtOAc. The organic layer was washed with water, brine, dried over Na_2SO_4 and concentrated. The crude material was used in the next step without further purification (13.0 g, 86% yield).

Step 4. 3-bromo-2-(2-hydroxyethyl)-6-methoxyphenol [3d]

TBAF (1.0 M in THF) (18.8 g, 72 mmol) was added to a mixture of 2-(6-bromo-2-((tert-butyldimethylsilyl)oxy)-3-methoxyphenyl)ethan-1-ol (13.0 g, 36 mmol) in THF (150 mL) and the resulting mixture was stirred at room temperature for 1 h. The mixture was concentrated under vacuum and the remaining material was purified by silica gel column chromatography, EtOAc/ hexane 20-70%, to give product (5.0 g, 56%). LCMS (m/z): 230 [M-18]⁺. ¹H NMR (400 MHz, CDCl₃): 7.087-7.109 (d, 1H), 6.66-6.68 (m, 1H) 6.214 (s, 1H), 3.92 (5H, s), 3.143-3.175 (m, 2H).

Step 5. 4-bromo-7-methoxy-2,3-dihydrobenzofuran [3e]


DEAD (2.89 g, 16 mmol) and PPh₃ (4.7 g, 17mmol) were added to a solution of 3-bromo-2-(2-hydroxyethyl)-6-methoxyphenol (5.0 g, 13 mmol) in DCM (70mL) at 0 °C and the resulting mixture was stirred at room temperature for 15 hours. The mixture was then concentrated and diethyl ether was added. The mixture was filtered and the filtrate was concentrated. The residue was dissolved in ethanol and 1.0 mL H₂O₂ was added. After 30 min, the solution was concentrated and the residue was purified by silica gel column chromatography, EtOAc/hexane 20-70% to give product. ¹H NMR (400 MHz, CDCl₃): 6.93-6.95 (d, 1H), 6.64-6.66 (m, 1H), 4.70-4.66 (m, 3H), 3.895 (s, 3H), 3.281-3.236 (m, 2H).

Step 6. Synthesis of 6-methoxy-12,12-dimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [3]

Compound **3** was synthesized from **3e** by the method described in the synthesis of **1.1** Step 5-11. The material was separated by chiral SFC, AD column, 100 mL/min, CO₂/IPA 65/35 to give four isomers. **3-I**: t_R 3.5 min, **3-II**: t_R 4.8 min, **3-III**: t_R 6.7 min, **3-IV**: t_R 9.2 min.


3-III: ^1H NMR (500 MHz, CDCl_3): 0.61 (s, 3 H), 1.26 (s, 3 H), 1.61 -1.65 (m, 2 H), 1.93 - 2.18 (m, 1 H), 2.47 (m, 1 H), 3.17 (m, 1 H), 3.43 (m, 1 H), 3.83 - 3.90 (m, 1 H), 3.93 - 3.99 (s, 3 H), 4.17 (m, 1 H), 4.66 (m, 1 H), 4.78 - 4.96 (m, 1 H), 7.06 - 7.14 (m, 1 H), 7.18 (d, $J=17.97$ Hz, 2 H), 8.42 (s, 1 H), 8.81 (s, 1 H), 15.83 - 16.39 (m, 1 H). LCMS (m/z): 382.2 $[\text{M}+\text{H}]^+$.

Example 4. 6-methoxy-8-(methoxymethyl)-8,12,12-trimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-




carboxylic acid [4]

Step-1. 4-bromo-1-methoxy-2-((2-methylallyl)oxy)-benzene [4a]




To the solution of 5-bromo-2-methoxyphenol (10.0 g, 49.2 mmol) in acetone was added K_2CO_3 (13.59 g, 98.4 mmol) and 3-bromo-2-methylprop-1-ene (8.64 g, 64.0 mmol). The resulting mixture was heated to reflux for overnight. The mixture was then filtered and the filtrate was concentrated. The residue was dissolved in DCM and washed with water, then brine. The separated organic layer was dried over Na_2SO_4 and concentrated to give product (11.0 g). ^1H NMR (400 MHz, CDCl_3): 7.28–6.99 (m, 2H), 6.77–6.41 (m, 1H), 5.11 (s, 1H), 5.02 (s, 1H), 4.505 (s, 2H), 3.864 (s, 3H), 1.84 (s, 3H)

Step 2. ethyl 6-methoxy-8-(methoxymethyl)-8,12,12-trimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylate [4b]



Compound **4b** was synthesized from **4a** by the method described in the synthesis of 1.1 Step 2-9. LCMS (m/z): 468.5 [M+H]⁺. The material was separated via chiral SFC to give isomers **4b-1** and **4b-2**. **4b-1**: AD column, t_R 8.02 min. **4b-2**: t_R 9.22 min.

Step 3. 6-methoxy-8-(methoxymethyl)-8,12,12-trimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [4]




To a solution of **4b-1** (16 mg, 0.034 mmol) in THF (0.17 mL) was added LiOH (1.0 M in water, 0.05 mL) and the mixture was stirred at rt for 1 hour. The solution was then acidified by adding 1.0 N aq HCl solution and extracted with DCM. The organic layers were combined and concentrated. The residue was purified by chiral SFC, OD column, 100 mL/min, CO₂/MeOH 70/30 to give isomers **4-I** and **4-II**.


4-I: t_R 2.05 min.

4-II: t_R 5.33 min. LCMS (m/z): 440.3 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): 0.58 (s, 3 H), 1.16 - 1.28 (s, 3 H), 1.42 (s, 3 H), 1.51 - 1.71 (m, 2 H), 1.95 - 2.14 (m, 1 H), 2.42 (m, 1 H), 2.76 (m, 1 H), 3.44 (s, 3 H), 3.47 - 3.56 (m, 2 H), 3.67 (m, 1 H), 3.77 - 3.86 (m, 1 H), 3.92 (s, 3 H), 4.13 (m, 1 H), 7.13 (s, 1 H), 7.15 (s, 1 H), 8.38 (s, 1 H).

Example 5. Synthesis of 6-methoxy-8,12,12-trimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [5]




Step 1: 4-bromo-7-methoxy-2-methyl-2,3-dihydrobenzofuran [5a]



To a solution of **1.1b** (10 g, 0.041 mol) in dry toluene (100mL) was added p-TSA (31.5 g, 0.165 mol) and the mixture was stirred at 110 °C for 12 h. The mixture was then filtered and the filtrate was concentrated. The residue was purified by silica gel column

chromatography (0-5% EtOAc/hexane 0-5%) to give product (8.5 g, 85%). $^1\text{H NMR}$ (400 MHz, CDCl_3): 6.93 (d, $J = 8.6$ Hz, 1H), 6.73 – 6.57 (m, 1H), 5.05 (m, 1H), 3.91 (s, 3H), 3.36 (m, 1H), 2.87 (m, 1H), 1.54 – 1.18 (m, 3H).

Step 2. 6-methoxy-8,12,12-trimethyl-3-oxo-3,8,9,9c,10,11,12,12a-octahydrocyclopenta[c]furo[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [5]




Compound **5** was synthesized from **5a** by the method described in the synthesis of 1.1 Step 4-11.


The product was separated by chiral SFC, AD column, flow 100ml/min, $\text{CO}_2/\text{EtOH}=60/40$ to give four isomers. **5-I**: t_R 2.3 min, **5-II**: t_R 3.8 min, **5-III**: t_R 6.9 min, **5-IV**: t_R 8.7 min.

5-II: LCMS (m/z): 396.3 $[\text{M}+\text{H}]^+$. $^1\text{H NMR}$ (500 MHz, CDCl_3): 0.61 (s, 3 H), 1.26 (s, 3 H), 1.67 (m, 3 H), 1.57 - 1.78 (m, 2 H), 1.96 - 2.10 (m, 1 H), 2.43 (m, 1 H), 3.00 (m, 1 H), 3.26 (m, 1 H), 3.86 (br s, 1 H), 3.96 (s, 3 H), 4.16 (m, 1 H), 4.96 - 5.08 (m, 1 H), 7.15 (s, 1 H), 7.19 (s, 1 H), 8.41 (s, 1 H), 15.97 - 16.34 (m, 1 H)

Example 6. Synthesis of 6-methoxy-9-(methoxymethyl)-13,13-dimethyl-3-oxo-8,9,10,10c,11,12,13,13a-octahydro-3H-cyclopenta[c]pyrano[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [6]




Step 1. 2-((5-bromo-2-methoxyphenoxy)methyl)prop-2-en-1-ol [6.1]




To a solution of 5-bromo-2-methoxyphenol (9.08 g, 44.7 mmol) in DMF (89 ml) at 0 °C was added NaH (2.1 g, 60 % in mineral oil, 53.7 mmol,) and the solution was then stirred at rt for 30 mins. 5-methylene-1,3,2-dioxathiane 2-oxide (6.0g, 44.7 mmol) was then added and the mixture was heated at 55 °C for 4 hours. The reaction was quenched by adding sat. aq. NH₄Cl solution, and the mixture was extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated. The residue was purified by silica gel column chromatography, EtOAc/heptane 0 to 60%, to give product (8.5 g, 69.6 % yield). ¹H NMR (400 MHz, CDCl₃): 4.29 (s, 2 H), 4.65 (s, 2 H), 5.33 (d, J=1.03 Hz, 2 H), 7.01 - 7.10 (m, 2 H), 7.22 (d, J=8.41 Hz, 1 H)

Step 2. 4-bromo-1-methoxy-2-((2-(methoxymethyl)allyl)oxy)benzene [6.2]




To a solution of **6.1** (8.5 g, 31.1 mmol) in DMF at 0 °C was added NaH (2.489 g, 62.2 mmol). After stirring at rt for 20 mins, MeI (5.84 ml, 93 mmol) was added and the resulting mixture was stirred at rt for 2 hours. The reaction was then quenched by adding sat. aq. NH₄Cl solution and the mixture was extracted with EtOAc. The organic layer was washed with brine, dried over MgSO₄ and concentrated. The residue was purified by silica gel column chromatography, EtOAc/heptane 0 to 40% to give product 8.2 g (92% yield). ¹H NMR (400 MHz, CDCl₃): 3.35 (s, 3 H), 3.76 - 3.89 (m, 4 H), 4.02 (s, 2 H), 4.50 - 4.71 (m, 1 H), 4.50 - 4.71 (m, 1 H), 4.59 (s, 1 H), 5.19 - 5.46 (m, 2 H), 6.67 - 6.78 (m, 1 H), 6.67 - 6.78 (m, 1 H), 6.67 - 6.78 (m, 1 H), 6.97 - 7.09 (m, 2 H)

Step 3. 3-bromo-6-methoxy-2-((2-(methoxymethyl)allyl)phenol [6.3]




A solution of **6.2** (8.2 g, 28.6 mmol) in DMF (14 ml) was heated at 200 °C for 3 hours. The solvent was then removed under vacuum and the residue was purified by silica gel column chromatography, EtOAc/heptane 0 to 40%, to give product (7.2 g, 88% yield). ¹H NMR (400 MHz, CDCl₃): 3.38 (s, 3 H), 3.55 (s, 2 H), 3.87 (s, 3 H), 3.97 (s, 2 H), 4.73 (s, 1 H), 5.05 (m, 1 H), 6.21 (s, 1 H), 6.65 (m, 1 H), 7.07 (m, 1 H)

Step 4. ethyl 6-methoxy-9-(methoxymethyl)-13,13-dimethyl-3-oxo-8,9,10,10c,11,12,13,13a-octahydro-3H-cyclopenta[c]pyrano[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylate [6.4]




Compound **6.4** was synthesized from **6.3** by the method described in the synthesis of **1.1** Step 3-10. LCMS (m/z): 468.4 [M+H]⁺. The material was separated by chiral SFC, OD column, flow 100 ml/min, CO₂/MeOH=70/30, to give isomers. **6.4-I**: t_R 4.51 min, **6.4-II**: t_R 5.36 min, **6.4-III**: t_R 7.69 min, **6.4-IV**: t_R 19.0 min.

Step 5. 6-methoxy-9-(methoxymethyl)-13,13-dimethyl-3-oxo-8,9,10,10c,11,12,13,13a-octahydro-3H-cyclopenta[c]pyrano[3,2-f]pyrido[2,1-a]isoquinoline-2-carboxylic acid [6]



To a solution of **6.4-IV** (25 mg, 0.053 mmol) in THF (0.27 mL) was added LiOH (1.0 M in water, 0.11 mL, 0.11 mmol). After stirring at rt for 1 hours, the reaction was acidified by adding 1.0 N HCl aq. solution. The mixture was extracted with DCM. The organic layer was concentrated and purified by reverse phase HPLC to give product **6** (19 mg, 80%). LCMS (m/z): 440.2 [M+H]⁺. ¹H NMR (500 MHz, CDCl₃): 0.61 (s, 3 H), 1.25 (s, 3 H), 1.60 - 1.69 (m, 2 H), 1.87 - 1.97 (m, 1 H), 2.60 - 2.75 (m, 4 H), 3.41 (s, 3 H), 3.50 (m, 2 H), 3.87 - 3.97 (m, 2 H), 3.98 (s, 3 H), 4.24 (m, 1 H), 4.58 (m, 1 H), 7.21 (s, 1 H), 7.47 - 7.53 (m, 1 H), 8.48 (s, 1 H)

The following compounds can be made by similar methods to those illustrated above using starting materials that are known in the art:



BIOLOGICAL EXAMPLES

HBV Cell Line

HepG2-Clone42, a Tet-inducible HBV-expressing cell line with a stably integrated 1.3mer copy of the HBV ayw strain, was generated based on the Tet-inducible HepAD38 cell line with slight modifications. Ladner SK, et al., *Antimicrobial Agents and Chemotherapy*. 41(8):1715-1720 (1997). HepG2-Clone42 cells were cultured in DMEM/F-12 + Glutamax™ (Life Technologies, Carlsbad, CA, USA), supplemented with 10% fetal bovine serum (Life Technologies), G-418 (Corning, Manassas, VA, USA) at a final concentration of 0.5 mg/mL, and 5 µg/mL Doxycycline (Sigma, St. Louis, MO, USA) and maintained in 5% CO₂ at 37°C.

HBsAg Assay

HepG2-Clone42 cells were seeded into black clear-bottom 96-well plates at a concentration of 6.0×10^4 cells/well. 24 hours post-seeding, the cells were treated with 200 μ L/well of media containing five-fold serial dilutions of compounds in DMSO. DMSO alone was used as the no drug control. The final DMSO concentration in all wells was 0.5%.

The HBsAg ELISA kit (Alpha Diagnostic International, San Antonio, TX, USE, Catalog # 4110) was used to determine the level (semi-quantitative) of secreted HBV sAg. The HBsAg ELISA assay was performed following the manufacturer's protocol as described.

Step 1. Pipet 100 μ L each of compound or DMSO treated samples into HBsAg ELISA plates. Seal plates and incubate at room temp for 60 minutes.

Step 2. Aspirate samples and wash three times with Wash Buffer. Dispense 100 μ L of antibody-HRP conjugate to each well. Incubate at room temp for 30 minutes.

Step 3. Aspirate samples and wash three times with Wash Buffer. Add 100 μ L of TMB Substrate to all wells and incubate 15 minutes at room temp.

Step 4. Dispense 100 μ L of Stop Solution to each well. Measure absorbance of ELISA plate at 450 nm.

Dose Response Curves

Dose-response curves were generated and the EC_{50} value was defined as the compound concentration at which HBsAg secretion was reduced 50% compared to the DMSO control.

EC_{50} values were determined as follows:

1. Determine the percent of HBsAg secretion inhibition. Calculate the percent inhibition on of HBsAg secretion inhibition using the following equation:

$$100 \times (X_C - M_B) / (M_D - M_B)$$

where X_C is the absorbance signal from compound-treated well; M_B is average absorbance signal (background signal) for column 12 (no cells + HBsAg ELISA sample buffer) and M_D is average absorbance signal from DMSO-treated wells. Then calculate EC_{50} values by non-linear regression using a four parameter curve logistic equation.


The curve fit model employed is XLFit Dose Response One Site Model 204: $y = (A + ((B - A) / (1 + (10^{((C - x) * D))})))$ where A is the minimum y value, B is the maximum y value, C is the log EC_{50} value, and D is the slope factor.

For the compound 1 of Example 1, EC_{50} in the HBsAg assay was 0.2 nM.

For the compound **1.2-I** of Example 1.2, EC₅₀ in the HBsAg assay was 12 nM.
For the compound **1.2-II** of Example 1.2, EC₅₀ in the HBsAg assay was 17 nM.
For the compound **1.2-III** of Example 1.2, EC₅₀ in the HBsAg assay was 1.4 nM.
For the compound **2** of Example 2, EC₅₀ in the HBsAg assay was 0.5 nM
For the compound **3-III** of Example 3, EC₅₀ in the HBsAg assay was 0.7 nM.
For the compound **4-II** of Example 4, EC₅₀ in the HBsAg assay was 4.3 nM.
For the compound **5-II** of Example 5, EC₅₀ in the HBsAg assay was 0.2 nM.
For the compound **6** of Example 6, EC₅₀ in the HBsAg assay was 10 nM.

CLAIMS

1. A compound of formula (I):



(I) wherein:

R¹ is H, halo, or C₁-C₃ alkyl;

R² is H, halo, CN, C₁-C₃ alkyl or C₁-C₃ haloalkyl, C₁-C₃ alkoxy;

R³ is H, OH, halo, CN, C₁-C₃ alkyl, C₃-C₆ cycloalkyl, C₁-C₃ haloalkyl, C₁-C₃ alkoxy, or C₁-C₃ haloalkoxy;

R⁴ and R⁵ taken together form a 5 or 6 membered ring that is fused to the phenyl ring to which R⁴ and R⁵ are attached, wherein the 5 or 6 membered ring contains one or two oxygen atoms as ring members and is optionally substituted by up to four groups independently selected from -CN, -OR, -SR, -SO₂R, and C₁-C₄ alkyl optionally substituted with one or two groups selected from halo, -OR, -SR, -SO₂R, C₁-C₃ haloalkoxy, CN, and -N(R)₂;

R⁶ is H, halo, C₁-C₃ alkoxy, or C₁-C₆ alkyl, or is taken together with R⁹ to form a ring;

R⁷ is H, halo, C₁-C₃ alkoxy, or C₁-C₆ alkyl, or is taken together with R⁹ to form a ring;

R⁸ is H or C₁-C₆ alkyl, or is taken together with R⁹ to form a ring;

R⁹ taken together with exactly one group selected from R⁶, R⁷ and R⁸ forms a 3-7 membered cycloalkyl ring or a 3-7 membered heterocyclic ring containing N, O or S as a ring member; wherein the cycloalkyl or heterocyclic ring is optionally substituted with up to three groups selected from R, -OR, -NR₂, halo, CN, COOR, CONR₂, and oxo;

W is -COOR*, -C(O)NH-SO₂R, -C(O)NH-SO₂NR₂, 5-tetrazolyl, or 1,2,4-oxadiazol-3-yl-5(4H)-one;

R* is H or C₁-C₆ alkyl that is optionally substituted with one or two groups selected from halo, -OR, oxo, CN, -NR₂, COOR, and CONR₂;

R is independently selected at each occurrence from H and C₁-C₄ alkyl optionally substituted with one to three groups selected from halo, -OH, C₁-C₃ alkoxy, oxo, CN, -NH₂, -NH(C₁-C₃ alkyl), -N(C₁-C₃ alkyl)₂, and cyclopropyl;

and two R groups directly attached to the same atom, which may be C or N, can optionally be taken together to form a 3-6 membered ring that can optionally contain an added heteroatom selected from N, O and S as a ring member, and can be substituted by up to two groups selected from -OH, oxo, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

or a pharmaceutically acceptable salt thereof.


2. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein R¹ is H or F.

3. The compound according to any one of the preceding claims, or a pharmaceutically acceptable salt thereof, wherein R² is H or halo.

4. A compound according to any one of claims 1 to 3, or a pharmaceutically acceptable salt thereof, wherein R³ is C₁-C₃ alkoxy or halo.

5. The compound of any of claims 1-4, wherein W is -COOR*;
or a pharmaceutically acceptable salt thereof.

6. The compound of any of claims 1-5, which is of the formula (II):



wherein each R¹⁰ is independently selected from H, R¹¹, -OR, -SR, -SO₂R, and -N(R)₂;


R¹¹ is C₁-C₄ alkyl, optionally substituted with one or two groups selected from CN, -OR, C₁-C₃ haloalkoxy, -NR₂, and -SO₂R;

n is 1 or 2;

Z is O or CR₂;

or a pharmaceutically acceptable salt thereof.

7. The compound of claim 6, which is of the formula (III), or a pharmaceutically acceptable salt thereof:



wherein R¹⁰ is H, R¹¹, -OR, -SR, -SO₂R, or -N(R)₂;

R¹¹ is C₁-C₄ alkyl optionally substituted with -OH or OMe;


R¹² and R¹³ are each selected from H and C₁-C₃ alkyl,

or R¹² and R¹³ taken together form a cyclopropyl or cyclobutyl ring;

and Z is O or CH₂.


8. A compound according to any of claims 6-7, or a pharmaceutically acceptable salt thereof, wherein R¹⁰ is -CH₂OMe, -(CH₂)₂-OMe, or -OR.

9. A compound according to any of claims 1-8, or a pharmaceutically acceptable salt thereof, which is of the formula:



wherein R¹⁰ is -CH₂OMe, -(CH₂)₂-OMe, or -OR.

10. A compound according to claim 9, which is of the formula:




wherein R^{10} is $-\text{CH}_2\text{OMe}$, $-(\text{CH}_2)_2\text{-OMe}$, or $-\text{OR}$;

or a pharmaceutically acceptable salt thereof.

11. A compound according to any of claims 6-10, or a pharmaceutically acceptable salt thereof, wherein one R^{10} group present represents $-\text{OR}'$, wherein R' is C_1 - C_3 alkyl substituted with up to two groups selected from OH and OMe.

12. A compound according to any of claims 6-10, or a pharmaceutically acceptable salt thereof, wherein one R^{10} is selected from $-\text{CH}_2\text{OMe}$, $-\text{CH}_2\text{CH}_2\text{OMe}$, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{OMe}$.

13. A compound according to any one of claims 1-9, which is of the formula:



















wherein R^{10} is $-\text{CH}_2\text{OMe}$, $-(\text{CH}_2)_2\text{-OMe}$, or $-\text{OR}$;



or a pharmaceutically acceptable salt thereof.

14. The compound of claim 13, wherein R^{10} is selected from $-\text{OMe}$, $-\text{CH}_2\text{OMe}$, $-\text{CH}_2\text{CH}_2\text{OMe}$, and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{OMe}$;

or a pharmaceutically acceptable salt thereof.


15. The compound according to claim 1, which is selected from:





or a pharmaceutically acceptable salt thereof.

16. The compound of claim 1, which is:



or a pharmaceutically acceptable salt thereof.

17. A pharmaceutical composition, comprising a compound of any of the preceding claims admixed with at least one pharmaceutically acceptable carrier.

18. A method to treat a subject having a hepatitis B infection, which comprises administering to the subject a compound of any of claims 1-16 or a pharmaceutical composition of claim 17.

19. The method of claim 18, wherein the compound of any one of claims 1-16 or the pharmaceutical composition of claim 17 is used in combination with an additional therapeutic agent selected from an interferon or peginterferon, an HBV polymerase inhibitor, a viral entry inhibitor, a viral maturation inhibitor, a capsid assembly inhibitor, an HBV core modulator, a reverse transcriptase inhibitor, a TLR-agonist, or an immunomodulator.

20. A method to inhibit replication of hepatitis B virus, which comprises contacting the hepatitis B virus, either in vitro or in vivo, with a compound according to any one of claims 1-16.

21. A pharmaceutical combination, comprising a compound of any of claims 1-16 and at least one additional therapeutic agent.

INTERNATIONAL SEARCH REPORT

International application No PCT/IB2017/053365

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D491/14 A61K31/4741 A61P31/12 ADD.				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED				
Minimum documentation searched (classification system followed by classification symbols) C07D				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
A	WO 2015/113990 A1 (HOFFMANN LA ROCHE [CH]; HOFFMANN LA ROCHE [US]) 6 August 2015 (2015-08-06) see structure of the compounds of claim 1 as HBV antivirals -----	1-21		
A,P	WO 2017/017042 A1 (F HOFFMANN-LA ROCHE AG [CH]; HOFFMANN-LA ROCHE INC [US]) 2 February 2017 (2017-02-02) see structure of the compounds of claim 1 as HBV antivirals -----	1-21		
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.				
* Special categories of cited documents : <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none; vertical-align: top;"> "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed </td> <td style="width: 50%; border: none; vertical-align: top;"> "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family </td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family			
Date of the actual completion of the international search 17 July 2017	Date of mailing of the international search report 28/07/2017			
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Traegler-Goeldel, M			

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/IB2017/053365

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2015113990	A1	06-08-2015	AU 2015212903 A1 19-05-2016 CA 2931329 A1 06-08-2015 CN 105899508 A 24-08-2016 CR 20160241 A 15-07-2016 EA 201691261 A1 30-11-2016 EP 3099685 A1 07-12-2016 JP 2017504633 A 09-02-2017 PE 06862016 A1 23-07-2016 PH 12016501254 A1 15-08-2016 SG 112016059700 A 30-08-2016 TW 201540715 A 01-11-2015 US 2015210682 A1 30-07-2015 US 2016296515 A1 13-10-2016 WO 2015113990 A1 06-08-2015

WO 2017017042	A1	02-02-2017	NONE
