

(19) **DANMARK**

(10)

DK 177529 B1



(12)

PATENTSKRIFT

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 9/127 (2006.01)** **A 61 K 31/00 (2006.01)** **A 61 K 33/24 (2006.01)**
- (21) Ansøgningsnummer: **PA 2009 01150**
- (22) Indleveringsdato: **2009-10-23**
- (24) Løbedag: **2009-10-23**
- (41) Alm. tilgængelig: **2011-04-24**
- (45) Patentets meddelelse bkg. den: **2013-09-08**
- (73) Patenthaver: **BIO-BEDST ApS, Dandyvej 19, 7100 Vejle, Danmark**
- (72) Opfinder: **Anders Falk Vikbjerg, Vesthegnet 52, 2670 Greve, Danmark**
Sune Allan Petersen, Hovgårdsparken 57, 2670 Greve, Danmark
Mogens Winkel Madsen, Bakkevej 53 A, 2830 Virum, Danmark
- (74) Fuldmægtig: **Chas. Hude A/S, H.C. Andersens Boulevard 33, 1780 København V, Danmark**
- (54) Benævnelse: **Liposomes with improved storage stability**
- (56) Fremdragne publikationer:
US A 5213804 A
EP A1 0565 36
WO A1 8601102
- (57) Sammendrag:
The present invention provides a composition comprising a sPLA2 hydrolysable liposome, an exterior solution, and an interior solution within the liposome, - wherein the concentration of osmolytes is higher in the interior solution than in the exterior solution. The composition improves storage stability of sPLA2 hydrolysable liposomes, in particular at when stored at 2-8 degrees Celsius. The liposome preferably encapsulates cispiatin. The invention also provides methods of preparing the composition of the invention.

Fortsættes ...




Figure 1

Liposomes with improved storage stability

Field of the invention

The present invention relates to sPLA2 hydrolysable liposomal drug delivery systems with improved storage stability, in particular improved storage stability at
5 2-8 degrees Celsius.

Background

Liposomes for drug delivery

Liposomes are microscopic spheres which were developed as drug delivery
10 vehicles/systems in the 1980s. The first liposome-based pharmaceuticals were approved for commercial use in the 1990s.

Liposomes have three distinct compartments that can be used to carry various compounds such as drugs: The interior aqueous compartment; the hydrophobic
15 bilayer; and the polar inter-phase of the inner and outer leaflet. Depending on the chemical nature of the compound to be encapsulated it will be localised to either of the compartments.

Currently, there are several parenteral liposome-drug formulations available on
20 the market. Water soluble drugs tend to be localised in the aqueous compartment of liposomes, and examples of drugs encapsulated in liposome's are, e.g. doxorubicin (Doxil), doxorubicin (Myocet) and daunorubicin (DaunoXone). Examples of drugs intercalated in the liposome membrane are, e.g. amphotericin B (AmBisome), amphotericin (Albelcet B), benzoporphyrin (Visudyne) and
25 muramyltripeptide-phosphatidylethanolamine (Junovan).

Liposomes are considered a promising drug delivery system since they passively target tumor tissue by using the pathophysiological characteristics of solid tumors such as hyperplasia and increased vascular permeability but also a defect in
30 lymphatic drainage. These features facilitate extravasation of nanoparticles and the liposomes can be retained in the tissue for longer time due to the enhanced permeability and retention effect (EPR).

The property of liposomes as drug delivery vehicles is crucially dependent on their surface charge, permeability, solubility, stability etc. which is significantly influenced by the lipids comprised in the liposome composition. In addition, the
5 drug to be encapsulated in the liposome may need further requirements to be considered in preparing a stable liposome formulation.

Considerations regarding safety and drug efficacy require that liposome formulations maintain their properties, i.e. remain stable, from the time of
10 preparation until administration.

Furthermore, it is desirable that such formulations are intact during the transport in the treated subject until they reach the target site where the drug is specifically released.
15

Various targeting strategies for liposomes have been described, e.g. conjugation to cell specific ligands such as antibodies.

sPLA2 hydrolysable liposomes

20 Another approach has been suggested based upon elevated levels of secretory phospholipase A2 (sPLA2) in cancerous tissue and also at sites of inflammation. The basic idea is that liposomes can be prepared which are hydrolysable by sPLA2 and that hydrolysis by sPLA2 leads to release of the drug encapsulated within the liposome. Moreover, the products of sPLA2 hydrolysis, a lysolipid and a fatty acid
25 act as permeabilizers of cell membranes leading to increased cell uptake of the drug. Since sPLA2 levels are elevated in cancerous tissues and at sites of inflammation, sPLA2 activated liposomes may be used to preferentially deliver encapsulated drugs to such sites.

30 Storage stability of sPLA2 hydrolysable liposomes

sPLA2 hydrolysable liposomes pose special challenges with regards to storage stability. These challenges are based in the particular lipid composition necessary for effective sPLA2 hydrolysis.

In general, many parameters influence storage stability, and it is difficult to predict the consequences of altering buffer composition on storage stability as this affects not only the osmolarity, but also the membrane stability.

5 Prior art

A number of documents have described sPLA2 activated liposomes, and various documents have studied storage stability of liposomes. However, none of the documents have studied the particular storage problems posed by sPLA2 hydrolysable liposomes.

10

WO0158910 described sPLA2 activated liposomes comprising prodrugs of mono-ether lyso-phospholipids. This document also described encapsulation of additional bioactive compounds.

15

WO0176555 suggested the use of a lipid-based drug delivery system for treatment of diseases or conditions associated with a localized increase in extracellular sPLA2 in cutaneous or subcutaneous tissue of a mammal, for administration of a prodrug of an ether-lysolipid that is activated by sPLA2. The system further comprised a so-called edge active compound.

20

WO0176556 suggested the use of a lipid-based drug delivery system for treatment or prevention of a parasitic infection selected from Leishmaniasis, Trypanosomiasis, malaria, Entamoeba, Histolyticasis and "Oriental liver fluke *Clonorchis sinensis*", wherein the system comprised prodrugs in the form of lipid derivatives that are activated by sPLA2. The liposomes may contain an additional bioactive compound.

25

WO06048017 and WO07107161 did also describe sPLA2 activated liposomes.

30

US5094854 disclosed temperature sensitive liposomes optimized with regards to stability and release by heating. The liposomes described in this document preferably has an interior solution with an osmotic pressure 1,2 to 2,5 times higher than that of body fluid of warm blooded animals and have a phase transition temperature of 40-45 degrees Celsius. The liposomes disclosed in this

document are not substrates of sPLA2, because they do not have the appropriate lipid composition.

US5213804 describes liposomes for delivery of compounds to solid tumours from
5 the bloodstream. Preparation of the liposome comprises generation of a transmembrane ammonium ion gradient. However, no osmolyte gradient is used or disclosed. US5213804 also does not disclose sPLA2 hydrolysable liposomes.

Most liposomal formulations do not have problems with leakage during storage.
10 Rather the formulations are so stable that it is difficult to get a release of encapsulated compounds at the intended sites.

Summary of the invention

In a first aspect, the present invention provides a composition comprising
15 a sPLA2 hydrolysable liposome characterized by a content of anionic lipid between 20% and 45%, a content of polymer conjugated lipid between 3% and 6% and a content of neutral lipids between 40% and 75%, wherein the sPLA2 hydrolysable liposome comprises

- 20
- cisplatin encapsulated within the liposome,
 - an exterior solution,
 - and an interior solution within the liposome,
 - wherein the difference in osmolyte concentration between the interior and the exterior solution (exterior concentration of osmolyte subtracted
25 from the interior concentration of osmolyte) is between 280 and 320 mM.

The composition improves storage stability of sPLA2 hydrolysable liposomes, in particular when stored at 2-8 degrees Celsius. The liposome preferably
30 encapsulates cisplatin.

The invention also provides methods of preparing the composition of the invention.

Brief description of the figures

Figure 1.

Contour plot illustrating the influence of sucrose concentration and homogenization pressure used during liposome preparation on particle size.

5 Numbers inside the contour plot indicate particle size

Figure 2.

Total Pt concentrations in liposomes as a function of particle size.

10 Figure 3.

Change in particle size during storage at 2-8°C for LPB0057-LPB0073

Figure 4.

Change in PDI during storage at 2-8°C for LPB0057-LPB0073

15

Figure 5.

Contour plot illustrating the influence of interior NaCl concentration and homogenization pressure used during liposome preparation on initial DOE%. Numbers inside the contour plot indicate DOE%.

20

Figure 6.

Pt concentration during storage for LPB0057-LPB0073.

Figure 7.

25 Pt exterior (%) during storage for LPB0057-LPB0073.

Figure 8.

Leakage from LPB0057-LPB0073 during storage at 2-8°C.

30 Figure 9.

Contour plot illustrating the influence of interior NaCl concentration and sucrose concentration on DOE% after 56d storage at 2-8°C with homogenization pressure maintained at 12,500 KPa. Numbers inside the contour plot indicate DOE% after 56d storage at 2-8°C.

Figure 10.

Contour plot illustrating the influence of interior NaCl concentration and sucrose concentration on %Leakage after 56d storage at 2-8°C with homogenization pressure maintained at 12,500 KPa. Numbers inside the contour plot indicate %Leakage after 56d storage at 2-8°C.

Figure 11.

Leakage from liposomal cisplatin formulation during storage at 2-8°C. Formulations were prepared with varying osmotic gradient. For further details refer to table 3.

Figure 12.

Particle size of liposomal cisplatin formulations during storage at 2-8°C. For detailed description about the buffer composition refer to table 3.

Figure 13.

Table 1 showing actual experimental settings of factors and the responses.

Figure 14.

Table 2 showing regression coefficients and significance (P) values of the model after back ward elimination.

Figure 15.

Table 3 showing overview of liposomal cisplatin formulations prepared with varying osmotic gradient.

Figure 16.

Table 4 showing overview of different liposomal drug formulations prepared with varying molecular weight.

Disclosure of the invention

The present invention is based on the discovery that leakage from sPLA2 hydrolysable liposomes may be reduced by storage in an exterior solution that has a lower concentration of osmolytes than the interior solution of the liposome.

5

The leakage problem was observed in relation to storage of cisplatin encapsulated in sPLA2 hydrolysable liposomes to be used for a phase 1 study of an iso-osmotic formulation of cisplatin encapsulated in sPLA2 hydrolysable liposomes. In this formulation, the interior solution was 0,9% NaCl and the exterior solution was 10 mM phosphate buffer at pH 6,5, 1 mM NaCl and 10% sucrose. Because of a high degree of leakage when stored 2-8 degrees Celsius, the iso-osmotic formulation had to be stored at minus 80 degrees Celsius, which is not an optimal solution for a commercial product. Thus, not all hospitals have facilities for storage at minus 80 degrees. Moreover, transportation at minus 80 degrees Celsius is cumbersome.

15

Additionally, when thawing the vials before use, problems with breakage of the used vials were encountered and thawing caused significant leakage from the liposomes.

20 As mentioned, it is believed that the leakage problem is a consequence of the specific requirements with regards to lipid composition of the sPLA2 hydrolysable liposomes. In particular it is believed that the leakage is caused by the absence or low amounts of stabilizing agents such as cholesterol in sPLA2 hydrolysable liposomes. Moreover, the anionic nature of sPLA2 hydrolysable liposomes may
25 influence leakage.

Thus, there is a need for a composition comprising sPLA2 hydrolysable liposomes with reduced leakage when stored at 2-8 degrees Celsius.

30 The terms "reduced leakage" and "improved stability" with respect to sPLA2 liposomes are used interchangeably herein.

In a first aspect, the present invention provides a composition comprising

a sPLA2 hydrolysable liposome characterized by a content of anionic lipid between 20% and 45%, a content of polymer conjugated lipid between 3% and 6% and a content of neutral lipids between 40% and 75%, wherein the sPLA2 hydrolysable liposome comprises

5

- cisplatin encapsulated within the liposome,
 - an exterior solution,
 - and an interior solution within the liposome,
 - wherein the difference in osmolyte concentration between the interior and the exterior solution (exterior concentration of osmolyte subtracted from the interior concentration of osmolyte) is between 280 and 320 mM.
- 10

The composition will normally be a pharmaceutical composition.

15

In a preferred embodiment, the only therapeutic agent in sPLA2 hydrolysable liposome is cisplatin. Thus, the interior solution only comprises osmolytes and cisplatin and optionally a buffer.

20

Chloride ions prevent cisplatin from forming an undesired highly toxic hydration product. Therefore, it is preferred that the interior solution comprises NaCl or KCl at a concentration of at least 0,2 % or 0,4%, more preferably 0,2-2,5 % w/w, or 0,4-1,8% and most preferably between 0,8-1%. In some embodiments, other chloride salts may be used. NaCl normally is preferred over KCl.

25

In some embodiments, it is preferred that the exterior solution comprises NaCl or KCl at a concentration of at least 0,2 % or 0,4%, more preferably 0,2-2,5 % w/w, or 0,4-1,8% and most preferably between 0,8-1%. The typical concentration is 0,9%. . Thus, if minor amounts of cisplatin leaks out of the liposome, the presence of chloride ions will minimize the formation of toxic hydration products. Again, NaCl is normally preferred over KCl.

30

In one embodiment, there are no divalent cations, such as Calcium ions, in the exterior solution.

In another embodiment, divalent cations may be included in the exterior solution.

- 5 The interior solution does typically not include a buffer. Moreover, sulfates are not normally included in the interior solution.

In one embodiment of the composition, the interior solution comprises at least 1,4% NaCl or KCl and more preferably 1,8% NaCl or KCl and does not comprise a
10 viscosity enhancer, preferably selected from the group consisting of sugars such as lactose, sucrose, maltose, galactose, glucose and hydrophilic polymers such as starch, dextran, polyvinyl alcohol, polyvinyl-pyrrolidone, dextrin, xanthan and partly hydrolyzed cellulose oligomers and proteins and polypeptides.

- 15 In another embodiment of the composition, the interior solution comprises at least 5 % of a viscosity enhancer. Preferably, the viscosity enhancer is selected from the group consisting of sugars such as lactose, sucrose, maltose, galactose, glucose and hydrophilic polymers like starch, dextran, polyvinyl alcohol, polyvinyl-pyrrolidone, dextrin, xanthan and partly hydrolyzed cellulose oligomers and
20 proteins and polypeptides. It should be recognized that viscosity enhancers will also function as osmolytes.

When the interior solution comprises a viscosity enhancer, it is preferred that the concentration is more than 5% w/w. Even more preferred is a concentration of
25 more than 6, 7, 8, 9 or 10 %%. In a preferred embodiment, the viscosity enhancer selected from the group consisting of sugars such as lactose, sucrose, maltose, galactose, glucose and hydrophilic polymers like starch, dextran, polyvinyl alcohol, polyvinyl-pyrrolidone, dextrin, xanthan and partly hydrolyzed cellulose oligomers and proteins and polypeptides. Most preferred are sugars, in
30 particular sucrose.

In general, it is preferred that the exterior solution has an osmolyte concentration around 300 mM since this the physiological osmolytes concentration. Thus, in embodiment, it is preferred that the exterior solution has an osmolytes

concentration between 200 and 400 mM, more preferably 250-350 mM and most preferred between 275 and 325 mM.

The composition of the invention is characterized in that there is a difference in
5 osmolyte concentration between the interior and the exterior solution. The difference is herein also termed the osmotic gradient.

As demonstrated in the examples section, an osmotic gradient (exterior
concentration of osmolyte subtracted from the interior concentration of osmolyte)
10 of 27 mM cannot reduce leakage sufficiently. A gradient of 59 mM reduces leakage further as do also a gradient of 91 mM. It should be noted that the osmolyte concentration is the total concentration of all solute particles, which is often referred to as the osmolarity (Osm) of a solution.

15 In one embodiment, the osmolarity of the solutions are corrected using the osmotic coefficients of the solute particles.

In one embodiment, the concentration of osmolytes in the interior solution is
20 assumed to be the same as the concentration of the osmolytes in the hydration solvent used when preparing the liposomes. I.e. when reference is made to the osmolytes concentration in the interior solution, this concentration can easily be determined since it is the concentration of osmolytes in the hydration solvent. The actual concentration of osmolytes in the interior solution may not be identical
25 to the concentration of osmolytes in the hydration solvent, because water may move into the liposome after formation and hence decrease the concentration of osmolytes (while increasing the osmotic pressure within the liposome)

If the osmotic gradient too high, it is believed that the liposomes will initially be
30 leaky, such that osmolytes will pass the membrane until a maximal tolerable gradient is established, where after leakage will be at a minimum.

Thus, the osmotic gradient is between 280 and 320 mM.

The pH of the exterior solution typically has a pH between 5 and 7 and the interior solution typically has a pH between 5 and 7. Appropriate buffers may be used to keep the pH at the desired value.

- 5 However, in one embodiment, the interior solution and/or the exterior solution do not comprise a buffer, preferably phosphate buffer. In some embodiments, phosphate buffers are undesired, because they promote conversion of cisplatin to byproducts.
- 10 Particular preferred compositions where cisplatin is encapsulated in the liposome comprise the following:
- 1) Interior solution: 0,8-1,0% NaCl, such as 0.9% NaCl and 9-11 % sucrose, such as 10% sucrose.
 - 15 Exterior solution: 8-12 mM Phosphate buffer (pH6.5), such as 10 mM Phosphate buffer (pH6.5) + 9-11 sucrose %, such as 10% sucrose.
 - 2) Interior solution: 1,6-2,0 % NaCl, such as 1.8% NaCl
Exterior solution: 8-12 mM Phosphate buffer (pH6.5), such as 10 mM
20 Phosphate buffer (pH6.5) + 9-11 % sucrose, such as 10% sucrose.
 - 3) Interior solution: 0,8-1,0% NaCl, such as 0.9% NaCl and 9-11 % sucrose, such as 10% sucrose.
Exterior solution: 8-12 mM Phosphate buffer (pH6.5), such as 10 mM
25 Phosphate buffer (pH6.5) + 0,35% - 0,55% such as 0.45% NaCl+ 4-6 % sucrose, such as 5% sucrose.
 - 4) Interior solution: 1,6-2,0 % NaCl, such as 1.8% NaCl
Exterior solution: 8-12 mM Phosphate buffer (pH6.5), such as 10 mM
30 Phosphate buffer (pH 6.5) + 0,8-1,0% NaCl, such as 0.9 % NaCl.

The concentration of the therapeutic agent in the final composition is typically between 0,1 mg/ml to 15 mg/ml. When the therapeutic agent is cisplatin, it is preferred that the concentration is between 0,5 mg/ml and 1,5 mg/ml. Such
35 concentrations can be achieved by using a cisplatin concentration of 8 mg/ml in

the hydration solution during liposome preparation. A concentration of 8 mg/ml can be achieved by heating the hydration solution to a temperature of 65% degrees Celsius.

- 5 The liposomes of the composition preferably have an average size (diameter) between 50 and 200 nm, more preferably between 75 and 160 nm.

The liposomes of the composition should have a leakage of less than 20 %, 15 % or 10 %, more preferably less than 9 %, 8% or 7 % leakage after 56 days of
10 storage at a temperature between 2 and 8 degrees Celsius. As demonstrated in the examples section, this can be achieved by adjusted the osmotic gradient.

When referring to the percentage of leakage, reference is to the amount of cisplatin in the liposomes after a step of dialysis or ultrafiltration compared to the
15 amount of cisplatin in the liposomes after storage. After a step of dialysis or ultrafiltration, the exterior solution may comprise 5% of the total amount of cisplatin in the composition. Thus, the internal solution comprises 95% of the cisplatin in the composition. After storage, the internal solution may comprise 85,5% of the total amount of cisplatin and the exterior solution consequently
20 14,5%. The leakage has then been $9,5 / 95 = 10\%$.

In a preferred embodiment, the composition of the first aspect is prepared by the method of the second aspect.

25 **sPLA2 hydrolysable liposomes**

sPLA2 hydrolysable liposomes for use in the composition of the present invention are defined in more detail in the following embodiments. In its broadest
embodiment, the term sPLA2 hydrolysable liposomes refer to liposomes that are hydrolysable under physiological conditions, particular in cancerous tissue.

30

Preferably, the sPLA2 hydrolysable liposomes comprises between 20% and 45% (mol/mol) of an anionic lipid. The content of anionic lipid affects important characteristics of the liposome, such as the rate of sPLA2 mediated lipid hydrolysis of the liposome and also the immune response toward the liposome.

As the content of anionic lipid increases, so does the rate of lipid hydrolysis by sPLA₂ (and the release of drug). It has been demonstrated that a reasonable rate of hydrolysis can be achieved by anionic lipid content between 20% and 45%.

5 Thus, in one embodiment, the content of anionic lipid is at least 20%. In another embodiment, the content of anionic lipid is no more than 45%. In yet another embodiment, the anionic lipid content of the liposome is selected from the group consisting of between 20% and 45%, between 25% and 45%, between 28% and 42%, between 30% and 40%, between 32% and 38% and between 34% and
10 36%.

As mentioned, also the immune response toward the liposomes is affected by the content of anionic lipid. Thus, the clearance rate of the liposome in body may be reduced by keeping the content of the anionic lipid in the liposome below a certain
15 level and the present inventors have recognized that the content of anionic lipid in the liposome can be used to strike a balance between hydrolysis rate of sPLA₂ and clearance by the reticuloendothelial system.

Preferably the anionic lipid is a phospholipid and preferably, the phospholipid is
20 selected from the group consisting of PI (phosphatidyl inositol), PS (phosphatidyl serine), DPG (bisphosphatidyl glycerol), PA (phosphatidic acid), PEOH (phosphatidyl alcohol), and PG (phosphatidyl glycerol). More preferably, the anionic phospholipid is PG. Preferably, the lipids comprise stearyl chains. Thus preferably PG is DSPG.

25

Hydrophilic polymers

In a preferred embodiment, the sPLA₂ hydrolysable liposome for use in the present invention further comprises a hydrophilic polymer selected from the group consisting of PEG [poly(ethylene glycol)], PACM [poly(N-acryloylmorpholine)], PVP
30 [poly(vinylpyrrolidone)], PLA [poly(lactide)], PG [poly(glycolide)], POZO [poly(2-methyl-2-oxazoline)], PVA [poly(vinyl alcohol)], HPMC (hydroxypropylmethylcellulose), PEO [poly(ethylene oxide)], chitosan [poly(D-glucosamine)], PAA [poly(aminoacid)], polyHEMA [Poly(2-hydroxyethylmethacrylate)] and co-polymers thereof.

35

Most preferably the polymer is PEG with a molecular weight between 100 Da and 10 kDa. Particular preferred are PEG sizes of 2-5 kDa (PEG2000 to PEG5000), and most preferred is PEG2000.

- 5 The inclusion of polymers on liposomes is well known to the skilled artisan and can be used to increase the half-life of the liposomes in the bloodstream, presumably by reducing clearance by the reticuloendothelial system.

Preferably, the polymer is conjugated to the head group of phosphatidyl
10 ethanolamine. Another option is conjugation to ceramide (even though this lipid is not hydrolyzable by sPLA₂). When the polymer is conjugated to phosphatidyl ethanolamine, a negative charge is introduced and hence DSPE-PEG is regarded as an anionic lipid (contrary to DSPE which is regarded as a neutral lipid).

The polymer-conjugated lipid is preferably present at an amount of at least 2%.
15 More preferably, the amount is at least 5% and no more than 15%. Even more preferably, the amount of polymer-conjugated lipid is at least 3% and no more than 6%. Liposomes containing anionic phospholipids and ≤ 2.5 % DSPE-PEG2000 have increased tendency to aggregate in the presence of calcium. This can usually be observed by formation of high viscous gel. Liposomes containing anionic
20 phospholipids and >7.5 % DSPE-PEG2000 causes the liposomes to sediment or phase separate.

Neutrally charged lipid components in the liposome

Preferably, the liposome to be used in the present invention also comprises an
25 uncharged phospholipid selected from the group consisting of zwitterionic phospholipids comprising PC (phosphatidyl choline) and PE (phosphatidylethanolamine). Most preferably, the zwitterionic phospholipid is PC.

In contrast to anionic phospholipid, zwitterionic phospholipid serves as a charge
30 neutral sPLA₂-hydrolyzable lipid component in the liposome. By combining zwitterionic- and anionic phospholipid in the same liposome, it is possible to adjust to a desired surface charge density which complies with both sufficiently high sPLA₂ hydrolysis and a low clearance rate in the blood.

The amount of zwitterionic phospholipid in the liposome is preferably between 40% and 75% and more preferably between 50 and 70%.

Preferably, the lipids (anionic lipids, neutral lipids and polymer conjugated lipids) 5 comprise stearyl chains). Thus preferably PG is DSPG, PE is preferably DSPE.

Ether-phospholipids

Some or all of the phospholipids may be ether-phospholipids.

10 Thus, they may harbour an ether-bond instead of an ester-bond at the *sn*-1 position of the glycerol backbone of the phospholipid. When sPLA₂ hydrolyze this particular type of phospholipids, mono-ether lyso-phospholipids are produced and these are toxic to e.g. cancer cells. I.e. ether phospholipids may be seen as pro-
15 drugs of mono-ether lyso-phospholipids and liposomes of the invention can be used to deliver such pro-drugs to the sPLA₂-enhanced environment of cancer cells, where the pro-drugs are activated by sPLA₂ hydrolysis. Ether-phospholipids have been described in EP 1254143 and WO 2006/048017.

In one embodiment, the sPLA₂ activated liposomes as used in the present 20 invention does not comprise ether-phospholipids.

Other pro-drugs

The moiety released from the lipid by sPLA₂ to create a lysolipid may also be a drug. Thus, a liposome may comprise pro-drugs of mono-ether lysolipids, pro-
25 drugs released from the lipid by sPLA₂ and other therapeutic agents, as further outlined below.

In one embodiment, the sPLA₂ activated liposomes as used in the present invention does not comprise prodrugs released from the lipid by sPLA₂.

30

Stabilizing agent

The liposome may also be stabilized by the inclusion of cholesterol as membrane component in the liposome. However, high amounts of cholesterol in the liposome have a negative effect on hydrolysis by PLA₂ and therefore it is preferred that the
35 liposome comprises no more than 10% cholesterol. Even more preferably, the

liposome comprises less than 1% cholesterol, less than 0,1 % or does not comprise any cholesterol at all.

The alkyl chain length of the lipids comprising the liposome may be adjusted for
5 optimal PLA₂ hydrolysis rate and minimum leakage of entrapped compound out of the liposome. Preferably, the alkyl chains are C18 or C16 saturated chains.

As described above, the liposomes may comprise pro-drugs of mono-ether lyso-
lipids and/or of the moiety released from the lipid by sPLA₂ to create the lysolipid.
10

Physical-chemical characteristics of the liposomes

The liposome can be unilamellar or multilamellar. Most preferably, the liposome is unilamellar. The diameter of the liposome should be between 50 and 400 nm, preferably between 80 and 160 nm and most preferable between 90 and 120nm.
15

Preferably, the Poly Dispersity Index (PDI) of the liposomal formulation of the second aspect of the invention should not exceed 0.2 and more preferable is 0.10 or less. A PDI value in this range expresses a relatively narrow particle size-distribution in the formulation.

20 As will be clear from the above, it is preferred that at least one of the lipids comprising the liposome is a substrate for sPLA₂ when present in the liposome.

In one embodiment, the liposome comprises lipids which are hydrolysed by sPLA₂
25 at the sn-3 position instead of at the sn-2 position. Such unnatural lipids and liposomes comprising unnatural lipids have been disclosed in WO 2006/048017.

In a most preferred embodiment, the liposomes to be used in the present invention comprise 70% DSPC, 25% DSPG and 5% DSPE-PEG.

30

Method of preparation

Provided is also a method comprising the steps

- a) Preparing a lipid mixture by dissolving selected lipids in an organic solvent

- b) Hydrating the product of step a) with an aqueous hydration solvent so as to form liposomes
- c) Removing the organic solvent of step a) either before addition of the aqueous hydration solvent or after the addition of the aqueous hydration solvent
- d) exchanging the hydration solvent with an exterior solution, which has a lower osmolyte concentration than the hydration solution
- e) Thereby forming a composition as described in the first aspect of the invention.

5

10

The hydration solvent may be exchanged by centrifugation, ultrafiltration, dialysis or similar. After changing the hydration solvent to the exterior solution, it is preferred that less than 15%, less 10 % or more preferably less than 8 or 6% of therapeutic agent is present in the exterior solution.

15

Preferably, the drug encapsulation in the liposomes should be >70%, more preferably >95% and most preferably >99%. The degree of drug encapsulation is the ratio of drug encapsulated to the total amount of drug in the formulation.

20 Preferably, the organic solvent is removed before addition of hydration solvent.

The method may further comprise high sheer mixing to reduce the size of the liposomes.

25 The method may further comprise a step of extruding the liposomes produced in step c) through a filter to produce liposomes of a certain mean size.

The method may also comprise a step of sonicating the liposomal formulation to produce liposomes of a certain size.

30

The method may also comprise homogenization at a pressure between 5000 and 20000 KPa.

35 Preferably, the liposome is a liposome as described in the first aspect of the invention, i.e. lipids are selected accordingly.

Liposomes may be loaded with at least one therapeutic agent by solubilizing the drug in the organic solvent or hydration solvent used to prepare the liposomes. Preferably, the therapeutic agent is solubilised in the hydration solvent and the
5 therapeutic agent is preferably cisplatin.

Alternatively, ionisable therapeutic agent can be loaded into liposomes by first forming the liposomes, establishing an electrochemical potential, e.g. by way of a pH gradient, across the outermost liposome layer, and then adding the ionisable
10 therapeutic agent to the aqueous medium external to the liposome.

Examples

Example 1, Optimization of storage stability of sPLA2 hydrolysable 15 liposomes

Abstract

It has been observed that upon storage at 2-8°C LiPlaCis (liposomal cisplatin) may have up to 30% leakage during the first couple of months. Leakage is most extensive during the first two months, and beyond this period only minor leakage
20 occurs.

A factorial design was set up to test stability of LiPlacis at 2-8°C. Formulations were prepared with varying interior buffer composition, and homogenized at different pressures according to composite face design. 3 factors were tested at 3
25 levels:

- Interior NaCl concentration (0.9, 1.4, and 1.9%)
- Interior Sucrose concentration (0, 5, and 10%)
- Homogenization pressure used during preparation (5000, 12500, and
30 20000 KPa).

Leakage from formulation was demonstrated to be influenced by all parameters tested. Increasing the sucrose and NaCl concentration on liposome interior and having low homogenization pressure during liposome preparation results in higher

drug retention during storage. The formulations having the highest degree of encapsulation after two months of storage was prepared with 1.9% NaCl and 10% sucrose on the liposome interior, and homogenized at 5000 KPa. No significant changes were seen on particle size during storage for any of the formulations
5 prepared.

Materials and methods

Preparation of sPLA2 liposomes (LiPlaCis)

1,2-di(octadecanoyl)-sn-glycero-3-phosphocholine (DSPC), 1,2-di(octadecanoyl)-
10 sn-glycero-3-phosphoglycerol (DSPG), and 1,2-
di(octadecanoyl)phosphatidylethanolamine-methoxy poly (ethylene glycol) 2000 (DSPE-PEG 2000) were all purchased from Lipoid (Ludvigshafen, Germany). Sucrose was purchased from Sigma-Aldrich (St. Louis, MO). Sodium chloride and sodium dihydrogen phosphate were purchased from Merck (Darmstadt, Germany)
15 and J.T.Baker (Deventer, Holland), respectively. Iridium atomic absorption standard solution was purchased from Sigma-Aldrich (St. Louis, Mo). Slide-A-Lyzer 10K MWCO dialysis casttes were purchased from Thermo Scientific (Rockford, IL). Amicon® Ultra-4 Ultracel-30k centrifuge filter devices were purchased from Millipore (Bedford, MA).

20

Preparation of LiPlaCis

Homogeneous lipid mixture (DSPC/DSPG/DSPE-PEG2000 70:25:5 mol-%) was prepared by dissolving lipids in chloroform/methanol 9:1 (v/v). The solvent was evaporated from the lipid mixtures at a temperature of 50oC, using a gentle
25 stream of nitrogen gas. Residual solvent was evaporated by storing under vacuum overnight in a Christ Epsilon 2-4 freeze-dryer from Martin Christ Freeze Dryers GmbH (Harz, Germany). Multilamellar vesicles (MLVs) were prepared, by dispersing the lipid films in a 65 oC hot hydration media containing 8 mg/mL cisplatin and varying concentrations of sucrose and sodium chloride according to
30 table 1. The lipids were hydrated for ½ h at 65 oC. During hydration samples were vortexed each 5 min. Hydrated lipid suspension was homogenized in EmulsiFlex-C3 from Avestin, Inc. (Ontartio, Canada) at varying pressure (5000-20,000 KPa) according to table 1. All steps of homogenization were carried out at a temperature of 65°C. Untrapped cisplatin was removed from the formulation by

precipitation and dialysis. Precipitation was carried out in two steps; first at 25°C for 1 h, and then at 5 °C for 1 h. Dialyzed against 100x volume 2 times in 10mM Phosphate buffer containing 10% sucrose pH 6.5 for 18 and 24h, respectively. Immediately after preparation the formulations were divided into glass vials with
5 500 µl sample in each vial. Vials were sealed with cap and placed in a refrigerator (2-8°C). The day of placement in refrigerator is considered as the initiation date. Samples are taken continually during storage, and analyzed by ICP-MS and Zetasizer nano for determination of Pt concentrations and particle size, respectively.

10

Determination of particle sizes

Two drops of LiPlaCis was mixed into 10 mL Milli-Q water. 1 mL mixture was transferred into a disposable cuvette and particle size was measured on Zetasizer Nano ZS Dynamic Light Scattering device from Malvern Inc. (Worcestershire,
15 United Kingdom). Particle sizes were determined three times at a temperature of 25 oC using water as the internal reference substance. A mean Z-average particle size (i.e. diameter) was calculated based on the three measurements.

Determination of degree of encapsulation

20 The liposomes were diluted by a factor 100 in dialysis solution. Subsequently, the diluted liposomes were equilibrated for 1h at 25 oC. Part of the diluted liposomes was then centrifuged at 2500g for 30 min. at 15 oC in an Amicon® Ultra centrifugal filter devices (30K MWCO) from Millipore (Billerica, MA). Samples before centrifugation (i.e. 100 times diluted liposomes) and after centrifugation
25 (permeate) were diluted 100 fold in Milli-Q water. 1 vol.% iridium atomic absorption standard solution (1.05 ppm) was added to all samples as internal standard. A Perkin-Elmer PESCIEX Elan 6000 inductively coupled plasma device (Ontario, Canada) equipped with a cross flow nebuliser and pt conus was used for measuring platinum levels in samples of LiPlaCis.

30

Experimental design and statistical analysis

Experiment was conducted using a central composite face design to investigate the linear, quadratic, and cross product effects of three factors, each varied at three levels and also includes three center points for replication. Factors and their
35 levels are shown in table 1. The three factors chosen were interior NaCl

concentration (w/v%), interior sucrose concentration (w/v%), and homogenization pressure (KPa).

The design of the experiment employed is presented in table 1. A software package (Modde 8.0, Umetri, Umeå, Sweden) was used to fit the second order model to the independent variables.

The response surface model was fitted to the following equation:

$$Y = \beta_0 + \sum_{i=1}^n \beta_i X_i + \sum_{i=1}^n \beta_{ii} X_i^2 + \sum_{i=1}^{n-1} \sum_{j=i+1}^n \beta_{ij} X_i X_j + \varepsilon$$

10

Where Y is the response variables, X_i the *i*th independent variable, β_0 the intercept, β_i the first-order model coefficient, β_{ii} the quadric coefficient for variable *i* and β_{ij} is the model coefficient for the interaction between factor *i* and *j*, ε is the combination of the experimental error for the factors. The quadratic term make it possible to obtain information about curvatures in the response.

The coefficient of determination (R^2) and lack-of-fit test were used to determine whether the constructed model was adequate to describe the observed data. $R^2 > 0.8$ indicates that the model has acceptable qualities. Where it is possible, the model is simplified by dropping terms, which were not significant ($P > 0.05$) by analysis of variance. Terms were however not removed from the model if R^2 become below 0.8.

RSM was used to evaluate the effect of selected factors on initial particle size, DOE% immediately after preparation, DOE% and leakage after 56d storage at 2-8°C,

Results and discussion

Model fitting

30 The best-fitting models by multiple regression and backward elimination were determined. The observed and predicted values were sufficiently correlated. The statistics for model coefficients and probability (P) values for the response variables were calculated (Table 2). The models generated were generally satisfactory for the evaluations, as the observed and the predicted results were

well correlated. According to analysis of variance there was no lack of fit for any of the generated model.

Particle size

5 Data shows that the homogenization pressure has the most significant influence on the particle size. With a higher homogenization pressure used during preparation the particle size decrease. Sucrose concentration used during preparation was also demonstrated to have significant influence on the particle size. Increasing sucrose concentration results in a larger particle size. Contour plot
10 illustrating the influence of interior sucrose concentration and homogenization pressure used during liposome preparation on particle is presented in figure 1.

From figure 2 it can be seen that with larger particle size there is usually seen a higher amount of encapsulated cisplatin. With larger particle size it would also be
15 expected that the interior volume would increase, allowing more Pt to be encapsulated.

Changes in particle size and PDI during storage are presented in figure 3-4. No significant changes in particle size were observed during cold storage for any of
20 the liposomal formulations prepared.

Initial degree of encapsulation (%)

Formulations had an initial degree of encapsulation ranging from 84 to 92%. It was demonstrated that NaCl had the most significant influence on the initial
25 degree of encapsulation (DOE). With a higher salt concentration used in the hydration solution a higher initial DOE was obtained. Furthermore higher homogenization pressure used during preparation resulted in lower DOE. This is probably due to the fact that smaller liposomes are obtained at higher homogenization pressure, and thus more platinum can be expected on the
30 liposome exterior. Contour illustrating the influence of interior NaCl concentration and homogenization pressure used during liposome preparation on initial DOE% is presented in figure 5.

Influence of parameters on DOE% and leakage after 56 d storage at 2-8°C Pt analysis (total Pt content, liposome exterior Pt content and leakage) during
35 storage are presented in figure 6-8.

It was demonstrated that interior NaCl and sucrose concentrations have significant influence on degree of encapsulation (DOE) after 56 d storage. With a higher NaCl and sucrose concentration on the liposome interior a higher DOE was
5 obtained. No main effect was observed for homogenization pressure; however interaction between homogenization pressure and sucrose was seen. At high interior sucrose concentration and a low homogenization pressure results in higher DOE compared, as compared to having higher homogenization pressure.

10 The amount of cisplatin leaking from the formulation during 56d storage depends on all factors examined. However the factor having the most significant effect on leakage was sucrose concentration. Increasing interior NaCl and sucrose concentrations were observed to increase drug retention during storage. More leakage could be observed from liposomes prepared at low pressure compared to
15 those prepared at higher pressure.

Lowest leakage was observed for formulations prepared with hydration solution containing 10% sucrose and 1.9% NaCl, and homogenized at 20000 KPa. The formulation having the highest DOE after 56d storage at 2-8°C was prepared with
20 hydration solution containing 10% sucrose and 1.9% NaCl, and homogenized at 5000 KPa.

Conclusion

The response models satisfactorily expressed the relationship between the
25 selected parameters and the responses. The results demonstrate that interior NaCl and sucrose concentrations have significant influence on the leakage from LiPlacis formulation during storage at 2-8°C. During storage the factor having the most significant influence on drug retention was sucrose. Increasing NaCl also improves drug retention. Highest encapsulation degree after 56d was seen for
30 formulations having high NaCl and sucrose concentration on the liposome interior. DOE was demonstrated to be improved by increasing NaCl concentration in the hydration media.

Leakage was most pronounced when liposomes were prepared at low pressure, as compared to liposomes prepared at high pressure. Initial DOE is also higher for liposomes prepared at lower pressures. Since the homogenization pressure has a major influence on the particle size the results indicate that drug retention is to
5 some extent influenced by particle size. i

The formulations having the highest degree of encapsulation after two months of storage was prepared with 1.9% NaCl and 10% sucrose on the liposome interior, and homogenized at low pressure.

10

No significant changes in particle size were seen during storage for any of the formulations prepared.

15 **Example 2**

Liposomal cisplatin formulations were prepared by film hydration followed by extrusion. Hydration solution was prepared with varying osmolyte concentrations. After extrusion the liposomes were dialyzed in different media. The formulations prepared had different osmotic gradient (difference in osmolarity between interior
20 and exterior) as outlined in table 3. For comparison liposomal formulation of other chemotherapeutics were prepared as outlined in table 4. All Formulations were placed in refrigerator at 2-8°C, and samples were taken continually during storage.

25 Preparation of Liposomal formulations

Phospholipids (DSPC/DSPG/DSPE-PEG2000, 70:25:5 mol%) were dissolved in 9:1 (v/v) chloroform/methanol. The solvent of the dissolved lipid mixtures were then evaporated in a 50°C hot water bath until visual dryness, under a stream of nitrogen gas. The samples were further dried under vacuum overnight.

30

Hydration liquids (varying Salt and sucrose concentrations according to table 3 and 4) containing a chemotherapeutic were added to the dried lipid mixtures at a temperature of 65°C for the preparation of multilamellar vesicles (MLV). The lipid suspensions were kept at 65°C for at least 30 min. in order to ensure complete

hydration. During this period, the lipid suspensions were vortex every 5 min. Large unilamellar vesicles (LUV) were prepared by extrusion through membranes with defined pore size (100 nm) at 65°C. LUV were subsequently transferred to dialysis cassettes (MWCO: 10 kDa) in order to remove un-entrapped drug.

- 5 Liposomal formulations were dialyzed in different buffer solutions as outlined in table 3 and 4.

Immediately after preparation the formulations were divided into glass vials. Glass vials were sealed with cap and placed in refrigerator (2-8°C). The day of
10 placement in refrigerator is considered as the initiation date. Samples were taken continually during storage, and particle size and drug concentration (Exterior and total) was determined.

Results:

- 15 Liposomal formulations were prepared with either cisplatin, oxaliplatin, MTX, Bleomycin, or 5FU. Formulations containing small molecular weight drug (<350 g/mol) such as cisplatin and 5FU was observed to be leaky when osmotic gradient between interior and exterior was low. Liposomal formulations containing drugs with higher molecular weight (>350 g/mol) were demonstrated to be practically
20 non-leaky during storage at 2-8°C.

Trials to increase the osmotic gradient in cisplatin formulations were demonstrated to be an effective way to minimize the leakage during storage. When the difference between the interior and exterior osmotic gradient was >282
25 mOsM, the leakage observed during storage was maintained at acceptable levels (see table 3 and figure 11).

No changes in particle size were observed in particle size during storage at 2-8°C (Figure 12). The particle size of the formulation is thus not influenced by higher
30 osmotic gradient in the cisplatin formulation.

Krav

- 1) En farmaceutisk komposition omfattende et sPLA2 hydrolyserbart liposom karakteriseret ved et indhold af anionisk lipid mellem 20 % og 45 %, et indhold af polymer konjugeret lipid mellem 3 % og 6 % og et indhold af neutrale lipider mellem 40 % og 75 %, hvori det sPLA2 hydrolyserbare liposome omfatter
- cisplatin indkapslet i liposomet,
 - en ydre opløsning,
 - en indre opløsning indeni liposomet,
 - hvori forskellen i osmolyt koncentration mellem den indre opløsning og the ydre opløsning (den ydre koncentration af osmolyte subtraheret fra den indre koncentration af osmolyt) er mellem 280 og 320 mM.
- 2) Den farmaceutiske komposition ifølge krav 1, hvori det eneste terapeutiske middel er cisplatin.
- 3) Den farmaceutiske komposition ifølge hvilket som helst af de foregående krav, hvor den indre opløsning omfatter NaCl eller KCl i en koncentration mellem 0,2-2,5 % vægt/vægt.
- 4) Den farmaceutiske komposition ifølge hvilket som helst af de foregående krav, hvor den ydre opløsning omfatter NaCl eller KCl på en koncentration mellem 0,2-2,5 % vægt/vægt.
- 5) Den farmaceutiske komposition ifølge hvilket som helst af de foregående krav, hvor den indre og ydre opløsning er udvalgt fra gruppen bestående af
- a. Indre opløsning af 0,8-1,0 % NaCl og 9-11 % sukrose og ydre opløsning på 8-12 mM fosfat buffer (pH 6,5) + 9-11 % sukrose,
 - b. Indre opløsning af 1,6-2,0 % NaCl og ydre opløsning på 8-12 mM fosfat buffer (pH 6,5) + 9-11 % sukrose,

c. Indre opløsning af 0,8-1,0 % NaCl og 9-11 % sukrose og ydre opløsning på 8-12 mM fosfat buffer (pH 6,5) + 0,35 % - 0,55 % NaCl + 4-6 % sukrose,

5

d. Indre opløsning af 1,6-2,0 % NaCl og ydre opløsning på 8-12 mM fosfat buffer (pH 6,5) + 0,8-1,0 % NaCl.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7






Figure 8



Pressure = 12500

Figure 9



Pressure = 12500

Figure 10




Figure 11

12/16




Figure 12

Table 1: Actual experimental settings of factors and the responses

Exp no.	Batch no	NaCl conc. (w/v%)	Sucrose concentration (w/v%)	Pressure (KPa)	Initial particle size (nm)	DOE% Day 0	DOE% Day 56	Leakage % Day0- 56
1	LPB0057	0.9	0	5000	82.7	86.0	71.5	14.5
2	LPB0058	1.9	0	5000	86.5	90.1	78.4	11.7
3	LPB0059	0.9	10	5000	93.9	87.0	78.0	9.1
4	LPB0060	1.9	10	5000	88.1	91.5	84.9	6.6
5	LPB0061	0.9	0	20000	67.0	85.1	74.8	10.2
6	LPB0062	1.9	0	20000	65.7	86.8	77.2	9.6
7	LPB0063	0.9	10	20000	76.9	83.9	76.1	7.8
8	LPB0064	1.9	10	20000	67.3	86.4	80.4	6.0
9	LPB0065	0.9	5	12500	71.9	84.8	75.4	9.4
10	LPB0066	1.9	5	12500	75.5	89.3	81.2	8.1
11	LPB0067	1.4	0	12500	70.8	87.6	74.4	13.2
12	LPB0068	1.4	10	12500	80.8	88.5	80.8	7.7
13	LPB0069	1.4	5	5000	97.4	89.6	79.5	10.1
14	LPB0070	1.4	5	20000	65.5	85.4	75.8	9.6
15	LPB0071	1.4	5	12500	76.8	87.7	77.7	10.0
16	LPB0072	1.4	5	12500	76.4	88.1	79.4	8.6
17	LPB0073	1.4	5	12500	67.4	86.3	73.6	12.7

Figure 13

Table 2. Regression coefficients and significance (P) values of the model after back ward elimination

Terms	Initial particle size		Initial DOE%		DOE% 56days		Leakage (%)	
	Regression coefficient	P	Regression coefficient	P	Regression coefficient	P	Regression coefficient	P
Constant	74.2	7.71×10^{-16}	87.3	2.79×10^{-30}	77.6	2.12×10^{-22}	10.3	1.78×10^{-11}
NaCl			1.73	1.25×10^{-5}	2.63	2.67×10^{-4}	-0.90	2.89×10^{-2}
Suc	3.43	2.34×10^{-2}	-	-	2.39	5.85×10^{-4}	-2.19	5.51×10^{-5}
Pre	-10.62	2.40×10^{-6}	-1.66	1.95×10^{-5}	-0.80	1.48×10^{-1}	-0.87	3.31×10^{-2}
NaCl*NaCl			-	-	-	-	-0.97	1.12×10^{-1}
Suc*Suc			-	-	-	-	-	-
Pre*Pre	4.87	3.59×10^{-2}	-	-	-	-	-	-
NaCl*Suc	-	-	-	-	-	-	-	-
NaCl*Pre	-	-	-	-	-	-	-	-
Suc*Pre	-	-	-	-	-1.06	9.08×10^{-2}	-	-
R2	0.852		0.855		0.816		0.812	
Q2	0.749		0.778		0.661		0.657	

Figure 14

Table 3. Overview of liposomal cisplatin formulations prepared with varying osmotic gradient

Batch No	Interior water phase		Exterior water phase		Osmotic gradient Interior-Exterior	Leaky? Yes/No [†]
	Composition [‡]	Osmolarity ^Δ	Composition [‡]	Osmolarity ^Δ		
LPB0021	8mg/mL Cis 0.9 wt% NaCl	313 mOsm	10 wt% Suc 10mM Phos	317 mOsm	- 4 mOsm	Yes
LPB0022	8mg/mL Cis 1.0 wt% NaCl	345 mOsm	10 wt% Suc 10mM Phos	317 mOsm	+ 28 mOsm	Yes
LPB0023	8mg/mL Cis 0.9 wt% NaCl	313 mOsm	0.9 wt% NaCl	286 mOsm	+ 27 mOsm	Yes
LPB0024	8mg/mL Cis 1.1 wt% NaCl	377 mOsm	10 wt% Suc 10mM Phos	317 mOsm	+ 59 mOsm	Yes
LPB0025	8mg/mL Cis 1.2 wt% NaCl	409 mOsm	10 wt% Suc 10mM Phos	317 mOsm	+ 91 mOsm	Yes
LPB0026	8mg/mL Cis 1.8 wt% NaCl	600 mOsm	10 wt% Suc 10mM Phos	317 mOsm	+ 282 mOsm	No
LPB0027	8mg/mL Cis 0.9 wt% NaCl 10 wt% Suc	611 mOsm	10 wt% Suc 10mM Phos	317 mOsm	+ 294 mOsm	No
LPB0110	8mg/mL Cis 0.9 wt% NaCl 10 wt% Suc	611 mOsm	0.9 wt% NaCl 10mM Phos	306 mOsm	+ 306 mOsm	No
LPB0111	8mg/mL Cis 0.9 wt% NaCl 10 wt% Suc	611 mOsm	5 wt% Suc 0.45 wt% NaCl	292 mOsm	+ 319 mOsm	No

[‡] Abbreviations: Cis = cisplatin, Suc = Sucrose, Phos = Sodium phosphate, NaCl = sodium chloride

^Δ Assumes: 1) All material in solution,
2) Negligible increases in interior volumes upon dialysis, and
3) Use of available osmotic coefficients, otherwise assumed to be 1 corresponding to ideality.

[†] Definitions: Yes = More than (>) 10 % relative platinum leakage after 150 days of storage.
No = Less or equal (≤) 10 % relative platinum leakage after 150 days of storage.

Figure 15

Table 4. Overview of different liposomal drug formulations prepared with varying molecular weight

	Liposomal oxaliplatin	Liposomal MTX	Liposomal bleomycin	Liposomal 5FU
Active substance	Oxaliplatin	Methotrexate	Bleomycin	5FU
Hydration media (Interior)	15 mg/mL Oxaliplatin /10% sucrose /1mM calcium gluconate	40mg/mL MTX/10mM Na-Phosphate	25mg/mL Bleomycin /10% Sucrose	30mg/mL 5-FU/ 0.9%NaCl
Dialysis solution (Exterior)	10% sucrose / 1mM calcium gluconate	10% sucrose /10mM Na-Phosphate	10% sucrose /1mM Calcium gluconate	10% sucrose /10mM Na-Phosphate
Interior osmolarity (mOsm)	338	270	315	517
Interior osmolarity (mOsm)	300	317	3300	317
Osmotic gradient (mOsm)	38	-47	15	200
Molecular weight active substance (g/mol)	397	454	1516	130
Leakage (>10%)	No	No	No	Yes

Figure 16