



US 20140286902A1

(19) **United States**

(12) **Patent Application Publication**  
**Proia**

(10) **Pub. No.: US 2014/0286902 A1**

(43) **Pub. Date: Sep. 25, 2014**

(54) **COMBINATION THERAPY OF HSP90  
INHIBITORS WITH  
PLATINUM-CONTAINING AGENTS**

USPC ..... 424/85.7; 514/384; 514/80; 424/649;  
514/266.22; 424/133.1; 514/171; 514/34;  
514/110; 424/85.1; 514/266.4; 514/64; 514/49;  
514/274; 514/249

(71) Applicant: **Synta Pharmaceuticals Corp.,**  
Lexington, MA (US)

(72) Inventor: **David Proia,** Newton, MA (US)

(21) Appl. No.: **14/355,689**

(22) PCT Filed: **Nov. 1, 2012**

(86) PCT No.: **PCT/US2012/063039**

§ 371 (c)(1),

(2), (4) Date: **May 1, 2014**

(57) **ABSTRACT**

A pharmaceutical combination comprising a platinum-containing anticancer agent, and an Hsp90 inhibitor according to the following formulae (I) & (Ia), a tautomer, or a pharmaceutically acceptable salt thereof, wherein the variables structural formulae are defined herein. Also provided is a method of treating a proliferative disorder such as cancer in a subject in need thereof, using the pharmaceutical combination described herein.

**Related U.S. Application Data**

(60) Provisional application No. 61/554,706, filed on Nov. 2, 2011.

**Publication Classification**

(51) **Int. Cl.**

**A61K 31/4196** (2006.01)

**A61K 45/06** (2006.01)

**A61K 33/24** (2006.01)



**A61K 31/675** (2006.01)

(52) **U.S. Cl.**

CPC ..... **A61K 31/4196** (2013.01); **A61K 31/675**

(2013.01); **A61K 45/06** (2013.01); **A61K 33/24**

(2013.01)






Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6

**COMBINATION THERAPY OF HSP90  
INHIBITORS WITH  
PLATINUM-CONTAINING AGENTS**

CROSS-REFERENCE TO RELATED PATENTS

**[0001]** This application claims the benefit of priority to U.S. Provisional Patent Application Nos. 61/554,706, filed on Nov. 2, 2011. The contents of the above application are incorporated herein by reference in their entirety.

BACKGROUND OF THE INVENTION

**[0002]** Although tremendous advances have been made in elucidating the genomic abnormalities that cause malignant cancer cells, currently available chemotherapy remains unsatisfactory, and the prognosis for the majority of patients diagnosed with cancer remains dismal. Most chemotherapeutic agents act on a specific molecular target thought to be involved in the development of the malignant phenotype. However, a complex network of signaling pathways regulate cell proliferation and the majority of malignant cancers are facilitated by multiple genetic abnormalities in these pathways. Therefore, it is less likely that a therapeutic agent that acts on one molecular target will be fully effective in curing a patient who has cancer.

**[0003]** Heat shock proteins (HSPs) are a class of chaperone proteins that are up-regulated in response to elevated temperature and other environmental stresses, such as ultraviolet light, nutrient deprivation and oxygen deprivation. HSPs act as chaperones to other cellular proteins (called client proteins), facilitate their proper folding and repair and aid in the refolding of mis-folded client proteins. There are several known families of HSPs, each having its own set of client proteins. The Hsp90 family is one of the most abundant HSP families, accounting for about 1-2% of proteins in a cell that is not under stress and increasing to about 4-6% in a cell under stress. Inhibition of Hsp90 results in the degradation of its client proteins via the ubiquitin proteasome pathway. Unlike other chaperone proteins, the client proteins of Hsp90 are mostly protein kinases or transcription factors involved in signal transduction, and a number of its client proteins have been shown to be involved in the progression of cancer.

SUMMARY OF THE INVENTION

**[0004]** It has been found that certain triazolone Hsp90 inhibitor and platinum-containing anti-cancer agent combinations are surprisingly effective at treating subjects with certain cancers without further increasing the side effects profile of the individual agents. The particular combination therapies disclosed herein demonstrate surprising biological activity by demonstrating significant anticancer effects.

**[0005]** In an embodiment, the invention provides a method of utilizing Hsp90 inhibitors according to formulae (I) or (Ia), or a compound in Tables 1 or 2 for the treatment of proliferative disorders such as cancer, in combination with a platinum-containing anti-cancer agent. In an embodiment, a method of treating a subject with cancer includes administering to the subject an Hsp90 inhibitor according to formulae (I) or (Ia), or a compound in Tables 1 or 2, and a platinum-containing anti-cancer agent. In an embodiment, the administration of the Hsp90 inhibitor and the platinum-containing agent are done concurrently. In another embodiment, the administration of the Hsp90 inhibitor and the platinum-containing agent are done sequentially. In another embodiment, the adminis-

tration of the Hsp90 inhibitor and the platinum-containing agent are dosed independently. In any one of these embodiments, the platinum-containing agent may be cisplatin, carboplatin, oxaliplatin, or their analogues. In any one of these embodiments, the Hsp90 inhibitor may be a compound represented in Tables 1 or 2. In any one of these embodiments, the platinum-containing agent may be cisplatin.

**[0006]** In some embodiments, the cancer may have mutations or translocations in the EGFR, K-Ras, c-Met-c-Met, c-Kit, HER2, B-Raf, PI3K and/or ALK proteins. In some embodiments, the cancer may express wild-type EGFR and K-Ras. In some embodiments, the cancer may express mutated EGFR and wild-type K-Ras. In some embodiments, the cancer may express wild-type EGFR and mutated K-Ras protein. In some embodiments, the cancer may be ALK positive ("ALK+"). In some embodiments, the cancer may have an EML4-ALK translocation. In some embodiments, the cancer may have a HER2 mutation. In some embodiments, the cancer may have a PI3K mutation. In some embodiments, the cancer may have a B-Raf protein mutation.

**[0007]** In an embodiment, kits for administration of the combination therapy are provided. In an embodiment, the kit includes separate pharmaceutical compositions containing the Hsp90 inhibitor according to formulae (I) or (Ia) or a compound in Tables 1 or 2, and a platinum-containing anti-cancer agent. In another embodiment, the kit includes one pharmaceutical composition containing both the Hsp90 inhibitor and the platinum-containing anti-cancer agent. In any of these embodiments, each pharmaceutical composition includes one or more pharmaceutically acceptable carrier or diluent. In any one of these embodiments, the platinum-containing agent may be cisplatin, carboplatin, oxaliplatin, or their analogues. In any one of these embodiments, the Hsp90 inhibitor may be a compound represented in Tables 1 or 2. In any one of these embodiments, the platinum-containing agent may be cisplatin.

**[0008]** In an embodiment, the invention provides the use of an Hsp90 inhibitor according to formulae (I) or (Ia) or a compound in Tables 1 or 2 for the manufacture of a medication for treating cancer in combination with a platinum-containing anti-cancer agent.

**[0009]** In an embodiment, a method of treating drug-resistant cancer in a subject includes administering to the subject an effective amount of the pharmaceutical combination including an Hsp90 compound according to formulae (I) or (Ia) or a compound in Tables 1 or 2 and a platinum-containing anti-cancer agent. In an embodiment, the method further includes the administration of one or more therapeutic agents. In certain embodiments, the combination treatment utilizes an Hsp90 compound according to formulae (I) or (Ia) or a compound in Tables 1 or 2 with a platinum-containing anti-cancer agent to help to arrest, partially or fully, or reduce the development of drug resistant cancer in a subject. The combinations described herein may allow a reduced dose of the platinum-containing agent given to a subject, because the Hsp90 inhibitor should inhibit the development of multidrug-resistant cancer cells. In an embodiment, the platinum-containing anti-cancer agent may be cisplatin, carboplatin, oxaliplatin, or their derivatives. In another embodiment, the platinum-containing agent may be cisplatin.

BRIEF DESCRIPTION OF THE DRAWINGS

**[0010]** The foregoing and other objects, features and advantages of the invention will be apparent from the follow-

ing more particular description of some embodiments of the invention, as illustrated in the accompanying drawings. The drawings are not necessarily to scale, emphasis instead being placed upon illustrating the principles of the invention.

**[0011]** FIG. 1 shows a dose-dependent curve with the  $IC_{50}$  of ganetespib, at about 32 nM.

**[0012]** FIG. 2 shows a dose-dependent curve with the  $IC_{50}$  of cisplatin at about 5.7  $\mu$ M.

**[0013]** FIG. 3 shows significant killing of HCT-116 cells by ganetespib in combination with cisplatin. Cells were exposed to the indicated single agent or combination, concurrently, for 3 days.

**[0014]** FIG. 4 shows significant killing of HCT-116 cells by the sequential combination of ganetespib with cisplatin. Cells were exposed to ganetespib for 1 hour, washed and then treated with vehicle (DMSO) or indicated chemotherapeutic for 3 days. Single agent chemotherapeutic was dosed for 3 days.

**[0015]** FIG. 5 shows substantial treatment improvement by a combination of ganetespib with cisplatin in an animal model of HCT-116 colon cancer xenograft. Ganetespib (150 mg/kg) was given as a single bolus injection 6 hours prior to cisplatin (2 mg/kg) administration.

**[0016]** FIG. 6 shows substantial treatment improvement by a combination of ganetespib with cisplatin in an animal model of MDA-MB-231 breast cancer tumor xenograft. Ganetespib (150 mg/kg) was given as a single bolus injection 6 hours prior to cisplatin (2 mg/kg) administration.

## DETAILED DESCRIPTION OF THE INVENTION

### Definitions

**[0017]** Unless otherwise specified, the below terms used herein are defined as follows:

**[0018]** As used herein, the term “alkyl” means a saturated or unsaturated, straight chain or branched, non-cyclic hydrocarbon having from 1 to 10 carbon atoms. Representative straight chain alkyls include methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl and n-decyl; while representative branched alkyls include isopropyl, sec-butyl, isobutyl, tert-butyl, isopentyl, 2-methylbutyl, 3-methylbutyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2-methylhexyl, 3-methylhexyl, 4-methylhexyl, 5-methylhexyl, 2,3-dimethylbutyl, 2,3-dimethylpentyl, 2,4-dimethylpentyl, 2,3-dimethylhexyl, 2,4-dimethylhexyl, 2,5-dimethylhexyl, 2,2-dimethylpentyl, 2,2-dimethylhexyl, 3,3-dimethylpentyl, 3,3-dimethylhexyl, 4,4-dimethylhexyl, 2-ethylpentyl, 3-ethylpentyl, 2-ethylhexyl, 3-ethylhexyl, 4-ethylhexyl, 2-methyl-2-ethylpentyl, 2-methyl-3-ethylpentyl, 2-methyl-4-ethylpentyl, 2-methyl-2-ethylhexyl, 2-methyl-3-ethylhexyl, 2-methyl-4-ethylhexyl, 2,2-diethylpentyl, 3,3-diethylhexyl, 2,2-diethylhexyl, 3,3-diethylhexyl, and the like. The term “(C<sub>1</sub>-C<sub>6</sub>)alkyl” means a saturated, straight chain or branched, non-cyclic hydrocarbon having from 1 to 6 carbon atoms. Alkyl groups included in compounds described herein may be optionally substituted with one or more substituents. Examples of unsaturated alkyls include vinyl, allyl, 1-butenyl, 2-butenyl, isobutylenyl, 1-pentenyl, 2-pentenyl, 3-methyl-1-butenyl, 2-methyl-2-butenyl, 2,3-dimethyl-2-butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 1-heptenyl, 2-heptenyl, 3-heptenyl, 1-octenyl, 2-octenyl, 3-octenyl, 1-nonenyl, 2-nonenyl, 3-nonenyl, 1-decenyl, 2-decenyl, 3-decenyl, acetylenyl, propynyl, 1-butyne, 2-butyne, 1-pentynyl, 2-pentynyl, 3-methyl-1-butyne, 4-pentynyl, 1-hexy-

nyl, 2-hexynyl, 5-hexynyl, 1-heptynyl, 2-heptynyl, 6-heptynyl, 1-octynyl, 2-octynyl, 7-octynyl, 1-nonynyl, 2-nonynyl, 8-nonynyl, 1-decynyl, 2-decynyl, 9-decynyl, and the like. Alkyl groups included in compounds described herein may be optionally substituted with one or more substituents.

**[0019]** As used herein, the term “cycloalkyl” means a saturated or unsaturated, mono- or polycyclic, non-aromatic hydrocarbon having from 3 to 20 carbon atoms. Representative cycloalkyls include cyclopropyl, 1-methylcyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, octahydroptaleny, cyclohexenyl, cyclooctenyl, cyclohexynyl, and the like. Cycloalkyl groups included in compounds described herein may be optionally substituted with one or more substituents.

**[0020]** As used herein, the term “alkylene” refers to an alkyl group that has two points of attachment. The term “(C<sub>1</sub>-C<sub>6</sub>)alkylene” refers to an alkylene group that has from one to six carbon atoms. Straight chain (C<sub>1</sub>-C<sub>6</sub>)alkylene groups are preferred. Non-limiting examples of alkylene groups include methylene (—CH<sub>2</sub>—), ethylene (—CH<sub>2</sub>CH<sub>2</sub>—), n-propylene (—CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>—), isopropylene (—CH<sub>2</sub>CH(CH<sub>3</sub>)—), and the like. Alkylene groups may be saturated or unsaturated, and may be optionally substituted with one or more substituents.

**[0021]** As used herein, the term “lower” refers to a group having up to four atoms. For example, a “lower alkyl” refers to an alkyl radical having from 1 to 4 carbon atoms, “lower alkoxy” refers to “—O—(C<sub>1</sub>-C<sub>4</sub>)alkyl”.

**[0022]** As used herein, the term “haloalkyl” means an alkyl group, in which one or more, including all, the hydrogen radicals are replaced by a halo group(s), wherein each halo group is independently selected from —F, —Cl, —Br, and —I. For example, the term “halomethyl” means a methyl in which one to three hydrogen radical(s) have been replaced by a halo group. Representative haloalkyl groups include trifluoromethyl, bromomethyl, 1,2-dichloroethyl, 4-iodobutyl, 2-fluoropentyl, and the like.

**[0023]** As used herein, an “alkoxy” is an alkyl group which is attached to another moiety via an oxygen linker. Alkoxy groups included in compounds described herein may be optionally substituted with one or more substituents.

**[0024]** As used herein, a “haloalkoxy” is a haloalkyl group which is attached to another moiety via an oxygen linker.

**[0025]** As used herein, the term an “aromatic ring” or “aryl” means a mono- or polycyclic hydrocarbon, containing from 6 to 15 carbon atoms, in which at least one ring is aromatic. Examples of suitable aryl groups include phenyl, tolyl, anthracenyl, fluorenyl, indenyl, azuleny, and naphthyl, as well as benzo-fused carbocyclic moieties such as 5,6,7,8-tetrahydronaphthyl. Aryl groups included in compounds described herein may be optionally substituted with one or more substituents. In an embodiment, the aryl group is a monocyclic ring, wherein the ring comprises 6 carbon atoms, referred to herein as “(C<sub>6</sub>)aryl.”

**[0026]** As used herein, the term “aralkyl” means an aryl group that is attached to another group by a (C<sub>1</sub>-C<sub>6</sub>)alkylene group. Representative aralkyl groups include benzyl, 2-phenyl-ethyl, naphth-3-yl-methyl and the like. Aralkyl groups included in compounds described herein may be optionally substituted with one or more substituents.

**[0027]** As used herein, the term “heterocyclyl” means a monocyclic or a polycyclic, saturated or unsaturated, non-aromatic ring or ring system which typically contains 5- to 20-members and at least one heteroatom. A heterocyclic ring

system can contain saturated ring(s) or unsaturated non-aromatic ring(s), or a mixture thereof. A 3- to 10-membered heterocycle can contain up to 5 heteroatoms, and a 7- to 20-membered heterocycle can contain up to 7 heteroatoms. Typically, a heterocycle has at least one carbon atom ring member. Each heteroatom is independently selected from nitrogen, which can be oxidized (e.g., N(O)) or quaternized, oxygen and sulfur, including sulfoxide and sulfone. The heterocycle may be attached via any heteroatom or carbon atom. Representative heterocycles include morpholinyl, thiomorpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, hydantoinyl, valerolactamyl, oxiranyl, oxetanyl, tetrahydrofuranlyl, tetrahydropyranlyl, tetrahydropyridinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranlyl, and the like. A heteroatom may be substituted with a protecting group known to those of ordinary skill in the art, for example, a nitrogen atom may be substituted with a tert-butoxycarbonyl group. Furthermore, the heterocycle included in compounds described herein may be optionally substituted with one or more substituents. Only stable isomers of such substituted heterocyclic groups are contemplated in this definition.

**[0028]** As used herein, the term “heteroaryl”, or like terms, means a monocyclic or a polycyclic, unsaturated radical containing at least one heteroatom, in which at least one ring is aromatic. Polycyclic heteroaryl rings must contain at least one heteroatom, but not all rings of a polycyclic heteroaryl moiety must contain heteroatoms. Each heteroatom is independently selected from nitrogen, which can be oxidized (e.g., N(O)) or quaternized, oxygen and sulfur, including sulfoxide and sulfone. Representative heteroaryl groups include pyridyl, 1-oxo-pyridyl, furanyl, benzo[1,3]dioxolyl, benzo[1,4]dioxinyl, thienyl, pyrrolyl, oxazolyl, imidazolyl, thiazolyl, an isoxazolyl, quinolinyl, pyrazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, a triazinyl, triazolyl, thiadiazolyl, isoquinolinyl, indazolyl, benzoxazolyl, benzofuryl, indoliziny, imidazopyridyl, tetrazolyl, benzimidazolyl, benzothiazolyl, benzothiadiazolyl, benzoxadiazolyl, indolyl, tetrahydroindolyl, azaindolyl, imidazopyridyl, quinazoliny, purinyl, pyrrolo[2,3]pyrimidinyl, pyrazolo[3,4]pyrimidinyl, imidazo[1,2-a]pyridyl, and benzothienyl. In one embodiment, the heteroaromatic ring may be a 5-8 membered monocyclic heteroaryl ring. The point of attachment of a heteroaromatic or heteroaryl ring may be at either a carbon atom or a heteroatom. Heteroaryl groups included in compounds described herein may be optionally substituted with one or more substituents. As used herein, the term “(C<sub>5</sub>) heteroaryl” means an heteroaromatic ring of 5 members, wherein at least one carbon atom of the ring is replaced with a heteroatom, such as, for example, oxygen, sulfur or nitrogen. Representative (C<sub>5</sub>)heteroaryls include furanyl, thienyl, pyrrolyl, oxazolyl, imidazolyl, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, pyrazinyl, triazolyl, thiadiazolyl, and the like. As used herein, the term “(C<sub>6</sub>)heteroaryl” means an aromatic heterocyclic ring of 6 members, wherein at least one carbon atom of the ring is replaced with a heteroatom such as, for example, oxygen, nitrogen or sulfur. Representative (C<sub>6</sub>) heteroaryls include pyridyl, pyridazinyl, pyrazinyl, triazinyl, tetrazinyl, and the like.

**[0029]** As used herein, the term “heteroaralkyl” means a heteroaryl group that is attached to another group by a (C<sub>1</sub>-C<sub>6</sub>)alkylene. Representative heteroaralkyls include 2-(pyridin-4-yl)-propyl, 2-(thien-3-yl)-ethyl, imidazol-4-yl-methyl,

and the like. Heteroaralkyl groups included in compounds described herein may be optionally substituted with one or more substituents.

**[0030]** As used herein, the term “halogen” or “halo” means —F, —Cl, —Br or —I.

**[0031]** Suitable substituents for an alkyl, alkylene, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, aralkyl, heteroaryl, and heteroaralkyl groups include are those substituents which form a stable compound described herein without significantly adversely affecting the reactivity or biological activity of the compound described herein. Examples of substituents for an alkyl, alkylene, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, aralkyl, heteroaryl, and heteroaralkyl include an alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, heteroaryl, aralkyl, heteraralkyl, heteroalkyl, alkoxy, (each of which can be optionally and independently substituted), —C(O)NR<sup>28</sup>R<sup>29</sup>, —C(S)NR<sup>28</sup>R<sup>29</sup>, —C(NR<sup>32</sup>)NR<sup>28</sup>R<sup>29</sup>, —NR<sup>33</sup>C(O)R<sup>31</sup>, —NR<sup>33</sup>C(S)R<sup>31</sup>, —NR<sup>33</sup>C(NR<sup>32</sup>)R<sup>31</sup>, halo, —OR<sup>33</sup>, cyano, nitro, —C(O)R<sup>33</sup>, —C(S)R<sup>33</sup>, —C(NR<sup>32</sup>)R<sup>33</sup>, —NR<sup>28</sup>R<sup>29</sup>, —C(O)OR<sup>33</sup>, —C(S)OR<sup>33</sup>, —C(NR<sup>32</sup>)OR<sup>33</sup>, —OC(O)R<sup>33</sup>, —OC(S)R<sup>33</sup>, —OC(NR<sup>32</sup>)R<sup>33</sup>, —NR<sup>30</sup>C(O)NR<sup>28</sup>R<sup>29</sup>, —NR<sup>33</sup>C(S)NR<sup>28</sup>R<sup>29</sup>, —NR<sup>33</sup>C(NR<sup>32</sup>)NR<sup>28</sup>R<sup>29</sup>, —OC(O)NR<sup>28</sup>R<sup>29</sup>, —OC(S)NR<sup>28</sup>R<sup>29</sup>, —OC(NR<sup>32</sup>)NR<sup>28</sup>R<sup>29</sup>, —OC(NR<sup>32</sup>)NR<sup>28</sup>R<sup>29</sup>, —NR<sup>33</sup>C(O)OR<sup>31</sup>, —NR<sup>33</sup>C(S)OR<sup>31</sup>, —NR<sup>33</sup>C(NR<sup>32</sup>)OR<sup>31</sup>, —S(O)<sub>k</sub>R<sup>33</sup>, —OS(O)<sub>k</sub>R<sup>33</sup>, —NR<sup>33</sup>S(O)<sub>k</sub>R<sup>33</sup>, —S(O)<sub>k</sub>NR<sup>28</sup>R<sup>29</sup>, —OS(O)<sub>k</sub>NR<sup>28</sup>R<sup>29</sup>, —NR<sup>33</sup>S(O)<sub>k</sub>NR<sup>28</sup>R<sup>29</sup>, guanidino, —C(O)SR<sup>31</sup>, —C(S)SR<sup>31</sup>, —C(NR<sup>32</sup>)SR<sup>31</sup>, —OC(O)OR<sup>31</sup>, —OC(S)OR<sup>31</sup>, —OC(NR<sup>32</sup>)OR<sup>31</sup>, —SC(O)R<sup>33</sup>, —SC(O)OR<sup>31</sup>, —SC(NR<sup>32</sup>)OR<sup>31</sup>, —SC(S)R<sup>33</sup>, —SC(S)OR<sup>31</sup>, —SC(O)NR<sup>28</sup>R<sup>29</sup>, —SC(NR<sup>32</sup>)NR<sup>28</sup>R<sup>29</sup>, —SC(S)NR<sup>28</sup>R<sup>29</sup>, —SC(NR<sup>32</sup>)R<sup>33</sup>, —OS(O)<sub>k</sub>OR<sup>31</sup>, —S(O)<sub>k</sub>OR<sup>31</sup>, —NR<sup>30</sup>S(O)<sub>k</sub>OR<sup>31</sup>, —SS(O)<sub>k</sub>R<sup>33</sup>, —SS(O)<sub>k</sub>OR<sup>31</sup>, —SS(O)<sub>k</sub>NR<sup>28</sup>R<sup>29</sup>, —OP(O)(OR<sup>31</sup>)<sub>2</sub>, or —SP(O)(OR<sup>31</sup>)<sub>2</sub>. In addition, any saturated portion of an alkyl, cycloalkyl, alkylene, heterocyclyl, alkenyl, cycloalkenyl, alkynyl, aralkyl and heteroaralkyl groups, may also be substituted with =O, =S, or =N—R<sup>32</sup>. Each R<sup>28</sup> and R<sup>29</sup> is independently H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, heteroaryl, aralkyl, or heteraralkyl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, heteroaryl, aralkyl, or heteroalkyl represented by R<sup>28</sup> or R<sup>29</sup> is optionally and independently substituted. Each R<sup>30</sup>, R<sup>31</sup> and R<sup>33</sup> is independently H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, heteroaryl, aralkyl, or heteraralkyl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, heteroaryl, aralkyl, and heteraralkyl represented by R<sup>30</sup> or R<sup>31</sup> or R<sup>33</sup> is optionally and independently unsubstituted. Each R<sup>32</sup> is independently H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, heteroaryl, aralkyl, heteraralkyl, —C(O)R<sup>33</sup>, —C(O)NR<sup>28</sup>R<sup>29</sup>, —S(O)<sub>k</sub>R<sup>33</sup>, or —S(O)<sub>k</sub>NR<sup>28</sup>R<sup>29</sup>, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, heteroaryl, aralkyl and heteraralkyl represented by R<sup>32</sup> is optionally and independently substituted. The variable k is 0, 1 or 2. In some embodiments, suitable substituents include C1-C4 alkyl, C1-C4 haloalkyl, C1-C4 alkoxy, C1-C4 haloalkoxy, C1-C4 hydroxyalkyl, halo, or hydroxyl.

**[0032]** When a heterocyclyl, heteroaryl or heteroaralkyl group contains a nitrogen atom, it may be substituted or unsubstituted. When a nitrogen atom in the aromatic ring of a heteroaryl group has a substituent, the nitrogen may be oxidized or a quaternary nitrogen.

**[0033]** Unless indicated otherwise, the compounds described herein containing reactive functional groups, such as, for example, carboxy, hydroxy, thiol and amino moieties, also include corresponding protected derivatives thereof. "Protected derivatives" are those compounds in which a reactive site or sites are blocked with one or more protecting groups. Examples of suitable protecting groups for hydroxyl groups include benzyl, methoxymethyl, allyl, trimethylsilyl, tert-butyldimethylsilyl, acetate, and the like. Examples of suitable amine protecting groups include benzyloxycarbonyl, tert-butoxycarbonyl, tert-butyl, benzyl and fluorenylmethoxycarbonyl (Fmoc). Examples of suitable thiol protecting groups include benzyl, tert-butyl, acetyl, methoxymethyl and the like. Other suitable protecting groups are well known to those of ordinary skill in the art and include those found in T. W. GREENE, *PROTECTING GROUPS IN ORGANIC SYNTHESIS*, (John Wiley & Sons, Inc., 1981).

**[0034]** As used herein, the term "compound(s) described herein" or similar terms refers to a compound of formulae (I), or (Ia) or a compound in Tables 1 or 2 or a tautomer or pharmaceutically acceptable salt thereof. Also included in the scope of the embodiments are a solvate, clathrate, hydrate, polymorph, prodrug, or protected derivative of a compound of formulae (I), or (Ia), or a compound in Tables 1 or 2.

**[0035]** The compounds described herein may contain one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as double-bond isomers (i.e., geometric isomers), enantiomers or diastereomers. Each chemical structure shown herein, including the compounds described herein, encompass all of the corresponding compound's enantiomers, diastereomers and geometric isomers, that is, both the stereochemically pure form (e.g., geometrically pure, enantiomerically pure, or diastereomerically pure) and isomeric mixtures (e.g., enantiomeric, diastereomeric and geometric isomeric mixtures). In some cases, one enantiomer, diastereomer or geometric isomer will possess superior activity or an improved toxicity or kinetic profile compared to other isomers. In those cases, such enantiomers, diastereomers and geometric isomers of compounds described herein are preferred.

**[0036]** When a disclosed compound is named or depicted by structure, it is to be understood that solvates (e.g., hydrates) of the compound or a pharmaceutically acceptable salt thereof is also included. "Solvates" refer to crystalline forms wherein solvent molecules are incorporated into the crystal lattice during crystallization. Solvates may include water or nonaqueous solvents such as ethanol, isopropanol, DMSO, acetic acid, ethanolamine and ethyl acetate. When water is the solvent molecule incorporated into the crystal lattice of a solvate, it is typically referred to as a "hydrate". Hydrates include stoichiometric hydrates as well as compositions containing variable amounts of water.

**[0037]** When a disclosed compound is named or depicted by structure, it is to be understood that the compound, including solvates thereof, may exist in crystalline forms, non-crystalline forms or a mixture thereof. The compounds or solvates may also exhibit polymorphism (i.e., the capacity to occur in different crystalline forms). These different crystalline forms are typically known as "polymorphs." It is to be understood that when named or depicted by structure, the disclosed compounds and solvates (e.g., hydrates) also include all polymorphs thereof. Polymorphs have the same chemical composition but differ in packing, geometrical arrangement and other descriptive properties of the crystal-

line solid state. Polymorphs, therefore, may have different physical properties such as shape, density, hardness, deformability, stability and dissolution properties. Polymorphs typically exhibit different melting points, IR spectra and X-ray powder diffraction patterns, which may be used for identification. One of ordinary skill in the art will appreciate that different polymorphs may be produced, for example, by changing or adjusting the conditions used in crystallizing the compound. For example, changes in temperature, pressure or solvent may result in different polymorphs. In addition, one polymorph may spontaneously convert to another polymorph under certain conditions.

**[0038]** When a disclosed compound is named or depicted by structure, it is to be understood that clathrates ("inclusion compounds") of the compound or its pharmaceutically acceptable salt, solvate or polymorph, are also included. "Clathrate" means a compound described herein, or a salt thereof, in the form of a crystal lattice that contains spaces (e.g., channels) that have a guest molecule trapped within (e.g., a solvent or water).

**[0039]** As used herein, and unless otherwise indicated, the term "prodrug" means a derivative of a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (in vitro or in vivo) to provide a compound described herein. Prodrugs may become active upon such reaction under biological conditions, or they may have activity in their unreacted forms. Examples of prodrugs contemplated herein include analogs or derivatives of compounds of formulae (I) or (Ia) or a compound in Tables 1 or 2 that comprise biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides and phosphate analogues. Prodrugs can typically be prepared using well-known methods, such as those described by BURGER'S MEDICINAL CHEMISTRY AND DRUG DISCOVERY, (Manfred E. Wolff Ed., 5th ed. (1995)) 172-178, 949-982.

**[0040]** The articles "a", "an" and "the" are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article unless otherwise clearly indicated by contrast. By way of example, "an element" means one element or more than one element.

**[0041]** The term "including" is used herein to mean, and is used interchangeably with, the phrase "including but not limited to".

**[0042]** The term "or" is used herein to mean, and is used interchangeably with, the term "and/or," unless context clearly indicates otherwise.

**[0043]** The term "such as" is used herein to mean, and is used interchangeably, with the phrase "such as but not limited to".

**[0044]** Unless specifically stated or obvious from context, as used herein, the term "about" is understood as within a range of normal tolerance in the art, for example within 2 standard deviations of the mean. About can be understood as within 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.1%, 0.05%, or 0.01% of the stated value. Unless otherwise clear from context, all numerical values provided herein can be modified by the term about.

**[0045]** As used herein, the terms "subject", "patient" and "mammal" are used interchangeably. The terms "subject" and "patient" refer to an animal (e.g., a bird such as a chicken, quail or turkey, or a mammal), preferably a mammal including a non-primate (e.g., a cow, pig, horse, sheep, rabbit, guinea pig, rat, cat, dog, and mouse) and a primate (e.g., a



monkey, chimpanzee and a human), and more preferably a human. In an embodiment, the subject is a non-human animal such as a farm animal (e.g., a horse, cow, pig or sheep), or a pet (e.g., a dog, cat, guinea pig or rabbit). In another embodiment, the subject is a human.

**[0046]** As used herein, “Hsp90” includes each member of the family of heat shock proteins having a mass of about 90-kiloDaltons. For example, in humans the highly conserved Hsp90 family includes the cytosolic Hsp90 $\alpha$  and Hsp90 $\beta$  isoforms, as well as GRP94, which is found in the endoplasmic reticulum, and HSP75/TRAP1, which is found in the mitochondrial matrix.

**[0047]** The term “c-Kit” or “c-Kit kinase” refers to a membrane receptor protein tyrosine kinase which is preferably activated upon binding Stem Cell Factor (SCF) to its extracellular domain. Yarden, et al., *Embo. J.*, (1987) 11:3341-3351; Qiu, et al., *Embo. J.*, (1988) 7:1003-1011. The full length amino acid sequence of a c-Kit kinase preferably is as set forth in Yarden, et al.; and Qiu, et al., which are incorporated by reference herein in their entirety. Mutant versions of c-Kit kinase are encompassed by the term “c-Kit” or “c-Kit kinase” and include those that fall into two classes: (1) having a single amino acid substitution at codon 816 of the human c-Kit kinase, or its equivalent position in other species (Ma, et al., *J. Invest Dermatol.*, (1999) 112:165-170), and (2) those which have mutations involving the putative juxtamembrane  $\alpha$ -helix of the protein (Ma, et al., *J. Biol. Chem.*, (1999) 274:13399-13402). Both of these publications are incorporated by reference herein in their entirety, including any drawings.

**[0048]** As used herein, “BCR-ABL” is a fusion protein that results from the translocation of gene sequences from c-ABL protein tyrosine kinase on chromosome 9 into BCR sequences on chromosome 22 producing the Philadelphia chromosome. A schematic representation of human BCR, ABL and BCR-ABL can be seen in FIG. 1 of U.S. patent application Ser. No. 10/193,651, filed on Jul. 9, 2002. Depending on the breaking point in the BCR gene, BCR-ABL fusion proteins can vary in size from 185-230 kD but they must contain at least the OLI domain from BCR and the TK domain from ABL for transforming activity. The most common BCR-ABL gene products found in humans are P230 BCR-ABL, P210 BCR-ABL and P190 BCR-ABL. P210 BCR-ABL is characteristic of CML and P190 BCR-ABL is characteristic of ALL.

**[0049]** FLT3 kinase is a tyrosine kinase receptor involved in the regulation and stimulation of cellular proliferation. Gilliland, et al., *Blood* (2002), 100:1532-42. The FLT3 kinase has five immunoglobulin-like domains in its extracellular region, as well as an insert region of 75-100 amino acids in the middle of its cytoplasmic domain. FLT3 kinase is activated upon the binding of the FLT3 ligand which causes receptor dimerization. Dimerization of the FLT3 kinase by FLT3 ligand activates the intracellular kinase activity as well as a cascade of downstream substrates including Stat5, Ras, phosphatidylinositol-3-kinase (PI3K), Erk2, Akt, MAPK, SHC, SHP2 and SHIP. Rosnet, et al., *Acta Haematol.* (1996), 95:218; Hayakawa, et al., *Oncogene* (2000), 19:624; Mizuki, et al., *Blood* (2000), 96:3907; Gilliland, et al., *Curr. Opin. Hematol.* (2002), 9: 274-81. Both membrane-bound and soluble FLT3 ligand bind, dimerize, and subsequently activate the FLT3 kinase.

**[0050]** Normal cells that express FLT3 kinase include immature hematopoietic cells, typically CD34+ cells, pla-

centa, gonads and brain. Rosnet, et al., *Blood* (1993), 82:1110-19; Small, et al., *Proc. Natl. Acad. Sci. U.S.A.* (1994), 91:459-63; Rosnet, et al., *Leukemia* (1996), 10:238-48. However, efficient stimulation of proliferation via FLT3 kinase typically requires other hematopoietic growth factors or interleukins. FLT3 kinase also plays a critical role in immune function through its regulation of dendritic cell proliferation and differentiation. McKenna, et al., *Blood* (2000), 95:3489-497. Numerous hematologic malignancies express FLT3 kinase, the most prominent of which is AML. Yokota, et al., *Leukemia* (1997), 11:1605-09. Other FLT3 expressing malignancies include B-precursor cell acute lymphoblastic leukemias, myelodysplastic leukemias, T-cell acute lymphoblastic leukemias, and chronic myelogenous leukemias. Rasko, et al., *Leukemia* (1995), 9:2058-66.

**[0051]** FLT3 kinase mutations associated with hematologic malignancies are activating mutations. In other words, the FLT3 kinase is constitutively activated without the need for binding and dimerization by FLT3 ligand, and therefore stimulates the cell to grow continuously. Two types of activating mutations have been identified: internal tandem duplications (ITDs) and point mutation in the activating loop of the kinase domain. As used herein, the term “FLT3 kinase” refers to both wild type FLT3 kinase and mutant FLT3 kinases, such as FLT3 kinases that have activating mutations. Compounds provided herein are useful in treating conditions characterized by inappropriate FLT3 activity, such as proliferative disorders. Inappropriate FLT3 activity includes enhanced FLT3 activity resulting from increased or de novo expression of FLT3 in cells, increased FLT3 expression or activity, and FLT3 mutations resulting in constitutive activation. The existence of inappropriate or abnormal FLT3 ligand and FLT3 levels or activity can be determined using well-known methods in the art. For example, abnormally high FLT3 levels can be determined using commercially available ELISA kits. FLT3 levels can also be determined using flow cytometric analysis, immunohistochemical analysis and in situ hybridization techniques.

**[0052]** “Epidermal growth factor receptor” or “EGFR”, as used herein, means any epidermal growth factor receptor (EGFR) protein, peptide, or polypeptide having EGFR or EGFR family activity (e.g., Her1, Her2, Her3 and/or Her4), such as encoded by EGFR Genbank Accession Nos. shown in Table I of U.S. patent application Ser. No. 10/923,354, filed on Aug. 20, 2004, or any other EGFR transcript derived from a EGFR gene and/or generated by EGFR translocation. The term “EGFR” is also meant to include other EGFR proteins, peptides, or polypeptides derived from EGFR isoforms (e.g., Her1, Her2, Her3 and/or Her4), mutant EGFR genes, splice variants of EGFR genes, and EGFR gene polymorphisms.

**[0053]** EGFR is a member of the type 1 subgroup of receptor tyrosine kinase family of growth factor receptors which play critical roles in cellular growth, differentiation and survival. Activation of these receptors typically occurs via specific ligand binding which results in hetero- or homodimerization between receptor family members, with subsequent autophosphorylation of the tyrosine kinase domain. Specific ligands which bind to EGFR include epidermal growth factor (EGF), transforming growth factor  $\alpha$  (TGF $\alpha$ ), amphiregulin and some viral growth factors. Activation of EGFR triggers a cascade of intracellular signaling pathways involved in both cellular proliferation (the ras/raf/MAP kinase pathway) and

survival (the PI3 kinase/Akt pathway). Members of this family, including EGFR and HER2, have been directly implicated in cellular transformation.

**[0054]** A number of human malignancies are associated with aberrant or overexpression of EGFR and/or overexpression of its specific ligands. Gullick, *Br. Med. Bull.* (1991), 47:87-98; Modjtahedi & Dean, *Int. J. Oncol.* (1994), 4:277-96; Salomon, et al., *Crit. Rev. Oncol. Hematol.* (1995), 19:183-232. Aberrant or overexpression of EGFR has been associated with an adverse prognosis in a number of human cancers, including head and neck, breast, colon, prostate, lung (e.g., NSCLC, adenocarcinoma and squamous lung cancer), ovarian, gastrointestinal cancers (gastric, colon, pancreatic), renal cell cancer, bladder cancer, glioma, gynecological carcinomas and prostate cancer. In some instances, overexpression of tumor EGFR has been correlated with both chemoresistance and a poor prognosis. Lei, et al., *Anti-cancer Res.* (1999), 19:221-28; Veale, et al., *Br. J. Cancer* (1993); 68:162-65. Mutations in EGFR are associated with many types of cancer as well. For example, EGFR mutations are highly prevalent in non-mucinous BAC patients. Finberg, et al., *J. Mol. Diagnostics* (2007) 9(3):320-26.

**[0055]** c-Met is a receptor tyrosine kinase that is encoded by the Met protooncogene and transduces the biological effects of hepatocyte growth factor (HGF), which is also referred to as scatter factor (SF). Jiang et al., *Crit. Rev. Oncol. Hematol.* 29: 209-248 (1999), the entire teachings of which are incorporated herein by reference. c-Met and HGF are expressed in numerous tissues, although their expression is normally confined predominantly to cells of epithelial and mesenchymal origin, respectively. c-Met and HGF are required for normal mammalian development and have been shown to be important in cell migration, cell proliferation and survival, morphogenic differentiation, and organization of 3-dimensional tubular structures (e.g., renal tubular cells, gland formation, etc.). The c-Met receptor has been shown to be expressed in a number of human cancers. c-Met and its ligand, HGF, have also been shown to be co-expressed at elevated levels in a variety of human cancers (particularly sarcomas). However, because the receptor and ligand are usually expressed by different cell types, c-Met signaling is most commonly regulated by tumor-stroma (tumor-host) interactions. Furthermore, c-Met gene amplification, mutation, and rearrangement have been observed in a subset of human cancers. Families with germline mutations that activate c-Met kinase are prone to multiple kidney tumors as well as tumors in other tissues. Numerous studies have correlated the expression of c-Met and/or HGF/SF with the state of disease progression of different types of cancer (including lung, colon, breast, prostate, liver, pancreas, brain, kidney, ovaries, stomach, skin, and bone cancers). Furthermore, the overexpression of c-Met or HGF have been shown to correlate with poor prognosis and disease outcome in a number of major human cancers including lung, liver, gastric, and breast.

**[0056]** The anaplastic lymphoma kinase (ALK) tyrosine kinase receptor is an enzyme that, in humans, is encoded by the ALK gene. The 2;5 chromosomal translocation is frequently associated with anaplastic large cell lymphomas (ALCLs). The translocation creates a fusion gene consisting of the ALK (anaplastic lymphoma kinase) gene and the nucleophosmin (NPM) gene: the 3' half of ALK, derived from chromosome 2, is fused to the 5' portion of NPM from chromosome 5. The product of the NPM-ALK fusion gene is

oncogenic. Other possible translocations of the ALK gene, such as the em14 translocation, are also implicated in cancer.

**[0057]** The general role of ALK in cancer has been described. See, e.g., Pulford et al., *J. Cell Physiol.* 199(3): 330-358 (2004). Abnormalities in the anaplastic lymphoma kinase (ALK) gene have an established pathogenic role in many pediatric and adult cancers, including non-small cell lung cancer (NSCLC), diffuse large B-cell lymphoma (DLBCL), anaplastic large cell lymphoma (ALCL), neuroblastoma (NBL), and inflammatory myofibroblastic tumors (IMT), non-Hodgkin's lymphoma (NHL), and esophageal squamous cell carcinoma (ESCC). These diseases account for more than 250,000 new cancer diagnoses each year in the United States alone.

**[0058]** More particularly, EML4-ALK and KIF5B-ALK translocations have been found in non-small cell lung cancer. See, e.g. Mano H., *Cancer Sci.* 2008 December;99(12):2349-55; Takeuchi K et al., *Clin Cancer Res.* 2009 May 1; 15(9): 3143-9. CLTC-ALK mutation has been found in DLBCL. See e.g. Rudzki Z et al., *Pol J Pathol.* 2005; 56 (1):37-45. NPM-ALK, MSN-ALK, and other mutations have been found in ALCL. See e.g. Lamant L et al., *Genes Chromosomes Cancer.* 2003 August; 37 (4):427-32; Webb T R et al. *Expert Rev Anticancer Ther* 2009 March; 9(3):331-56. TPM4-ALK mutation has been found in esophageal squamous cell carcinoma (ESCC). See e.g. Li R, Morris S W., *Med Res Rev.* 2008 May; 28 (3):372-412. F1174L, R1275Q, and other point mutations have been found in NBL. See e.g. van Roy N et al. *Genome Med* 2009 July 27; 1 (7):74. TPM3-ALK, TPM4-ALK, CLTC-ALK, RanBP2-ALK, and TPM4-ALK mutations have been found in IMT. See e.g. Gleason B C, Hornick J L. *J Clin Pathol* 2008 April; 61(4):428-37. The methods of detection and identification of these alterations, mutations or rearrangements in an ALK gene or gene product can be found in the above-identified references and references cited therein.

**[0059]** The KRAS oncogene (the cellular homolog of the Kirsten rat sarcoma virus gene) is a critical gene in the development of a variety of cancers, and the mutation status of this gene is an important characteristic of many cancers. Mutation status of the gene can provide diagnostic, prognostic and predictive information for several cancers. The KRAS gene is a member of a family of genes (KRAS, NRAS and HRAS). KRAS is a member of the RAS family of oncogenes, a collection of small guanosine triphosphate (GTP)-binding proteins that integrate extracellular cues and activate intracellular signaling pathways to regulate cell proliferation, differentiation, and survival. Gain-of-function mutations that confer transforming capacity are frequently observed in KRAS, predominantly arising as single amino acid substitutions at amino acid residues G12, G13 or Q61. Constitutive activation of KRAS leads to the persistent stimulation of downstream signaling pathways that promote tumorigenesis, including the RAF/MEK/ERK and PI3K/AKT/mTOR cascades. In NSCLC, KRAS mutations are highly prevalent (20-30%) and are associated with unfavorable clinical outcomes. Mutations in KRAS appear mutually exclusive with those in EGFR in NSCLC tumors; more importantly, they can account for primary resistance to targeted EGFR TKI therapies. Mutations in the KRAS gene are common in many types of cancer, including pancreatic cancer (~65%), colon cancer (~40%), lung cancer (~20%) and ovarian cancer (~15%).

**[0060]** The methods and procedures for the detections and/or identifications of EGFR, KRAS, and/or ALK over-express-

sions and/or mutations are known in the literature and can be easily carried out by a skilled person. See, e.g., U.S. Pat. Nos. 7,700,339; 5,529,925; 5,770,421; U.S. Patent Application Publication No. US2011/0110923; Palmer et al, *Biochem. J.* (2009), 345-361; Koivunen et al, *Clin. Can. Res.*, 2008, 14, 4275-4283; Anderson, *Expert Rev. Mol. Diagn.* 11(6), 635-642 (2011); Pinto et al, *Cancer Genetics* 204 (2011), 439-446; Rekhman et al; *Clin Cancer Res* 2012; 18:1167-1176; Massarelli et al, *Clin Cancer Res* 2007; 13:2890-2896; Lamy et al, *Modern Pathology* (2011) 24, 1090-1100; Balschun et al, *Expert Rev. Mol. Diagn.* 11(8), 799-802 (2011); Vakiani et al, *J Pathol* 2011; 223, 219-229; Okudela et al, *Pathology International* 2010; 60: 651-660; John et al, *Oncogene* (2009) 28, S14-S23; Jimeno et al, *J. Clin. Oncol.* 27, 1130-1135 (2009); *Van Krieken et al, Virchows Archiv.* 453, 417-431 (2008); and the references cited in the-above identified references. Thresholds of increased expression that constitute an EGFR mutation or an ALK mutation are well known in the art. Moreover, it is generally recognized that once an EGFR mutation is detected in a cancer, the KRAS mutation will be eliminated in the same cancer. Put reversely, if a KRAS mutation is positively identified in a cancer from a subject, it is then not necessary to engage in any further EGFR-related identification. Similar principle can be applied to an ALK mutation in a cancer. That is if there is an ALK mutation detected in a cancer, it is extremely rare that an EGFR or KRAS mutation will be implicated. Stated another way, once an ALK mutation is positively identified in a cancer, no further identification is necessary either for EGFR mutation or for KRAS mutation in the same cancer.

**[0061]** As used herein, a “proliferative disorder” or a “hyperproliferative disorder,” and other equivalent terms, means a disease or medical condition involving pathological growth of cells. Proliferative disorders include cancer, smooth muscle cell proliferation, systemic sclerosis, cirrhosis of the liver, adult respiratory distress syndrome, idiopathic cardiomyopathy, lupus erythematosus, retinopathy, (e.g., diabetic retinopathy or other retinopathies), cardiac hyperplasia, reproductive system associated disorders such as benign prostatic hyperplasia and ovarian cysts, pulmonary fibrosis, endometriosis, fibromatosis, hamartomas, lymphangiomatosis, sarcoidosis and desmoid tumors. Non-cancerous proliferative disorders also include hyperproliferation of cells in the skin such as psoriasis and its varied clinical forms, Reiter’s syndrome, pityriasis rubra pilaris, hyperproliferative variants of disorders of keratinization (e.g., actinic keratosis, senile keratosis), scleroderma, and the like. In an embodiment, the proliferative disorder is cancer.

**[0062]** In an embodiment, the invention provides a method of treating a proliferative disorder in a subject, comprising administering to the subject an effective amount of the combination of Hsp90 inhibitor and platinum-containing anti-cancer agent as described herein. In an embodiment, the proliferative disorder is cancer. In an embodiment, the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer. In an embodiment, the cancer is solid cancer, gastric cancer, bladder cancer, ovarian cancer, or colorectal cancer. In an embodiment, the cancer is colon cancer. In an embodiment,

the cancer is metastatic colorectal cancer. In an embodiment, the cancer is bladder cancer. In an embodiment, the cancer is solid cancer. In an embodiment, the cancer is gastric cancer. In an embodiment, the cancer may have a mutation or translocation in EGFR, K-Ras, PI3K, ALK, HER2 and/or B-Raf proteins.

**[0063]** Some of the disclosed methods can be particularly effective at treating subjects whose cancer has become “drug resistant” or “multi-drug resistant”. A cancer which initially responded to an anti-cancer drug becomes resistant to the anti-cancer drug when the anti-cancer drug is no longer effective in treating the subject with the cancer. For example, many tumors will initially respond to treatment with an anti-cancer drug by decreasing in size or even going into remission, only to develop resistance to the drug. “Drug resistant” tumors are characterized by a resumption of their growth and/or reappearance after having seemingly gone into remission, despite the administration of increased dosages of the anti-cancer drug. Cancers that have developed resistance to two or more anti-cancer drugs are said to be “multi-drug resistant”. For example, it is common for cancers to become resistant to three or more anti-cancer agents, often five or more anti-cancer agents and at times ten or more anti-cancer agents.

**[0064]** Other anti-proliferative or anti-cancer therapies may be combined with the pharmaceutical combination of this invention to treat proliferative diseases and cancer. Other therapies or anti-cancer agents that may be used in combination with the inventive anti-cancer agents of the present invention include surgery, radiotherapy (including, but not limited to, gamma-radiation, neutron beam radiotherapy, electron beam radiotherapy, proton therapy, brachytherapy, and systemic radioactive isotopes), endocrine therapy, biologic response modifiers (including, but not limited to, interferons, interleukins, and tumor necrosis factor (TNF)), hyperthermia and cryotherapy, agents to attenuate any adverse effects (e.g., antiemetics), and other approved chemotherapeutic drugs. In an embodiment, the pharmaceutical combination of the invention is administered with one or more therapeutic agent selected from DFMO, vandetanib, trastuzumab, temodar, dexamethasone, epirubicin, ifosfamide, mitoxantrone, vorinostat, interferon alpha, rituximab, prednisone, cyclophosphamide, bendamustine, adriamycin, valproate, celecoxib, thalidomide, nelarabine, methotrexate, filgrastim, gemtuzumab, ozogamicin, testosterone, clofarabine, cytarabine, everolimus, rituxumab, busulfan, capecitabine, pegfilgrastim, mesna, amrubicin, obatoclax, gefitinib, cyclosporine, dasatinib, temozolomide, thiotepa, plerixafor, mitotane, vincristine, doxorubicin, cixutumumab, endostar, fenofibrate, melphalan, sunitinib, rubitecan, enoxaparin, isotretinoin, tariquidar, pomalidomide, sorafenib, altretamine, idarubicin, rapamycin, zevalin, everolimus, pravastatin, carmustine, nelfinavir, streptozocin, tirapazamine, aprepitant, lenalidomide, G-CSF, procarbazine, alemtuzumab, amifostine, valspodar, lomustine, oblimersen, temsirolimus, vinblastine, figitumumab, belinostat, niacinamide, tipifarnib, estramustine, erlotinib, bevacizumab, paclitaxel, docetaxel, Abraxane®, pemetrexed, bortezomib, cetuximab, gemcitabine, 5-fluorouracil, leucovorin and tetracycline. In an embodiment, the one or more therapeutic agent is selected from erlotinib, bevacizumab, bortezomib, paclitaxel, doxorubicin, docetaxel, mitoxantrone, cytarabine, 5-fluorouracil, leucovorin, pemetrexed and vincristine.

**[0065]** As used herein, the term “pharmaceutically acceptable salt” refers to a salt prepared from a compound of for-

mulae (I) or (Ia) or a compound in Tables 1 or 2 having an acidic functional group, such as a carboxylic acid functional group, and a pharmaceutically acceptable inorganic or organic base. Suitable bases include hydroxides of alkali metals such as sodium, potassium, and lithium; hydroxides of alkaline earth metal such as calcium and magnesium; hydroxides of other metals, such as aluminum and zinc; ammonia, and organic amines, such as unsubstituted or hydroxy-substituted mono-, di-, or trialkylamines; dicyclohexylamine; tributyl amine; pyridine; N-methyl,N-ethylamine; diethylamine; triethylamine; mono-, bis-, or tris-(2-hydroxy-lower alkyl amines), such as mono-, bis-, or tris-(2-hydroxyethyl)amine, 2-hydroxy-tert-butylamine, or tris-(hydroxymethyl)methylamine, N,N-di-lower alkyl-N-(hydroxy lower alkyl)-amines, such as N,N-dimethyl-N-(2-hydroxyethyl)amine, or tri-(2-hydroxyethyl)amine; N-methyl-D-glucamine; and amino acids such as arginine, lysine, and the like. The term “pharmaceutically acceptable salt” also refers to a salt prepared from a compound of formulae (I) or (Ia) or a compound in Tables 1 or 2 having a basic functional group, such as an amine functional group, and a pharmaceutically acceptable inorganic or organic acid. Suitable acids include hydrogen sulfate, citric acid, acetic acid, oxalic acid, hydrochloric acid (HCl), hydrogen bromide (HBr), hydrogen iodide (HI), nitric acid, hydrogen bisulfide, phosphoric acid, isonicotinic acid, oleic acid, tannic acid, pantothenic acid, saccharic acid, lactic acid, salicylic acid, tartaric acid, bitartronic acid, ascorbic acid, succinic acid, maleic acid, besylic acid, fumaric acid, gluconic acid, glucaronic acid, formic acid, benzoic acid, glutamic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, pantoic acid and p-toluenesulfonic acid.

**[0066]** As used herein, the term “pharmaceutically acceptable solvate,” is a solvate formed from the association of one or more pharmaceutically acceptable solvent molecules to one of the compounds of formulae (I) or (Ia) or a compound in Tables 1 or 2. The term “solvate” includes hydrates, e.g., hemihydrate, monohydrate, dihydrate, trihydrate, tetrahydrate, and the like.

**[0067]** A pharmaceutically acceptable carrier may contain inert ingredients which do not unduly inhibit the biological activity of the compound(s) described herein. The pharmaceutically acceptable carriers should be biocompatible, i.e., non-toxic, non-inflammatory, non-immunogenic and devoid of other undesired reactions upon the administration to a subject. Standard pharmaceutical formulation techniques can be employed, such as those described in REMINGTON, J. P., REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., 17th ed., 1985). Suitable pharmaceutical carriers for parenteral administration include, for example, sterile water, physiological saline, bacteriostatic saline (saline containing about 0.9% mg/ml benzyl alcohol), phosphate-buffered saline, Hank's solution, Ringer's-lactate, and the like. Methods for encapsulating compositions, such as in a coating of hard gelatin or cyclodextran, are known in the art. See BAKER, ET AL., CONTROLLED RELEASE OF BIOLOGICAL ACTIVE AGENTS, (John Wiley and Sons, 1986).

**[0068]** As used herein, the term “effective amount” refers to an amount of a compound described herein which is sufficient to reduce or ameliorate the severity, duration, progression, or onset of a disease or disorder, delay onset of a disease or disorder, retard or halt the advancement of a disease or disorder, cause the regression of a disease or disorder, prevent or delay the recurrence, development, onset or progression of a

symptom associated with a disease or disorder, or enhance or improve the therapeutic effect(s) of another therapy. In an embodiment of the invention, the disease or disorder is a proliferative disorder. The precise amount of compound administered to a subject will depend on the mode of administration, the type and severity of the disease or condition and on the characteristics of the subject, such as general health, age, sex, body weight and tolerance to drugs. For example, for a proliferative disease or disorder, determination of an effective amount will also depend on the degree, severity and type of cell proliferation. The skilled artisan will be able to determine appropriate dosages depending on these and other factors. When co-administered with other therapeutic agents, e.g., when co-administered with an anti-cancer agent, an “effective amount” of any additional therapeutic agent(s) will depend on the type of drug used. Suitable dosages are known for approved therapeutic agents and can be adjusted by the skilled artisan according to the condition of the subject, the type of condition(s) being treated and the amount of a compound of the invention being used. In cases where no amount is expressly noted, an effective amount should be assumed. Non-limiting examples of an effective amount of a compound described herein are provided herein below. In a specific embodiment, the invention provides a method of treating, managing, or ameliorating a disease or disorder, e.g. a proliferative disorder, or one or more symptoms thereof, the method comprising administering to a subject in need thereof a dose of the Hsp90 inhibitor at least 150  $\mu\text{g}/\text{kg}$ , at least 250  $\mu\text{g}/\text{kg}$ , at least 500  $\mu\text{g}/\text{kg}$ , at least 1 mg/kg, at least 5 mg/kg, at least 10 mg/kg, at least 25 mg/kg, at least 50 mg/kg, at least 75 mg/kg, at least 100 mg/kg, at least 125 mg/kg, at least 150 mg/kg, or at least 200 mg/kg or more of one or more compounds described herein once every day, once every 2 days, once every 3 days, once every 4 days, once every 5 days, once every 6 days, once every 7 days, once every 8 days, once every 10 days, once every two weeks, once every three weeks, or once a month.

**[0069]** The dosage of the individual platinum-containing anti-cancer agent used in the pharmaceutical combination may be equal to or lower than the dose of an individual therapeutic agent when given independently to treat, manage, or ameliorate a disease or disorder, or one or more symptoms thereof. In an embodiment of the invention, the disease or disorder being treated with a combination therapy is a proliferative disorder. In an embodiment, the proliferative disorder is cancer. In an embodiment, the platinum-containing anti-cancer agent cisplatin is administered IV at a dose of between about 20  $\text{mg}/\text{m}^2$  to about 150  $\text{mg}/\text{m}^2$ . In an embodiment, cisplatin is administered once daily. In an embodiment, cisplatin is administered IV at 20  $\text{mg}/\text{m}^2$  once daily. In an embodiment, cisplatin is administered IV at a dose of between about 75  $\text{mg}/\text{m}^2$  to about 100  $\text{mg}/\text{m}^2$ . A treatment cycle can last between one and 6 weeks. The recommended dosages of therapeutic agents currently used for the treatment, management, or amelioration of a disease or disorder, or one or more symptoms thereof, can be obtained from any reference in the art. For a more in depth review of dosage and treatment schedules for various disorders, see, e.g., GOODMAN & GILMAN'S THE PHARMACOLOGICAL BASIS OF THERAPEUTICS 9<sup>TH</sup> ED, (Hardman, et al., Eds., NY:Mc-Graw-Hill (1996)); PHYSICIAN'S DESK REFERENCE 57<sup>TH</sup> ED. (Medical Economics Co., Inc., Montvale, N.J. (2003)).

**[0070]** As used herein, the terms “treat”, “treatment”, “inhibiting”, and “treating” refer to the reduction or amelio-

ration of the progression, severity and/or duration of a disease or disorder, delay of the onset of a disease or disorder, or the amelioration of one or more symptoms (preferably, one or more discernible symptoms) of a disease or disorder, resulting from the administration of one or more therapies (e.g., one or more therapeutic agents such as a compound of the invention). The terms “treat”, “treatment” and “treating” also encompass the reduction of the risk of developing a disease or disorder, and the delay or inhibition of the recurrence of a disease or disorder. In an embodiment, the disease or disorder being treated is a proliferative disorder such as cancer. In specific embodiments, the terms “treat”, “treatment” and “treating” refer to the amelioration of at least one measurable physical parameter of a disease or disorder, such as growth of a tumor, not necessarily discernible by the patient. In other embodiments the terms “treat”, “treatment” and “treating” refer to the inhibition of the progression of a disease or disorder, e.g., a proliferative disorder, either physically by the stabilization of a discernible symptom, physiologically by the stabilization of a physical parameter, or both. In another embodiment, the terms “treat”, “treatment” and “treating” of a proliferative disease or disorder refers to the reduction or stabilization of tumor size or cancerous cell count, and/or delay of tumor formation. In another embodiment, the terms “treat”, “treating” and “treatment” also encompass the administration of a compound described herein as a prophylactic measure to patients with a predisposition (genetic or environmental) to any disease or disorder described herein.

**[0071]** As used herein, the terms “therapeutic agent” and “therapeutic agents” refer to any agent(s) that can be used in the treatment of a disease or disorder, e.g. a proliferative disorder, or one or more symptoms thereof. In certain embodiments, the term “therapeutic agent” refers to a compound described herein. In certain other embodiments, the term “therapeutic agent” does not refer to a compound described herein. Preferably, a therapeutic agent is an agent that is known to be useful for, or has been or is currently being used for the treatment of a disease or disorder, e.g., a proliferative disorder, or one or more symptoms thereof.

**[0072]** As used herein, the term “synergistic” refers to a combination of a compound described herein and another therapeutic agent, which, when taken together, is more effective than the additive effects of the individual therapies. A synergistic effect of a combination of therapies (e.g., a combination of therapeutic agents) permits the use of lower dosages of one or more of the therapeutic agent(s) and/or less frequent administration of the agent(s) to a subject with a disease or disorder, e.g., a proliferative disorder. The ability to utilize lower the dosage of one or more therapeutic agent and/or to administer the therapeutic agent less frequently reduces the toxicity associated with the administration of the agent to a subject without reducing the efficacy of the therapy in the treatment of a disease or disorder. In addition, a synergistic effect can result in improved efficacy of agents in the prevention, management or treatment of a disease or disorder, e.g. a proliferative disorder. Finally, a synergistic effect of a combination of therapies may avoid or reduce adverse or unwanted side effects associated with the use of either therapeutic agent alone.

**[0073]** As used herein, the phrase “side effects” encompasses unwanted and adverse effects of a therapeutic agent. Side effects are always unwanted, but unwanted effects are not necessarily adverse. An adverse effect from a therapeutic agent might be harmful or uncomfortable or risky to a subject.

Side effects include fever, chills, lethargy, gastrointestinal toxicities (including gastric and intestinal ulcerations and erosions), nausea, vomiting, neurotoxicities, nephrotoxicities, renal toxicities (including such conditions as papillary necrosis and chronic interstitial nephritis), hepatic toxicities (including elevated serum liver enzyme levels), myelotoxicities (including leukopenia, myelosuppression, thrombocytopenia and anemia), dry mouth, metallic taste, prolongation of gestation, weakness, somnolence, pain (including muscle pain, bone pain and headache), hair loss, asthenia, dizziness, extra-pyramidal symptoms, akathisia, cardiovascular disturbances and sexual dysfunction.

**[0074]** As used herein, the term “in combination” refers to the use of more than one therapeutic agent. The use of the term “in combination” does not restrict the order in which the therapeutic agents are administered to a subject with a disease or disorder, e.g., a proliferative disorder. A first therapeutic agent, such as a compound described herein, can be administered prior to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapeutic agent, such as an anti-cancer agent, to a subject with a disease or disorder, e.g. a proliferative disorder, such as cancer. In an embodiment, the Hsp90 inhibitor and the platinum-containing anti-cancer agent are dosed on independent schedules. In another embodiment, the Hsp90 inhibitor and the platinum-containing anti-cancer agent are dosed on approximately the same schedule. In another embodiment, the Hsp90 inhibitor and the platinum-containing anticancer agent are dosed concurrently or sequentially on the same day.

**[0075]** As used herein, the terms “therapies” and “therapy” can refer to any protocol(s), method(s), and/or agent(s) that can be used in the prevention, treatment, management, or amelioration of a disease or disorder, e.g., a proliferative disorder, or one or more symptoms thereof.

**[0076]** As used herein, a “protocol” includes dosing schedules and dosing regimens. The protocols herein are methods of use and include therapeutic protocols.

**[0077]** As used herein, a composition that “substantially” comprises a compound means that the composition contains more than about 80% by weight, more preferably more than about 90% by weight, even more preferably more than about 95% by weight, and most preferably more than about 97% by weight of the compound.

**[0078]** As used herein, a “racemic mixture” means about 50% of one enantiomer and about 50% of its corresponding enantiomer of the molecule. The combination encompasses all enantiomerically-pure, enantiomerically-enriched, diastereomerically pure, diastereomerically enriched, and racemic mixtures of the compounds described herein. Enantiomeric and diastereomeric mixtures can be resolved into their component enantiomers or diastereomers by well-known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Enantiomers and diastereomers can also be obtained from diastereomerically- or enantiomerically-

cally-pure intermediates, reagents, and catalysts by well-known asymmetric synthetic methods.


**[0079]** The compounds described herein are defined by their chemical structures and/or chemical names. Where a compound is referred to by both a chemical structure and a chemical name, and the chemical structure and the chemical name conflict, the chemical structure is determinative of the compound's identity.

**[0080]** When administered to a subject (e.g., a non-human animal for veterinary use or for improvement of livestock or to a human for clinical use), the compounds described herein are administered in an isolated form, or as the isolated form in a pharmaceutical composition. As used herein, "isolated" means that the compounds described herein are separated from other components of either: (a) a natural source, such as a plant or cell, preferably bacterial culture, or (b) a synthetic organic chemical reaction mixture. Preferably, the compounds described herein are purified via conventional techniques. As used herein, "purified" means that when isolated, the isolate contains at least 95%, preferably at least 98%, of a compound described herein by weight of the isolate either as a mixture of stereoisomers, or as a diastereomerically or enantiomerically pure isolate.

**[0081]** Only those choices and combinations of substituents that result in a stable structure are contemplated. Such choices and combinations will be apparent to those of ordinary skill in the art and may be determined without undue experimentation.

**[0082]** The invention can be understood more fully by reference to the following detailed description and illustrative examples, which are intended to exemplify non-limiting embodiments of the invention.

**[0083]** The methods described herein utilize triazolone compounds listed in Tables 1 or 2, or a compound represented by Formulae (I) or (Ia):



**[0084]** or a tautomer, or a pharmaceutically acceptable salt thereof, wherein:

**[0085]** Z is OH, SH, or NH<sub>2</sub>;

**[0086]** X is CR<sub>4</sub> or N;

**[0087]** R<sub>1</sub> is —H, —OH, —SH, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted

cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro, guanidino, a haloalkyl, a heteroalkyl, an alkoxy or cycloalkoxy, a haloalkoxy, —NR<sub>10</sub>R<sub>11</sub>, —OR<sub>7</sub>, —C(O)R<sub>7</sub>, —C(O)OR<sub>7</sub>, —C(S)R<sub>7</sub>, —C(O)SR<sub>7</sub>, —C(S)SR<sub>7</sub>, —C(S)OR<sub>7</sub>, —C(S)NR<sub>10</sub>R<sub>11</sub>, —C(NR<sub>8</sub>)OR<sub>7</sub>, —C(NR<sub>8</sub>)R<sub>7</sub>, —C(NR<sub>8</sub>)NR<sub>10</sub>R<sub>11</sub>, —C(NR<sub>8</sub>)SR<sub>7</sub>, —OC(O)R<sub>7</sub>, —OC(O)OR<sub>7</sub>, —OC(S)OR<sub>7</sub>, —OC(NRS)OR<sub>7</sub>, —SC(O)R<sub>7</sub>, —SC(O)OR<sub>7</sub>, —SC(NR<sub>8</sub>)OR<sub>7</sub>, —OC(S)R<sub>7</sub>, —SC(S)R<sub>7</sub>, —SC(S)OR<sub>7</sub>, —OC(O)NR<sub>10</sub>R<sub>11</sub>, —OC(S)NR<sub>10</sub>R<sub>11</sub>, —OC(NR<sub>8</sub>)NR<sub>10</sub>R<sub>11</sub>, —SC(O)NR<sub>10</sub>R<sub>11</sub>, —SC(NR<sub>8</sub>)NR<sub>10</sub>R<sub>11</sub>, —SC(S)NR<sub>10</sub>R<sub>11</sub>, —OC(NR<sub>8</sub>)R<sub>7</sub>, —SC(NR<sub>8</sub>)R<sub>7</sub>, —C(O)NR<sub>10</sub>R<sub>11</sub>, —NR<sub>8</sub>C(O)R<sub>7</sub>, —NR<sub>2</sub>C(S)R<sub>7</sub>, —NR<sub>7</sub>C(S)OR<sub>7</sub>, —NR<sub>7</sub>C(NR<sub>8</sub>)R<sub>7</sub>, —NR<sub>7</sub>C(O)OR<sub>7</sub>, —NR<sub>7</sub>C(NR<sub>8</sub>)OR<sub>7</sub>, —NR<sub>7</sub>C(O)NR<sub>10</sub>R<sub>11</sub>, —NR<sub>7</sub>C(S)NR<sub>10</sub>R<sub>11</sub>, —NR<sub>7</sub>C(NR<sub>8</sub>)NR<sub>10</sub>R<sub>11</sub>, —SR<sub>7</sub>, —S(O)<sub>p</sub>R<sub>7</sub>, —OS(O)<sub>p</sub>R<sub>7</sub>, —OS(O)<sub>p</sub>OR<sub>7</sub>, —OS(O)<sub>p</sub>NR<sub>10</sub>R<sub>11</sub>, —S(O)<sub>p</sub>OR<sub>7</sub>, —NR<sub>8</sub>S(O)<sub>p</sub>R<sub>7</sub>, —NR<sub>7</sub>S(O)<sub>p</sub>NR<sub>10</sub>R<sub>11</sub>, —NR<sub>7</sub>S(O)<sub>p</sub>OR<sub>7</sub>, —S(O)<sub>p</sub>NR<sub>10</sub>R<sub>11</sub>, —SS(O)<sub>p</sub>R<sub>7</sub>, —SS(O)<sub>p</sub>OR<sub>7</sub>, —SS(O)<sub>p</sub>NR<sub>10</sub>R<sub>11</sub>, —OP(O)(OR<sub>7</sub>)<sub>2</sub>, or —SP(O)(OR<sub>7</sub>)<sub>2</sub>;

**[0088]** R<sub>2</sub> is —H, —OH, —SH, —NR<sub>7</sub>H, —OR<sub>15</sub>, —SR<sub>15</sub>, —NHR<sub>15</sub>, —O(CH<sub>2</sub>)<sub>m</sub>OH, —O(CH<sub>2</sub>)<sub>m</sub>SH, —O(CH<sub>2</sub>)<sub>m</sub>NR<sub>7</sub>H, —S(CH<sub>2</sub>)<sub>m</sub>OH, —S(CH<sub>2</sub>)<sub>m</sub>H, —S(CH<sub>2</sub>)<sub>m</sub>NR<sub>7</sub>H, —OC(O)NR<sub>10</sub>R<sub>11</sub>, —C(O)NR<sub>10</sub>R<sub>11</sub>, —NR<sub>7</sub>C(O)NR<sub>10</sub>R<sub>11</sub>, —OC(O)R<sub>7</sub>, —SC(O)R<sub>7</sub>, —NR<sub>7</sub>C(O)R<sub>7</sub>, —OC(O)OR<sub>7</sub>, —SC(O)OR<sub>7</sub>, —NR<sub>7</sub>C(O)OR<sub>7</sub>, —OCH<sub>2</sub>C(O)R<sub>7</sub>, —SCH<sub>2</sub>C(O)R<sub>7</sub>, —NR<sub>7</sub>CH<sub>2</sub>C(O)R<sub>7</sub>, —OCH<sub>2</sub>C(O)OR<sub>7</sub>, —SCH<sub>2</sub>C(O)OR<sub>7</sub>, —NR<sub>7</sub>CH<sub>2</sub>C(O)OR<sub>7</sub>, —OCH<sub>2</sub>C(O)NR<sub>10</sub>R<sub>11</sub>, —SCH<sub>2</sub>C(O)NR<sub>10</sub>R<sub>11</sub>, —NR<sub>7</sub>CH<sub>2</sub>C(O)NR<sub>10</sub>R<sub>11</sub>, —OS(O)<sub>p</sub>R<sub>7</sub>, —SS(O)<sub>p</sub>R<sub>7</sub>, —NR<sub>7</sub>S(O)<sub>p</sub>R<sub>7</sub>, —OS(O)<sub>p</sub>NR<sub>10</sub>R<sub>11</sub>, —SS(O)<sub>p</sub>NR<sub>10</sub>R<sub>11</sub>, —NR<sub>7</sub>S(O)<sub>p</sub>NR<sub>10</sub>R<sub>11</sub>, —OS(O)<sub>p</sub>OR<sub>7</sub>, —SS(O)<sub>p</sub>OR<sub>7</sub>, —NR<sub>7</sub>S(O)<sub>p</sub>OR<sub>7</sub>, —OC(S)R<sub>7</sub>, —SC(S)R<sub>7</sub>, —NR<sub>7</sub>C(S)R<sub>7</sub>, —OC(S)OR<sub>7</sub>, —SC(S)OR<sub>7</sub>, —NR<sub>7</sub>C(S)OR<sub>7</sub>, —OC(S)NR<sub>10</sub>R<sub>11</sub>, —SC(S)NR<sub>10</sub>R<sub>11</sub>, —NR<sub>7</sub>C(S)NR<sub>10</sub>R<sub>11</sub>, —OC(NR<sub>8</sub>)R<sub>7</sub>, —SC(NR<sub>8</sub>)R<sub>7</sub>, —NR<sub>7</sub>C(NR<sub>8</sub>)R<sub>7</sub>, —OC(NR<sub>8</sub>)OR<sub>7</sub>, —SC(NR<sub>8</sub>)OR<sub>7</sub>, —NR<sub>7</sub>C(NR<sub>8</sub>)OR<sub>7</sub>, —OC(NR<sub>8</sub>)NR<sub>10</sub>R<sub>11</sub>, —SC(NR<sub>8</sub>)NR<sub>10</sub>R<sub>11</sub>, or —NR<sub>7</sub>C(NR<sub>8</sub>)NR<sub>10</sub>R<sub>11</sub>;

**[0089]** R<sub>3</sub> is —H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, a haloalkyl, a heteroalkyl, —C(O)R<sub>7</sub>, —(CH<sub>2</sub>)<sub>m</sub>C(O)OR<sub>7</sub>, —C(O)OR<sub>7</sub>, —OC(O)R<sub>7</sub>, —C(O)NR<sub>10</sub>R<sub>11</sub>, —S(O)<sub>p</sub>R<sub>7</sub>, —S(O)<sub>p</sub>OR<sub>7</sub>, or —S(O)<sub>p</sub>NR<sub>10</sub>R<sub>11</sub>;

**[0090]** R<sub>4</sub> is —H, —OH, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, halo, cyano, nitro, guanidino, a haloalkyl, a heteroalkyl, —C(O)R<sub>7</sub>, —C(O)OR<sub>7</sub>, —OC(O)R<sub>7</sub>, —C(O)NR<sub>10</sub>R<sub>11</sub>, —NR<sub>8</sub>C(O)R<sub>7</sub>, —SR<sub>7</sub>,

- S(O)<sub>p</sub>R<sub>7</sub>, —OS(O)<sub>p</sub>R<sub>7</sub>, —S(O)<sub>p</sub>OR<sub>7</sub>, —NR<sub>8</sub>S(O)<sub>p</sub>R<sub>7</sub>, —S(O)<sub>p</sub>NR<sub>10</sub>R<sub>11</sub>, or R<sub>3</sub> and R<sub>4</sub> taken together with the carbon atoms to which they are attached form an optionally substituted cycloalkenyl, an optionally substituted aryl, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;
- [0091]** R<sub>7</sub> and R<sub>8</sub>, for each occurrence, are independently, —H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkenyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;
- [0092]** R<sub>10</sub> and R<sub>11</sub>, for each occurrence, are independently —H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R<sub>10</sub> and R<sub>11</sub>, taken together with the nitrogen to which they are attached, form an optionally substituted heterocyclyl or an optionally substituted heteroaryl;
- [0093]** R<sub>15</sub>, for each occurrence, is independently, a lower alkyl;
- [0094]** p, for each occurrence, is, independently, 1 or 2; and
- [0095]** m, for each occurrence, is independently, 1, 2, 3, or 4.
- [0096]** In an embodiment, in formula (I) or (Ia), X is CR<sub>4</sub>. In another embodiment, in formula (I) or (Ia), X is N.
- [0097]** In another embodiment, in formula (I) or (Ia), R<sub>1</sub> may be —H, lower alkyl, lower alkoxy, lower cycloalkyl, or lower cycloalkoxy.
- [0098]** In another embodiment, in formula (I) or (Ia), R<sub>1</sub> may be —H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, or cyclopropoxy.
- [0099]** In another embodiment, in formula (I) or (Ia), R<sub>3</sub> may be —H, a lower alkyl, a lower cycloalkyl, —C(O)N(R<sub>27</sub>)<sub>2</sub>, or —C(O)OH, wherein R<sub>27</sub> is —H or a lower alkyl.
- [0100]** In another embodiment, in formula (I) or (Ia), R<sub>3</sub> may be —H, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, sec-butyl, tert-butyl, n-pentyl, n-hexyl, —C(O)OH, —(CH<sub>2</sub>)<sub>m</sub>C(O)OH, —CH<sub>2</sub>OCH<sub>3</sub>, —CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, or —C(O)N(CH<sub>3</sub>)<sub>2</sub>.
- [0101]** In another embodiment, R<sub>4</sub> may be H or a lower alkyl. In another embodiment, in formula (I) or (Ia), R<sub>4</sub> may be —H, methyl, ethyl, propyl, isopropyl or cyclopropyl.
- [0102]** In another embodiment, in formula (I) or (Ia), R<sub>1</sub> may be —H, —OH, —SH, —NH<sub>2</sub>, a lower alkoxy or a lower alkyl amino. In another embodiment, in formula (I) or (Ia), R<sub>1</sub> may be —H, —OH, methoxy or ethoxy.
- [0103]** In another embodiment, in formula (I) or (Ia), Z is —OH.
- [0104]** In another embodiment, in formula (I) or (Ia), Z is —SH.
- [0105]** In another embodiment, in formula (I) or (Ia), R<sub>2</sub> may be —H, —OH, —SH, —NH<sub>2</sub>, a lower alkoxy or a lower alkyl amino.
- [0106]** In another embodiment, in formula (I) or (Ia), R<sub>2</sub> may be —H, —OH, methoxy, or ethoxy.
- [0107]** In another embodiment, in formula (I) or (Ia), R<sub>1</sub> may be —H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, or cyclopropoxy; R<sub>3</sub> may be —H, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, sec-butyl, tert-butyl, n-pentyl, n-hexyl, —C(O)OH, —(CH<sub>2</sub>)<sub>m</sub>C(O)OH, —CH<sub>2</sub>OCH<sub>3</sub>, —CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, or —C(O)N(CH<sub>3</sub>)<sub>2</sub>; R<sub>4</sub> may be —H, methyl, ethyl, propyl, isopropyl or cyclopropyl; R<sub>2</sub> may be —H, —OH, —SH, —NH<sub>2</sub>, a lower alkoxy or a lower alkyl amino; and Z is OH.
- [0108]** In another embodiment, in formula (I) or (Ia), R<sub>1</sub> may be —H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, or cyclopropoxy; R<sub>3</sub> may be of —H, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, sec-butyl, tert-butyl, n-pentyl, n-hexyl, —C(O)OH, —(CH<sub>2</sub>)<sub>m</sub>C(O)OH, —CH<sub>2</sub>OCH<sub>3</sub>, —CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, or —C(O)N(CH<sub>3</sub>)<sub>2</sub>; R<sub>4</sub> may be —H, methyl, ethyl, propyl, isopropyl or cyclopropyl; R<sub>2</sub> may be of —H, —OH, —SH, —NH<sub>2</sub>, a lower alkoxy or a lower alkyl amino; and Z is SH.
- [0109]** In another embodiment, the compound may be:
- [0110]** 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole,
- [0111]** 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole,
- [0112]** 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole,
- [0113]** 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-hydroxy-[1,2,4]triazole,
- [0114]** 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indazol-5-yl)-5-mercapto-[1,2,4]triazole,
- [0115]** 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indazol-6-yl)-5-mercapto-[1,2,4]triazole,
- [0116]** 3-(2,4-dihydroxyphenyl)-4-(1-ethyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0117]** 3-(2,4-dihydroxyphenyl)-4-(1-isopropyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0118]** 3-(2,4-dihydroxyphenyl)-4-(indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0119]** 3-(2,4-dihydroxyphenyl)-4-(1-methoxyethyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0120]** 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0121]** 3-(2,4-dihydroxyphenyl)-4-(1-dimethylcarbamoyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0122]** 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-propyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0123]** 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,2,3-trimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- [0124]** 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- [0125]** 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-acetyl-2,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- [0126]** 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-propyl-2,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- [0127]** 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-n-butyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0128]** 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-n-pentyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0129]** 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-n-hexyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0130]** 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-(1-methylcyclopropyl)-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- [0131]** 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1,2,3-trimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,

- [0132] 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-methyl-3-ethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0133] 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0134] 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-methyl-3-isopropyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0135] 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,2-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0136] 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(N-methyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0137] 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0138] 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0139] 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0140] 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1H-indol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0141] 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,2-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0142] 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-ethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole, or  
 [0143] 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-propyl-indol-5-yl)-5-mercapto-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof.  
 [0144] In another embodiment, the compound may be:  
 [0145] 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-benzimidazol-4-yl)-5-mercapto-[1,2,4]triazole,  
 [0146] 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-benzimidazol-4-yl)-5-mercapto-[1,2,4]triazole HCL salt,  
 [0147] 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2-methyl-3-ethyl-benzimidazol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0148] 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-2-methyl-benzimidazol-5-yl)-5-mercapto-[1,2,4]triazole, or  
 [0149] 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-2-trifluoromethyl-benzimidazol-5-yl)-5-mercapto-[1,2,4]triazole,  
 [0150] or a tautomer, or a pharmaceutically acceptable salt thereof.  
 [0151] In another embodiment, the compound may be:  
 [0152] 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate,  
 [0153] sodium 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl phosphate,  
 [0154] 2-(3,4-dimethoxyphenethyl)-5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)phenyl dihydrogen phosphate,  
 [0155] 5-hydroxy-2-isopropyl-4-(5-mercapto-4-(4-methoxybenzyl)-4H-1,2,4-triazol-3-yl)phenyl dihydrogen phosphate,  
 [0156] 5-hydroxy-4-(5-hydroxy-4-(4-methoxybenzyl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or  
 [0157] 4-(4-(1,3-dimethyl-1H-indol-5-yl)-5-hydroxy-4H-1,2,4-triazol-3-yl)-2-ethyl-5-hydroxyphenyl dihydrogen phosphate,  
 [0158] or a tautomer, or a pharmaceutically acceptable salt thereof.  
 [0159] Hsp90 inhibitory compounds, as well as tautomers or pharmaceutically acceptable salts thereof that may be used in the methods described herein are depicted in Tables 1 or 2.

TABLE 1

	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
1			3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(1-METHYL-INDOL-5-YL)-5-HYDROXY-[1,2,4]TRIAZOLE (GANETESPIB)
2			3-(2,4-DIHYDROXY-PHENYL)-4-(1-ETHYL-INDOL-4-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE



TABLE 1-continued

	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
3			3-(2,4-DIHYDROXY-PHENYL)-4-(2,3-DIMETHYL-1H-INDOL-4-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
4			3-(2,4-DIHYDROXYPHENYL)-4-(1-ISOPROPYL-INDOL-4-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
5			3-(2,4-DIHYDROXY-PHENYL)-4-(INDOL-4-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
6			3-(2,4-DIHYDROXY-PHENYL)-4-[1-(2-METHOXYETHOXY)-INDOL-4-YL]-5-MERCAPTO-[1,2,4] TRIAZOLE
7			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ISOPROPYL-INDOL-4-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE

TABLE 1-continued

	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
8			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-[1-(DIMETHYL-CARBAMOYL)-INDOL-4-YL]-5-MERCAPTO-[1,2,4] TRIAZOLE
9			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ETHYL-BENZIMIDAZOL-4-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
10			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1,2,3-TRIMETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
11			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ISOPROPYL-INDOL-3-YL)-5-HYDROXY-[1,2,4] TRIAZOLE

TABLE 1-continued

	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
12			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ISOPROPYL-INDOL-4-yl)-5-AMINO-[1,2,4] TRIAZOLE
15			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ISOPROPYL-INDOL-4-yl)-5-UREIDO-[1,2,4] TRIAZOLE
16			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-METHYL-INDOL-4-yl)-5-CARBAMOYLOXY-[1,2,4] TRIAZOLE
17			3-(2,4-DIHYDROXY-PHENYL)-4-(1-METHYL-2-CHLORO-INDOL-4-yl)-5-CARBAMOYLOXY-[1,2,4] TRIAZOLE
18			3-(2,4-DIHYDROXY-5-METHOXY-PHENYL)-4-(1-ISOPROPYL-BENZIMIDAZOL-4-yl)-5-(SULFAMOYLAMINO)-[1,2,4] TRIAZOLE

TABLE 1-continued

	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
20			3-(2,4-DIHYDROXY-5-METHOXY-PHENYL)-4-(1-ISOPROPYL-BENZOIMIDAZOL-4-yl)-5-(SULFAMOYLOXY)-[1,2,4]TRIAZOLE
21			3-(2-HYDROXY-4-ETHOXYCARBOXYOXY-5-METHOXY-PHENYL)-4-(1-ISOPROPYL-BENZOIMIDAZOL-4-YL)-5-HYDROXY-[1,2,4]TRIAZOLE
22			3-[2-HYDROXY-4-ISOBUTYRYLOXY-5-ETHYL-PHENYL]-4-(1-METHYL-BENZOIMIDAZOL-4-YL)-5-HYDROXY-[1,2,4]TRIAZOLE
23			3-(2,4-DIHYDROXY-PHENYL)-4-(1-DIMETHYLCARBAMOYL-INDOL-4-yl)-5-MERCAPTO-[1,2,4]TRIAZOLE

TABLE 1-continued

24	STRUCTURE	TAUTOMERIC STRUCTURE	NAME	
			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(2,3-DIMETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE	
	25			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ETHYL-1H-BENZOIMIDAZOL-4-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE, HCL SALT
	26			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ISOPROPYL-7-METHOXY-INDOL-4-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE
	27			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-PROPYL-INDOL-4-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE

TABLE 1-continued

	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
28			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ACETYL-2,3-DIMETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE
29			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(2-METHYL-3-ETHYL-BENZIMIDAZOL-5-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE
30			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ETHYL-2-METHYL-BENZIMIDAZOL-5-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE
31			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-PROPYL-2,3-DIMETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE

TABLE 1-continued









STRUCTURE	TAUTOMERIC STRUCTURE	NAME
<p>34</p> 		<p>3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-N-BUTYL-INDOL-4-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE</p>
<p>35</p> 		<p>3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-N-PENTYL-INDOL-4-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE</p>
<p>36</p> 		<p>3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-N-HEXYL-INDOL-4-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE</p>
<p>37</p> 		<p>3-(2,4-DIHYDROXY-5-CYCLOPROPYL-PHENYL)-4-(1-(1-METHYLCYCLOPROPYL)-INDOL-4-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE</p>

TABLE 1-continued

	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
38			3-(2,4-DIHYDROXY-5-CYCLOPROPYL-PHENYL)-4-(1-ISOPROPYL-7-METHOXY-INDOL-4-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
39			3-(2,4-DIHYDROXY-5-CYCLOPROPYL-PHENYL)-4-(1,2,3-TRIMETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
40			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ISOPROPYL-7-METHOXY-INDOL-4-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE DISODIUM SALT
41			3-(2,4-DIHYDROXY-5-TERT-BUTYL-PHENYL)-4-(1-ISOPROPYL-7-METHOXY-INDOL-4-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE



TABLE 1-continued











	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
42			3-(2,4-DIHYDROXY-5-CYCLOPROPYL-PHENYL)-4-(1-PROPYL-7-METHOXY-INDOL-4-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE
43			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-METHYL-3-ETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE
44			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1,3-DIMETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE
45			3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(1-ISOPROPYL-7-METHOXY-INDOL-4-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE
46			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-METHYL-3-ISOPROPYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE

TABLE 1-continued









	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
48			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ISOPROPYL-7-HYDROXY-INDOL-4-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
49			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1-ISOPROPYL-7-ETHOXY-INDOL-4-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
50			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1,2-DIMETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
51			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(N-METHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE

TABLE 1-continued











	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
55			3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(1,3-DIMETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
56			3-(2,4-DIHYDROXY-5-CYCLOPROPYL-PHENYL)-4-(1,3-DIMETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
57			3-(2,4-DIHYDROXY-5-ETHYL-PHENYL)-4-(1,3-DIMETHYL-INDOL-5-YL)-5-HYDROXY-[1,2,4] TRIAZOLE
58			3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(N-METHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE
59			3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(1,2-DIMETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE

TABLE 1-continued

STRUCTURE	TAUTOMERIC STRUCTURE	NAME
<p>60</p>		<p>3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(1,3-DIMETHYL-INDOL-5-YL)-5-HYDROXY-[1,2,4] TRIAZOLE</p>
<p>62</p>		<p>3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(1H-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE</p>
<p>63</p>		<p>3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(1-ETHYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE</p>
<p>64</p>		<p>3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(1-PROPYL-INDOL-5-YL)-5-MERCAPTO-[1,2,4] TRIAZOLE</p>

TABLE 1-continued

	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
65			3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(1-METHYL-2-TRIFLUOROMETHYL-BENZIMIDAZOL-5-YL)-5-MERCAPTO-[1,2,4]TRIAZOLE
66			3-(2,4-DIHYDROXY-5-ISOPROPYL-PHENYL)-4-(1-ISOPROPYL-INDOL-4-YL)-5-HYDROXY-[1,2,4]TRIAZOLE

TABLE 2

Compounds according to Formula (Ia)


No.	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
1A			5-HYDROXY-4-(5-HYDROXY-4-(1-METHYL-1H-INDOL-5-YL)-4H-1,2,4-TRIAZOL-3-YL)-2-ISOPROPYLPHENYL DIHYDROGEN PHOSPHATE
2A			SODIUM 5-HYDROXY-4-(5-HYDROXY-4-(1-METHYL-1H-INDOL-5-YL)-4H-1,2,4-TRIAZOL-3-YL)-2-ISOPROPYLPHENYL PHOSPHATE

TABLE 2-continued

Compounds according to Formula (Ia)			
No.	STRUCTURE	TAUTOMERIC STRUCTURE	NAME
3A			2-(3,4-DIMETHOXYPHENETHYL)-5-HYDROXY-4-(5-HYDROXY-4-(1-METHYL-1H-INDOL-5-YL)-4H-1,2,4-TRIAZOL-3-YL)PHENYL DIHYDROGEN PHOSPHATE
4A			4-(4-(1,3-DIMETHYL-1H-INDOL-5-YL)-5-HYDROXY-4H-1,2,4-TRIAZOL-3-YL)-2-ETHYL-5-HYDROXYPHENYL DIHYDROGEN PHOSPHATE

[0160] The Hsp90 inhibitory compounds used in the disclosed combination methods can be prepared according to the procedures disclosed in U.S. Patent Publication No. 2006/0,167,070, and WO2009/023,211.

[0161] These triazolone compounds typically can form a tautomeric structure as shown below and as exemplified by the tautomeric structures shown in Tables 1 and 2:



[0162] In an embodiment, the present invention provides a pharmaceutical combination for the treatment, prophylaxis, and amelioration of proliferative disorders, such as cancer. In a specific embodiment, the combination comprises one or more Hsp90 inhibitors according to formulae (I) or (Ia), or a compound in Tables 1 or 2, or a tautomer or a pharmaceutically acceptable salt thereof in addition to a platinum-containing anti-cancer agent.

[0163] In an embodiment, the combination includes a pharmaceutical composition or a single unit dosage form containing both an Hsp90 inhibitor and a platinum-containing anti-cancer agent. Pharmaceutical combinations and dosage forms described herein comprise the two active ingredients in relative amounts and formulated in such a way that a given

pharmaceutical combination or dosage form can be used to treat proliferative disorders, such as cancer. Preferred pharmaceutical combinations and dosage forms comprise a compound of formulae (I) or (Ia), or a compound in Tables 1 or 2, or a tautomer or pharmaceutically acceptable salt thereof, in combination with a platinum-containing anticancer agent. In other embodiments, the Hsp90 inhibitor and the platinum-containing anticancer agent may be in individual or separate pharmaceutical compositions, depending on the dosing schedules, preferred routes of administration, and available formulations of the two compounds. Optionally, these embodiments can also contain one or more additional therapeutic agents.

[0164] The pharmaceutical combinations described herein are formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral, intranasal (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. In a specific embodiment, the combination is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous, subcutaneous, intramuscular, oral, intranasal or topical administration to human beings. In an embodiment, the combination is formulated in accordance with routine procedures for subcutaneous administration to human beings.

[0165] In a specific embodiment, the combination therapies described herein comprise one or more compounds and at least one other therapy which has the same mechanism of action as the compounds. In another specific embodiment, the

combination therapies described herein comprise one or more compounds described herein and at least one other therapy which has a different mechanism of action than the compounds. In certain embodiments, the combination therapies described herein improve the therapeutic effect of one or more triazolone compounds described herein by functioning together with the platinum-containing anticancer agent to have an additive or synergistic effect. In certain embodiments, the combination therapies described herein reduce the side effects associated with the therapies. In certain embodiments, the combination therapies described herein reduce the effective dosage of one or more of the therapies.

**[0166]** In a specific embodiment, the combination comprising one or more triazolone compounds described herein is administered to a subject, preferably a human, to prevent, treat, manage, or ameliorate cancer, or one or more symptom thereof. In accordance with the invention, the pharmaceutical combinations described herein may also comprise one or more other agents being used, have been used, or are known to be useful in the treatment or amelioration of cancer, particularly breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, or colon cancer. The pharmaceutical combinations described herein utilize pharmaceutical compositions and dosage forms which comprise one or more excipients. Suitable excipients are well known to those skilled in the art of pharmacy.

**[0167]** The triazolone compounds described herein can be also formulated into or administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include those described in U.S. Pat. Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566.

**[0168]** In an embodiment, the present invention also provides a method of treating a proliferative disorder in a subject, comprising administering to the subject an effective amount of the combination of an Hsp90 inhibitor and a platinum-containing anticancer agent as described herein. In an embodiment, the proliferative disorder is cancer. In one aspect of this embodiment, the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer.

**[0169]** Other anti-proliferative or anti-cancer therapies may be combined with the compounds described herein to treat proliferative diseases and cancer. Other therapies or anti-cancer agents that may be used in combination with the inventive anti-cancer agents described herein include surgery, radiotherapy (including gamma-radiation, neutron beam radiotherapy, electron beam radiotherapy, proton therapy, brachytherapy, and systemic radioactive isotopes), endocrine therapy, biologic response modifiers (including interferons, interleukins, and tumor necrosis factor (TNF)), hyperthermia and cryotherapy, agents to attenuate any adverse effects (e.g., antiemetics), and other approved chemotherapeutic drugs.

**[0170]** The therapeutic agents of the combination therapies described herein can be administered sequentially or concurrently. In an embodiment, the administration of the Hsp90 inhibitor and the platinum-containing anticancer agent are done concurrently. In another embodiment, the administration of the Hsp90 inhibitor and the platinum-containing anticancer agent are done separately. In another embodiment, the administration of the Hsp90 inhibitor and the platinum-containing anticancer agent are done sequentially. In an embodiment, the administration of the Hsp90 inhibitor and the platinum-containing anticancer agent are done until the cancer is cured or stabilized or improved.

**[0171]** In a specific embodiment, the present method includes treating, managing, or ameliorating cancer, or one or more symptoms thereof, comprising administering to a subject in need thereof one or more compounds represented by the structural formulae (I) or (Ia) or a compound in Table 1 or Table 2, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin, wherein the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer.

**[0172]** In another embodiment, the method of treating a subject with cancer includes administering to the subject an effective amount of a triazolone compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with an effective amount of a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0173]** In another embodiment, the method of treating a subject with cancer includes administering to the subject an effective amount of a triazolone compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with an effective amount of cisplatin. In another embodiment, the method of treating a subject with cancer includes administering to the subject an effective amount of a triazolone compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a synergistic amount of cisplatin. In an embodiment, cisplatin is administered IV at a dose of between about 20 mg/m<sup>2</sup> to about 150 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered at a dose of about 20 mg/m<sup>2</sup> once daily. In an embodiment, cisplatin is administered IV at about 20 mg/m<sup>2</sup> once daily for 5 days per cycle. In an embodiment, cisplatin is administered IV at a dose of between about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered IV at a dose of between about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of between about 50 mg/m<sup>2</sup> to about 70 mg/m<sup>2</sup> once every 3 or 4 weeks per cycle. In an embodiment, the amount of the

Hsp90 inhibitor is from about 2 mg/m<sup>2</sup> to about 260 mg/m<sup>2</sup>. In an embodiment, the amount of the Hsp90 inhibitor is about 75 mg/m<sup>2</sup>, about 85 mg/m<sup>2</sup>, about 100 mg/m<sup>2</sup>, about 110 mg/m<sup>2</sup>, about 115 mg/m<sup>2</sup>, about 120 mg/m<sup>2</sup>, about 145 mg/m<sup>2</sup>, about 150 mg/m<sup>2</sup>, about 175 mg/m<sup>2</sup>, about 180 mg/m<sup>2</sup>, about 200 mg/m<sup>2</sup>, about 215 mg/m<sup>2</sup> or about 260 mg/m<sup>2</sup>. In an embodiment, the Hsp90 inhibitor is administered IV once weekly or twice weekly. In any one of the above embodiments, the cancer may have a KRAS mutation. In any one of the above embodiments, the cancer may have an ALK mutation. In any one of the above embodiments, the cancer may have a BRAF mutation.

**[0174]** In another embodiment, the method of treating a subject with cancer includes administering to the subject an effective amount of a triazolone compound of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with an effective amount of a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0175]** In another embodiment, the method of treating a subject with cancer includes administering to the subject an effective amount of a triazolone compound of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with an effective amount of cisplatin. In another embodiment, the method of treating a subject with cancer includes administering to the subject an effective amount of a triazolone compound of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a synergistic amount of cisplatin. In an embodiment, cisplatin is administered IV at a dose of between about 20 mg/m<sup>2</sup> to about 150 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered at a dose of about 20 mg/m<sup>2</sup> once daily. In an embodiment, cisplatin is administered IV at about 20 mg/m<sup>2</sup> once daily for 5 days per cycle. In an embodiment, cisplatin is administered IV at a dose of between about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered IV at a dose of between about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of between about 50 mg/m<sup>2</sup> to about 70 mg/m<sup>2</sup> once every 3 or 4 weeks per cycle. In an embodiment, the amount of the Hsp90 inhibitor is from about 2 mg/m<sup>2</sup> to about 260 mg/m<sup>2</sup>. In an embodiment, the amount of the Hsp90 inhibitor is about 75 mg/m<sup>2</sup>, about 85 mg/m<sup>2</sup>, about 100 mg/m<sup>2</sup>, about 110 mg/m<sup>2</sup>, about 115 mg/m<sup>2</sup>, about 120 mg/m<sup>2</sup>, about 145 mg/m<sup>2</sup>, about 150 mg/m<sup>2</sup>, about 175 mg/m<sup>2</sup>, about 180 mg/m<sup>2</sup>, about 200 mg/m<sup>2</sup>, about 215 mg/m<sup>2</sup> or about 260 mg/m<sup>2</sup>. In an embodiment, the Hsp90 inhibitor is administered IV once weekly or twice weekly. In any one of the above embodiments, the cancer may have a KRAS mutation. In any one of the above embodiments, the cancer may have an ALK mutation. In any one of the above embodiments, the cancer may have a BRAF mutation.

**[0176]** In another embodiment, the method of treating a subject with cancer includes administering to the subject an

effective amount of a triazolone compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin, wherein the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0177]** In another embodiment, the method of treating a subject with cancer includes administering to the subject an effective amount of a triazolone compound of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin, wherein the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or cancer. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0178]** In yet another embodiment, the method of treating a subject with cancer, wherein the subject is being or has been treated with a chemotherapeutic agent, includes administering to the subject an effective amount of a triazolone compound represented by the structural formulae (I) or (Ia) or a compound in Table 1 or Table 2, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0179]** In an embodiment, the method of treating a subject with cancer, wherein the subject is being or has been treated with a chemotherapeutic agent, includes administering to the subject an effective amount of a triazolone compound represented by the structural formulae (I) or (Ia) or a compound in Table 1 or Table 2, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin, wherein the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0180]** In another embodiment, the method of treating a subject with cancer, wherein the subject is being or has been treated with a chemotherapeutic agent, includes administer-



ing to the subject an effective amount of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0181]** In another embodiment, the method of treating a subject with cancer, wherein the subject is being or has been treated with a chemotherapeutic agent, includes administering to the subject an effective amount of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with cisplatin. In an embodiment, cisplatin is administered IV at a dose of between about 20 mg/m<sup>2</sup> to about 150 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered at a dose of about 20 mg/m<sup>2</sup> once daily. In an embodiment, cisplatin is administered IV at about 20 mg/m<sup>2</sup> once daily for 5 days per cycle. In an embodiment, cisplatin is administered IV at a dose of between about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered IV at a dose of between about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of between about 50 mg/m<sup>2</sup> to about 70 mg/m<sup>2</sup> once every 3 or 4 weeks per cycle. In an embodiment, the amount of the Hsp90 inhibitor is from about 2 mg/m<sup>2</sup> to about 260 mg/m<sup>2</sup>. In an embodiment, the amount of the Hsp90 inhibitor is about 75 mg/m<sup>2</sup>, about 85 mg/m<sup>2</sup>, about 100 mg/m<sup>2</sup>, about 110 mg/m<sup>2</sup>, about 115 mg/m<sup>2</sup>, about 120 mg/m<sup>2</sup>, about 145 mg/m<sup>2</sup>, about 150 mg/m<sup>2</sup>, about 175 mg/m<sup>2</sup>, about 180 mg/m<sup>2</sup>, about 200 mg/m<sup>2</sup>, about 215 mg/m<sup>2</sup> or about 260 mg/m<sup>2</sup>. In an embodiment, the Hsp90 inhibitor is administered IV once weekly or twice weekly. In any one of the above embodiments, the cancer may have a KRAS mutation. In any one of the above embodiments, the cancer may have an ALK mutation. In any one of the above embodiments, the cancer may have a BRAF mutation.

**[0182]** In another embodiment, the method of treating a subject with cancer, wherein the subject is being or has been treated with a chemotherapeutic agent, includes administering to the subject an effective amount of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0183]** In another embodiment, the method of treating a subject with cancer, wherein the subject is being or has been treated with a chemotherapeutic agent, includes administering to the subject an effective amount of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with cisplatin. In an embodiment, cisplatin is administered IV at a dose of between about 20 mg/m<sup>2</sup> to about 150 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered at a dose of about 20 mg/m<sup>2</sup> once daily. In an embodiment, cisplatin is administered IV at about 20 mg/m<sup>2</sup> once daily for 5 days per cycle. In

an embodiment, cisplatin is administered IV at a dose of between about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered IV at a dose of between about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of between about 50 mg/m<sup>2</sup> to about 70 mg/m<sup>2</sup> once every 3 or 4 weeks per cycle. In an embodiment, the amount of the Hsp90 inhibitor is from about 2 mg/m<sup>2</sup> to about 260 mg/m<sup>2</sup>. In an embodiment, the amount of the Hsp90 inhibitor is about 75 mg/m<sup>2</sup>, about 85 mg/m<sup>2</sup>, about 100 mg/m<sup>2</sup>, about 110 mg/m<sup>2</sup>, about 115 mg/m<sup>2</sup>, about 120 mg/m<sup>2</sup>, about 145 mg/m<sup>2</sup>, about 150 mg/m<sup>2</sup>, about 175 mg/m<sup>2</sup>, about 180 mg/m<sup>2</sup>, about 200 mg/m<sup>2</sup>, about 215 mg/m<sup>2</sup> or about 260 mg/m<sup>2</sup>. In an embodiment, the Hsp90 inhibitor is administered IV once weekly or twice weekly. In any one of the above embodiments, the cancer may have a KRAS mutation. In any one of the above embodiments, the cancer may have an ALK mutation. In any one of the above embodiments, the cancer may have a BRAF mutation.

**[0184]** In an embodiment, the method of treating a subject with cancer, wherein the subject is being or has been treated with a chemotherapeutic agent, includes administering to the subject an effective amount of a triazolone compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin, wherein the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0185]** In an embodiment, the method of treating a subject with cancer, wherein the subject is being or has been treated with a chemotherapeutic agent, includes administering to the subject an effective amount of a triazolone compound of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin, wherein the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0186]** In an embodiment, the method of treating a subject with cancer, wherein the subject has proven refractory to other therapies but is no longer on these therapies, includes administering to the subject an effective amount of a triazolone compound represented by the structural formulae (I) or

(Ia) or a compound in Table 1 or Table 2, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin, wherein the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0187]** In another embodiment, the method of treating a subject with cancer, wherein the subject has proven refractory to other therapies but is no longer on these therapies, includes administering to the subject an effective amount of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0188]** In another embodiment, the method of treating a subject with cancer, wherein the subject has proven refractory to other therapies but is no longer on these therapies, includes administering to the subject an effective amount of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with cisplatin. In an embodiment, cisplatin is administered IV at a dose of between about 20 mg/m<sup>2</sup> to about 150 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered at a dose of about 20 mg/m<sup>2</sup> once daily. In an embodiment, cisplatin is administered IV at about 20 mg/m<sup>2</sup> once daily for 5 days per cycle. In an embodiment, cisplatin is administered IV at a dose from about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered IV at a dose from about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of between about 50 mg/m<sup>2</sup> to about 70 mg/m<sup>2</sup> once every 3 or 4 weeks per cycle. In an embodiment, the amount of the Hsp90 inhibitor is from about 2 mg/m<sup>2</sup> to about 260 mg/m<sup>2</sup>. In an embodiment, the amount of the Hsp90 inhibitor is about 75 mg/m<sup>2</sup>, about 85 mg/m<sup>2</sup>, about 100 mg/m<sup>2</sup>, about 110 mg/m<sup>2</sup>, about 115 mg/m<sup>2</sup>, about 120 mg/m<sup>2</sup>, about 145 mg/m<sup>2</sup>, about 150 mg/m<sup>2</sup>, about 175 mg/m<sup>2</sup>, about 180 mg/m<sup>2</sup>, about 200 mg/m<sup>2</sup>, about 215 mg/m<sup>2</sup> or about 260 mg/m<sup>2</sup>. In an embodiment, the Hsp90 inhibitor is administered IV once weekly or twice weekly. In any one of the above embodiments, the cancer may have a KRAS mutation. In any one of the above embodiments, the cancer may have an ALK mutation. In any one of the above embodiments, the cancer may have a BRAF mutation.

**[0189]** In another embodiment, the method of treating a subject with cancer, wherein the subject has proven refractory to other therapies but is no longer on these therapies, includes administering to the subject an effective amount of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a platinum-containing anticancer agent

such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0190]** In another embodiment, the method of treating a subject with cancer, wherein the subject has proven refractory to other therapies but is no longer on these therapies, includes administering to the subject an effective amount of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with cisplatin. In an embodiment, cisplatin is administered IV at a dose of between about 20 mg/m<sup>2</sup> to about 150 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered at a dose of about 20 mg/m<sup>2</sup> once daily. In an embodiment, cisplatin is administered IV at about 20 mg/m<sup>2</sup> once daily for 5 days per cycle. In an embodiment, cisplatin is administered IV at a dose from about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup>. In an embodiment, cisplatin is administered IV at a dose from about 75 mg/m<sup>2</sup> to about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose of about 100 mg/m<sup>2</sup> once every 4 weeks per cycle. In an embodiment, cisplatin is administered IV at a dose from about 50 mg/m<sup>2</sup> to about 70 mg/m<sup>2</sup> once every 3 or 4 weeks per cycle. In an embodiment, the amount of the Hsp90 inhibitor is from about 2 mg/m<sup>2</sup> to about 260 mg/m<sup>2</sup>. In an embodiment, the amount of the Hsp90 inhibitor is about 75 mg/m<sup>2</sup>, about 85 mg/m<sup>2</sup>, about 100 mg/m<sup>2</sup>, about 110 mg/m<sup>2</sup>, about 115 mg/m<sup>2</sup>, about 120 mg/m<sup>2</sup>, about 145 mg/m<sup>2</sup>, about 150 mg/m<sup>2</sup>, about 175 mg/m<sup>2</sup>, about 180 mg/m<sup>2</sup>, about 200 mg/m<sup>2</sup>, about 215 mg/m<sup>2</sup> or about 260 mg/m<sup>2</sup>. In an embodiment, the Hsp90 inhibitor is administered IV once weekly or twice weekly. In any one of the above embodiments, the cancer may have a KRAS mutation. In any one of the above embodiments, the cancer may have an ALK mutation. In any one of the above embodiments, the cancer may have a BRAF mutation.

**[0191]** In an embodiment, the method of treating a subject with cancer, wherein the subject has proven refractory to other therapies but is no longer on these therapies, includes administering to the subject an effective amount of a triazolone compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin, wherein the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0192]** In an embodiment, the method of treating a subject with cancer, wherein the subject has proven refractory to other therapies but is no longer on these therapies, includes administering to the subject an effective amount of a triazolone compound of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, in combination with a platinum-con-

taining anticancer agent such as cisplatin, carboplatin, or oxaliplatin, wherein the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, melanoma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0193]** In a further embodiment, the method includes inhibiting the growth of a cancer or tumor cell comprising the steps of: (a) contacting the cell with an effective amount of a compound of formulae (I) or (Ia) or a compound in Table (1) or Table (2), or tautomer or a pharmaceutically acceptable salt thereof; and (b) exposing the cell to an effective amount of a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin.

**[0194]** In a further embodiment, the method includes inhibiting the growth of a cancer or tumor cell comprising the steps of: (a) contacting the cell with an effective amount of a compound of -(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof; and (b) exposing the cell to an effective amount of a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0195]** In another embodiment, the invention also provides a method of inhibiting the growth of a cancer or tumor cell, comprising the steps of: (a) contacting the cell with an effective amount of a compound of -(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof; and (b) exposing the cell to an effective amount of cisplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0196]** In another embodiment, the invention also provides a method of inhibiting the growth of a cancer or tumor cell, comprising the steps of: (a) contacting the cell with an effective amount of a compound of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or tautomer or a pharmaceutically acceptable salt thereof; and (b) exposing the cell to an effective amount of a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0197]** In another embodiment, the invention also provides a method of inhibiting the growth of a cancer or tumor cell, comprising the steps of: (a) contacting the cell with an effective amount of a compound of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or tautomer or a pharmaceutically acceptable salt thereof; and (b) exposing the cell to an effective amount of cisplatin. In an embodiment, the cancer has a KRAS mutation. In an embodiment, the cancer has an ALK mutation. In an embodiment, the cancer has a BRAF mutation.

**[0198]** In general, the recommended daily dose range of a triazolone compound for the conditions described herein lie within the range of from about 0.01 mg to about 1000 mg per day, given as a single once-a-day dose preferably as divided doses throughout a day. In an embodiment, the daily dose is administered twice daily in equally divided doses. Specifically, a daily dose range should be from about 5 mg to about 500 mg per day, more specifically, between about 10 mg and about 200 mg per day. In managing the patient, the therapy should be initiated at a lower dose, perhaps about 1 mg to about 25 mg, and increased if necessary up to about 200 mg to about 1000 mg per day as either a single dose or divided doses, depending on the patient's global response. It may be necessary to use dosages of the active ingredient outside the ranges disclosed herein in some cases, as will be apparent to those of ordinary skill in the art. Furthermore, it is noted that the clinician or treating physician will know how and when to interrupt, adjust, or terminate therapy in conjunction with individual patient response.

**[0199]** Different therapeutically effective amounts may be applicable for different cancers, as will be readily known by those of ordinary skill in the art. Similarly, amounts sufficient to prevent, manage, treat or ameliorate such cancers, but insufficient to cause, or sufficient to reduce, adverse effects associated with the triazolone compounds described herein are also encompassed by the above described dosage amounts and dose frequency schedules. Further, when a patient is administered multiple dosages of a triazolone compound described herein, not all of the dosages need be the same. For example, the dosage administered to the patient may be increased to improve the prophylactic or therapeutic effect of the compound or it may be decreased to reduce one or more side effects that a particular patient is experiencing.

**[0200]** In a specific embodiment, the dosage of the composition comprising a triazolone compound described herein administered to prevent, treat, manage, or ameliorate cancer, or one or more symptoms thereof in a patient is 150 µg/kg, preferably 250 µg/kg, 500 µg/kg, 1 mg/kg, 5 mg/kg, 10 mg/kg, 25 mg/kg, 50 mg/kg, 75 mg/kg, 100 mg/kg, 125 mg/kg, 150 mg/kg, or 200 mg/kg or more of a patient's body weight. In another embodiment, the dosage of the composition comprising a compound described herein administered to prevent, treat, manage, or ameliorate cancer, or one or more symptoms thereof in a patient is a unit dose of 0.1 mg to 20 mg, 0.1 mg to 15 mg, 0.1 mg to 12 mg, 0.1 mg to 10 mg, 0.1 mg to 8 mg, 0.1 mg to 7 mg, 0.1 mg to 5 mg, 0.1 to 2.5 mg, 0.25 mg to 20 mg, 0.25 to 15 mg, 0.25 to 12 mg, 0.25 to 10 mg, 0.25 to 8 mg, 0.25 mg to 7 mg, 0.25 mg to 5 mg, 0.5 mg to 2.5 mg, 1 mg to 20 mg, 1 mg to 15 mg, 1 mg to 12 mg, 1 mg to 10 mg, 1 mg to 8 mg, 1 mg to 7 mg, 1 mg to 5 mg, or 1 mg to 2.5 mg. The unit dose can be administered 1, 2, 3, 4 or more times daily, or once every 2, 3, 4, 5, 6 or 7 days, or once weekly, once every two weeks, once every three weeks or once monthly.

**[0201]** In certain embodiments, when the triazolone compounds described herein are administered in combination with a platinum-containing anticancer agent, the therapies are administered less than 5 minutes apart, less than 30 minutes apart, 1 hour apart, at about 1 hour apart, at about 1 to about 2 hours apart, at about 2 hours to about 3 hours apart, at about 3 hours to about 4 hours apart, at about 4 hours to about 5 hours apart, at about 5 hours to about 6 hours apart, at about 6 hours to about 7 hours apart, at about 7 hours to about 8 hours apart, at about 8 hours to about 9 hours apart, at about 9 hours to about 10 hours apart, at about 10 hours to about 11

hours apart, at about 11 hours to about 12 hours apart, at about 12 hours to 18 hours apart, 18 hours to 24 hours apart, 24 hours to 36 hours apart, 36 hours to 48 hours apart, 48 hours to 52 hours apart, 52 hours to 60 hours apart, 60 hours to 72 hours apart, 72 hours to 84 hours apart, 84 hours to 96 hours apart, or 96 hours to 120 hours part. In one embodiment, two or more therapies are administered within the same patient visit.

**[0202]** In certain embodiments, one or more compounds described herein and one or more other the therapies (e.g., therapeutic agents) are cyclically administered. Cycling therapy involves the administration of a first therapy (e.g., a first prophylactic or therapeutic agents) for a period of time, followed by the administration of a second therapy (e.g., a second prophylactic or therapeutic agents) for a period of time, followed by the administration of a third therapy (e.g., a third prophylactic or therapeutic agents) for a period of time and so forth, and repeating this sequential administration, i.e., the cycle in order to reduce the development of resistance to one of the agents, to avoid or reduce the side effects of one of the agents, and/or to improve the efficacy of the treatment.

**[0203]** In certain embodiments, administration of the same compound described herein may be repeated and the administrations may be separated by at least 1 day, 2 days, 3 days, 5 days, 10 days, 15 days, 30 days, 45 days, 2 months, 75 days, 3 months, or 6 months. In other embodiments, administration of the same prophylactic or therapeutic agent may be repeated and the administration may be separated by at least 1 day, 2 days, 3 days, 5 days, 10 days, 15 days, 30 days, 45 days, 2 months, 75 days, 3 months, or 6 months.

**[0204]** In a specific embodiment, a method of preventing, treating, managing, or ameliorating a proliferative disorders, such as cancer, or one or more symptoms thereof, the methods comprising administering to a subject in need thereof a dose of at least 150  $\mu\text{g}/\text{kg}$ , preferably at least 250  $\mu\text{g}/\text{kg}$ , at least 500  $\mu\text{g}/\text{kg}$ , at least 1 mg/kg, at least 5 mg/kg, at least 10 mg/kg, at least 25 mg/kg, at least 50 mg/kg, at least 75 mg/kg, at least 100 mg/kg, at least 125 mg/kg, at least 150 mg/kg, or at least 200 mg/kg or more of one or more compounds described herein once every day, preferably, once every 2 days, once every 3 days, once every 4 days, once every 5 days, once every 6 days, once every 7 days, once every 8 days, once every 10 days, once every two weeks, once every three weeks, or once a month. Alternatively, the dose can be divided into portions (typically equal portions) administered two, three, four or more times a day.

**[0205]** In an embodiment, the invention also provides the use of a compound of formulae (I) or (Ia), or a compound in Tables 1 or 2, or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of a subject with cancer. In an embodiment, the invention further provides the use of a compound of formulae (I) or (Ia), or a compound in Tables 1 or 2, or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of a subject with a cancer, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the invention further provides the use of a compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the

treatment of a subject with a cancer, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the invention further provides the use of a compound of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a subject with a cancer, in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the invention further provides the use of the compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a subject with a cancer, in combination with cisplatin. In an embodiment, the invention further provides the synergistic use of the compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a subject with a cancer, in combination with cisplatin.

**[0206]** In an embodiment, the invention also provides a compound of formulae (I) or (Ia) or a pharmaceutically acceptable salt thereof for use in treating a subject with a cancer. In an embodiment, the invention also provides a compound of formulae (I) or (Ia) or a pharmaceutically acceptable salt thereof for use in treating a subject with cancer in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the invention also provides a compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, for use in treating a subject with cancer in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the invention also provides a compound of 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, for use in treating a subject with cancer in combination with a platinum-containing anticancer agent such as cisplatin, carboplatin, or oxaliplatin. In an embodiment, the invention also provides a compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, for use in treating a subject with cancer in combination with cisplatin. In an embodiment, the invention also provides a compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer, or a pharmaceutically acceptable salt thereof, for synergistic use in treating a subject with cancer in combination with cisplatin.

## EXAMPLES

**[0207]** In Vitro combination analysis of ganetespib with chemotherapy in colorectal cancer cells

### A. Materials and Methods

**[0208]** Cell Lines

**[0209]** Human HCT-116 colorectal cancer cells (CRC) and MDA-MB-231 breast cancer cells were purchased from the American Type Culture Collection (Manassas, Va.) and grown following ATCC recommendations, in the presence of

fetal bovine serum (10%), 2 mL-glutamine and antibiotics (100 IU/ml penicillin and 100 µg/ml streptomycin, Sigma). Cells were maintained at 37° C., 5% CO<sub>2</sub> atmosphere.

**[0210]** Cell Viability Assays

**[0211]** Cell viability was measured using the alamarBlue assay (Invitrogen). In brief, cells were plated in 96-well plates in triplicate at 5K cells per well and incubated at 37° C., 5% CO<sub>2</sub> atmosphere for 24 hr prior to the addition of drug or vehicle (0.3% DMSO) to the culture medium. After 72 hr, 10 µl/well alamarBlue was added to the wells and incubated for an additional 3 hr at 37° C., 5% CO<sub>2</sub> atmosphere. Fluorescence (560<sub>EX</sub>/590<sub>EM</sub> nm) was measured with a SpectraMax microplate reader (Molecular Devices) and the resulting data were used to calculate cell viability, normalized to vehicle control.

**[0212]** Mouse Studies

**[0213]** Six to seven week old, female CB17/Icr-Prkdc<sup>scid</sup>/Crl (SCID) mice were obtained from Charles River Laboratories (Wilmington, Mass., USA). Animals were housed 4-5/cage in micro-isolators, with a 12 hr/12 hr light/dark cycle, acclimated for at least 1 week prior to use and fed normal laboratory chow ad libitum. Animals were between seven to eight weeks of age at implantation. To implant HCT-116 or MDA-MB-231 tumor cells into SCID mice, cells were harvested by trypsinization, washed in PBS and re-suspended at a concentration of 5×10<sup>7</sup> cells/mL in 50% non-supplemented medium and 50% Matrigel Basement Membrane Matrix (BD Biosciences; Bedford, Mass., USA). Using a 27 gauge needle and 1 cc syringe, 5×10<sup>6</sup> cells in 0.1 mL of a cell suspension were injected subcutaneously into the flanks of SCID mice.

**[0214]** Tumors were then permitted to develop in vivo until the majority reached 95-195 mm<sup>3</sup> in tumor volume. Animals with oblong, very small or large tumors were discarded and only animals carrying tumors that displayed consistent growth rates were selected for studies. Tumor volumes (V) were calculated by caliper measurement of the width (W), length (L) and thickness (T) of tumors using the following formula: V=0.5236×(L×W×T). Animals were randomized into treatment groups so that the average tumor volumes of each group were similar at the start of dosing.

**[0215]** Ganetespiib was prepared by dissolving the appropriate amounts of the compound in dimethyl sulfoxide (DMSO) by sonication in an ultrasonic water bath. Stock solutions were prepared weekly, stored at -20° C. and diluted fresh each day for dosing. A solution of 20% Cremophor RH40 (polyoxyl 40 hydrogenated castor oil; BASF Corp., Aktiengesellschaft, Ludwigshafen, Germany) in 5% dextrose in water (Abbott Laboratories, North Chicago, Ill., USA) was also prepared by first heating 100% Cremophor RH40 at 50-60° C. until liquefied and clear, diluting 1:5 with 100% D5W, reheating again until clear and then mixing well. This solution can be stored at room temperature for up to 3 months prior to use. To prepare DRD formulations for daily dosing, DMSO stock solutions were diluted 1:10 with 20% Cremophor RH40. The final DRD formulation for dosing contained 10% DMSO, 18% Cremophor RH40, 3.6% dextrose, 68.4% water and the appropriate amount of test article. Animals were intravenously (i.v.) injected with this formulation at 10 mL per kg body weight 1 day each week. Cisplatin was prepared fresh in saline and given intravenously 6 hours after ganetespiib injection, once per week.

B. Combination Studies with Ganetespiib and Cisplatin

**[0216]** The half maximal inhibitory concentration (IC<sub>50</sub>) for ganetespiib (synthesized at Synta Pharmaceuticals) and cisplatin (purchased from Sigma) were first determined using a 1.5-fold serial dilution series of compound. After HCT-116 cells were exposed to drug for 72 hr, cell viability was measured and results were fit to a four parameter logistic model (XLFit, ID Business Solutions) shown in FIGS. 1 and 2. The IC<sub>50</sub> for ganetespiib was calculated at approximately 32 nM, and 5.7 µM for cisplatin.

**[0217]** Combinations between ganetespiib and cisplatin were then performed in HCT-116 cells concurrently based on the IC<sub>50</sub> for each agent in matrix format with 54 combination pairs for each drug. The combined drugs, as well as each drug alone, were incubated with the cells for 3 days and the surviving fraction of cells relative to control was determined using the alamarBlue assay. Representative figures are shown in FIGS. 3 and 4. The combination of ganetespiib with cisplatin displayed enhanced cytotoxicity relative to single agent drugs alone. Similar results were observed when cells were exposed to ganetespiib for just one hour, washed and then treated with cisplatin for 3 days.


**[0218]** Combinations between ganetespiib and cisplatin were further performed in animal models, using HCT-116 colon cancer or MDA-MB-231 breast cancer tumor xenografts. Ganetespiib (150 mg/kg) was given as a single bolus injection 6 hours prior to cisplatin (2 mg/kg) administration. Representative figures are shown in FIGS. 5 and 6. In both cancer xenograft models, the combination of ganetespiib with cisplatin displayed greater antitumor activity relative to monotherapy.

**[0219]** In conclusion, these data support the use of ganetespiib in combination with cisplatin in solid cancers such as gastric, bladder, colorectal, breast, ovarian and lung. See also Acquaviva et al, Mol Cancer Ther. 2012, September issue.

**[0220]** All publications, patent applications, patents, and other documents cited herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples throughout the specification are illustrative only and not intended to be limiting in any way.


What is claimed is:

1. A pharmaceutical composition comprising a platinum-containing anti-cancer agent and an Hsp90 inhibitor according to the following formulae:



-continued

(1a)



or a tautomer, or a pharmaceutically acceptable salt thereof, wherein:

Z is OH, SH, or  $\text{NH}_2$ ;

X is  $\text{CR}_4$  or N;

$\text{R}_1$  is —H, —OH, —SH, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro, guanidino, a haloalkyl, a heteroalkyl, an alkoxy or cycloalkoxy, a haloalkoxy,  $-\text{NR}_{10}\text{R}_{11}$ ,  $-\text{O}_7$ ,  $-\text{C}(\text{O})\text{R}_7$ ,  $-\text{C}(\text{O})\text{OR}_7$ ,  $-\text{C}(\text{S})\text{R}_7$ ,  $-\text{C}(\text{O})\text{SR}_7$ ,  $-\text{C}(\text{S})\text{SR}_7$ ,  $-\text{C}(\text{S})\text{OR}_7$ ,  $-\text{C}(\text{S})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{C}(\text{NR}_8)$ ,  $\text{OR}_7$ ,  $-\text{C}(\text{NR}_8)\text{R}_7$ ,  $-\text{C}(\text{NR}_8)\text{NR}_{10}\text{R}_{11}$ ,  $-\text{C}(\text{NR}_8)\text{SR}_7$ ,  $-\text{OC}(\text{O})\text{R}_7$ ,  $-\text{OC}(\text{O})\text{OR}_7$ ,  $-\text{OC}(\text{S})\text{OR}_7$ ,  $-\text{OC}(\text{NR}_8)$ ,  $\text{OR}_7$ ,  $-\text{SC}(\text{O})\text{R}_7$ ,  $-\text{SC}(\text{O})\text{OR}_7$ ,  $-\text{SC}(\text{NR}_8)\text{OR}_7$ ,  $-\text{OC}(\text{S})\text{R}_7$ ,  $-\text{SC}(\text{S})\text{R}_7$ ,  $-\text{SC}(\text{S})\text{OR}_7$ ,  $-\text{OC}(\text{O})$ ,  $\text{NR}_{10}\text{R}_{11}$ ,  $-\text{OC}(\text{S})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{OC}(\text{NR}_8)\text{NR}_{10}\text{R}_{11}$ ,  $-\text{SC}(\text{O})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{SC}(\text{NR}_8)\text{NR}_{10}\text{R}_{11}$ ,  $-\text{SC}(\text{S})$ ,  $\text{NR}_{10}\text{R}_{11}$ ,  $-\text{OC}(\text{NR}_8)\text{R}_7$ ,  $-\text{SC}(\text{NR}_8)\text{R}_7$ ,  $-\text{C}(\text{O})$ ,  $\text{NR}_{10}\text{R}_{11}$ ,  $-\text{NR}_8\text{C}(\text{O})\text{R}_7$ ,  $-\text{NR}_7\text{C}(\text{S})\text{R}_7$ ,  $-\text{NR}_7\text{C}(\text{S})$ ,  $\text{OR}_7$ ,  $-\text{NR}_7\text{C}(\text{NR}_8)\text{R}_7$ ,  $-\text{NR}_7\text{C}(\text{O})\text{OR}_7$ ,  $-\text{NR}_7\text{C}(\text{NR}_8)\text{OR}_7$ ,  $-\text{NR}_7\text{C}(\text{O})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{NR}_7\text{C}(\text{S})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{NR}_7\text{C}(\text{NR}_8)\text{NR}_{10}\text{R}_{11}$ ,  $-\text{SR}_7$ ,  $-\text{S}(\text{O})_p\text{R}_7$ ,  $-\text{OS}(\text{O})_p\text{R}_7$ ,  $-\text{OS}(\text{O})_p\text{OR}_7$ ,  $-\text{OS}(\text{O})_p\text{NR}_{10}\text{R}_{11}$ ,  $-\text{S}(\text{O})_p\text{OR}_7$ ,  $-\text{NR}_8\text{S}(\text{O})_p\text{R}_7$ ,  $-\text{NR}_7\text{S}(\text{O})_p\text{NR}_{10}\text{R}_{11}$ ,  $-\text{NR}_7\text{S}(\text{O})_p\text{OR}_7$ ,  $-\text{S}(\text{O})_p\text{NR}_{10}\text{R}_{11}$ ,  $-\text{SS}(\text{O})_p\text{R}_7$ ,  $-\text{SS}(\text{O})_p\text{OR}_7$ ,  $-\text{SS}(\text{O})_p\text{NR}_{10}\text{R}_{11}$ ,  $-\text{OP}(\text{O})(\text{OR}_7)_2$ , or  $-\text{SP}(\text{O})(\text{OR}_7)_2$ ;

$\text{R}_2$  is —H, —OH, —SH,  $-\text{NR}_7\text{H}$ ,  $-\text{OR}_{15}$ ,  $-\text{SR}_{15}$ ,  $-\text{NHR}_{15}$ ,  $-\text{O}(\text{CH}_2)_m\text{OH}$ ,  $-\text{O}(\text{CH}_2)_m\text{SH}$ ,  $-\text{O}(\text{CH}_2)_m\text{NR}_7\text{H}$ ,  $-\text{S}(\text{CH}_2)_m\text{OH}$ ,  $-\text{S}(\text{CH}_2)_m\text{SH}$ ,  $-\text{S}(\text{CH}_2)_m\text{NR}_7\text{H}$ ,  $-\text{OC}(\text{O})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{SC}(\text{O})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{NR}_7\text{C}(\text{O})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{OC}(\text{O})\text{R}_7$ ,  $-\text{SC}(\text{O})\text{R}_7$ ,  $-\text{NR}_7\text{C}(\text{O})\text{R}_7$ ,  $-\text{OC}(\text{O})\text{OR}_7$ ,  $-\text{SC}(\text{O})\text{OR}_7$ ,  $-\text{NR}_7\text{C}(\text{O})\text{OR}_7$ ,  $-\text{OCH}_2\text{C}(\text{O})\text{R}_7$ ,  $-\text{SCH}_2\text{C}(\text{O})\text{R}_7$ ,  $-\text{NR}_7\text{CH}_2\text{C}(\text{O})\text{R}_7$ ,  $-\text{OCH}_2\text{C}(\text{O})\text{OR}_7$ ,  $-\text{SCH}_2\text{C}(\text{O})\text{OR}_7$ ,  $-\text{NR}_7\text{CH}_2\text{C}(\text{O})\text{OR}_7$ ,  $-\text{OCH}_2\text{C}(\text{O})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{SCH}_2\text{C}(\text{O})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{NR}_7\text{CH}_2\text{C}(\text{O})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{OS}(\text{O})_p\text{R}_7$ ,  $-\text{SS}(\text{O})_p\text{R}_7$ ,  $-\text{NR}_7\text{S}(\text{O})_p\text{R}_7$ ,  $-\text{OS}(\text{O})_p\text{NR}_{10}\text{R}_{11}$ ,  $-\text{SS}(\text{O})_p\text{NR}_{10}\text{R}_{11}$ ,  $-\text{NR}_7\text{S}(\text{O})_p\text{NR}_{10}\text{R}_{11}$ ,  $-\text{OS}(\text{O})_p\text{OR}_7$ ,  $-\text{SS}(\text{O})_p\text{OR}_7$ ,  $-\text{NR}_7\text{S}(\text{O})_p\text{OR}_7$ ,  $-\text{OC}(\text{S})\text{R}_7$ ,  $-\text{SC}(\text{S})\text{R}_7$ ,  $-\text{NR}_7\text{C}(\text{S})\text{R}_7$ ,  $-\text{OC}(\text{S})\text{OR}_7$ ,  $-\text{SC}(\text{S})\text{OR}_7$ ,  $-\text{NR}_7\text{C}(\text{S})\text{OR}_7$ ,  $-\text{OC}(\text{S})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{SC}(\text{S})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{NR}_7\text{C}(\text{S})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{OC}(\text{NR}_8)\text{R}_7$ ,  $-\text{SC}(\text{NR}_8)\text{R}_7$ ,  $-\text{NR}_7\text{C}(\text{NR}_8)\text{R}_7$ ,  $-\text{OC}(\text{NR}_8)\text{OR}_7$ ,  $-\text{SC}(\text{NR}_8)\text{OR}_7$ ,  $-\text{NR}_7\text{C}(\text{NR}_8)\text{OR}_7$ ,  $-\text{OC}(\text{NR}_8)\text{NR}_{10}\text{R}_{11}$ ,  $-\text{SC}(\text{NR}_8)\text{NR}_{10}\text{R}_{11}$ , or  $-\text{NR}_7\text{C}(\text{NR}_8)\text{NR}_{10}\text{R}_{11}$ ;

$\text{R}_3$  is —H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, a haloalkyl, a heteroalkyl,  $-\text{C}(\text{O})\text{R}_7$ ,  $-(\text{CH}_2)_m\text{C}(\text{O})\text{OR}_7$ ,  $-\text{C}(\text{O})\text{OR}_7$ ,  $-\text{OC}(\text{O})\text{R}_7$ ,  $-\text{C}(\text{O})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{S}(\text{O})_p\text{R}_7$ ,  $-\text{S}(\text{O})_p\text{OR}_7$ , or  $-\text{S}(\text{O})_p\text{NR}_{10}\text{R}_{11}$ ;

$\text{R}_4$  is —H, —OH, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, halo, cyano, nitro, guanidino, a haloalkyl, a heteroalkyl,  $-\text{C}(\text{O})\text{R}_7$ ,  $-\text{C}(\text{O})\text{OR}_7$ ,  $-\text{OC}(\text{O})\text{R}_7$ ,  $-\text{C}(\text{O})\text{NR}_{10}\text{R}_{11}$ ,  $-\text{NR}_8\text{C}(\text{O})\text{R}_7$ ,  $-\text{SR}_7$ ,  $-\text{S}(\text{O})_p\text{R}_7$ ,  $-\text{OS}(\text{O})_p\text{R}_7$ ,  $-\text{S}(\text{O})_p\text{OR}_7$ ,  $-\text{NR}_8\text{S}(\text{O})_p\text{R}_7$ ,  $-\text{S}(\text{O})_p\text{NR}_{10}\text{R}_{11}$ , or  $\text{R}_3$  and  $\text{R}_4$  taken together with the carbon atoms to which they are attached form an optionally substituted cycloalkenyl, an optionally substituted aryl, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

$\text{R}_7$  and  $\text{R}_8$ , for each occurrence, are independently, —H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

$\text{R}_{10}$  and  $\text{R}_{11}$ , for each occurrence, are independently —H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or  $\text{R}_{10}$  and  $\text{R}_{11}$ , taken together with the nitrogen to which they are attached, form an optionally substituted heterocyclyl or an optionally substituted heteroaryl;

$\text{R}_{15}$ , for each occurrence, is independently, a lower alkyl;  $p$ , for each occurrence, is, independently, 1 or 2; and  $m$ , for each occurrence, is, independently, 1, 2, 3, or 4.

2. The pharmaceutical composition of claim 1, wherein the Hsp90 inhibitor is 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole or a tautomer or a pharmaceutically acceptable salt thereof.

3. The pharmaceutical composition of claim 1, wherein the Hsp90 inhibitor is 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof.

4. The pharmaceutical composition of claim 1, wherein the platinum-containing anticancer agent is selected from the group consisting of cisplatin, carboplatin, and oxaliplatin.

5. The pharmaceutical composition of claim 4, wherein the platinum-containing anticancer agent is cisplatin.

6. The pharmaceutical composition of claim 1, wherein the Hsp90 inhibitor is 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer or a pharmaceutically acceptable salt thereof, and the platinum-containing anticancer agent is cisplatin.

7. The pharmaceutical composition of claim 1, wherein the Hsp90 inhibitor is 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, and the platinum-containing anticancer agent is cisplatin.

8. The composition of claim 1, further comprising one or more additional therapeutic agents selected from the group consisting of vandetanib, trastuzumab, temodar, dexamethasone, epirubicin, Ifosfamide, mitoxantrone, vorinostat, interferon alpha, rituximab, prednisone, cyclophosphamide, bendamustine, adriamycin, valproate, celecoxib, thalidomide, nelarabine, methotrexate, filgrastim, gemtuzumab ozogamicin, testosterone, clofarabine, cytarabine, everolimus, rituximab, busulfan, capecitabine, pegfilgrastim, mesna, amrubicin, obatoclast, gefitinib, cyclosporine, dasatinib, temozolomide, thiotepa, plerixafor, mitotane, vincristine, doxorubicin, cixutumumab, endostar, fenofibrate, melphalan, sunitinib, rubitecan, enoxaparin, isotretinoin, tariquidar, pomalidomide, sorafenib, altretamine, idarubicin, rapamycin, zevalin, everolimus, pravastatin, carmustine, nelfinavir, streptozocin, tirapazamine, aprepitant, lenalidomide, G-CSF, procarbazine, alemtuzumab, amifostine, valspodar, lomustine, oblimersen, temsirolimus, vinblastine, figitumumab, belinostat, niacinamide, tipifarnib, estramustine, erlotinib, bevacizumab, paclitaxel, docetaxel, Abraxane®, pemetrexed, bortezomib, cetuximab, gemcitabine, 5-fluorouracil, leucovorin, and tetracycline.

9. The composition of claim 8, wherein the one or more therapeutic agents is selected from the group consisting of erlotinib, bevacizumab, bortezomib, paclitaxel, doxorubicin, docetaxel, mitoxantrone, cytarabine, 5-fluorouracil, leucovorin, and vincristine.

10. A method of treating cancer, comprising administering to the subject an effective amount of the pharmaceutical composition of claim 1.

11. The method of claim 10, wherein the cancer is breast cancer, gastric cancer, colorectal cancer, pancreatic cancer, ocular melanoma, prostate cancer, gastrointestinal stromal tumors (GIST), advanced esophagogastric cancer, mela-

noma, hepatocellular cancer, solid tumor, liver cancer, head and neck cancer, small cell lung cancer, non-small cell lung cancer, bladder cancer, testicular tumor, ovarian cancer, lymphoma, leukemia, multiple myeloma, or colon cancer.

12. The method of claim 11, wherein the cancer is gastric cancer, bladder cancer, pancreatic cancer, non-small cell lung cancer, ovarian cancer, or colorectal cancer.

13. (canceled)

14. The method of claim 10, wherein the method comprising administering a compound of 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer or a pharmaceutically acceptable salt thereof, and cisplatin.

15. The method of claim 14, wherein cisplatin is administered at a dose from about 20 mg/m<sup>2</sup> to about 150 mg/m<sup>2</sup>.

16. The method of claim 14, wherein the amount of the triazole compound is from about 2 mg/m<sup>2</sup> to about 260 mg/m<sup>2</sup>.

17. The method of claim 16, wherein the amount of the triazole compound is about 75 mg/m<sup>2</sup>, about 85 mg/m<sup>2</sup>, about 100 mg/m<sup>2</sup>, about 110 mg/m<sup>2</sup>, about 115 mg/m<sup>2</sup>, about 120 mg/m<sup>2</sup>, about 145 mg/m<sup>2</sup>, about 150 mg/m<sup>2</sup>, about 175 mg/m<sup>2</sup>, about 180 mg/m<sup>2</sup>, about 200 mg/m<sup>2</sup>, about 215 mg/m<sup>2</sup> or about 260 mg/m<sup>2</sup>.

18. The method of claim 16, wherein the Hsp90 inhibitor is administered IV once weekly or twice weekly.

19. A method of inhibiting the growth of a cancer or tumor cell in a subject, the method comprising the steps of: (a) contacting the cell with an effective amount of a compound of formulae (I) or (Ia) as defined in claim 1, and (b) exposing the cell to an effective amount of a platinum-containing anticancer agent, wherein the platinum-containing anticancer agent is selected from the group consisting of cisplatin, carboplatin, and oxaliplatin.

20. The method of claim 19, wherein the compound is 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, or a tautomer or a pharmaceutically acceptable salt thereof and the platinum-containing anticancer agent is cisplatin.

21. The method of claim 19, wherein the compound is 5-hydroxy-4-(5-hydroxy-4-(1-methyl-1H-indol-5-yl)-4H-1,2,4-triazol-3-yl)-2-isopropylphenyl dihydrogen phosphate, or a tautomer, or a pharmaceutically acceptable salt thereof, and the platinum-containing anticancer agent is cisplatin.

\* \* \* \* \*