

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 October 2002 (10.10.2002)

PCT

(10) International Publication Number
WO 02/078437 A2

- (51) International Patent Classification⁷: **A01N**
- (21) International Application Number: PCT/US02/10264
- (22) International Filing Date: 1 April 2002 (01.04.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/280,025 30 March 2001 (30.03.2001) US
60/336,657 4 December 2001 (04.12.2001) US
- (71) Applicant (for all designated States except US): **SYNGENTA PARTICIPATIONS AG** [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **MILES, Paul** [US/US]; 3054 Cornwallis Road, Research Triangle Park, NC 27709 (US). **KRAMER, Vance** [US/US]; 3054 Cornwallis Road, Research Triangle Park, NC 27709 (US). **SHEN, Zhicheng** [CN/US]; 3054 Cornwallis Road, Research Triangle Park, NC 27709 (US). **SHOTKOSKI, Frank** [US/US]; 3054 Cornwallis Road, Research Triangle Park, NC 27709 (US). **WARREN, Gregory, W.** [US/US]; 3054 Cornwallis Road, Research Triangle Park, NC 27709 (US).
- (74) Agent: **WARREN, Gregory, W.**; 3054 Cornwallis Road, Research Triangle Park, NC 27709 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/078437 A2

(54) Title: NOVEL PESTICIDAL TOXINS

(57) Abstract: A novel pesticidal toxin that is highly active against a wide range of lepidopteran insect pests is disclosed. The DNA encoding the pesticidal toxin can be used to transform various prokaryotic and eukaryotic organisms to express the pesticidal toxin. These recombinant organisms can be used to control lepidopteran insects in various environment.

NOVEL PESTICIDAL TOXINS

FIELD OF THE INVENTION

[0001] The present invention relates to novel Vip3 toxins from *Bacillus thuringiensis*, nucleic acid sequences whose expression results in said toxins, and methods of making and methods of using the toxins and corresponding nucleic acid sequences to control insects.

BACKGROUND

[0002] Plant pests are a major factor in the loss of the world's important agricultural crops. About \$8 billion are lost every year in the U.S. alone due to infestations of non-mammalian pests including insects. In addition to losses in field crops, insect pests are also a burden to vegetable and fruit growers, to producers of ornamental flowers, and to home gardeners.

[0003] Insect pests are mainly controlled by intensive applications of chemical pesticides, which are active through inhibition of insect growth, prevention of insect feeding or reproduction, or cause death. Good insect control can thus be reached, but these chemicals can sometimes also affect other, beneficial insects. Another problem resulting from the wide use of chemical pesticides is the appearance of resistant insect varieties. This has been partially alleviated by various resistance management practices, but there is an increasing need for alternative pest control agents. Biological pest control agents, such as *Bacillus thuringiensis* strains expressing pesticidal toxins like δ -endotoxins, have also been applied to crop plants with satisfactory results, offering an alternative or complement to chemical pesticides. The genes coding for some of these δ -endotoxins have been isolated and their expression in heterologous hosts have been shown to provide another tool for the control of economically important insect pests. In particular, the expression of insecticidal toxins in transgenic plants, such as *Bacillus thuringiensis* δ -endotoxins, has provided efficient protection against selected insect pests, and transgenic plants expressing

such toxins have been commercialized, allowing farmers to reduce applications of chemical insect control agents.

[0004] Other, non-endotoxin genes and the proteins they encode have now been identified. United States Patents 5,877,012, 6,107,279, 6,137,033, and 6,291,156, as well as Estruch *et al.* (1996, Proc. Natl. Acad. Sci. 93:5389-5394) and Yu *et al.* (1997, Appl. Environ. Microbiol. 63:532-536), all herein incorporated by reference, describe a new class of insecticidal proteins called Vip3. Vip3 coding sequences encode approximately 88 kDa proteins that are produced and secreted by *Bacillus* during its vegetative stages of growth (vegetative insecticidal proteins, VIP). The Vip3A protein possesses insecticidal activity against a wide spectrum of lepidopteran pests, including, but not limited to, black cutworm (BCW, *Agrotis ipsilon*), fall armyworm (FAW, *Spodoptera frugiperda*), tobacco budworm (TBW, *Heliothis virescens*), and corn earworm (CEW, *Helicoverpa zea*). More recently, plants expressing the Vip3A protein have been found to be resistant to feeding damage caused by hemipteran insect pests. Thus, the Vip3A protein displays a unique spectrum of insecticidal activities. Other disclosures, WO 98/18932, WO 98/33991, WO 98/00546, and WO 99/57282, have also now identified homologues of the Vip3 class of proteins.

[0005] The continued use of chemical and biological control methods heightens the chance for insects to develop resistance to such control measures. Also, only a few specific insect pests are controllable with current measures.

[0006] Therefore, there remains a need to discover new and effective pest control agents that provide an economic benefit to farmers and that are environmentally acceptable. Particularly needed are control agents that are targeted to a wider spectrum of economically important insect pests and that efficiently control insect strains that are or could become resistant to existing insect control agents. Furthermore, agents whose application minimizes the burden on the environment are desirable.

SUMMARY

[0007] The present invention addresses the need for novel pest control agents by providing new genes and toxins that are distinct from those disclosed in U.S. Patents

5,877,012, 6,107,279, and 6,137,033, and Estruch *et al.* (1996), and Yu *et al.* (1997), as well as WO 98/18932, WO 99/33991, WO 99/5782, and WO 98/00546.

[0008] Within the present invention, compositions and methods for controlling plant pests are provided. In particular, novel *vip3* nucleic acid sequences isolated from *Bacillus thuringiensis*, and sequences substantially identical thereto, whose expression results in pesticidal toxins with high specific toxicity to economically important insect pests, particularly insect pests that infest plants, are provided. The invention is further drawn to the novel pesticidal toxins resulting from the expression of the nucleic acid sequences, and to compositions and formulations containing the pesticidal toxins, which are capable of inhibiting the ability of insect pests to survive, grow and reproduce, or of limiting insect-related damage or loss to crop plants. The invention is also drawn to methods of using the nucleic acid sequences, for example in making hybrid toxins with enhanced pesticidal activity or in a recombinogenic procedure such as DNA shuffling. The invention is further drawn to a method of making the toxins and to methods of using the nucleic acid sequences, for example in microorganisms to control insects or in transgenic plants to confer protection from insect damage, and to a method of using the pesticidal toxins, and compositions and formulations comprising the pesticidal toxins, for example applying the pesticidal toxins or compositions or formulations to insect-infested areas, or to prophylactically treat insect-susceptible areas or plants to confer protection against the insect pests.

[0009] The novel pesticidal toxins described herein are highly active against insects. For example, a number of economically important insect pests, such as the lepidopterans *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Diatraea grandiosella* (southwestern corn borer), *Diatraea saccharalis* (sugarcane borer), *Sesamia nonagroides* (mediterranean corn borer), *Helicoverpa punctigera* (native budworm) and *Helicoverpa armigera* (cotton bollworm) can be controlled by the pesticidal toxins. The pesticidal toxins can be used singly or in combination with other insect control strategies to confer maximal pest control efficiency with minimal environmental impact.

[0010] According to one aspect, the present invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a toxin that is active against

insects, wherein the nucleotide sequence: (a) has at least 92% sequence identity with SEQ ID NO: 1; or (b) is isocoding with the nucleotide sequence of (a); or (c) encodes an amino acid sequence that has at least 91% sequence identity with SEQ ID NO: 3.

- [0011] In one embodiment of this aspect, the isolated nucleic acid molecule comprises a nucleotide sequence that has at least 92% sequence identity with SEQ ID NO: 1.
- [0012] In another embodiment of this aspect, the isolated nucleic acid molecule comprises a nucleotide sequence that is isocoding with a nucleotide sequence that has at least 92% sequence identity with SEQ ID NO: 1.
- [0013] In a further embodiment, the isolated nucleic acid molecule comprises the nucleotide sequence set forth in SEQ ID NO: 1 or SEQ ID NO: 3.
- [0014] In another embodiment of this aspect, the isolated nucleic acid molecule comprises a nucleotide sequence that encodes an amino acid sequence with at least 91% sequence identity with SEQ ID NO: 2. In a further embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence that encodes the amino acid sequence set forth in SEQ ID NO: 3.
- [0015] In one embodiment, the isolated nucleic acid molecule comprises the approximately 2.4 kb DNA fragment comprised in pNOV1325 harbored in *E. coli* strain DH5 α , designated as ATCC PTA-3868. In another embodiment, the isolated nucleic acid molecule comprises the approximately 2.4 kb DNA fragment comprised in pNOV1328 harbored in *E. coli* DH5 α , designated as ATCC PTA-3869.
- [0016] According to one embodiment of the invention, the isolated nucleic acid molecule encodes a toxin that is active against a lepidopteran insect. In a further embodiment, the lepidopteran insect is selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth).
- [0017] The present invention also provides a chimeric gene comprising a heterologous promoter sequence operatively linked to the nucleic acid molecule of the invention. Further, the present invention provides a recombinant vector comprising such a chimeric

gene. The present invention also provides a virus comprising such a chimeric gene. A virus according to this aspect of the invention may be an animal virus or a plant virus. Still further, the present invention provides a transgenic host cell comprising such a chimeric gene. A transgenic host cell according to this aspect of the invention may be an animal cell, a bacterial cell, a yeast cell or a plant cell, preferably, a plant cell. Even further, the present invention provides a transgenic plant comprising such a plant cell. A transgenic plant according to this aspect of the invention may be sorghum, wheat, sunflower, tomato, cole crops, cotton, rice, soybean, sugar beet, sugarcane, tobacco, barley, oilseed rape or maize, preferably maize. Still further, the present invention provides seed from the group of transgenic plants consisting of sorghum, wheat, sunflower, tomato, cole crops, cotton, rice, soybean, sugar beet, sugarcane, tobacco, barley, oilseed rape and maize. In a further embodiment, the seed is from a transgenic maize plant.

[0018] Also provided by the present invention are transgenic plants further comprising a second nucleic acid sequence or groups of nucleic acid sequences that encode a second pesticidal principle. Particularly preferred second nucleic acid sequences are those that encode a δ -endotoxin, those that encode another Vegetative Insecticidal Protein toxin or those that encode a pathway for the production of a non-proteinaceous pesticidal principle.

[0019] According to one aspect, the present invention provides an isolated toxin that is active against insects, wherein the toxin comprises an amino acid sequence that: (a) has at least 91% sequence identity with SEQ ID NO: 2; or (b) is produced by the expression of a nucleic acid molecule comprising a nucleotide sequence that has at least 92% sequence identity with SEQ ID NO: 1.

[0020] In one embodiment of this aspect, the isolated toxin comprises an amino acid sequence that has at least 91% sequence identity with SEQ ID NO: 2.

[0021] In a further embodiment, the isolated toxin comprises the amino acid sequence set forth in SEQ ID NO: 2.

[0022] In another embodiment of this aspect, the isolated toxin is produced by the expression of a nucleic acid molecule comprising a nucleotide sequence that has at least 92% sequence identity with SEQ ID NO: 1.

[0023] In yet another embodiment, the isolated toxin is produced by the expression of a nucleic acid molecule comprising the nucleotide sequence set forth in SEQ ID NO: 1 or SEQ ID NO: 3.

[0024] In another embodiment, the toxins of the invention are active against lepidopteran insects. In a further embodiment, the toxins are active against *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth).

[0025] In a further embodiment, the toxins are produced by the *E. coli* strain deposited as ATCC accession PTA-3868 or the *E. coli* strain deposited as ATCC accession PTA-3869.

[0026] The present invention also provides a composition comprising an effective insect-controlling amount of a toxin according to the invention.

[0027] In another aspect, the present invention provides a method of producing a toxin that is active against insects, comprising: (a) obtaining a transgenic host cell comprising a chimeric gene, which itself comprises a heterologous promoter sequence operatively linked to the nucleic acid molecule of the invention; and (b) expressing the nucleic acid molecule in the transgenic cell, which results in at least one toxin that is active against insects.

[0028] In a further aspect, the present invention provides a method of producing an insect-resistant transgenic plant, comprising introducing a nucleic acid molecule of the invention into the transgenic plant, wherein the nucleic acid molecule is expressible in the transgenic plant in an effective amount to control insects. According to one embodiment, the insects are lepidopteran insects. In a further embodiment, the lepidopteran insects are selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth).

[0029] In a still further aspect, the present invention provides a method of controlling insects comprising delivering to the insects an effective amount of a toxin of the present invention. According to one embodiment, the insects are lepidopteran insects. In a further

embodiment, the lepidopteran insects are selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth). In another embodiment, the toxin is delivered to the insects orally. In a still further embodiment, the toxin is delivered orally through a transgenic plant comprising a nucleic acid sequence that expresses a toxin of the present invention.

[0030] The present invention also provides hybrid toxins active against insects, wherein the hybrid toxins are encoded by a nucleic acid molecule comprising a nucleotide sequence according to the invention.

[0031] In one embodiment, the hybrid toxins of the invention are active against lepidopteran insects. In a further embodiment, the lepidopteran insects are selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth)

[0032] In another embodiment, the hybrid toxin is encoded by the nucleotide sequence set forth in SEQ ID NO: 6.

[0033] The present invention also provides a composition comprising an insecticidally effective amount of a hybrid toxin according to the invention.

[0034] In another aspect, the present invention provides a method of producing a hybrid toxin active against insects, comprising: (a) obtaining a transgenic host cell comprising a chimeric gene, which itself comprises a heterologous promoter sequence operatively linked to the nucleic acid molecule of the invention; and (b) expressing the nucleic acid molecule in the transgenic cell, which results in at least one hybrid toxin that is active against insects.

[0035] In a further aspect, the present invention provides a method of producing an insect-resistant transgenic plant, comprising introducing a nucleic acid molecule of the invention into the plant, wherein the nucleic acid molecule encodes a hybrid toxin and wherein the hybrid toxin is expressible in the transgenic plant in an effective amount to control an insect. According to one embodiment, the insects are lepidopteran insects. In another embodiment, the lepidopteran insect is selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Pectinophora gossypiella* (pink boll worm), *Trichoplusia ni* (cabbage looper), *Cochyles hospes* (banded sunflower moth), and *Homoeosoma electellum* (sunflower head moth).

[0036] In a still further aspect, the present invention provides a method of controlling an insect comprising delivering to the insects an effective amount of a hybrid toxin of the present invention. According to one embodiment, the insects are lepidopteran insects. In a further embodiment, the lepidopteran insects are selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Pectinophora gossypiella* (pink boll worm), *Trichoplusia ni* (cabbage looper), *Cochyles hospes* (banded sunflower moth), and *Homoeosoma electellum* (sunflower head moth). In another embodiment, the hybrid toxin is delivered to the insects orally. In a further embodiment, the hybrid toxin is delivered orally through a transgenic plant comprising a nucleic acid sequence that expresses a hybrid toxin of the present invention.

[0037] The present invention also provides a hybrid toxin active against insects, comprising a carboxy-terminal region of a Vip3 toxin joined in the amino to carboxy direction to an amino-terminal region of a different Vip3 toxin, wherein the carboxy-terminal region comprises an amino acid sequence which has at least 75% identity with amino acids 579-787 of SEQ ID NO: 2; and wherein the amino-terminal region has at least 75% identity with amino acids 1-578 of SEQ ID NO: 4. In a further embodiment, the carboxy-terminal region comprises amino acids 578-787 of SEQ ID NO: 2, and the amino-terminal region comprises amino acids 1-579 of SEQ ID NO: 5. In a still further embodiment, the hybrid toxin comprises amino acids 1-787 of SEQ ID NO: 7.

- [0038] The hybrid toxin, according to this aspect of the invention, is active against lepidopteran insects. In a further embodiment, the lepidopteran insects are selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Pectinophora gossypiella* (pink boll worm), *Trichoplusia ni* (cabbage looper), *Cochyles hospes* (banded sunflower moth), and *Homoeosoma electellum* (sunflower head moth).
- [0039] Also encompassed by this aspect of the invention is a nucleic acid molecule comprising a nucleotide sequence that encodes the hybrid toxin of this aspect.
- [0040] Also provided by the invention is a method of controlling insects wherein the transgenic plant further comprises a second nucleic acid sequence or groups of nucleic acid sequences that encode a second pesticidal principle. Particularly preferred second nucleic acid sequences are those that encode a δ -endotoxin, those that encode another Vegetative Insecticidal Protein toxin or those that encode a pathway for the production of a non-proteinaceous pesticidal principle.
- [0041] Yet another aspect of the present invention is the provision of a method for mutagenizing a nucleic acid molecule according to the present invention, wherein the nucleic acid molecule has been cleaved into populations of double-stranded random fragments of a desired size, comprising: (a) adding to the population of double-stranded random fragments one or more single- or double-stranded oligonucleotides, wherein the oligonucleotides each comprise an area of identity and an area of heterology to a double-stranded template polynucleotide; (b) denaturing the resultant mixture of double-stranded random fragments and oligonucleotides into single-stranded fragments; (c) incubating the resultant population of single-stranded fragments with polymerase under conditions which result in the annealing of the single-stranded fragments at the areas of identity to form pairs of annealed fragments, the areas of identity being sufficient for one member of the pair to prime replication of the other, thereby forming a mutagenized double-stranded polynucleotide; and (d) repeating the second and third steps for at least two further cycles, wherein the resultant mixture in the second step of a further cycle includes the mutagenized double-stranded polynucleotide from the third step of the previous cycle, and wherein the further cycle forms a further mutagenized double-stranded polynucleotide.

[0042] Other aspects and advantages of the present invention will become apparent to those skilled in the art from a study of the following description of the invention and non-limiting examples.

BRIEF DESCRIPTION OF THE SEQUENCES IN THE SEQUENCE LISTING

SEQ ID NO: 1 is the coding sequence of the native *vip3B* gene.

SEQ ID NO: 2 is the amino acid sequence encoded SEQ ID NO: 1.

SEQ ID NO: 3 is the coding sequence of the maize optimized *vip3B* gene.

SEQ ID NO: 4 is the coding sequence of the native *vip3A* gene.

SEQ ID NO: 5 is the amino acid sequence encoded by SEQ ID NO: 4.

SEQ ID NO: 6 is the coding sequence of the hybrid *vip3A-B* gene.

SEQ ID NO: 7 is the amino acid sequence encoded by SEQ ID NO: 6.

SEQ ID NO: 8-13 are primers useful in the present invention.

DEPOSITS

The following material was deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA, under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. All restrictions on the availability of the deposited material will be irrevocably removed upon granting of the patent.

<u>Clone</u>	<u>Accession Number</u>	<u>Date of Deposit</u>
pNOV1325	ATCC No. PTA-3868	November 16, 2001
pNOV1328	ATCC No. PTA-3869	November 16, 2001

DEFINITIONS

[0043] "Activity" of the toxins of the invention is meant that the toxins function as orally active insect control agents, have a toxic effect, or are able to disrupt or deter insect feeding, which may or may not cause death of the insect. When a toxin of the invention is

delivered to the insect, the result is typically death of the insect, or the insect does not feed upon the source that makes the toxin available to the insect.

[0044] "Associated with / operatively linked" refer to two nucleic acid sequences that are related physically or functionally. For example, a promoter or regulatory DNA sequence is said to be "associated with" a DNA sequence that codes for an RNA or a protein if the two sequences are operatively linked, or situated such that the regulator DNA sequence will affect the expression level of the coding or structural DNA sequence.

[0045] A "chimeric gene" is a recombinant nucleic acid sequence in which a promoter or regulatory nucleic acid sequence is operatively linked to, or associated with, a nucleic acid sequence that codes for an mRNA or which is expressed as a protein, such that the regulator nucleic acid sequence is able to regulate transcription or expression of the associated nucleic acid sequence. The regulator nucleic acid sequence of the chimeric gene is not normally operatively linked to the associated nucleic acid sequence as found in nature.

[0046] A "coding sequence" is a nucleic acid sequence that is transcribed into RNA such as mRNA, rRNA, tRNA, snRNA, sense RNA or antisense RNA. Preferably the RNA is then translated in an organism to produce a protein.

[0047] To "control" insects means to inhibit, through a toxic effect, the ability of insect pests to survive, grow, feed, and/or reproduce, or to limit insect-related damage or loss in crop plants. To "control" insects may or may not mean killing the insects, although it preferably means killing the insects.

[0048] To "deliver" a toxin means that the toxin comes in contact with an insect, resulting in toxic effect and control of the insect. The toxin can be delivered in many recognized ways, e.g., orally by ingestion by the insect or by contact with the insect via transgenic plant expression, formulated protein composition(s), sprayable protein composition(s), a bait matrix, or any other art-recognized toxin delivery system.

[0049] "Effective insect-controlling amount" means that concentration of toxin that inhibits, through a toxic effect, the ability of insects to survive, grow, feed and/or reproduce, or to limit insect-related damage or loss in crop plants. "Effective insect-controlling amount" may or may not mean killing the insects, although it preferably means killing the insects.

[0050] "Expression cassette" as used herein means a nucleic acid sequence capable of directing expression of a particular nucleotide sequence in an appropriate host cell,

comprising a promoter operably linked to the nucleotide sequence of interest which is operably linked to termination signals. It also typically comprises sequences required for proper translation of the nucleotide sequence. The expression cassette comprising the nucleotide sequence of interest may be chimeric, meaning that at least one of its components is heterologous with respect to at least one of its other components. The expression cassette may also be one that is naturally occurring but has been obtained in a recombinant form useful for heterologous expression. Typically, however, the expression cassette is heterologous with respect to the host, i.e., the particular nucleic acid sequence of the expression cassette does not occur naturally in the host cell and must have been introduced into the host cell or an ancestor of the host cell by a transformation event. The expression of the nucleotide sequence in the expression cassette may be under the control of a constitutive promoter or of an inducible promoter that initiates transcription only when the host cell is exposed to some particular external stimulus. In the case of a multicellular organism, such as a plant, the promoter can also be specific to a particular tissue, or organ, or stage of development.

[0051] A "gene" is a defined region that is located within a genome and that, besides the aforementioned coding nucleic acid sequence, comprises other, primarily regulatory, nucleic acid sequences responsible for the control of the expression, that is to say the transcription and translation, of the coding portion. A gene may also comprise other 5' and 3' untranslated sequences and termination sequences. Further elements that may be present are, for example, introns.

[0052] "Gene of interest" refers to any gene which, when transferred to a plant, confers upon the plant a desired characteristic such as antibiotic resistance, virus resistance, insect resistance, disease resistance, or resistance to other pests, herbicide tolerance, improved nutritional value, improved performance in an industrial process or altered reproductive capability. The "gene of interest" may also be one that is transferred to plants for the production of commercially valuable enzymes or metabolites in the plant.

[0053] A "heterologous" nucleic acid sequence is a nucleic acid sequence not naturally associated with a host cell into which it is introduced, including non-naturally occurring multiple copies of a naturally occurring nucleic acid sequence.

[0054] A "homologous" nucleic acid sequence is a nucleic acid sequence naturally associated with a host cell into which it is introduced.

- [0055] "Homologous recombination" is the reciprocal exchange of nucleic acid fragments between homologous nucleic acid molecules.
- [0056] "Hybrid toxin" as used herein is an insecticidal toxin made by the hand of man which comprises amino acid regions or fragments of one toxin joined with amino acid regions or fragments from a different toxin, For example, without limitation, joining the C-terminal region of Vip3B, from amino acid number 579 to amino acid number 787 of SEQ ID NO: 2, with the N-terminal region of Vip3A, from amino acid number 1 to amino acid number 578 of SEQ ID NO: 5.
- [0057] "Insecticidal" is defined as a toxic biological activity capable of controlling insects, preferably by killing them.
- [0058] A nucleic acid sequence is "isocoding with" a reference nucleic acid sequence when the nucleic acid sequence encodes a polypeptide having the same amino acid sequence as the polypeptide encoded by the reference nucleic acid sequence.
- [0059] An "isolated" nucleic acid molecule or an isolated enzyme is a nucleic acid molecule or enzyme that, by the hand of man, exists apart from its native environment and is therefore not a product of nature. An isolated nucleic acid molecule or enzyme may exist in a purified form or may exist in a non-native environment such as, for example, a recombinant host cell.
- [0060] A "nucleic acid molecule" or "nucleic acid sequence" is a linear segment of single- or double-stranded DNA or RNA that can be isolated from any source. In the context of the present invention, the nucleic acid molecule is preferably a segment of DNA.
- [0061] A "plant" is any plant at any stage of development, particularly a seed plant.
- [0062] A "plant cell" is a structural and physiological unit of a plant, comprising a protoplast and a cell wall. The plant cell may be in form of an isolated single cell or a cultured cell, or as a part of higher organized unit such as, for example, plant tissue, a plant organ, or a whole plant.
- [0063] "Plant cell culture" means cultures of plant units such as, for example, protoplasts, cell culture cells, cells in plant tissues, pollen, pollen tubes, ovules, embryo sacs, zygotes and embryos at various stages of development.
- [0064] "Plant material" refers to leaves, stems, roots, flowers or flower parts, fruits, pollen, egg cells, zygotes, seeds, cuttings, cell or tissue cultures, or any other part or product of a plant.

- [0065] A "plant organ" is a distinct and visibly structured and differentiated part of a plant such as a root, stem, leaf, flower bud, or embryo.
- [0066] "Plant tissue" as used herein means a group of plant cells organized into a structural and functional unit. Any tissue of a plant in planta or in culture is included. This term includes, but is not limited to, whole plants, plant organs, plant seeds, tissue culture and any groups of plant cells organized into structural and/or functional units. The use of this term in conjunction with, or in the absence of, any specific type of plant tissue as listed above or otherwise embraced by this definition is not intended to be exclusive of any other type of plant tissue.
- [0067] A "promoter" is an untranslated DNA sequence upstream of the coding region that contains the binding site for RNA polymerase II and initiates transcription of the DNA. The promoter region may also include other elements that act as regulators of gene expression.
- [0068] A "protoplast" is an isolated plant cell without a cell wall or with only parts of the cell wall.
- [0069] "Regulatory elements" refer to sequences involved in controlling the expression of a nucleotide sequence. Regulatory elements comprise a promoter operably linked to the nucleotide sequence of interest and termination signals. They also typically encompass sequences required for proper translation of the nucleotide sequence.
- [0070] A "shuffled" nucleic acid is a nucleic acid produced by a shuffling procedure such as any shuffling procedure set forth herein. Shuffled nucleic acids are produced by recombining (physically or virtually) two or more nucleic acids (or character strings), e.g., in an artificial, and optionally recursive, fashion. Generally, one or more screening steps are used in shuffling processes to identify nucleic acids of interest; this screening step can be performed before or after any recombination step. In some (but not all) shuffling embodiments, it is desirable to perform multiple rounds of recombination prior to selection to increase the diversity of the pool to be screened. The overall process of recombination and selection are optionally repeated recursively. Depending on context, shuffling can refer to an overall process of recombination and selection, or, alternately, can simply refer to the recombinational portions of the overall process.
- [0071] Substantially identical: the phrase "substantially identical," in the context of two nucleic acid or protein sequences, refers to two or more sequences or subsequences that have at least 60%, preferably 80%, more preferably 90, even more preferably 95%, and

most preferably at least 99% nucleotide or amino acid residue identity, when compared and aligned for maximum correspondence, as measured using one of the following sequence comparison algorithms or by visual inspection. Preferably, the substantial identity exists over a region of the sequences that is at least about 50 residues in length, more preferably over a region of at least about 100 residues, and most preferably the sequences are substantially identical over at least about 150 residues. In an especially preferred embodiment, the sequences are substantially identical over the entire length of the coding regions. Furthermore, substantially identical nucleic acid or protein sequences perform substantially the same function.

[0072] For sequence comparison, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

[0073] Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2: 482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48: 443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (*see generally*, Ausubel *et al.*, *infra*).

[0074] One example of an algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul *et al.*, *J. Mol. Biol.* 215: 403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, 1990). These initial neighborhood word hits act as seeds

for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when the cumulative alignment score falls off by the quantity X from its maximum achieved value, the cumulative score goes to zero or below due to the accumulation of one or more negative-scoring residue alignments, or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (*see* Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89: 10915 (1989)).

[0075] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (*see, e.g.,* Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90: 5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a test nucleic acid sequence is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid sequence to the reference nucleic acid sequence is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

[0076] Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions. The phrase "hybridizing specifically to" refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent conditions when that sequence is present in a complex mixture (*e.g.,* total cellular) DNA or RNA. "Bind(s) substantially" refers to complementary hybridization between a probe nucleic acid and a target nucleic acid and embraces minor mismatches that can be accommodated by reducing the stringency of the hybridization media to achieve the desired detection of the target nucleic acid sequence.

[0077] "Stringent hybridization conditions" and "stringent hybridization wash conditions" in the context of nucleic acid hybridization experiments such as Southern and Northern hybridizations are sequence dependent, and are different under different environmental parameters. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes* part I chapter 2 "Overview of principles of hybridization and the strategy of nucleic acid probe assays" Elsevier, New York. Generally, highly stringent hybridization and wash conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. Typically, under "stringent conditions" a probe will hybridize to its target subsequence, but to no other sequences.

[0078] The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Very stringent conditions are selected to be equal to the T_m for a particular probe. An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is 50% formamide with 1 mg of heparin at 42°C, with the hybridization being carried out overnight. An example of highly stringent wash conditions is 0.1 5M NaCl at 72°C for about 15 minutes. An example of stringent wash conditions is a 0.2x SSC wash at 65°C for 15 minutes (*see, Sambrook, infra*, for a description of SSC buffer). Often, a high stringency wash is preceded by a low stringency wash to remove background probe signal. An example medium stringency wash for a duplex of, e.g., more than 100 nucleotides, is 1x SSC at 45°C for 15 minutes. An example low stringency wash for a duplex of, e.g., more than 100 nucleotides, is 4-6x SSC at 40°C for 15 minutes. For short probes (*e.g.*, about 10 to 50 nucleotides), stringent conditions typically involve salt concentrations of less than about 1.0 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3, and the temperature is typically at least about 30°C. Stringent conditions can also be achieved with the addition of destabilizing agents such as formamide. In general, a signal to noise ratio of 2x (or higher) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization. Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the proteins that they encode are substantially identical. This occurs, e.g., when

a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code.

[0079] The following are examples of sets of hybridization/wash conditions that may be used to clone homologous nucleotide sequences that are substantially identical to reference nucleotide sequences of the present invention: a reference nucleotide sequence preferably hybridizes to the reference nucleotide sequence in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 2X SSC, 0.1% SDS at 50°C, more desirably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 1X SSC, 0.1% SDS at 50°C, more desirably still in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.5X SSC, 0.1% SDS at 50°C, preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 50°C, more preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 65°C.

[0080] A further indication that two nucleic acid sequences or proteins are substantially identical is that the protein encoded by the first nucleic acid is immunologically cross reactive with, or specifically binds to, the protein encoded by the second nucleic acid. Thus, a protein is typically substantially identical to a second protein, for example, where the two proteins differ only by conservative substitutions.

[0081] "Synthetic" refers to a nucleotide sequence comprising structural characters that are not present in the natural sequence. For example, an artificial sequence that resembles more closely the G+C content and the normal codon distribution of dicot and/or monocot genes is said to be synthetic.

[0082] "Transformation" is a process for introducing heterologous nucleic acid into a host cell or organism. In particular, "transformation" means the stable integration of a DNA molecule into the genome of an organism of interest.

[0083] "Transformed / transgenic / recombinant" refer to a host organism such as a bacterium or a plant into which a heterologous nucleic acid molecule has been introduced. The nucleic acid molecule can be stably integrated into the genome of the host or the nucleic acid molecule can also be present as an extrachromosomal molecule. Such an extrachromosomal molecule can be auto-replicating. Transformed cells, tissues, or plants are understood to encompass not only the end product of a transformation process, but also transgenic progeny thereof. A "non-transformed", "non-transgenic", or "non-

recombinant" host refers to a wild-type organism, e.g., a bacterium or plant, which does not contain the heterologous nucleic acid molecule.

[0084] The "Vip3 class of proteins" comprises, for example, without limitation, Vip3A(a), Vip3A(b), Vip3A(c), Vip3B, Vip3C(a), Vip3C(b), Vip3Z, and their homologues. "Homologue" is used herein to mean that the indicated protein or polypeptide bears a defined relationship to other members of the Vip3 class of proteins. This defined relationship includes but is not limited to, 1) proteins which are at least 70%, more preferably at least 80% and most preferably at least 90% identical at the sequence level to another member of the Vip3 class of proteins while also retaining pesticidal activity, 2) proteins which are cross-reactive to antibodies which immunologically recognize another member of the Vip3 class of proteins, 3) proteins which are cross-reactive with a receptor to another member of the Vip3 class of proteins and retain the ability to induce programmed cell death, and 4) proteins which are at least 70%, more preferably at least 80% and most preferably at least 90% identical at the sequence level to the toxic core region of another member of the Vip3 class of proteins while also retaining pesticidal activity. Other Vip3 homologues have been disclosed in WO 98/18932, WO 98/33991, WO 98/00546, and WO 99/57282

[0085] Nucleotides are indicated by their bases by the following standard abbreviations: adenine (A), cytosine (C), thymine (T), and guanine (G). Amino acids are likewise indicated by the following standard abbreviations: alanine (Ala; A), arginine (Arg; R), asparagine (Asn; N), aspartic acid (Asp; D), cysteine (Cys; C), glutamine (Gln; Q), glutamic acid (Glu; E), glycine (Gly; G), histidine (His; H), isoleucine (Ile; I), leucine (Leu; L), lysine (Lys; K), methionine (Met; M), phenylalanine (Phe; F), proline (Pro; P), serine (Ser; S), threonine (Thr; T), tryptophan (Trp; W), tyrosine (Tyr; Y), and valine (Val; V).

DETAILED DESCRIPTION

[0086] This invention relates to nucleic acid sequences whose expression results in novel toxins, and to the making and using of the toxins to control insect pests. The nucleic acid sequences are derived from *Bacillus*, a gram-positive spore-forming microorganism. In particular, novel Vip3 proteins, useful as pesticidal agents, are provided.

[0087] For purposes of the present invention, insect pests include insects selected from the orders Coleoptera, Diptera, Hymenoptera, Lepidoptera, Mallophaga, Homoptera, Hemiptera, Orthoptera, Thysanoptera, Dermaptera, Isoptera, Anoplura, Siphonaptera, Trichoptera, etc., particularly Lepidoptera.

[0088] The expression of the nucleic acid sequences of the present invention results in toxins that can be used to control lepidopteran insects such as *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth).

[0089] In one embodiment, the invention encompasses an isolated nucleic acid molecule comprising a nucleotide sequence that has at least 92% sequence identity with SEQ ID NO: 1, wherein expression of the isolated nucleic acid molecule results in insect control activity. When expressed in a heterologous host, the nucleic acid molecule of SEQ ID NO: 1 results in insect control activity against *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth), showing that the nucleotide sequence set forth in SEQ ID NO: 1 is sufficient for such insect control activity. In a further embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence that has at least 93% sequence identity with SEQ ID NO: 1. In a still further embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence that has at least 94% sequence identity with SEQ ID NO: 1. In a still further embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence that has at least 95% sequence identity with SEQ ID NO: 1. In a still further embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence that has at least 96% sequence identity with SEQ ID NO: 1. In a still further embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence that has at least 97% sequence identity with SEQ ID NO: 1. In a still further

embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence that has at least 98% sequence identity with SEQ ID NO: 1. In a still further embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence that has at least 99% sequence identity with SEQ ID NO: 1. In a still further embodiment, the isolated nucleic acid molecule comprises the nucleotide sequence set forth in SEQ ID NO: 1.

[0090] In another embodiment, the invention encompasses a nucleic acid molecule comprised in pNOV1325, whose expression results in an insecticidal toxin, which is deposited in the *E. coli* strain DH5 α designated ATCC accession number PTA-3868.

[0091] In one embodiment, the invention encompasses an isolated nucleic acid molecule comprising a nucleotide sequence that is isocoding with a nucleotide sequence that has at least 92% sequence identity with SEQ ID NO: 1, wherein expression of the isolated nucleic acid molecule results in insect control activity. When expressed in a heterologous host, the nucleic acid molecule of SEQ ID NO: 3 results in insect control activity against *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm) and *Helicoverpa armigera* (cotton bollworm), showing that the nucleotide sequence set forth in SEQ ID NO: 3 is sufficient for such insect control activity. In a further embodiment, the invention encompasses a nucleic acid molecule comprised in pNOV1328, whose expression results in an insecticidal toxin, which is deposited in the *E. coli* strain DH5 α designated ATCC accession number PTA-3869.

[0092] In another embodiment, the isolated nucleic acid molecule encodes a toxin comprising an amino acid sequence which has at least 91% sequence identity to the amino acid sequence set forth in SEQ ID NO: 2. In a further embodiment, the isolated nucleic acid molecule encodes a toxin comprising an amino acid sequence which has at least 92% sequence identity to the amino acid sequence set forth in SEQ ID NO: 2. In a still further embodiment, the isolated nucleic acid molecule encodes a toxin comprising an amino acid sequence which has at least 93% sequence identity to the amino acid sequence set forth in SEQ ID NO: 2. In a still further embodiment, the isolated nucleic acid molecule encodes a toxin comprising an amino acid sequence which has at least 94% sequence identity to the amino acid sequence set forth in SEQ ID NO: 2. In a still further embodiment, the isolated

nucleic acid molecule encodes a toxin comprising an amino acid sequence which has at least 95% sequence identity to the amino acid sequence set forth in SEQ ID NO: 2. In a still further embodiment, the isolated nucleic acid molecule encodes a toxin comprising an amino acid sequence which has at least 96% sequence identity to the amino acid sequence set forth in SEQ ID NO: 2. In a still further embodiment, the isolated nucleic acid molecule encodes a toxin comprising an amino acid sequence which has at least 97% sequence identity to the amino acid sequence set forth in SEQ ID NO: 2. In a still further embodiment, the isolated nucleic acid molecule encodes a toxin comprising an amino acid sequence which has at least 98% sequence identity to the amino acid sequence set forth in SEQ ID NO: 2. In a still further embodiment, the isolated nucleic acid molecule encodes a toxin comprising an amino acid sequence which has at least 99% sequence identity to the amino acid sequence set forth in SEQ ID NO: 2. In a still further embodiment, the isolated nucleic acid molecule encodes a toxin comprising the amino acid sequence set forth in SEQ ID NO: 2.

[0093] The present invention also encompasses recombinant vectors comprising the nucleic acid sequences of this invention. In such vectors, the nucleic acid sequences are preferably comprised in expression cassettes comprising regulatory elements for expression of the nucleotide sequences in a host cell capable of expressing the nucleotide sequences. Such regulatory elements usually comprise promoter and termination signals and preferably also comprise elements allowing efficient translation of polypeptides encoded by the nucleic acid sequences of the present invention. Vectors comprising the nucleic acid sequences are usually capable of replication in particular host cells, preferably as extrachromosomal molecules, and are therefore used to amplify the nucleic acid sequences of this invention in the host cells. In one embodiment, host cells for such vectors are microorganisms, such as bacteria, in particular *E. coli* or *Bacillus*. In another embodiment, host cells for such recombinant vectors are endophytes or epiphytes. A preferred host cell for such vectors is a eukaryotic cell, such as a yeast cell, a plant cell, or an insect cell. Plant cells such as maize cells are most preferred host cells. In another preferred embodiment, such vectors are viral vectors and are used for replication of the nucleotide sequences in particular host cells, e.g. insect cells or plant cells. Recombinant vectors are also used for transformation of the nucleotide sequences of this invention into host cells, whereby the nucleotide sequences are stably integrated into the DNA of such host cells. In one, such host cells are prokaryotic cells. In a preferred embodiment, such

host cells are eukaryotic cells, such as yeast cells, insect cells, or plant cells. In a most preferred embodiment, the host cells are plant cells, such as maize cells.

[0094] The present invention also provides a method of producing a toxin that is active against insects, comprising (a) obtaining a transgenic host cell according to the invention and (b) expressing a nucleic acid molecule of the invention in the transgenic host cell which results in at least one toxin that is active against insects.

[0095] The present invention further provides a method of producing an insect-resistant transgenic plant, comprising introducing a nucleic acid molecule of the invention into the transgenic plant, wherein the nucleic acid molecule is expressible in the transgenic plant in an effective amount to control insects. In a still further embodiment the insects are lepidopteran insects. In a still further embodiment, the lepidopteran insects are selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth).

[0096] In another aspect, the present invention provides an isolated toxin that is active against insects, wherein the toxin comprises an amino acid sequence that: (a) has at least 91% sequence identity with SEQ ID NO: 2; or (b) is encoded by a nucleic acid molecule comprising a nucleotide sequence that has at least 92% sequence identity with SEQ ID NO: 1.

[0097] In one embodiment, the present invention provides an isolated toxin that is active against insects, wherein the toxin comprises an amino acid sequence that has at least 91% sequence identity with SEQ ID NO: 2. In a further embodiment, the toxin comprises an amino acid sequence that has at least 92% sequence identity with SEQ ID NO: 2. In a still further embodiment, the toxin has at least 93% sequence identity with SEQ ID NO: 2. In a still further embodiment, the toxin has at least 94% sequence identity with SEQ ID NO: 2. In a still further embodiment, the toxin has at least 95% sequence identity with SEQ ID NO: 2. In a still further embodiment, the toxin has at least 96% sequence identity with SEQ ID NO: 2. In a still further embodiment, the toxin has at least 97% sequence identity with SEQ ID NO: 2. In a still further embodiment, the toxin has at least 98% sequence

identity with SEQ ID NO: 2. In a still further embodiment, the toxin has at least 99% sequence identity with SEQ ID NO: 2. In a still further embodiment, the toxin comprises the amino acid sequence set forth in SEQ ID NO: 2.

[0098] In one embodiment, the present invention provides a toxin active against insects, wherein the toxin is produced by the expression of a nucleic acid molecule comprising a nucleotide sequence that has at least 92% sequence identity with SEQ ID NO: 1. In a further embodiment, the nucleotide sequence has at least 93% sequence identity with SEQ ID NO: 1. In a still further embodiment, the nucleotide sequence has at least 94% sequence identity with SEQ ID NO: 1. In a still further embodiment, the nucleotide sequence has at least 95% sequence identity with SEQ ID NO: 1. In a still further embodiment, the nucleotide sequence has at least 96% sequence identity with SEQ ID NO: 1. In a still further embodiment, the nucleotide sequence has at least 97% sequence identity with SEQ ID NO: 1. In a still further embodiment, the nucleotide sequence has at least 98% sequence identity with SEQ ID NO: 1. In a still further embodiment, the nucleotide sequence has at least 99% sequence identity with SEQ ID NO: 1. In a still further embodiment, the toxin is produced by the expression of a nucleotide sequence comprising nucleotides 1-2364 of SEQ ID NO: 1 or nucleotides 1-2364 of SEQ ID NO: 3.

[0099] In one embodiment, a toxin of the present invention is produced by the expression of a nucleotide sequence comprising the approximately 2.4 kb DNA fragment comprised in pNOV1325, deposited as ATCC accession number PTA-3868. In another embodiment, a toxin of the present invention is produced by the expression of a nucleotide sequence comprising the approximately 2.4 kb DNA fragment comprised in pNOV1328, deposited as ATCC accession number PTA-3869.

[00100] In another embodiment, a toxin of the present invention is produced by the *E. coli* strain designated as ATCC accession PTA-3868. In still another embodiment, the toxin is produced by the *E. coli* strain designated as ATCC accession PTA-3869.

[00101] The toxins of the present invention have insect control activity when tested against insect pests in bioassays. In one embodiment, the toxins of the invention are active against lepidopteran insects. In a further embodiment, the lepidopteran insects are selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm),

Helicoverpa armigera (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth). The insect controlling properties of the insecticidal toxins of the invention are further illustrated in Example 5 and Example 8.

[00102] The present invention also encompasses hybrid toxins which are active against insects, wherein the hybrid toxins are encoded by nucleic acid molecules comprising a nucleotide sequence that: (a) has a complement that hybridizes to nucleotides 1734-2364 of SEQ ID NO: 1 in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50° C. with washing in 0.1XSSC, 0.1% SDS at 65°C.; or (b) is isocoding with the nucleotide sequence of (a); or (c) comprises a consecutive 20 base pair nucleotide portion identical in sequence to a consecutive 20 base pair nucleotide portion of a nucleotide sequence of (a) or (b), wherein expression of the nucleic acid molecule results in insect control activity. Specifically exemplified herein is a hybrid toxin that is encoded by the nucleotide sequence set forth in SEQ ID NO: 6. When expressed in a heterologous host, the nucleic acid molecule of SEQ ID NO: 6 results in insect control activity against *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Pectinophora gossypiella* (pink boll worm), *Trichoplusia ni* (cabbage looper), *Cochylis hospes* (banded sunflower moth), and *Homoeosoma electellum* (sunflower head moth).

[00103] The present invention also encompasses hybrid toxins active against insects that comprise a carboxy-terminal region of a Vip3 toxin joined in the amino to carboxy direction to an amino-terminal region of a different Vip3 toxin, wherein the carboxy-terminal region comprises an amino acid sequence which has at least 75% identity with amino acids 579-787 of SEQ ID NO: 2; and wherein the amino-terminal region has at least 75% identity with amino acids 1-578 of SEQ ID NO: 5. In a further embodiment, the carboxy-terminal region comprises amino acids 579-787 of SEQ ID NO: 2, and the amino-terminal region comprises amino acids 1-578 of SEQ ID NO: 5. In a still further embodiment, the hybrid toxin comprises amino acids 1-787 of SEQ ID NO: 6.

Expression of the Nucleotide Sequences in Heterologous Microbial Hosts

[00104] As biological insect control agents, the insecticidal toxins are produced by expression of the nucleotide sequences in heterologous host cells capable of expressing the nucleotide sequences. In a first embodiment, *B. thuringiensis* cells comprising modifications of a nucleotide sequence of this invention are made. Such modifications encompass mutations or deletions of existing regulatory elements, thus leading to altered expression of the nucleotide sequence, or the incorporation of new regulatory elements controlling the expression of the nucleotide sequence. In another embodiment, additional copies of one or more of the nucleotide sequences are added to *Bacillus thuringiensis* cells either by insertion into the chromosome or by introduction of extrachromosomally replicating molecules containing the nucleotide sequences.

[00105] In another embodiment, at least one of the nucleotide sequences of the invention is inserted into an appropriate expression cassette, comprising a promoter and termination signals. Expression of the nucleotide sequence is constitutive, or an inducible promoter responding to various types of stimuli to initiate transcription is used. In a preferred embodiment, the cell in which the toxin is expressed is a microorganism, such as a virus, a bacteria, or a fungus. In a preferred embodiment, a virus, such as a baculovirus, contains a nucleotide sequence of the invention in its genome and expresses large amounts of the corresponding insecticidal toxin after infection of appropriate eukaryotic cells that are suitable for virus replication and expression of the nucleotide sequence. The insecticidal toxin thus produced is used as an insecticidal agent. Alternatively, baculoviruses engineered to include the nucleotide sequence are used to infect insects *in vivo* and kill them either by expression of the insecticidal toxin or by a combination of viral infection and expression of the insecticidal toxin.

[00106] Bacterial cells are also hosts for the expression of the nucleotide sequences of the invention. In a preferred embodiment, non-pathogenic symbiotic bacteria, which are able to live and replicate within plant tissues, so-called endophytes, or non-pathogenic symbiotic bacteria, which are capable of colonizing the phyllosphere or the rhizosphere, so-called epiphytes, are used. Such bacteria include bacteria of the genera *Agrobacterium*, *Alcaligenes*, *Azospirillum*, *Azotobacter*, *Bacillus*, *Clavibacter*, *Enterobacter*, *Erwinia*, *Flavobacter*, *Klebsiella*, *Pseudomonas*, *Rhizobium*, *Serratia*, *Streptomyces* and *Xanthomonas*. Symbiotic fungi, such as *Trichoderma* and *Gliocladium* are also possible hosts for expression of the inventive nucleotide sequences for the same purpose.

[00107] Techniques for these genetic manipulations are specific for the different available hosts and are known in the art. For example, the expression vectors pKK223-3 and pKK223-2 can be used to express heterologous genes in *E. coli*, either in transcriptional or translational fusion, behind the *tac* or *trc* promoter. For the expression of operons encoding multiple ORFs, the simplest procedure is to insert the operon into a vector such as pKK223-3 in transcriptional fusion, allowing the cognate ribosome binding site of the heterologous genes to be used. Techniques for overexpression in gram-positive species such as *Bacillus* are also known in the art and can be used in the context of this invention (Quax et al. In:Industrial Microorganisms:Basic and Applied Molecular Genetics, Eds. Baltz et al., American Society for Microbiology, Washington (1993)). Alternate systems for overexpression rely for example, on yeast vectors and include the use of *Pichia*, *Saccharomyces* and *Kluyveromyces* (Sreekrishna, In:Industrial microorganisms:basic and applied molecular genetics, Baltz, Hegeman, and Skatrud eds., American Society for Microbiology, Washington (1993); Dequin & Barre, *Biotechnology L2*:173- 177 (1994); van den Berg et al., *Biotechnology 8*:135-139 (1990)).

Plant transformation

[00108] In a particularly preferred embodiment, at least one of the insecticidal toxins of the invention is expressed in a higher organism, e.g., a plant. In this case, transgenic plants expressing effective amounts of the toxins protect themselves from insect pests. When the insect starts feeding on such a transgenic plant, it also ingests the expressed toxins. This will deter the insect from further biting into the plant tissue or may even harm or kill the insect. A nucleotide sequence of the present invention is inserted into an expression cassette, which is then preferably stably integrated in the genome of said plant. In another preferred embodiment, the nucleotide sequence is included in a non- pathogenic self-replicating virus. Plants transformed in accordance with the present invention may be monocots or dicots and include, but are not limited to, maize, wheat, barley, rye, sweet potato, bean, pea, chicory, lettuce, cabbage, cauliflower, broccoli, turnip, radish, spinach, asparagus, onion, garlic, pepper, celery, squash, pumpkin, hemp, zucchini, apple, pear, quince, melon, plum, cherry, peach, nectarine, apricot, strawberry, grape, raspberry, blackberry, pineapple, avocado, papaya, mango, banana, soybean, tomato, sorghum,

sugarcane, sugar beet, sunflower, rapeseed, clover, tobacco, carrot, cotton, alfalfa, rice, potato, eggplant, cucumber, Arabidopsis, and woody plants such as coniferous and deciduous trees.

[00109] Once a desired nucleotide sequence has been transformed into a particular plant species, it may be propagated in that species or moved into other varieties of the same species, particularly including commercial varieties, using traditional breeding techniques.

[00110] A nucleotide sequence of this invention is preferably expressed in transgenic plants, thus causing the biosynthesis of the corresponding toxin in the transgenic plants. In this way, transgenic plants with enhanced resistance to insects are generated. For their expression in transgenic plants, the nucleotide sequences of the invention may require modification and optimization. Although in many cases genes from microbial organisms can be expressed in plants at high levels without modification, low expression in transgenic plants may result from microbial nucleotide sequences having codons that are not preferred in plants. It is known in the art that all organisms have specific preferences for codon usage, and the codons of the nucleotide sequences described in this invention can be changed to conform with plant preferences, while maintaining the amino acids encoded thereby. Furthermore, high expression in plants is best achieved from coding sequences that have at least about 35% GC content, preferably more than about 45%, more preferably more than about 50%, and most preferably more than about 60%. Microbial nucleotide sequences that have low GC contents may express poorly in plants due to the existence of ATTTA motifs that may destabilize messages, and AATAAA motifs that may cause inappropriate polyadenylation. Although preferred gene sequences may be adequately expressed in both monocotyledonous and dicotyledonous plant species, sequences can be modified to account for the specific codon preferences and GC content preferences of monocotyledons or dicotyledons as these preferences have been shown to differ (Murray et al. Nucl. Acids Res. 17:477-498 (1989)). In addition, the nucleotide sequences are screened for the existence of illegitimate splice sites that may cause message truncation. All changes required to be made within the nucleotide sequences such as those described above are made using well known techniques of site directed mutagenesis, PCR, and synthetic gene construction using the methods described in the published patent applications EP 0 385 962, EP 0 359 4721, and WO 93/07278.

[00111] In one embodiment of the invention the *vip3B* gene is made according to the procedure disclosed in U.S. Patent 5,625,136, herein incorporated by reference. In this

procedure, maize preferred codons, i.e., the single codon which most frequently encodes that amino acid in maize, are used. The maize preferred codon for a particular amino acid may be derived, for example, from known gene sequences from maize. Maize codon usage for 28 genes from maize plants is found in Murray et al., *Nucleic Acids Research* 17:477-498 (1989), the disclosure of which is incorporated herein by reference. A synthetic sequence made with maize optimized codons is set forth in SEQ ID NO: 2.

[00112] In this manner, the nucleotide sequences can be optimized for expression in any plant. It is recognized that all or any part of the gene sequence may be optimized or synthetic. That is, synthetic or partially optimized sequences may also be used.

[00113] For efficient initiation of translation, sequences adjacent to the initiating methionine may require modification. For example, they can be modified by the inclusion of sequences known to be effective in plants. Joshi has suggested an appropriate consensus for plants (NAR 15:6643-6653 (1987)) and Clontech suggests a further consensus translation initiator (1993/1994 catalog, page 210). These consensus are suitable for use with the nucleotide sequences of this invention. The sequences are incorporated into constructions comprising the nucleotide sequences, up to and including the ATG (whilst leaving the second amino acid unmodified), or alternatively up to and including the GTC subsequent to the ATG (with the possibility of modifying the second amino acid of the transgene).

[00114] The novel *vip3* toxin genes of the present invention, either as their native sequence or as optimized synthetic sequences as described above, can be operably fused to a variety of promoters for expression in plants including constitutive, inducible, temporally regulated, developmentally regulated, chemically regulated, tissue-preferred and tissue-specific promoters to prepare recombinant DNA molecules, i.e., chimeric genes. The choice of promoter will vary depending on the temporal and spatial requirements for expression, and also depending on the target species. Thus, expression of the nucleotide sequences of this invention in leaves, in stalks or stems, in ears, in inflorescences (e.g. spikes, panicles, cobs, etc.), in roots, and/or seedlings is preferred. In many cases, however, protection against more than one type of insect pest is sought, and thus expression in multiple tissues is desirable. Although many promoters from dicotyledons have been shown to be operational in monocotyledons and vice versa, ideally dicotyledonous promoters are selected for expression in dicotyledons, and monocotyledonous promoters for expression in monocotyledons. However, there is no

restriction to the provenance of selected promoters; it is sufficient that they are operational in driving the expression of the nucleotide sequences in the desired cell.

- [00115] Preferred constitutive promoters include the CaMV 35S and 19S promoters (Fraley *et al.*, U.S. Pat. No. 5,352,605 issued Oct. 4, 1994). An additionally preferred promoter is derived from any one of several of the actin genes, which are expressed in most cell types. The promoter expression cassettes described by McElroy *et al.* (Mol. Gen. Genet. 231: 150-160 (1991)) can be easily modified for the expression of the novel toxin gene and are particularly suitable for use in monocotyledonous hosts.
- [00116] Yet another preferred constitutive promoter is derived from ubiquitin, which is another gene product known to accumulate in many cell types. A ubiquitin promoter has been cloned from several species for use in transgenic plants, for example, sunflower (Binet *et al.*, 1991. Plant Science 79: 87-94), maize (Christensen *et al.*, 1989. Plant Molec. Biol. 12: 619-632), and arabidopsis (Norris *et al.* 1993. Plant Molec. Biol. 21:895-906). The maize ubiquitin promoter (*UbiP*) has been developed in transgenic monocot systems and its sequence and vectors constructed for monocot transformation are disclosed in the patent publication EP 0 342 926. The ubiquitin promoter is suitable for the expression of the novel toxin gene in transgenic plants, especially monocotyledons.
- [00117] Tissue-specific or tissue-preferential promoters useful for the expression of the novel toxin genes of the invention in plants, particularly maize, are those that direct expression in root, pith, leaf or pollen. Such promoters are disclosed in WO 93/07278, herein incorporated by reference in its entirety. Other tissue specific promoters useful in the present invention include the cotton rubisco promoter disclosed in US Patent 6,040,504; the rice sucrose synthase promoter disclosed in US Patent 5,604,121; and the cestrum yellow leaf curling virus promoter disclosed in WO 01/73087, all incorporated by reference. Chemically inducible promoters useful for directing the expression of the novel toxin gene in plants are disclosed in US Patent 5,614,395 herein incorporated by reference in its entirety.
- [00118] The nucleotide sequences of this invention can also be expressed under the regulation of promoters that are chemically regulated. This enables the Vip3 toxins to be synthesized only when the crop plants are treated with the inducing chemicals. Preferred technology for chemical induction of gene expression is detailed in the published application EP 0 332 104 (to Ciba-Geigy) and U.S. Patent 5,614,395. A preferred promoter for chemical induction is the tobacco PR-1a promoter.

- [00119]** A preferred category of promoters is that which is wound inducible. Numerous promoters have been described which are expressed at wound sites and also at the sites of phytopathogen infection. Ideally, such a promoter should only be active locally at the sites of infection, and in this way the insecticidal toxins only accumulate in cells that need to synthesize the insecticidal toxins to kill the invading insect pest. Preferred promoters of this kind include those described by Stanford *et al.* *Mol. Gen. Genet.* 215:200-208 (1989), Xu *et al.* *Plant Molec. Biol.* 22:573-588 (1993), Logemann *et al.* *Plant Cell* 1:151-158 (1989), Rohrmeier & Lehle, *Plant Molec. Biol.* 22:783-792 (1993), Firek *et al.* *Plant Molec. Biol.* 22:129-142 (1993), and Warner *et al.* *Plant J.* 3:191-201 (1993).
- [00120]** Preferred tissue specific expression patterns include green tissue specific, root specific, stem specific, and flower specific. Promoters suitable for expression in green tissue include many that regulate genes involved in photosynthesis and many of these have been cloned from both monocotyledons and dicotyledons. A preferred promoter is the maize PEPC promoter from the phosphoenol carboxylase gene (Hudspeth & Grula, *Plant Molec. Biol.* 12:579-589 (1989)). A preferred promoter for root specific expression is that described by de Framond (*FEBS* 290:103-106 (1991); EP 0 452 269 to Ciba-Geigy). A preferred stem specific promoter is that described in U.S. Patent 5,625,136 (to Ciba-Geigy) and which drives expression of the maize *trpA* gene.
- [00121]** Further preferred embodiments are transgenic plants expressing the nucleotide sequences in a wound-inducible or pathogen infection-inducible manner.
- [00122]** In addition to the selection of a suitable promoter, constructions for expression of an insecticidal toxin in plants require an appropriate transcription terminator to be attached downstream of the heterologous nucleotide sequence. Several such terminators are available and known in the art (e.g. *tml* from CaMV, *E9* from *rbcS*). Any available terminator known to function in plants can be used in the context of this invention.
- [00123]** Numerous other sequences can be incorporated into expression cassettes described in this invention. These include sequences that have been shown to enhance expression such as intron sequences (e.g. from *Adhl* and *bronzel*) and viral leader sequences (e.g. from TMV, MCMV and AMV).
- [00124]** It may be preferable to target expression of the nucleotide sequences of the present invention to different cellular localizations in the plant. In some cases, localization in the cytosol may be desirable, whereas in other cases, localization in some subcellular organelle may be preferred. Subcellular localization of transgene-encoded enzymes is

undertaken using techniques well known in the art. Typically, the DNA encoding the target peptide from a known organelle-targeted gene product is manipulated and fused upstream of the nucleotide sequence. Many such target sequences are known for the chloroplast and their functioning in heterologous constructions has been shown. The expression of the nucleotide sequences of the present invention is also targeted to the endoplasmic reticulum or to the vacuoles of the host cells. Techniques to achieve this are well known in the art.

[00125] Vectors suitable for plant transformation are described elsewhere in this specification. For *Agrobacterium*-mediated transformation, binary vectors or vectors carrying at least one T-DNA border sequence are suitable, whereas for direct gene transfer any vector is suitable and linear DNA containing only the construction of interest may be preferred. In the case of direct gene transfer, transformation with a single DNA species or co-transformation can be used (Schocher et al. *Biotechnology* 4:1093- 1096 (1986)). For both direct gene transfer and *Agrobacterium*-mediated transfer, transformation is usually (but not necessarily) undertaken with a selectable marker that may provide resistance to an antibiotic (kanamycin, hygromycin or methotrexate) or a herbicide (basta). The choice of selectable marker is not, however, critical to the invention.

[00126] In another preferred embodiment, a nucleotide sequence of the present invention is directly transformed into the plastid genome. A major advantage of plastid transformation is that plastids are generally capable of expressing bacterial genes without substantial modification, and plastids are capable of expressing multiple open reading frames under control of a single promoter. Plastid transformation technology is extensively described in U.S. Patent Nos. 5,451,513, 5,545,817, and 5,545,818, in PCT application no. WO 95/16783, and in McBride et al (1994) *Proc. Natl. Acad. Sci. USA* 91, 7301-7305. The basic technique for chloroplast transformation involves introducing regions of cloned plastid DNA flanking a selectable marker together with the gene of interest into a suitable target tissue, e.g., using biolistics or protoplast transformation (e.g., calcium chloride or PEG mediated transformation). The 1 to 1.5 kb flanking regions, termed targeting sequences, facilitate homologous recombination with the plastid genome and thus allow the replacement or modification of specific regions of the plastome. Initially, point mutations in the chloroplast 16S rRNA and rps12 genes conferring resistance to spectinomycin and/or streptomycin are utilized as selectable markers for transformation (Svab, Z., Hajdukiewicz, P., and Maliga, P. (1990) *Proc. Natl. Acad. Sci. USA* 87, 8526-

8530; Staub, J. M., and Maliga, P. (1992) *Plant Cell* 4, 39-45). This resulted in stable homoplasmic transformants at a frequency of approximately one per 100 bombardments of target leaves. The presence of cloning sites between these markers allowed creation of a plastid targeting vector for introduction of foreign genes (Staub, J.M., and Maliga, P. (1993) *EMBO J.* 12, 601-606). Substantial increases in transformation frequency are obtained by replacement of the recessive rRNA or r-protein antibiotic resistance genes with a dominant selectable marker, the bacterial *aadA* gene encoding the spectinomycin-cletoxyfying enzyme aminoglycoside- 3'- adenylyltransferase (Svab, Z., and Maliga, P. (1993) *Proc. Natl. Acad. Sci. USA* 90, 913-917). Previously, this marker had been used successfully for high-frequency transformation of the plastid genome of the green alga *Chlamydomonas reinhardtii* (Goldschmidt- Clermont, M. (1991) *Nucl. Acids Res.* 19:4083-4089). Other selectable markers useful for plastid transformation are known in the art and encompassed within the scope of the invention. Typically, approximately 15-20 cell division cycles following transformation are required to reach a homoplastidic state. Plastid expression, in which genes are inserted by homologous recombination into all of the several thousand copies of the circular plastid genome present in each plant cell, takes advantage of the enormous copy number advantage over nuclear- expressed genes to permit expression levels that can readily exceed 10% of the total soluble plant protein. In a preferred embodiment, a nucleotide sequence of the present invention is inserted into a plastid-targeting vector and transformed into the plastid genome of a desired plant host. Plants homoplasmic for plastid genomes containing a nucleotide sequence of the present invention are obtained, and are preferentially capable of high expression of the nucleotide sequence.

Combinations of Insect Control Principles

[00127] The pesticidal toxins of the invention can be used in combination with Bt δ -endotoxins or other pesticidal principles to increase pest target range. Furthermore, the use of the pesticidal toxins of the invention in combination with Bt δ -endotoxins or other pesticidal principles of a distinct nature has particular utility for the prevention and/or management of insect resistance.

- [00128] The various insecticidal crystal proteins from *Bacillus thuringiensis* have been classified based upon their spectrum of activity and sequence similarity. The classification put forth by Hofte and Whiteley, *Microbiol. Rev.* 53: 242-255 (1989) placed the then known insecticidal crystal proteins into four major classes. Generally, the major classes are defined by the spectrum of activity, with the Cry1 proteins active against Lepidoptera, Cry2 proteins active against both Lepidoptera and Diptera, Cry3 proteins active against Coleoptera, and Cry4 proteins active against Diptera.
- [00129] Within each major class, the δ -endotoxins are grouped according to sequence similarity. The Cry1 proteins are typically produced as 130-140 kDa protoxin proteins that are proteolytically cleaved to produce active toxins that are about 60-70 kDa. The active portion of the δ -endotoxin resides in the NH₂-terminal portion of the full-length molecule. Hofte and Whiteley, *supra*, classified the then known Cry1 proteins into six groups, 1Aa, 1Ab, 1Ac, 1B, 1C, and 1D. Since then, proteins classified as Cry1Ea, Cry1Fa, Cry9A, Cry9C and Cry9B, as well as others, have also been characterized.
- [00130] The spectrum of insecticidal activity of an individual δ -endotoxin from *Bacillus thuringiensis* tends to be quite narrow, with a given δ -endotoxin being active against only a few insects. Specificity is the result of the efficiency of the various steps involved in producing an active toxin protein and its subsequent ability to interact with the epithelial cells in the insect digestive tract. In one preferred embodiment, expression of the nucleic acid molecules of the invention in transgenic plants is accompanied by the expression of one or more Bt δ -endotoxins. Particularly preferred Bt δ -endotoxins are those disclosed in U.S. Patent 5,625,136, herein incorporated by reference.
- [00131] It is well known that many δ -endotoxin proteins from *Bacillus thuringiensis* are actually expressed as protoxins. These protoxins are solubilized in the alkaline environment of the insect gut and are proteolytically converted by proteases into a toxic core fragment (Hofte and Whiteley, *Microbiol. Rev.* 53: 242-255 (1989)). For δ -endotoxin proteins of the Cry1 class, the toxic core fragment is localized in the N-terminal half of the protoxin. It is within the scope of the present invention that genes encoding either the full-length protoxin form or the truncated toxic core fragment of the novel toxin proteins can be used in plant transformation vectors to confer insecticidal properties upon the host plant.

- [00132] Other insecticidal principles include protease inhibitors (both serine and cysteine types), lectins, α -amylase, peroxidase and cholesterol oxidase. Other Vip coding sequences, such as *vip1A(a)* and *vip2A(a)* as disclosed in U.S. Pat. No. 5,849,870 and herein incorporated by reference, are also useful in the present invention.
- [00133] This co-expression of more than one insecticidal principle in the same transgenic plant can be achieved by genetically engineering a plant to contain and express all the genes necessary. Alternatively, a plant, Parent 1, can be genetically engineered for the expression of genes of the present invention. A second plant, Parent 2, can be genetically engineered for the expression of a supplemental insect control principle. By crossing Parent 1 with Parent 2, progeny plants are obtained which express all the genes introduced into Parents 1 and 2.
- [00134] The present invention further encompasses variants of the disclosed nucleic acid molecules. Naturally occurring variant sequences can be identified and/or isolated with the use of well-known molecular biology techniques, as, for example, with PCR and hybridization techniques as outlined below.
- [00135] Variant *vip3* nucleotide sequences include synthetically derived nucleotide sequences, such as those generated, for example, by using site-directed mutagenesis or those made by whole domain swaps, but which still exhibit pesticidal activity. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel (1985) Proc. Natl. Acad. Sci. USA 82:488-492; Kunkel et al. (1987) Methods in Enzymol. 154:367-382; U.S. Pat. No. 4,873,192; Walker and Gaastra, eds. (1983) Techniques in Molecular Biology (MacMillan Publishing Company, New York) and the references cited therein. Generally, a nucleotide sequence of the invention will have at least 80%, preferably 85%, 90%, 95%, up to 98% or more sequence identity to its respective reference *vip3* nucleotide sequence, and have pesticidal activity.
- [00136] Variant *vip3* nucleotide sequences also encompass sequences derived from a mutagenic and recombinogenic procedure such as DNA shuffling. With such a procedure, one or more different *vip3* sequences of the present invention, for example, without limitation, *vip3B* and *vip3A-B* can be recombined together or with other *vip3* or related sequences, for example, and without limitation, *vip3A* (SEQ ID NO: 5), to create new *vip3* nucleic acid molecules encoding Vip3 toxins possessing the desired properties. In this manner, libraries of recombinant *vip3* polynucleotides are generated from a

population of sequence related *vip3* polynucleotides comprising sequence regions that have substantial sequence identity and can be homologously recombined *in vitro* or *in vivo*. Strategies for such DNA shuffling are known in the art. See, for example, Stemmer (1994) Proc. Natl. Acad. Sci. USA 91:10747-10751; Stemmer (1994) Nature 370:389-391; Cramer et al. (1997) Nature Biotech. 15:436-438; Moore et al. (1997) J. Mol. Biol. 272:336-347; Zhang et al. (1997) Proc. Natl. Acad. Sci. USA 94:4504-4509; Cramer et al. (1998) Nature 391:288-291; International Patent Application WO 99/57128, and U.S. Pat. Nos. 5,605,793, 5,837,458 and 6,335,179.

[00137] Mutagenesis methods as disclosed herein can be combined with high-throughput, screening methods to detect the pesticidal activity of cloned, mutagenized Vip3 polypeptides in host cells. Mutagenized DNA molecules that encode active Vip3 polypeptides (e.g., secreted and detected by antibodies; or insecticidal in an insect bioassay) can be recovered from the host cells and rapidly sequenced using standard art procedures. These methods allow the rapid determination of the importance of individual amino acid residues in a Vip3 polypeptide of interest, and can be applied to polypeptides of unknown structure.

[00138] The libraries of recombinant *vip3* genes that are produced using DNA shuffling methods are screened to identify those that exhibit improved properties for use in protecting plants against pests. Included among properties for which DNA shuffling is useful for obtaining improved *vip3* pest resistance genes are increased potency against a target pest, increased target pest range, decreased susceptibility to development of resistance by pests, increased expression level, increased resistance to protease degradation, increased stability in environmental conditions, and reduced toxicity to a host plant. By using an appropriate screening strategy, one can simultaneously or sequentially obtain *vip3* genes that are optimized for more than one property.

[00139] DNA shuffling is useful for obtaining *vip3* pest resistance genes that encode toxins that exhibit enhanced potency against a target pest. Once the shuffling is completed, the resulting library of shuffled *vip3* genes is screened to identify those that exhibit enhanced pesticidal activity. One way of performing this screening is to clone the protein-coding region of the shuffled *vip3* genes into an expression vector that is suitable for expressing the genes in a chosen host cell such as, for example, *E. coli* or a crystal minus strain of *Bacillus thuringiensis*. One skilled in the art will recognize the advantages and disadvantages of using either of these two expression systems. For example, *Bacillus*

thuringiensis would be more desirable in producing secreted Vip3 proteins. If desired, clones can be subjected to a preliminary screen, for example, by immunoassay, to identify those that produce a Vip3 protein of the correct size. Those that are positive in the preliminary screen are then tested in a functional screen to identify shuffled *vip3* genes that encode a toxin having the desired enhanced activity.

[00140] A whole insect assay can be used for determining toxicity. In these assays, the Vip3 toxins expressed from the shuffled *vip3* genes are placed on insect diet, for example, artificial diet or plant tissue, and consumed by the target insect. Those clones causing growth inhibition or mortality to the target insect can be tested in further bioassays to determine potency. Shuffled *vip3* genes encoding toxins with enhanced potency can be identified as those that have a decreased EC_{50} (concentration of toxin necessary to reduce insect growth by 50%) and/or LC_{50} (concentration of toxin necessary to cause 50% mortality).

[00141] *In vitro* assays can also be used for screening shuffled *vip3* gene libraries. Such assays typically involve the use of cultured insect cells that are susceptible to Vip3 toxins, and/or cells that express a receptor for the Vip3 toxins, either naturally or as a result of expression of a heterologous gene. Other *in vitro* assays can be used, for example, detection of morphological changes in cells, dyes and labels useful for detecting cell death, or detection of the release of ATPase by cells. One example of a suitable *in vitro* assay using cultured insect cells for Vip3 toxicity is Sf9 (*Spodoptera frugiperda*) cells. Sf9 is highly sensitive to Vip3 toxins. When Vip3 toxins are mixed with Sf9 cells, the cell membrane becomes highly permeable to small molecules. When a dye such as trypan blue is added to the cell suspension, those cells which are killed by the Vip3 toxin are stained blue. Thus, the cytotoxicity of the Vip3 toxin can be determined by image analysis.

[00142] Additional *in vitro* assays involve the use of receptors for the Vip3 toxins. One such receptor is disclosed in US Patent 6,291,156, herein incorporated by reference. The Vip3 receptor protein can be immobilized on a receiving surface, for example, without limitation, a 96-well plate or a nitrocellulose membrane, and exposed to clones comprising shuffled *vip3* genes. Thus, shuffled *vip3* genes that encode functional toxins can be identified on the basis of binding affinity to the Vip3 receptor. Further, the gene encoding the Vip3 receptor can be transformed into a non-Vip3 susceptible cell line, for example the Schneider 2 (S2) *Drosophila* cell line, using methods known in the art (see for example, Clem and Miller, 1194, Mol. Cel. Biol. 14:5212-522). The transformed S2

cells can then be exposed to clones comprising shuffled *vip3* genes. Thus, shuffled *vip3* genes that encode functional toxins can be identified on the basis of induction of cell death.

EXAMPLES

[00143] The invention will be further described by reference to the following detailed examples. These examples are provided for the purposes of illustration only, and are not intended to be limiting unless otherwise specified. Standard recombinant DNA and molecular cloning techniques used here are well known in the art and are described by Ansel (ed.) Current Protocols in Molecular Biology, John Wiley and Sons, Inc. (1994); T. Maniatis, E. F. Fritsch and J. Sambrook, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor laboratory, Cold Spring Harbor, NY (1998); and by T. J. Silhavy, M. L. Berman, and L. W. Enquist, *Experiments with Gene Fusions*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1984).

Example 1: Construction of Cosmid Library from *Bacillus thuringiensis* strain AB1183

[00144] Total DNA was isolated from a *Bacillus thuringiensis* strain, designated AB1183, by treating freshly grown cells resuspended in 100 mM Tris pH 8, 10 mM EDTA with 2 mg/ml lysozyme for 30 minutes at 37°C. Proteinase K was added to a final concentration of 100 µg/ml in 1% SDS, 50mM EDTA, 1M urea and incubated at 55°C. An equal volume of phenol-chloroform-isoamyl alcohol was added. The sample was gently mixed for 5 minutes and centrifuged at 3K. This was repeated twice. The aqueous phase was then mixed with 0.7 volumes isopropanol and centrifuged. The DNA pellet was washed three times with 70% ethanol and gently resuspended in 0.5X TE. 12 µg of DNA was treated with 0.3 unit of *Sau3A* per µg of DNA at 37°C in a volume of 100 µl. Samples were taken at 2-min intervals for 10 minutes. Then 1/10 volume 10X TE was added and samples were then heated for 30 minutes at 65°C to inactivate the enzyme. The samples were subjected to electrophoresis to determine which fraction was in the 40-kb range and this sample was used in the ligation.

[00145] SuperCos cosmid vector (Stratagene, La Jolla, CA) was prepared as described by the supplier utilizing the *Bam*HI cloning site. Prepared SuperCos at 100 ng/ml was ligated with the AB1183 DNA previously digested with *Sau*3A at a ratio of 2:1 in a 5 µl volume overnight at 6°C. The ligation mixture was packaged using Gigapack XL III (Stratagene) as described by the supplier. Packaged phages were infected into XL-1MR *E. coli* cells (Stratagene) as described by the supplier. The cosmid library was plated on L-agar with 50 µg/ml kanamycin and incubated 16 hours at 37°C. 1200 colonies were picked and grown for testing against insects.

Example 2. Bioassay of Cosmid Clones.

[00146] The 1200 colonies from Example 1 were screened for insecticidal activity against *Heliothis virescens* neonate larvae. The bioassay was performed using a surface contamination method on artificial diet. The bioassay was scored after 7 days. Eight clones were found to be insecticidal to *Heliothis virescens*.

Example 3: Analysis of Insecticidal Cosmid Clones

[00147] To identify *vip3* homologous sequences, PCR analysis was performed on the eight *Heliothis virescens*-positive clones using primers from the 5' prime region of the *vip3A* gene (SEQ ID NO: 4), using the methods of Carozzi *et al.* (1991, Appl. Env. Microbiol. 57: 3057-3061). The primers used for this analysis are:

Forward: 5'-GTGATCTAACCTAGACG-3' (SEQ ID NO: 8)

Reverse: 5'-GCTTTAGTTCCATTCCTCC-3' (SEQ ID NO: 9).

[00148] One clone produced a DNA band of the size expected for genes related to the *vip3* class. A 3.8 kb EcoRI fragment from this clone was subcloned into pBluescript (Stratagene) and transformed into *E. coli*. This *E. coli* clone was confirmed to comprise a *vip3* homologous coding sequence using PCR analysis. The *vip3* homologous coding sequence was designated *vip3B*. Bioassay results of the *E. coli* clone comprising the *vip3B* coding sequence demonstrated that the Vip3B toxin was responsible for the *Heliothis virescens* activity. The plasmid comprised in this clone, and the clone, were designated pCIB9400.

Example 4. Cloning and Sequencing the Full-length *vip3B* Gene.

[00149] pCIB9400 was cut with *Bgl*II and *Eco*RV to remove approximately 800 bp of flanking sequence near the 3' end of the *vip3B* coding sequence. The ends of the resulting fragment were filled in using Klenow polymerase (New England Biolabs, Beverly, Mass.) and then ligated together using T4 ligase (New England Biolabs, Beverly, Mass.) This ligation mixture was transformed into *E. coli* DH5 α cells. Plasmid DNA was isolated from single colonies using a standard alkaline lysis procedure and several restriction digests were done to ensure that the 800 bp flanking region was deleted. The resulting plasmid was designated pNOV1325 and deposited in *E. coli* as ATCC accession no. PTA-3868.

[00150] Sequencing was performed using the dideoxy chain-termination method and was completed using Applied Biosystems Inc. model 3700 automated DNA sequencer (Foster City, CA). The sequence was assembled using Sequencher 4.05 from Gene Codes Corporation (Ann Arbor, Michigan).

[00151] Sequence analysis identified a 2364 bp coding sequence (SEQ ID NO: 1) coding for a 787 amino acid protein (SEQ ID NO: 2) with an approximate molecular weight of 88 kDa. The *vip3B* nucleotide sequence has 86% identity to the nucleotide sequence of the *vip3A* gene. The amino acid sequence of the Vip3B protein is 81% identical to the amino acid sequence of the Vip3A protein.

Example 5: Bioassay of the Vip3B Protein Expressed in pNOV1325 and pCIB9400 .

[00152] Molten black cutworm diet (BioServ, Frenchtown, NJ) was poured into 50 mm petri dishes and allowed to solidify. A 200 μ l suspension of Top10 (Invitrogen,) *E. coli* cells comprising either pNOV1325 or pNOV9400 (each of which comprise the *vip3B* coding sequence) was pipetted onto the diet surface. The solution was uniformly spread with a bacterial loop so that the suspension covered the entire surface of the diet. The surface was allowed to dry thoroughly. First instar larvae of a lepidopteran species listed in the table below were placed on the diet with a fine tip brush. Each species was tested separately. Larval mortality was recorded at 3 days and 5 days after larval infestation of the diet. A sample containing Top10 *E. coli* cells without a vector was used for the

negative control. Vip3A protein can also be tested in the same bioassay for comparative purposes. For this example, the Vip3B data obtained in this example were compared with the known activity spectrum data of Vip3A.

[00153] The bioassay results are shown in Table 1. Data shown in table 1 are from the day 5 post-infestation recording. Little or no activity was observed in the *E. coli* negative control. Results indicate that the Vip3B toxin has a broader spectrum of activity than the Vip3A toxin since Vip3B was active against *Ostrinia nubilalis* and *Plutella xylostella*. Results also suggest that Vip3B has a higher specific activity to *Helicoverpa zea* than the Vip3A toxin.

Table 1.

Insect	Percent Mortality		<i>E.coli</i> Control	Activity Spectrum of Vip3A ^b
	pCIB9400	pNOV1325		
<i>Manduca sexta</i>	100	100	0	+
<i>Agrotis ipsilon</i>	100	100	5	+
<i>Helicoverpa zea</i>	100	100	10	+
<i>Heliothis virescens</i>	100	100	10	+
<i>Spodoptera exguia</i>	100	90	10	+
<i>Spodoptera frugiperda</i>	100	100	0	+
<i>Ostrinia nubilalis</i>	100	100	0	-
<i>Plutella xylostella</i>	100	100	0	-
<i>Trichplusia ni</i>	100	100	0	+
<i>Pectinophora gossypiella</i>	50 ^a	50 ^a	0	+
<i>Cochyles hospes</i>	60 ^a	80 ^a	0	+

^aSurviving insects were observed to have severe feeding and growth inhibition.

^bA “+” indicates an insect species that is susceptible to Vip3A. A “-“ indicates an insect species with little or no susceptibility of Vip3A.

Example 6. Construction of Maize Optimized *vip3B* Coding Sequence

[00154] A synthetic maize-optimized *vip3B* coding sequence was made according to the procedure disclosed in U.S. Patent 5,625,136. In this procedure, maize preferred codons, i.e., the single codon that most frequently encodes that amino acid in maize, are used. The maize preferred codon for a particular amino acid is derived from known gene sequences

from maize. Maize codon usage for 28 genes from maize plants is found in Murray *et al.*, Nucleic Acids Research 17:477-498 (1989). The synthetic *vip3B* coding sequence (SEQ ID NO: 3) was cloned into a pET101/D-Topo expression vector. The resulting vector, designated pNOV1328, was transformed into *E. coli* DH5 α cells and deposited as ATCC accession number PTA-3869.

Example 7. Creation of Transgenic Maize Plants Comprising a *vip3B* Gene.

[00155] The synthetic maize-optimized *vip3B* (SEQ ID NO: 3) coding sequence was chosen for transformation into maize plants. An expression cassette comprising the synthetic *vip3B* coding sequence was transferred to a suitable vector for *Agrobacterium*-mediated maize transformation. Three vectors were constructed for this example: (a) a vector comprising two *vip3B* expression cassettes, the first expression cassette comprising MTL:*vip3B*, and the second expression cassette comprising PEPC:*vip3B*, (b) a vector comprising CMP:*vip3B*, and (c) a vector comprising UbiP:*vip3B*. All vectors used in this example also comprise the phosphomannose isomerase (PMI) gene for selection of transgenic lines (Negrotto *et al.* (2000) Plant Cell Reports 19: 798-803).

[00156] All three vectors were individually transformed into maize. Transformation of immature maize embryos was performed essentially as described in Negrotto *et al.*, 2000, Plant Cell Reports 19: 798-803. For this example, all media constituents were as described in Negrotto *et al.*, *supra*. However, various media constituents known in the art may be substituted.

[00157] *Agrobacterium* strain LBA4404 (pSB1) containing the plant transformation plasmid was grown on YEP (yeast extract (5 g/L), peptone (10g/L), NaCl (5g/L), 15g/l agar, pH 6.8) solid medium for 2 – 4 days at 28°C. Approximately 0.8×10^9 *Agrobacterium* were suspended in LS-inf media supplemented with 100 μ M As (Negrotto *et al.*, (2000) Plant Cell Rep 19: 798-803). Bacteria were pre-induced in this medium for 30-60 minutes.

[00158] Immature embryos from A188 were excised from 8 – 12 day old ears into liquid LS-inf + 100 μ M As. Immature embryos from other maize germplasm can also be used. Embryos were rinsed once with fresh infection medium. *Agrobacterium* solution was then added and embryos were vortexed for 30 seconds and allowed to settle with the bacteria for 5 minutes. The embryos were then transferred scutellum side up to LSAs

medium and cultured in the dark for two to three days. Subsequently, between 20 and 25 embryos per petri plate were transferred to LSDc medium supplemented with cefotaxime (250 mg/l) and silver nitrate (1.6 mg/l) and cultured in the dark for 28°C for approximately 10 days.

[00159] Immature embryos, producing embryogenic callus were transferred to LSD1M0.5S medium. The cultures were selected on this medium for approximately 6 weeks with a subculture step at approximately 3 weeks. Surviving calli were transferred to Reg1 medium supplemented with mannose. Following culturing in the light (16 hour light/ 8 hour dark regiment), green tissues were then transferred to Reg2 medium without growth regulators and incubated for approximately 1-2 weeks. Plantlets were transferred to Magenta GA-7 boxes (Magenta Corp, Chicago Ill.) containing Reg3 medium and grown in the light. After approximately 2-3 weeks, plants were tested for the presence of the PMI gene and the *vip3B* gene by PCR. Positive plants from the PCR assay were transferred to the greenhouse and tested for resistance to lepidopteran pests.

Example 8. Analysis of Transgenic Maize Plants Expressing Vip3B

Insect Bioassays

[00160] Plants were sampled as they were being transplanted from Magenta GA-7 boxes into soil. Sampling consisted of cutting two small pieces of leaf (ca. 2-4 cm long) and placing each piece in a small petri dish. Negative controls were either transgenic plants that were PCR negative for the *vip3B* gene from the same transformation experiment, or from non-transgenic plants (of a similar size to test plants) that were being grown under similar growth conditions as the transgenics.

[00161] Leaf samples from each plant were inoculated with either European corn borer (*Ostrinia nubilalis*) or fall armyworm (*Spodoptera frugiperda*) by placing 10 first instar larvae onto each leaf piece. Petri dishes are then tightly sealed. Other suitable insect pests can also be used.

[00162] At approximately 3-4 days post inoculation, data was collected. The percent mortality of the larvae was calculated. Also, a visual damage rating of the leaf can be ascertained at the same time. Feeding damage is rated as high, moderate, low, or absent and given a numerical value of 3, 2, 1 or 0, respectively.

[00163] Results of the bioassay of the transgenic plants are shown in Table 2. Results indicate that transgenic maize plants comprising the *vip3B* gene and expressing the Vip3B protein, are insecticidal to European corn borer and fall armyworm.

Table 2. Efficacy of Transgenic Maize Plants Expressing Vip3B.

Event	Promter: <i>vip3B</i> Construct	% FAW Mortality Per Plant ^a	% ECB Mortality Per Plant ^b
118A	MTL: <i>vip3B</i> / PEPC: <i>vip3B</i>	100,100,100,100,100,100,100,100,100, 100,100,100,100,100,100,100,100,100, 100,100,100,100,100,100,100,100,100, 100	90,100,100,90,100,100,80,100, 100,100,100,90,90,90,100,100, 90,100,70,80,70,100,100,100, 70,100,80,80
121A	MTL: <i>vip3B</i> / PEPC: <i>vip3B</i>	90,100,100,100,100,100,100,100,100, 100,100,100,100,100,100,100,100,100, 100,100,100,100,100,100,100,100,100, 100,100,100,100	70,80,70,90,80,80,90,80,80,90, 100,100,80,90,70,90,80,100,80, 100,70,80,70,70,90,100,100,90, 90,90,90
142C	MTL: <i>vip3B</i> / PEPC: <i>vip3B</i>	100,100,100,100,100,100,100,100,100	80,100,90,80,90,70,70,90,90
145B	MTL: <i>vip3B</i> / PEPC: <i>vip3B</i>	100,100,100,100,100,100,100,100,100, 100,100,100,100,100,100,100,100,100, 100	80,70,90,100,70,70,100,80,80, 70,100,100,100,100,100,70,100,90, 90,90
89A	CMP: <i>vip3B</i>	100,100,100,100,100,100,100,100	100,100,100,100,100,100,100,100
190A	CMP: <i>vip3B</i>	100,100,100,100,100,100,100,100,100	100,80,90,80,90,100,80,100,80

^aFAW= fall armyworm

^bECB = European corn borer

ELISA Assay

[00164] Levels of Vip3B protein in various transgenic maize tissues were determined using an ELISA. ELISA analysis was done according to the method disclosed in U.S. Patent No. 5,625,136. Results of the ELISA analysis are shown in Table 3.

Table 3. Vip3B Protein Levels in Transgenic Maize.

Event	Mean Vip3B Protein Level in Specified Tissue ($\mu\text{g}/\text{mg}$ soluble protein)			
	Leaf	Pith	Rind	Pollen
118A	20 \pm 12	.316 \pm .240	.686 \pm .645	0
121A	20 \pm 19	.523 \pm .387	.723 \pm .343	0
142C	18 \pm 17	1.12 \pm .770	3.75 \pm 1.30	0
145B	19 \pm 13	.720 \pm .720	2.23 \pm 2.0	0

Example 9. Hybrid Vip3 Toxins

[00165] Vip3B is toxic to *Ostrinia nubilalis* (European corn borer) and *Plutella xylostella* (diamond back moth), whereas a related Vip toxin, Vip3A, has little or no activity. Vip3B and Vip3A differ primarily in the C-terminal region of their respective amino acid sequences particularly in the region from amino acid 579 to amino acid 787 of SEQ ID NO: 2. In order to demonstrate that this C-terminal region of Vip3B is the portion of the Vip3B toxin that is sufficient for the activity against European corn borer and diamond back moth, a hybrid toxin comprising the C-terminal region of Vip3B, starting at amino acid number 579 and ending at amino acid number 787 of SEQ ID NO: 2, was joined in an amino to carboxy direction with the N-terminal region of Vip3A, starting at amino acid number 1 and ending at amino acid number 578 of SEQ ID NO: 5. The hybrid toxin was designated Vip3A-B (SEQ ID NO: 7).

[00166] The nucleic acid molecule encoding the Vip3A-B hybrid toxin was constructed using two steps of PCR with the following primers:

VIP3A-N: 5'-ATGACCAAGAATAATACTAAATTAAGCAC-3' (SEQ ID NO: 10)

VIPfus4: 5'-TCCTTATGAACATATAAAGCTTTAGTTCCATT-3' (SEQ ID NO: 11)

VIP3B-C:5'-GGCGAATTCTCACTTAATCGAAAAATCCGGAAATTTAT-3' (SEQ ID NO: 12)

VIPfus3: 5'-AATGGAACTAAAGCTTTATATGTTTCATAAGGA-3' (SEQ ID NO: 13)

[00167] In the first PCR step, primers Vip3A-N (SEQ ID NO: 10) and Vipfus4 (SEQ ID NO: 11) were used to generate an approximately 1.7 kb fragment of the 5' end of the *vip3A* gene, encoding the N-terminal region, and primers Vip3B-C (SEQ ID NO: 12) and Vipfus3 (SEQ ID NO: 13) were used to generate an approximately 0.7 kb fragment of the 3' end of the *vip3B* gene encoding the C-terminal region. In the second PCR step, these two fragments were combined as the templates for primers Vip3A-N (SEQ ID NO: 10) and Vip3B-C (SEQ ID NO: 12) to generate an approximately 2.4 kb hybrid *vip3A-vip3B* gene, designated *vip3A-B* (SEQ ID NO: 6).

[00168] An *E. coli* clone expressing the hybrid Vip3A-B toxin was tested for insecticidal activity against fall armyworm and European corn borer using the method outlined in Example 5. Results of some of the bioassays suggested that the C-terminal region of Vip3B is sufficient to confer European corn borer activity on the hybrid toxin.

Example 10. *In vitro* Recombination of *vip3* Genes by DNA Shuffling

[00169] One of the *vip3* genes of the present invention, for example SEQ ID NO: 1, 3, or 6 is amplified by PCR. The resulting DNA fragment is digested by DNaseI treatment essentially as described in Stemmer *et al.*, *PNAS* 91: 10747-10751 (1994), and the PCR primers are removed from the reaction mixture. A PCR reaction is carried out without primers and is followed by a PCR reaction with the primers, both as described in Stemmer *et al.* (1994). The resulting DNA fragments are cloned into pTRC99a (Pharmacia, Cat no: 27-5007-01) and transformed into *E. coli* strain SASX38 by electroporation using the Biorad Gene Pulser and the manufacturer's conditions. The transformed bacteria are grown on medium overnight and screened for insecticidal activity.

[00170] In a similar reaction, PCR-amplified DNA fragments comprising one of the *vip3* genes described herein (SEQ ID NO: 1, 3, or 6, or mutants thereof), and PCR-amplified DNA fragments comprising at least one other of the *vip3* genes described herein (or a mutant thereof) are recombined *in vitro* and resulting variants with improved insecticidal properties are recovered as described below.

[00171] In order to increase the diversity of the shuffled *vip3* gene library, a *vip3* gene or genes (called the primary genes) are shuffled using synthetic oligonucleotide shuffling. A plurality (e.g., 2, 5, 10, 20, 50, 75, or 100 or more) of oligonucleotides corresponding to at least one region of diversity are synthesized. These oligonucleotides can be shuffled directly, or can be recombined with one or more of the family of nucleic acids.

[00172] The oligonucleotide sequence can be taken from other *vip3* genes called secondary genes. The secondary genes have a certain degree of homology to the primary genes. There are several ways to select parts of the secondary gene for the oligonucleotide synthesis. For example, portions of the secondary gene can be selected at random. The DNA shuffling process will select those oligonucleotides, which can be incorporated into the shuffled genes.

- [00173] The selected portions can be any lengths as long as they are suitable to synthesize. The oligonucleotides can also be designed based on the homology between the primary and secondary genes. A certain degree of homology is necessary for crossover, which must occur among DNA fragments during the shuffling. At the same time, strong heterogeneity is desired for the diversity of the shuffled gene library. Furthermore, a specific portion of the secondary genes can be selected for the oligonucleotide synthesis based on the knowledge in the protein sequence and function relationship.
- [00174] The present invention has disclosed that the C-terminal domain of Vip3B is in part responsible for spectrum of activity of the Vip3 toxins. When the insecticidal spectrum is modified by the current invention utilizing the DNA shuffling technology, the C-terminal region of the nucleotide sequence of the secondary genes can be selected as a target region for synthesizing oligonucleotides used in an oligonucleotide shuffling procedure.
- [00175] Since the insecticidal activity of the Vip3 protein is dependent, at least in part, to the N-terminal region, the N-terminal region of the secondary genes can be selected for oligonucleotide shuffling for increased insecticidal activity.

Example 11. High-throughput Screen for Insecticidal Activity.

- [00176] Shuffled *vip3* gene libraries in either *E. coli* or *Bacillus thuringiensis* are screened for insecticidal activity. Colonies are picked with a Q-bot (Beckman), placed in growth media in a standard 96-well format and grown over night. Each clone is then layered onto the surface of an insect diet in 96-well format and the surface allowed to dry. Optionally, pools of transformed cells are added to each well to increase the number of clones that are tested in the initial screening round. For example, screening 100 clones per well and using 10,000 wells provides a screen of 10^6 clones.
- [00177] Several neonate larvae of a target insect, for example, *Heliothis virescens*, *Helicoverpa zea* or *Spodoptera frugiperda*, are added to each well. The plate is covered with an air permeable membrane that retains the larvae in the wells into which they were placed. After 5 days the wells are evaluated for amount of diet consumed and/or insect mortality. Clones in wells indicating that little or no diet is consumed and/or where high insect mortality is observed are chosen for further analysis. Several clones should be found to have enhanced activity against the target insect.

[00178] All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art that this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

[00179] It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and the scope of the appended claims.

What is claimed is:

1. An isolated toxin that is active against insects, wherein said toxin comprises an amino acid sequence that:
 - (a) has at least 91% sequence identity with SEQ ID NO: 2; or
 - (b) is produced by the expression of a nucleic acid molecule comprising a nucleotide sequence that has at least 92% sequence identity with SEQ ID NO: 1.
2. The isolated toxin according to claim 1, wherein said toxin comprises an amino acid sequence with at least 95% identity with the amino acid sequence set forth in SEQ ID NO: 2.
3. The isolated toxin according to claim 1, wherein said toxin comprises an amino acid sequence with at least 99% identity with the amino acid sequence set forth in SEQ ID NO: 2.
4. The isolated toxin according to claim 1, wherein said toxin comprises the amino acid sequence set forth in SEQ ID NO: 2.
5. The isolated toxin according to claim 1, wherein said toxin is produced by the expression of a nucleic acid molecule comprising a nucleotide sequence that has at least 95% sequence identity with SEQ ID NO: 1.
6. The isolated toxin according to claim 1, wherein said toxin is produced by the expression of a nucleic acid molecule comprising a nucleotide sequence that has at least 99% sequence identity with SEQ ID NO: 1.
7. The isolated toxin according to claim 1, wherein said toxin is produced by the expression of a nucleic acid molecule comprising a nucleotide sequence comprising nucleotides 1-2364 of SEQ ID NO: 1 or nucleotides 1-2364 of SEQ ID NO: 3.

8. The isolated toxin according to claim 7, wherein said toxin is produced by the expression of a nucleic acid molecule comprising a nucleotide sequence comprising nucleotides 1-2364 of SEQ ID NO: 1.
9. The isolated toxin according to claim 7, wherein said toxin is produced by the expression of a nucleic acid molecule comprising a nucleotide sequence comprising nucleotides 1-2364 of SEQ ID NO: 3.
10. The isolated toxin according claim 1, wherein said toxin is produced by the expression of a nucleotide sequence comprising the approximately 2.4 kb DNA fragment comprised in pNOV1325 (ATCC PTA-3868).
11. The isolated toxin according to claim 1, wherein said toxin is produced by the expression of a nucleotide sequence comprising the approximately 2.4 kb DNA fragment comprised in pNOV1328 (ATCC PTA-3869).
12. The isolated toxin according to claim 1, wherein said insect is a lepidopteran insect.
13. The isolated toxin according to claim 12, wherein said lepidopteran insect is selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth)..
14. A composition comprising an effective insect-controlling amount of the toxin according to claim 1.
15. A method of controlling insects, comprising delivering to the insects an effective amount of the toxin according to claim 1.

16. The method of claim 15, wherein said insects are lepidopteran insects.
17. The method of claim 16, wherein said lepidopteran insects are selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth).
18. The method of claim 15, wherein said toxin is delivered to the insects orally.
19. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes a toxin that is active against insects, wherein said nucleotide sequence:
- (a) has at least 92% sequence identity with SEQ ID NO: 1; or
 - (b) is isocoding with the nucleotide sequence of (a); or
 - (c) encodes an amino acid sequence that has at least 91% sequence identity with SEQ ID NO: 2.
20. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence has at least 92% sequence identity with the nucleotide sequence set forth in SEQ ID NO: 1.
21. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence has at least 95% sequence identity with the nucleotide sequence set forth in SEQ ID NO: 1.
22. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence has at least 99% sequence identity with the nucleotide sequence set forth in SEQ ID NO: 1.
23. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence comprises nucleotides 1-2364 of SEQ ID NO: 1.

24. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence is isocoding for a nucleotide sequence that has at least 92% sequence identity to SEQ ID NO: 1.
25. The isolated nucleic acid molecule according to claim 24, wherein said nucleotide sequence comprises nucleotides 1-2364 of SEQ ID NO: 3.
26. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence encodes an amino acid sequence with at least 91% sequence identity with SEQ ID NO: 2.
27. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence encodes an amino acid sequence with at least 95% sequence identity with SEQ ID NO: 2.
28. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence encodes an amino acid sequence with at least 99% sequence identity with SEQ ID NO: 2.
29. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence encodes the amino acid sequence set forth in SEQ ID NO: 2.
30. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence comprises the approximately 2.4 kd DNA fragment comprised in pNOV1325 (ATCC PTA-3868).
31. The isolated nucleic acid molecule according to claim 19, wherein said nucleotide sequence comprises the approximately 2.4 kd DNA fragment comprised in pNOV1328 (ATCC PTA-3869).
32. The isolated nucleic acid molecule according to claim 19, wherein said toxin has activity against a lepidopteran insect.

33. The isolated nucleic acid molecule according to claim 32, wherein said lepidopteran insect is selected from the group consisting of *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth)..
34. A chimeric gene comprising a heterologous promoter sequence operatively linked to the nucleic acid molecule of claim 19.
35. A recombinant vector comprising the chimeric gene of claim 34.
36. A transgenic host cell comprising the chimeric gene of claim 34.
37. An animal virus comprising the chimeric gene of claim 34.
38. A plant virus comprising the chimeric gene of claim 34.
39. The transgenic host cell according to claim 36, which is a bacterial cell.
40. The transgenic host cell according to claim 36, which is a yeast cell.
41. The transgenic host cell according to claim 36, which is a plant cell.
42. A transgenic plant comprising the transgenic plant cell of claim 41.
43. The transgenic plant according to claim 42, wherein said plant is selected from the group consisting of sorghum, wheat, sunflower, tomato, cole crops, cotton, rice, soybean, sugar beet, sugarcane, tobacco, barley, oilseed rape and maize.

44. The transgenic plant according to claim 43, wherein said plant is a maize plant.
45. Transgenic seed from the transgenic plant of claim 42.
46. Transgenic seed from the maize plant of claim 44.
47. A method of producing a toxin that is active against insects, comprising
- (a) obtaining the transgenic host cell according to claim 36; and
 - (b) expressing the nucleic acid molecule in said cell, which results in at least one toxin that is active against insects.
48. A method of producing an insect-resistant transgenic plant, comprising introducing a nucleic acid molecule according to claim 19 into said transgenic plant, wherein said nucleic acid molecule is expressible in said transgenic plant in an effective amount to control insects.
49. The method according to claim 48, wherein said insects are lepidopteran insects.
50. The method according to claim 49, wherein said lepidopteran insects are selected from the group consisting of: *Ostrinia nubilalis* (European corn borer), *Plutella xylostella* (diamondback moth), *Spodoptera frugiperda* (fall armyworm), *Agrotis ipsilon* (black cutworm), *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Spodoptera exigua* (beet armyworm), *Helicoverpa punctigera* (native budworm), *Helicoverpa armigera* (cotton bollworm), *Manduca sexta* (tobacco hornworm), *Trichoplusia ni* (cabbage looper), *Pectinophora gossypiella* (pink bollworm), and *Cochylis hospes* (banded sunflower moth).

SEQUENCE LISTING

<110> Miles, Paul
 Kramer, Vance
 Shen, Zhicheng
 Shotkoski, Frank
 Warren, Greg

<120> Novel Pesticidal Toxins

<130> S-60000P1

<160> 13

<170> PatentIn version 3.0

<210> 1

<211> 2364

<212> DNA

<213> Bacillus thuringiensis

<220>

<221> CDS

<222> (1)..(2364)

<223> Native vip3B coding sequence

<400> 1
 atg aac aag aat aat act aaa tta aac gca agg gcc tta ccg agt ttt 48
 Met Asn Lys Asn Asn Thr Lys Leu Asn Ala Arg Ala Leu Pro Ser Phe
 1 5 10 15
 att gat tat ttt aat ggc att tat gga ttt gcc act ggt atc aaa gac 96
 Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
 20 25 30

att atg aac atg att ttt aaa acg gat aca ggt gga aat cta acc cta Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asn Leu Thr Leu 35 40 45	144
gac gaa att tta aaa aat cag cag tta tta aat gag att tct ggt aaa Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Glu Ile Ser Gly Lys 50 55 60	192
ttg gat ggg gta aat ggg agc tta aac gat ctt atc gca cag gga aac Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn 65 70 75 80	240
tta aat aca gaa tta tct aag gaa atc tta aaa att gca aat gag cag Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln 85 90 95	288
aat caa gtc tta aat gat gtt aat aac aaa ctt aat gcg ata aat aca Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asn Ala Ile Asn Thr 100 105 110	336
atg ctt cac ata tat cta cct aaa att aca tct atg tta aat gat gta Met Leu His Ile Tyr Leu Pro Lys Ile Thr Ser Met Leu Asn Asp Val 115 120 125	384
atg aaa caa aat tat gca cta agt ctg caa ata gaa tac cta agt aaa Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys 130 135 140	432
caa ttg caa gaa att tcc gac aag tta gat gtc att aac gtg aat gta Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Val Ile Asn Val Asn Val 145 150 155 160	480
ctt att aac tct aca ctt act gaa att aca cct gcg tat caa cgg atg Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Met 165 170 175	528
aaa tat gta aat gaa aaa ttt gaa gat tta act ttt gct aca gaa acc Lys Tyr Val Asn Glu Lys Phe Glu Asp Leu Thr Phe Ala Thr Glu Thr 180 185 190	576
act tta aaa gta aaa aag aat agc tcc cct gca gat att ctt gat gag Thr Leu Lys Val Lys Lys Asn Ser Ser Pro Ala Asp Ile Leu Asp Glu 195 200 205	624
tta act gag tta act gaa cta gcg aaa agt gta aca aaa aat gac gtg Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val 210 215 220	672
gat ggt ttt gaa ttt tac ctt aat aca ttc cac gat gta atg gta gga Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly 225 230 235 240	720
aac aat tta ttc ggg cgt tca gct tta aaa act gct tcg gaa tta atc Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile 245 250 255	768
gct aaa gaa aat gtg aaa aca agt ggc agt gag gta gga aat gtt tat Ala Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr 260 265 270	816

aat ttc tta att gta tta aca gct ctg caa gca aaa gct ttt ctt act	864
Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Lys Ala Phe Leu Thr	
275 280 285	
tta aca aca tgc cgg aaa tta tta ggc tta gca gat att gat tat act	912
Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr	
290 295 300	
ttc att atg aat gaa cat tta gat aag gaa aaa gag gaa ttt aga gta	960
Phe Ile Met Asn Glu His Leu Asp Lys Glu Lys Glu Glu Phe Arg Val	
305 310 315 320	
aat atc ctt cct aca ctt tct aat act ttt tct aat cct aac tat gca	1008
Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala	
325 330 335	
aaa gct aaa gga agc aat gaa gat gca aag ata att gtg gaa gct aaa	1056
Lys Ala Lys Gly Ser Asn Glu Asp Ala Lys Ile Ile Val Glu Ala Lys	
340 345 350	
cca gga tat gct ttg gtt gga ttt gaa atg agc aat gat tca atc aca	1104
Pro Gly Tyr Ala Leu Val Gly Phe Glu Met Ser Asn Asp Ser Ile Thr	
355 360 365	
gta tta aaa gca tat cag gct aag cta aaa caa gat tat caa gtt gat	1152
Val Leu Lys Ala Tyr Gln Ala Lys Leu Lys Gln Asp Tyr Gln Val Asp	
370 375 380	
aaa gat tcg tta tca gaa att gtc tat ggt gat atg gat aaa tta ttg	1200
Lys Asp Ser Leu Ser Glu Ile Val Tyr Gly Asp Met Asp Lys Leu Leu	
385 390 395 400	
tgc ccg gat caa tct gaa caa ata tat tat aca aat aac att gct ttt	1248
Cys Pro Asp Gln Ser Glu Gln Ile Tyr Tyr Thr Asn Asn Ile Ala Phe	
405 410 415	
ccc aat gaa tat gta att act aaa att act ttt act aaa aaa atg aat	1296
Pro Asn Glu Tyr Val Ile Thr Lys Ile Thr Phe Thr Lys Lys Met Asn	
420 425 430	
agt tta aga tat gag gca aca gct aat ttt tat gat tct tct aca ggg	1344
Ser Leu Arg Tyr Glu Ala Thr Ala Asn Phe Tyr Asp Ser Ser Thr Gly	
435 440 445	
gat att gat cta aat aag aca aaa gta gaa tca agt gaa gca gag tat	1392
Asp Ile Asp Leu Asn Lys Thr Lys Val Glu Ser Ser Glu Ala Glu Tyr	
450 455 460	
agt acg cta agt gct agt act gat gga gtc tat atg ccg tta ggt att	1440
Ser Thr Leu Ser Ala Ser Thr Asp Gly Val Tyr Met Pro Leu Gly Ile	
465 470 475 480	
atc agt gaa aca ttt ttg act cca att aat ggg ttt gga atc gta gtc	1488
Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Ile Val Val	
485 490 495	
gat gaa aat tca aaa tta gta aat tta aca tgt aaa tca tat tta aga	1536
Asp Glu Asn Ser Lys Leu Val Asn Leu Thr Cys Lys Ser Tyr Leu Arg	
500 505 510	

gag gta tta tta gca aca gac tta agt aat aaa gaa act aaa ttg att	1584
Glu Val Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile	
515 520 525	
gtc cca cct att ggt ttt att agc aat att gta gaa aat ggg aac tta	1632
Val Pro Pro Ile Gly Phe Ile Ser Asn Ile Val Glu Asn Gly Asn Leu	
530 535 540	
gag gga gaa aac tta gag ccg tgg aaa gca aat aac aaa aat gcg tat	1680
Glu Gly Glu Asn Leu Glu Pro Trp Lys Ala Asn Asn Lys Asn Ala Tyr	
545 550 555 560	
gta gat cat aca ggc ggc gta aat gga act aaa gct tta tat gtt cat	1728
Val Asp His Thr Gly Gly Val Asn Gly Thr Lys Ala Leu Tyr Val His	
565 570 575	
aag gat ggt gag ttt tca caa ttt att gga gat aag ttg aaa tcg aaa	1776
Lys Asp Gly Glu Phe Ser Gln Phe Ile Gly Asp Lys Leu Lys Ser Lys	
580 585 590	
aca gaa tat gta att caa tat att gta aag gga aaa gct tct att ctt	1824
Thr Glu Tyr Val Ile Gln Tyr Ile Val Lys Gly Lys Ala Ser Ile Leu	
595 600 605	
ttg aaa gat gaa aaa aat ggt gat tgc att tat gaa gat aca aat aat	1872
Leu Lys Asp Glu Lys Asn Gly Asp Cys Ile Tyr Glu Asp Thr Asn Asn	
610 615 620	
ggt tta gaa gat ttt caa acc att act aaa agt ttt att aca gga acg	1920
Gly Leu Glu Asp Phe Gln Thr Ile Thr Lys Ser Phe Ile Thr Gly Thr	
625 630 635 640	
gat tct tca gga gtt cat tta ata ttt aat agt caa aat ggc gat gaa	1968
Asp Ser Ser Gly Val His Leu Ile Phe Asn Ser Gln Asn Gly Asp Glu	
645 650 655	
gca ttt ggg gaa aac ttt act att tca gaa att agg ctt tcc gaa gat	2016
Ala Phe Gly Glu Asn Phe Thr Ile Ser Glu Ile Arg Leu Ser Glu Asp	
660 665 670	
tta tta agt cca gaa ttg ata aat tca gat gct tgg gtt gga tct cag	2064
Leu Leu Ser Pro Glu Leu Ile Asn Ser Asp Ala Trp Val Gly Ser Gln	
675 680 685	
gga act tgg atc tca gga aat tca ctc act att aat agt aat gtg aat	2112
Gly Thr Trp Ile Ser Gly Asn Ser Leu Thr Ile Asn Ser Asn Val Asn	
690 695 700	
gga act ttt cga caa aac ctt tcg tta gaa agc tat tca act tat agt	2160
Gly Thr Phe Arg Gln Asn Leu Ser Leu Glu Ser Tyr Ser Thr Tyr Ser	
705 710 715 720	
atg aac ttt aat gtg aat gga ttt gcc aag gtg aca gta aga aat tcc	2208
Met Asn Phe Asn Val Asn Gly Phe Ala Lys Val Thr Val Arg Asn Ser	
725 730 735	
cgt gaa gta tta ttt gaa aaa aat tat ccg cag ctt tca cct aaa gat	2256
Arg Glu Val Leu Phe Glu Lys Asn Tyr Pro Gln Leu Ser Pro Lys Asp	
740 745 750	

att tct gaa aaa ttc aca act gca gcc aat aat acc ggg ttg tat gta 2304
 Ile Ser Glu Lys Phe Thr Thr Ala Ala Asn Asn Thr Gly Leu Tyr Val
 755 760 765

gag ctt tct cgt ttt aca tcg ggt ggc gct ata aat ttc cgg aat ttt 2352
 Glu Leu Ser Arg Phe Thr Ser Gly Gly Ala Ile Asn Phe Arg Asn Phe
 770 775 780

tcg att aag tga 2364
 Ser Ile Lys
 785

<210> 2

<211> 787

<212> PRT

<213> Bacillus thuringiensis

<400> 2

Met Asn Lys Asn Asn Thr Lys Leu Asn Ala Arg Ala Leu Pro Ser Phe
 1 5 10 15

Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
 20 25 30

Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asn Leu Thr Leu
 35 40 45

Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Glu Ile Ser Gly Lys
 50 55 60

Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
 65 70 75 80

Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
 85 90 95

Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asn Ala Ile Asn Thr
 100 105 110

Met Leu His Ile Tyr Leu Pro Lys Ile Thr Ser Met Leu Asn Asp Val
 115 120 125

Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys
 130 135 140

Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Val Ile Asn Val Asn Val
 145 150 155 160

Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Met
 165 170 175

Lys Tyr Val Asn Glu Lys Phe Glu Asp Leu Thr Phe Ala Thr Glu Thr
 180 185 190

Thr Leu Lys Val Lys Lys Asn Ser Ser Pro Ala Asp Ile Leu Asp Glu
 195 200 205

Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val
 210 215 220

Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly
 225 230 235 240

Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile
 245 250 255

Ala Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr
 260 265 270

Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Lys Ala Phe Leu Thr
 275 280 285

Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr
 290 295 300

Phe Ile Met Asn Glu His Leu Asp Lys Glu Lys Glu Glu Phe Arg Val
 305 310 315 320

Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala
 325 330 335

Lys Ala Lys Gly Ser Asn Glu Asp Ala Lys Ile Ile Val Glu Ala Lys
 340 345 350

Pro Gly Tyr Ala Leu Val Gly Phe Glu Met Ser Asn Asp Ser Ile Thr
 355 360 365

Val Leu Lys Ala Tyr Gln Ala Lys Leu Lys Gln Asp Tyr Gln Val Asp
 370 375 380

Lys Asp Ser Leu Ser Glu Ile Val Tyr Gly Asp Met Asp Lys Leu Leu
 385 390 395 400

Cys Pro Asp Gln Ser Glu Gln Ile Tyr Tyr Thr Asn Asn Ile Ala Phe
 405 410 415

Pro Asn Glu Tyr Val Ile Thr Lys Ile Thr Phe Thr Lys Lys Met Asn
 420 425 430

Ser Leu Arg Tyr Glu Ala Thr Ala Asn Phe Tyr Asp Ser Ser Thr Gly
 435 440 445

Asp Ile Asp Leu Asn Lys Thr Lys Val Glu Ser Ser Glu Ala Glu Tyr
 450 455 460

Ser Thr Leu Ser Ala Ser Thr Asp Gly Val Tyr Met Pro Leu Gly Ile
 465 470 475 480

Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Ile Val Val
 485 490 495

Asp Glu Asn Ser Lys Leu Val Asn Leu Thr Cys Lys Ser Tyr Leu Arg
 500 505 510

Glu Val Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile
 515 520 525

Val Pro Pro Ile Gly Phe Ile Ser Asn Ile Val Glu Asn Gly Asn Leu
 530 535 540

Glu Gly Glu Asn Leu Glu Pro Trp Lys Ala Asn Asn Lys Asn Ala Tyr
 545 550 555 560

Val Asp His Thr Gly Gly Val Asn Gly Thr Lys Ala Leu Tyr Val His
 565 570 575

Lys Asp Gly Glu Phe Ser Gln Phe Ile Gly Asp Lys Leu Lys Ser Lys
 580 585 590

Thr Glu Tyr Val Ile Gln Tyr Ile Val Lys Gly Lys Ala Ser Ile Leu
 595 600 605

Leu Lys Asp Glu Lys Asn Gly Asp Cys Ile Tyr Glu Asp Thr Asn Asn
 610 615 620

Gly Leu Glu Asp Phe Gln Thr Ile Thr Lys Ser Phe Ile Thr Gly Thr
625 630 635 640

Asp Ser Ser Gly Val His Leu Ile Phe Asn Ser Gln Asn Gly Asp Glu
645 650 655

Ala Phe Gly Glu Asn Phe Thr Ile Ser Glu Ile Arg Leu Ser Glu Asp
660 665 670

Leu Leu Ser Pro Glu Leu Ile Asn Ser Asp Ala Trp Val Gly Ser Gln
675 680 685

Gly Thr Trp Ile Ser Gly Asn Ser Leu Thr Ile Asn Ser Asn Val Asn
690 695 700

Gly Thr Phe Arg Gln Asn Leu Ser Leu Glu Ser Tyr Ser Thr Tyr Ser
705 710 715 720

Met Asn Phe Asn Val Asn Gly Phe Ala Lys Val Thr Val Arg Asn Ser
725 730 735

Arg Glu Val Leu Phe Glu Lys Asn Tyr Pro Gln Leu Ser Pro Lys Asp
740 745 750

Ile Ser Glu Lys Phe Thr Thr Ala Ala Asn Asn Thr Gly Leu Tyr Val
755 760 765

Glu Leu Ser Arg Phe Thr Ser Gly Gly Ala Ile Asn Phe Arg Asn Phe
770 775 780

Ser Ile Lys
785

<210> 3

<211> 2364

<212> DNA

<213> Synthetic DNA

<220>

<221> misc_feature

<222> (1) .. (2364)

<223> Maize optimized vip3B

<400> 3

```

atgaacatga acaacaccaa gctcaacgcc cgcgccctcc cgtccttcat cgactacttc      60
aacggcatct acggcttcgc caccggcatc aaggacatca tgaacatgat cttcaagacc     120
gacaccggcg gcaacctcac cctcgacgag atcctcaaga accagcagct cctcaacgag     180
atctccggca agctcgacgg cgtgaacggc tcctcaacg acctcatcgc ccagggcaac     240
ctcaacaccg agctgtccaa ggagatcctc aagatcgcca acgagcagaa ccaggtgctc     300
aacgacgtga acaacaagct caacgccatc aacaccatgc tccacatcta cctcccgaag     360
atcacctcca tgctcaacga cgtgatgaag cagaactacg ccctctccct ccagatcgag     420
tacctctcca agcagctcca ggagatttcc gacaagctcg acgtgatcaa cgtgaacgtg     480
ctcatcaact ccaccctcac cgagatcacc cggcctacc agcgcgatgaa gtacgtgaac     540
gagaagttcg aggacctcac cttcgccacc gagaccaccc tcaaggtgaa gaagaactcc     600
tccccggccg acatcctcga cgagctgacc gagctgaccg agctggccaa gtccgtgacc     660
aagaacgacg tggacggctt cgagttctac ctcaacacct tccacgacgt gatggtgggc     720
aacaacctct tcggccgctc cgcctcaag accgcctccg agctgatcgc caaggagaac     780
gtgaagacct ccggctccga ggtgggcaac gtgtacaact tcctcatcgt gctcaccgcc     840
ctccaggcca aggccttctt caccctcacc acctgcccga agctcctcgg cctcgccgac     900
atcgactaca cttcatcat gaacgagcac ctcgacaagg agaaggagga gttccgctg     960
aacatcctcc cgaccctctc caacaccttc tccaaccoga actacgcca ggccaagggc    1020
tccaacgagg acgccaagat catcgtggag gccaagccgg gctacgccct cgtgggcttc    1080
gagatgtcca acgactccat caccgtgctc aaggcctacc aggccaaagct caagcaggac    1140
taccaggtgg acaaggactc cctctccgag atcgtgtacg gcgacatgga caagctcctc    1200
tgcccggacc agtccgagca aatctactac accaacaaca tcgccttccc gaacgagtac    1260
gtgatcacca agatcacctt caccaagaag atgaactccc tccgctacga ggccaccgcc    1320
aacttctacg actcctccac cggcgacatc gacctcaaca agaccaaggt ggagtcctec    1380
gaggccgagt actccaccct ctccgcctcc accgacggcg tgtacatgcc gctcggcatc    1440
atctccgaga ccttctcac cccgatcaac ggcttcggca tcgtggtgga cgagaactcc    1500
aagctcgtga acctcacctg caagtccctc ctccgagagg tgctcctcgc caccgacctc    1560
tccaacaagg agaccaagct catcgtgccg ccgatoggct tcattctcaa catcgtggag    1620

```


aacggcaacc tggagggcga gaacctggag ccgtggaagg ccaacaacaa gaacgcctac 1680
 gtggaccaca cgggcggcgt gaacggcacc aagggccctct acgtgcacaa ggacggcgag 1740
 ttctcccagt tcatcggcga caagctcaag tccaagaccg agtacgtgat ccagtacatc 1800
 gtgaagggca aggcctccat cctcctcaag gacgagaaga acggcgactg catctacgag 1860
 gacaccaaca acggcctgga ggacttccag accatcacca agtccttcat caccggcacc 1920
 gactcctccg gcgtgcacct catcttcaac tcccagaacg gcgacgaggg cttcggcgag 1980
 aacttcacca tctccgagat ccgcctctcc gaggacctcc tctccccgga gctgatcaac 2040
 tccgacgcct ggggtgggctc ccagggcacc tggatctccg gcaactccct caccatcaac 2100
 tccaacgtga acggcacctt ccgccagaac ctctccctgg agtcctactc cacctactcc 2160
 atgaacttca acgtgaacgg cttcggccaag gtgaccgtgc gcaactcccg cgaggtgctc 2220
 ttcgagaaga actaccgca gctctccccg aaggacatct ccgagaagtt caccaccgcc 2280
 gccacaaca ccggcctcta cgtggagctg tcccgcttca cctccggcgg cgccatcaac 2340
 ttccgcaact tctccatcaa gtag 2364

<210> 4

<211> 2367

<212> DNA

<213> Bacillus thuringiensis

<220>

<221> CDS

<222> (1)..(2367)

<223> Native vip3A coding sequence

<400> 4

atg aac aag aat aat act aaa tta agc aca aga gcc tta cca agt ttt 48
 Met Asn Lys Asn Asn Thr Lys Leu Ser Thr Arg Ala Leu Pro Ser Phe
 1 5 10 15
 att gat tat ttt aat ggc att tat gga ttt gcc act ggt atc aaa gac 96
 Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
 20 25 30
 att atg aac atg att ttt aaa acg gat aca ggt ggt gat cta acc cta 144
 Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Thr Leu
 35 40 45

gac gaa att tta aag aat cag cag tta cta aat gat att tct ggt aaa 192
 Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Asp Ile Ser Gly Lys
 50 55 60

ttg gat ggg gtg aat gga agc tta aat gat ctt atc gca cag gga aac 240
 Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
 65 70 75 80

tta aat aca gaa tta tct aag gaa ata tta aaa att gca aat gaa caa 288
 Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
 85 90 95

aat caa gtt tta aat gat gtt aat aac aaa ctc gat gcg ata aat acg 336
 Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
 100 105 110

atg ctt cgg gta tat cta cct aaa att acc tct atg ttg agt gat gta 384
 Met Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
 115 120 125

atg aaa caa aat tat gcg cta agt ctg caa ata gaa tac tta agt aaa 432
 Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys
 130 135 140

caa ttg caa gag att tct gat aag ttg gat att att aat gta aat gta 480
 Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val
 145 150 155 160

ctt att aac tct aca ctt act gaa att aca cct gcg tat caa agg att 528
 Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile
 165 170 175

aaa tat gtg aac gaa aaa ttt gag gaa tta act ttt gct aca gaa act 576
 Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr
 180 185 190

agt tca aaa gta aaa aag gat ggc tct cct gca gat att ctt gat gag 624
 Ser Ser Lys Val Lys Lys Asp Gly Ser Pro Ala Asp Ile Leu Asp Glu
 195 200 205

tta act gag tta act gaa cta gcg aaa agt gta aca aaa aat gat gtg 672
 Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val
 210 215 220

gat ggt ttt gaa ttt tac ctt aat aca ttc cac gat gta atg gta gga 720
 Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly
 225 230 235 240

aat aat tta ttc ggg cgt tca gct tta aaa act gca tcg gaa tta att 768
 Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile
 245 250 255

act aaa gaa aat gtg aaa aca agt ggc agt gag gtc gga aat gtt tat 816
 Thr Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr
 260 265 270

aac ttc tta att gta tta aca gct ctg caa gca aaa gct ttt ctt act 864
 Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Lys Ala Phe Leu Thr
 275 280 285

tta aca aca tgc cga aaa tta tta ggc tta gca gat att gat tat act	912
Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr	
290 295 300	
tct att atg aat gaa cat tta aat aag gaa aaa gag gaa ttt aga gta	960
Ser Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe Arg Val	
305 310 315 320	
aac atc ctc cct aca ctt tct aat act ttt tct aat cct aat tat gca	1008
Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala	
325 330 335	
aaa gtt aaa gga agt gat gaa gat gca aag atg att gtg gaa gct aaa	1056
Lys Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu Ala Lys	
340 345 350	
cca gga cat gca ttg att ggg ttt gaa att agt aat gat tca att aca	1104
Pro Gly His Ala Leu Ile Gly Phe Glu Ile Ser Asn Asp Ser Ile Thr	
355 360 365	
gta tta aaa gta tat gag gct aag cta aaa caa aat tat caa gtc gat	1152
Val Leu Lys Val Tyr Glu Ala Lys Leu Lys Gln Asn Tyr Gln Val Asp	
370 375 380	
aag gat tcc tta tcg gaa gtt att tat ggt gat atg gat aaa tta ttg	1200
Lys Asp Ser Leu Ser Glu Val Ile Tyr Gly Asp Met Asp Lys Leu Leu	
385 390 395 400	
tgc cca gat caa tct gaa caa atc tat tat aca aat aac ata gta ttt	1248
Cys Pro Asp Gln Ser Glu Gln Ile Tyr Tyr Thr Asn Asn Ile Val Phe	
405 410 415	
cca aat gaa tat gta att act aaa att gat ttc act aaa aaa atg aaa	1296
Pro Asn Glu Tyr Val Ile Thr Lys Ile Asp Phe Thr Lys Lys Met Lys	
420 425 430	
act tta aga tat gag gta aca gcg aat ttt tat gat tct tct aca gga	1344
Thr Leu Arg Tyr Glu Val Thr Ala Asn Phe Tyr Asp Ser Ser Thr Gly	
435 440 445	
gaa att gac tta aat aag aaa aaa gta gaa tca agt gaa gcg gag tat	1392
Glu Ile Asp Leu Asn Lys Lys Lys Val Glu Ser Ser Glu Ala Glu Tyr	
450 455 460	
aga acg tta agt gct aat gat gat ggg gtg tat atg ccg tta ggt gtc	1440
Arg Thr Leu Ser Ala Asn Asp Asp Gly Val Tyr Met Pro Leu Gly Val	
465 470 475 480	
atc agt gaa aca ttt ttg act ccg att aat ggg ttt ggc ctc caa gct	1488
Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Leu Gln Ala	
485 490 495	
gat gaa aat tca aga tta att act tta aca tgt aaa tca tat tta aga	1536
Asp Glu Asn Ser Arg Leu Ile Thr Leu Thr Cys Lys Ser Tyr Leu Arg	
500 505 510	
gaa cta ctg cta gca aca gac tta agc aat aaa gaa act aaa ttg atc	1584
Glu Leu Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile	
515 520 525	

gtc ccg cca agt ggt ttt att agc aat att gta gag aac ggg tcc ata	1632
Val Pro Pro Ser Gly Phe Ile Ser Asn Ile Val Glu Asn Gly Ser Ile	
530 535 540	
gaa gag gac aat tta gag ccg tgg aaa gca aat aat aag aat gcg tat	1680
Glu Glu Asp Asn Leu Glu Pro Trp Lys Ala Asn Asn Lys Asn Ala Tyr	
545 550 555 560	
gta gat cat aca ggc gga gtg aat gga act aaa gct tta tat gtt cat	1728
Val Asp His Thr Gly Gly Val Asn Gly Thr Lys Ala Leu Tyr Val His	
565 570 575	
aag gac gga gga att tca caa ttt att gga gat aag tta aaa ccg aaa	1776
Lys Asp Gly Gly Ile Ser Gln Phe Ile Gly Asp Lys Leu Lys Pro Lys	
580 585 590	
act gag tat gta atc caa tat act gtt aaa gga aaa cct tct att cat	1824
Thr Glu Tyr Val Ile Gln Tyr Thr Val Lys Gly Lys Pro Ser Ile His	
595 600 605	
tta aaa gat gaa aat act gga tat att cat tat gaa gat aca aat aat	1872
Leu Lys Asp Glu Asn Thr Gly Tyr Ile His Tyr Glu Asp Thr Asn Asn	
610 615 620	
aat tta gaa gat tat caa act att aat aaa cgt ttt act aca gga act	1920
Asn Leu Glu Asp Tyr Gln Thr Ile Asn Lys Arg Phe Thr Thr Gly Thr	
625 630 635 640	
gat tta aag gga gtg tat tta att tta aaa agt caa aat gga gat gaa	1968
Asp Leu Lys Gly Val Tyr Leu Ile Leu Lys Ser Gln Asn Gly Asp Glu	
645 650 655	
gct tgg gga gat aac ttt att att ttg gaa att agt cct tct gaa aag	2016
Ala Trp Gly Asp Asn Phe Ile Ile Leu Glu Ile Ser Pro Ser Glu Lys	
660 665 670	
tta tta agt cca gaa tta att aat aca aat aat tgg acg agt acg gga	2064
Leu Leu Ser Pro Glu Leu Ile Asn Thr Asn Asn Trp Thr Ser Thr Gly	
675 680 685	
tca act aat att agc ggt aat aca ctc act ctt tat cag gga gga cga	2112
Ser Thr Asn Ile Ser Gly Asn Thr Leu Thr Leu Tyr Gln Gly Gly Arg	
690 695 700	
ggg att cta aaa caa aac ctt caa tta gat agt ttt tca act tat aga	2160
Gly Ile Leu Lys Gln Asn Leu Gln Leu Asp Ser Phe Ser Thr Tyr Arg	
705 710 715 720	
gtg tat ttt tct gtg tcc gga gat gct aat gta agg att aga aat tct	2208
Val Tyr Phe Ser Val Ser Gly Asp Ala Asn Val Arg Ile Arg Asn Ser	
725 730 735	
agg gaa gtg tta ttt gaa aaa aga tat atg agc ggt gct aaa gat gtt	2256
Arg Glu Val Leu Phe Glu Lys Arg Tyr Met Ser Gly Ala Lys Asp Val	
740 745 750	
tct gaa atg ttc act aca aaa ttt gag aaa gat aac ttt tat ata gag	2304
Ser Glu Met Phe Thr Thr Lys Phe Glu Lys Asp Asn Phe Tyr Ile Glu	
755 760 765	

ctt tct caa ggg aat aat tta tat ggt ggt cct att gta cat ttt tac 2352
 Leu Ser Gln Gly Asn Asn Leu Tyr Gly Gly Pro Ile Val His Phe Tyr
 770 775 780

gat gtc tct att aag 2367
 Asp Val Ser Ile Lys
 785

<210> 5

<211> 789

<212> PRT

<213> Bacillus thuringiensis

<400> 5

Met Asn Lys Asn Asn Thr Lys Leu Ser Thr Arg Ala Leu Pro Ser Phe
 1 5 10 15

Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
 20 25 30

Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Thr Leu
 35 40 45

Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Asp Ile Ser Gly Lys
 50 55 60

Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
 65 70 75 80

Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
 85 90 95

Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
 100 105 110

Met Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
 115 120 125

Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys
 130 135 140

Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val
 145 150 155 160

Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile
 165 170 175

Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr
 180 185 190

Ser Ser Lys Val Lys Lys Asp Gly Ser Pro Ala Asp Ile Leu Asp Glu
 195 200 205

Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val
 210 215 220

Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly
 225 230 235 240

Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile
 245 250 255

Thr Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr
 260 265 270

Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Lys Ala Phe Leu Thr
 275 280 285

Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr
 290 295 300

Ser Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe Arg Val
 305 310 315 320

Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala
 325 330 335

Lys Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu Ala Lys
 340 345 350

Pro Gly His Ala Leu Ile Gly Phe Glu Ile Ser Asn Asp Ser Ile Thr
 355 360 365

Val Leu Lys Val Tyr Glu Ala Lys Leu Lys Gln Asn Tyr Gln Val Asp
 370 375 380

Lys Asp Ser Leu Ser Glu Val Ile Tyr Gly Asp Met Asp Lys Leu Leu
 385 390 395 400

Cys Pro Asp Gln Ser Glu Gln Ile Tyr Tyr Thr Asn Asn Ile Val Phe
 405 410 415

Pro Asn Glu Tyr Val Ile Thr Lys Ile Asp Phe Thr Lys Lys Met Lys
 420 425 430

Thr Leu Arg Tyr Glu Val Thr Ala Asn Phe Tyr Asp Ser Ser Thr Gly
 435 440 445

Glu Ile Asp Leu Asn Lys Lys Lys Val Glu Ser Ser Glu Ala Glu Tyr
 450 455 460

Arg Thr Leu Ser Ala Asn Asp Asp Gly Val Tyr Met Pro Leu Gly Val
 465 470 475 480

Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Leu Gln Ala
 485 490 495

Asp Glu Asn Ser Arg Leu Ile Thr Leu Thr Cys Lys Ser Tyr Leu Arg
 500 505 510

Glu Leu Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile
 515 520 525

Val Pro Pro Ser Gly Phe Ile Ser Asn Ile Val Glu Asn Gly Ser Ile
 530 535 540

Glu Glu Asp Asn Leu Glu Pro Trp Lys Ala Asn Asn Lys Asn Ala Tyr
 545 550 555 560

Val Asp His Thr Gly Gly Val Asn Gly Thr Lys Ala Leu Tyr Val His
 565 570 575

Lys Asp Gly Gly Ile Ser Gln Phe Ile Gly Asp Lys Leu Lys Pro Lys
 580 585 590

Thr Glu Tyr Val Ile Gln Tyr Thr Val Lys Gly Lys Pro Ser Ile His
 595 600 605

Leu Lys Asp Glu Asn Thr Gly Tyr Ile His Tyr Glu Asp Thr Asn Asn
 610 615 620

Asn Leu Glu Asp Tyr Gln Thr Ile Asn Lys Arg Phe Thr Thr Gly Thr
 625 630 635 640

Asp Leu Lys Gly Val Tyr Leu Ile Leu Lys Ser Gln Asn Gly Asp Glu
 645 650 655

Ala Trp Gly Asp Asn Phe Ile Ile Leu Glu Ile Ser Pro Ser Glu Lys
 660 665 670

Leu Leu Ser Pro Glu Leu Ile Asn Thr Asn Asn Trp Thr Ser Thr Gly
 675 680 685

Ser Thr Asn Ile Ser Gly Asn Thr Leu Thr Leu Tyr Gln Gly Gly Arg
 690 695 700

Gly Ile Leu Lys Gln Asn Leu Gln Leu Asp Ser Phe Ser Thr Tyr Arg
 705 710 715 720

Val Tyr Phe Ser Val Ser Gly Asp Ala Asn Val Arg Ile Arg Asn Ser
 725 730 735

Arg Glu Val Leu Phe Glu Lys Arg Tyr Met Ser Gly Ala Lys Asp Val
 740 745 750

Ser Glu Met Phe Thr Thr Lys Phe Glu Lys Asp Asn Phe Tyr Ile Glu
 755 760 765

Leu Ser Gln Gly Asn Asn Leu Tyr Gly Gly Pro Ile Val His Phe Tyr
 770 775 780

Asp Val Ser Ile Lys
 785

<210> 6

<211> 2364

<212> DNA

<213> Artificial Sequence

<220>

<221> CDS

<222> (1)..(2364)

<223> Hybrid vip3A-B(1734)


```

<400> 6
atg aac aag aat aat act aaa tta agc aca aga gcc tta cca agt ttt      48
Met Asn Lys Asn Asn Thr Lys Leu Ser Thr Arg Ala Leu Pro Ser Phe
1          5          10          15

att gat tat ttt aat ggc att tat gga ttt gcc act ggt atc aaa gac      96
Ile Asp Tyr Phe Asn Gly Ile Tyr Gly Phe Ala Thr Gly Ile Lys Asp
          20          25          30

att atg aac atg att ttt aaa acg gat aca ggt ggt gat cta acc cta      144
Ile Met Asn Met Ile Phe Lys Thr Asp Thr Gly Gly Asp Leu Thr Leu
          35          40          45

gac gaa att tta aag aat cag cag tta cta aat gat att tct ggt aaa      192
Asp Glu Ile Leu Lys Asn Gln Gln Leu Leu Asn Asp Ile Ser Gly Lys
          50          55          60

ttg gat ggg gtg aat gga agc tta aat gat ctt atc gca cag gga aac      240
Leu Asp Gly Val Asn Gly Ser Leu Asn Asp Leu Ile Ala Gln Gly Asn
65          70          75          80

tta aat aca gaa tta tct aag gaa ata tta aaa att gca aat gaa caa      288
Leu Asn Thr Glu Leu Ser Lys Glu Ile Leu Lys Ile Ala Asn Glu Gln
          85          90          95

aat caa gtt tta aat gat gtt aat aac aaa ctc gat gcg ata aat acg      336
Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
          100          105          110

atg ctt cgg gta tat cta cct aaa att acc tct atg ttg agt gat gta      384
Met Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
          115          120          125

atg aaa caa aat tat gcg cta agt ctg caa ata gaa tac tta agt aaa      432
Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys
          130          135          140

caa ttg caa gag att tct gat aag ttg gat att att aat gta aat gta      480
Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val
145          150          155          160

ctt att aac tct aca ctt act gaa att aca cct gcg tat caa agg att      528
Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile
          165          170          175

aaa tat gtg aac gaa aaa ttt gag gaa tta act ttt gct aca gaa act      576
Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr
          180          185          190

agt tca aaa gta aaa aag gat ggc tct cct gca gat att ctt gat gag      624
Ser Ser Lys Val Lys Lys Asp Gly Ser Pro Ala Asp Ile Leu Asp Glu
          195          200          205

tta act gag tta act gaa cta gcg aaa agt gta aca aaa aat gat gtg      672
Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val
          210          215          220

gat ggt ttt gaa ttt tac ctt aat aca ttc cac gat gta atg gta gga      720

```

Asp 225	Gly	Phe	Glu	Phe	Tyr 230	Leu	Asn	Thr	Phe	His 235	Asp	Val	Met	Val	Gly 240	
aat	aat	tta	ttc	ggg	cgt	tca	gct	tta	aaa	act	gca	tcg	gaa	tta	att	768
Asn	Asn	Leu	Phe	Gly 245	Arg	Ser	Ala	Leu	Lys 250	Thr	Ala	Ser	Glu	Leu	Ile 255	
act	aaa	gaa	aat	gtg	aaa	aca	agt	ggc	agt	gag	gtc	gga	aat	gtt	tat	816
Thr	Lys	Glu	Asn	Val 260	Lys	Thr	Ser	Gly 265	Ser	Glu	Val	Gly	Asn	Val 270	Tyr	
aac	ttc	tta	att	gta	tta	aca	gct	ctg	caa	gca	aaa	gct	ttt	ctt	act	864
Asn	Phe	Leu	Ile	Val	Leu	Thr	Ala	Leu	Gln	Ala	Lys	Ala	Phe	Leu	Thr 285	
tta	aca	aca	tgc	cga	aaa	tta	tta	ggc	tta	gca	gat	att	gat	tat	act	912
Leu	Thr	Thr	Cys	Arg	Lys	Leu	Leu	Gly 295	Leu	Ala	Asp	Ile	Asp	Tyr	Thr 300	
tct	att	atg	aat	gaa	cat	tta	aat	aag	gaa	aaa	gag	gaa	ttt	aga	gta	960
Ser	Ile	Met	Asn	Glu	His	Leu	Asn	Lys	Glu	Lys	Glu	Glu	Phe	Arg	Val 320	
305					310					315						
aac	atc	ctc	cct	aca	ctt	tct	aat	act	ttt	tct	aat	cct	aat	tat	gca	1008
Asn	Ile	Leu	Pro	Thr	Leu	Ser	Asn	Thr	Phe	Ser	Asn	Pro	Asn	Tyr	Ala 335	
325										330						
aaa	gtt	aaa	gga	agt	gat	gaa	gat	gca	aag	atg	att	gtg	gaa	gct	aaa	1056
Lys	Val	Lys	Gly	Ser	Asp	Glu	Asp	Ala	Lys	Met	Ile	Val	Glu	Ala	Lys 350	
340									345							
cca	gga	cat	gca	ttg	att	ggg	ttt	gaa	att	agt	aat	gat	tca	att	aca	1104
Pro	Gly	His	Ala	Leu	Ile	Gly	Phe	Glu	Ile	Ser	Asn	Asp	Ser	Ile	Thr 365	
355																
gta	tta	aaa	gta	tat	gag	gct	aag	cta	aaa	caa	aat	tat	caa	gtc	gat	1152
Val	Leu	Lys	Val	Tyr	Glu	Ala	Lys	Leu	Lys	Gln	Asn	Tyr	Gln	Val	Asp 380	
370																
aag	gat	tcc	tta	tcg	gaa	gtt	att	tat	ggt	gat	atg	gat	aaa	tta	ttg	1200
Lys	Asp	Ser	Leu	Ser	Glu	Val	Ile	Tyr	Gly	Asp	Met	Asp	Lys	Leu	Leu 400	
385					390					395						
tgc	cca	gat	caa	tct	gaa	caa	atc	tat	tat	aca	aat	aac	ata	gta	ttt	1248
Cys	Pro	Asp	Gln	Ser	Glu	Gln	Ile	Tyr	Tyr	Thr	Asn	Asn	Ile	Val	Phe 415	
405										410						
cca	aat	gaa	tat	gta	att	act	aaa	att	gat	ttc	act	aaa	aaa	atg	aaa	1296
Pro	Asn	Glu	Tyr	Val	Ile	Thr	Lys	Ile	Asp	Phe	Thr	Lys	Lys	Met	Lys 430	
420									425							
act	tta	aga	tat	gag	gta	aca	gcg	aat	ttt	tat	gat	tct	tct	aca	gga	1344
Thr	Leu	Arg	Tyr	Glu	Val	Thr	Ala	Asn	Phe	Tyr	Asp	Ser	Ser	Thr	Gly 445	
435								440								
gaa	att	gac	tta	aat	aag	aaa	aaa	gta	gaa	tca	agt	gaa	gcg	gag	tat	1392
Glu	Ile	Asp	Leu	Asn	Lys	Lys	Lys	Val	Glu	Ser	Ser	Glu	Ala	Glu	Tyr 460	
450						455					460					
aga	acg	tta	agt	gct	aat	gat	gat	ggg	gtg	tat	atg	ccg	tta	ggt	gtc	1440

Arg	Thr	Leu	Ser	Ala	Asn	Asp	Asp	Gly	Val	Tyr	Met	Pro	Leu	Gly	Val	
465					470					475					480	
atc	agt	gaa	aca	ttt	ttg	act	ccg	att	aat	ggg	ttt	ggc	ctc	caa	gct	1488
Ile	Ser	Glu	Thr	Phe	Leu	Thr	Pro	Ile	Asn	Gly	Phe	Gly	Leu	Gln	Ala	
				485					490					495		
gat	gaa	aat	tca	aga	tta	att	act	tta	aca	tgt	aaa	tca	tat	tta	aga	1536
Asp	Glu	Asn	Ser	Arg	Leu	Ile	Thr	Leu	Thr	Cys	Lys	Ser	Tyr	Leu	Arg	
			500					505					510			
gaa	cta	ctg	cta	gca	aca	gac	tta	agc	aat	aaa	gaa	act	aaa	ttg	atc	1584
Glu	Leu	Leu	Leu	Ala	Thr	Asp	Leu	Ser	Asn	Lys	Glu	Thr	Lys	Leu	Ile	
		515					520					525				
gtc	ccg	cca	agt	ggg	ttt	att	agc	aat	att	gta	gag	aac	ggg	tcc	ata	1632
Val	Pro	Pro	Ser	Gly	Phe	Ile	Ser	Asn	Ile	Val	Glu	Asn	Gly	Ser	Ile	
	530					535					540					
gaa	gag	gac	aat	tta	gag	ccg	tgg	aaa	gca	aat	aat	aag	aat	gcg	tat	1680
Glu	Glu	Asp	Asn	Leu	Glu	Pro	Trp	Lys	Ala	Asn	Asn	Lys	Asn	Ala	Tyr	
545				550				555						560		
gta	gat	cat	aca	ggc	gga	gtg	aat	gga	act	aaa	gct	tta	tat	gtt	cat	1728
Val	Asp	His	Thr	Gly	Gly	Val	Asn	Gly	Thr	Lys	Ala	Leu	Tyr	Val	His	
			565				570							575		
aag	gat	ggg	gag	ttt	tca	caa	ttt	att	gga	gat	aag	ttg	aaa	tcg	aaa	1776
Lys	Asp	Gly	Glu	Phe	Ser	Gln	Phe	Ile	Gly	Asp	Lys	Leu	Lys	Ser	Lys	
			580				585					590				
aca	gaa	tat	gta	att	caa	tat	att	gta	aag	gga	aaa	gct	tct	att	ctt	1824
Thr	Glu	Tyr	Val	Ile	Gln	Tyr	Ile	Val	Lys	Gly	Lys	Ala	Ser	Ile	Leu	
		595					600					605				
ttg	aaa	gat	gaa	aaa	aat	ggg	gat	tgc	att	tat	gaa	gat	aca	aat	aat	1872
Leu	Lys	Asp	Glu	Lys	Asn	Gly	Asp	Cys	Ile	Tyr	Glu	Asp	Thr	Asn	Asn	
	610					615					620					
ggg	tta	gaa	gat	ttt	caa	acc	att	act	aaa	agt	ttt	att	aca	gga	acg	1920
Gly	Leu	Glu	Asp	Phe	Gln	Thr	Ile	Thr	Lys	Ser	Phe	Ile	Thr	Gly	Thr	
625					630				635					640		
gat	tct	tca	gga	ggt	cat	tta	ata	ttt	aat	agt	caa	aat	ggc	gat	gaa	1968
Asp	Ser	Ser	Gly	Val	His	Leu	Ile	Phe	Asn	Ser	Gln	Asn	Gly	Asp	Glu	
				645					650					655		
gca	ttt	ggg	gaa	aac	ttt	act	att	tca	gaa	att	agg	ctt	tcc	gaa	gat	2016
Ala	Phe	Gly	Glu	Asn	Phe	Thr	Ile	Ser	Glu	Ile	Arg	Leu	Ser	Glu	Asp	
			660					665					670			
tta	tta	agt	cca	gaa	ttg	ata	aat	tca	gat	gct	tgg	ggt	gga	tct	cag	2064
Leu	Leu	Ser	Pro	Glu	Leu	Ile	Asn	Ser	Asp	Ala	Trp	Val	Gly	Ser	Gln	
		675					680					685				
gga	act	tgg	atc	tca	gga	aat	tca	ctc	act	att	aat	agt	aat	gtg	aat	2112
Gly	Thr	Trp	Ile	Ser	Gly	Asn	Ser	Leu	Thr	Ile	Asn	Ser	Asn	Val	Asn	
	690					695					700					
gga	act	ttt	cga	caa	aac	ctt	tcg	tta	gaa	agc	tat	tca	act	tat	agt	2160

Gly	Thr	Phe	Arg	Gln	Asn	Leu	Ser	Leu	Glu	Ser	Tyr	Ser	Thr	Tyr	Ser	
705					710					715					720	
atg	aac	ttt	aat	gtg	aat	gga	ttt	gcc	aag	gtg	aca	gta	aga	aat	tcc	2208
Met	Asn	Phe	Asn	Val	Asn	Gly	Phe	Ala	Lys	Val	Thr	Val	Arg	Asn	Ser	
			725						730					735		
cgt	gaa	gta	tta	ttt	gaa	aaa	aat	tat	ccg	cag	ctt	tca	cct	aaa	gat	2256
Arg	Glu	Val	Leu	Phe	Glu	Lys	Asn	Tyr	Pro	Gln	Leu	Ser	Pro	Lys	Asp	
			740					745					750			
att	tct	gaa	aaa	ttc	aca	act	gca	gcc	aat	aat	acc	ggg	ttg	tat	gta	2304
Ile	Ser	Glu	Lys	Phe	Thr	Thr	Ala	Ala	Asn	Asn	Thr	Gly	Leu	Tyr	Val	
		755					760					765				
gag	ctt	tct	cgt	ttt	aca	tcg	ggg	ggc	gct	ata	aat	ttc	cgg	aat	ttt	2352
Glu	Leu	Ser	Arg	Phe	Thr	Ser	Gly	Gly	Ala	Ile	Asn	Phe	Arg	Asn	Phe	
	770					775					780					
tcg	att	aag	tga													2364
Ser	Ile	Lys														
785																
<210>	7															
<211>	787															
<212>	PRT															
<213>	Artificial Sequence															
<400>	7															
Met	Asn	Lys	Asn	Asn	Thr	Lys	Leu	Ser	Thr	Arg	Ala	Leu	Pro	Ser	Phe	
1			5						10					15		
Ile	Asp	Tyr	Phe	Asn	Gly	Ile	Tyr	Gly	Phe	Ala	Thr	Gly	Ile	Lys	Asp	
			20					25					30			
Ile	Met	Asn	Met	Ile	Phe	Lys	Thr	Asp	Thr	Gly	Gly	Asp	Leu	Thr	Leu	
		35					40					45				
Asp	Glu	Ile	Leu	Lys	Asn	Gln	Gln	Leu	Leu	Asn	Asp	Ile	Ser	Gly	Lys	
	50					55					60					
Leu	Asp	Gly	Val	Asn	Gly	Ser	Leu	Asn	Asp	Leu	Ile	Ala	Gln	Gly	Asn	
65					70					75					80	
Leu	Asn	Thr	Glu	Leu	Ser	Lys	Glu	Ile	Leu	Lys	Ile	Ala	Asn	Glu	Gln	
				85					90						95	

Asn Gln Val Leu Asn Asp Val Asn Asn Lys Leu Asp Ala Ile Asn Thr
 100 105 110

Met Leu Arg Val Tyr Leu Pro Lys Ile Thr Ser Met Leu Ser Asp Val
 115 120 125

Met Lys Gln Asn Tyr Ala Leu Ser Leu Gln Ile Glu Tyr Leu Ser Lys
 130 135 140

Gln Leu Gln Glu Ile Ser Asp Lys Leu Asp Ile Ile Asn Val Asn Val
 145 150 155 160

Leu Ile Asn Ser Thr Leu Thr Glu Ile Thr Pro Ala Tyr Gln Arg Ile
 165 170 175

Lys Tyr Val Asn Glu Lys Phe Glu Glu Leu Thr Phe Ala Thr Glu Thr
 180 185 190

Ser Ser Lys Val Lys Lys Asp Gly Ser Pro Ala Asp Ile Leu Asp Glu
 195 200 205

Leu Thr Glu Leu Thr Glu Leu Ala Lys Ser Val Thr Lys Asn Asp Val
 210 215 220

Asp Gly Phe Glu Phe Tyr Leu Asn Thr Phe His Asp Val Met Val Gly
 225 230 235 240

Asn Asn Leu Phe Gly Arg Ser Ala Leu Lys Thr Ala Ser Glu Leu Ile
 245 250 255

Thr Lys Glu Asn Val Lys Thr Ser Gly Ser Glu Val Gly Asn Val Tyr
 260 265 270

Asn Phe Leu Ile Val Leu Thr Ala Leu Gln Ala Lys Ala Phe Leu Thr
 275 280 285

Leu Thr Thr Cys Arg Lys Leu Leu Gly Leu Ala Asp Ile Asp Tyr Thr
 290 295 300

Ser Ile Met Asn Glu His Leu Asn Lys Glu Lys Glu Glu Phe Arg Val
 305 310 315 320

Asn Ile Leu Pro Thr Leu Ser Asn Thr Phe Ser Asn Pro Asn Tyr Ala
 325 330 335

Lys Val Lys Gly Ser Asp Glu Asp Ala Lys Met Ile Val Glu Ala Lys
 340 345 350

Pro Gly His Ala Leu Ile Gly Phe Glu Ile Ser Asn Asp Ser Ile Thr
 355 360 365

Val Leu Lys Val Tyr Glu Ala Lys Leu Lys Gln Asn Tyr Gln Val Asp
 370 375 380

Lys Asp Ser Leu Ser Glu Val Ile Tyr Gly Asp Met Asp Lys Leu Leu
 385 390 400

Cys Pro Asp Gln Ser Glu Gln Ile Tyr Tyr Thr Asn Asn Ile Val Phe
 405 410 415

Pro Asn Glu Tyr Val Ile Thr Lys Ile Asp Phe Thr Lys Lys Met Lys
 420 425 430

Thr Leu Arg Tyr Glu Val Thr Ala Asn Phe Tyr Asp Ser Ser Thr Gly
 435 440 445

Glu Ile Asp Leu Asn Lys Lys Lys Val Glu Ser Ser Glu Ala Glu Tyr
 450 455 460

Arg Thr Leu Ser Ala Asn Asp Asp Gly Val Tyr Met Pro Leu Gly Val
 465 470 475 480

Ile Ser Glu Thr Phe Leu Thr Pro Ile Asn Gly Phe Gly Leu Gln Ala
 485 490 495

Asp Glu Asn Ser Arg Leu Ile Thr Leu Thr Cys Lys Ser Tyr Leu Arg
 500 505 510

Glu Leu Leu Leu Ala Thr Asp Leu Ser Asn Lys Glu Thr Lys Leu Ile
 515 520 525

Val Pro Pro Ser Gly Phe Ile Ser Asn Ile Val Glu Asn Gly Ser Ile
 530 535 540

Glu Glu Asp Asn Leu Glu Pro Trp Lys Ala Asn Asn Lys Asn Ala Tyr
 545 550 555 560

Val Asp His Thr Gly Gly Val Asn Gly Thr Lys Ala Leu Tyr Val His
 565 570 575

Lys Asp Gly Glu Phe Ser Gln Phe Ile Gly Asp Lys Leu Lys Ser Lys
580 585 590

Thr Glu Tyr Val Ile Gln Tyr Ile Val Lys Gly Lys Ala Ser Ile Leu
595 600 605

Leu Lys Asp Glu Lys Asn Gly Asp Cys Ile Tyr Glu Asp Thr Asn Asn
610 615 620

Gly Leu Glu Asp Phe Gln Thr Ile Thr Lys Ser Phe Ile Thr Gly Thr
625 630 635 640

Asp Ser Ser Gly Val His Leu Ile Phe Asn Ser Gln Asn Gly Asp Glu
645 650 655

Ala Phe Gly Glu Asn Phe Thr Ile Ser Glu Ile Arg Leu Ser Glu Asp
660 665 670

Leu Leu Ser Pro Glu Leu Ile Asn Ser Asp Ala Trp Val Gly Ser Gln
675 680 685

Gly Thr Trp Ile Ser Gly Asn Ser Leu Thr Ile Asn Ser Asn Val Asn
690 695 700

Gly Thr Phe Arg Gln Asn Leu Ser Leu Glu Ser Tyr Ser Thr Tyr Ser
705 710 715 720

Met Asn Phe Asn Val Asn Gly Phe Ala Lys Val Thr Val Arg Asn Ser
725 730 735

Arg Glu Val Leu Phe Glu Lys Asn Tyr Pro Gln Leu Ser Pro Lys Asp
740 745 750

Ile Ser Glu Lys Phe Thr Thr Ala Ala Asn Asn Thr Gly Leu Tyr Val
755 760 765

Glu Leu Ser Arg Phe Thr Ser Gly Gly Ala Ile Asn Phe Arg Asn Phe
770 775 780

Ser Ile Lys
785

<210> 8

<211> 18

<212> DNA

<213> Synthetic DNA

<220>

<221> misc_feature

<223> Forward primer

<400> 8

gtgatctaac cctagacg

18

<210> 9

<211> 20

<212> DNA

<213> Synthetic DNA

<220>

<221> misc_feature

<223> Reverse primer

<400> 9

gctttagttc cattcactcc

20

<210> 10

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<221> misc_feature

<223> Forward primer Vip3A-N

<400> 10

atgaccaaga ataataactaa attaagcac

29

<210> 11

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<221> misc_feature

<223> Reverse primer Vipfus4

<400> 11

tccttatgaa catataaagc tttagttcca tt

32

<210> 12

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<221> misc_feature

<223> Reverse primer Vip3B-C

<400> 12

ggcgaattct cacttaatcg aaaaattccg gaaatttat

39

<210> 13

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<221> misc_feature

<223> Forward primer Vipfus3

<400> 13

aatggaacta aagctttata tgttcataag ga

32