



(51) International Patent Classification:
C12N 15/11 (2006.01)

(21) International Application Number:
PCT/US2015/065393

(22) International Filing Date:
11 December 2015 (11.12.2015)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/091,462 12 December 2014 (12.12.2014) US
62/096,324 23 December 2014 (23.12.2014) US
62/180,681 17 June 2015 (17.06.2015) US
62/237,496 5 October 2015 (05.10.2015) US

(71) Applicants: THE BROAD INSTITUTE INC. [US/US];
415 Main Street, Cambridge, MA 02142 (US). MAS-
SACHUSETTS INSTITUTE OF TECHNOLOGY
[US/US]; 77 Massachusetts Ave., Cambridge, MA 02142
(US).

(72) Inventors: ZHANG, Feng; 100 Pacific Street, Apt. 11,
Cambridge, MA 02139 (US). KONERMANN, Silvana;
Foerrlibuckstr. 226, 8005 Zurich (CH). DAHLMAN,
James; 86 Rice Street, Cambridge, MA 02140 (US).
ABUDAYYEH, Omar, O.; c/o Massachusetts Institute of
Technology, 77 Massachusetts Ave., Cambridge, MA
02142 (US).


(74) Agents: KOWALSKI, Thomas, J. et al.; Vedder Price
P.c., 1633 Broadway, New York, NY 10019 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

[Continued on next page]

(54) Title: DEAD GUIDES FOR CRISPR TRANSCRIPTION FACTORS

Fig. 15



(57) Abstract: The invention provides for systems, methods, and compositions for altering expression of target gene sequences and related gene products. Provided are structural information on the Cas protein of the CRISPR-Cas system, use of this information in generating modified components of the CRISPR complex, vectors and vector systems which encode one or more components or modified components of a CRISPR complex, as well as methods for the design and use of such vectors and components. Also provided are methods of directing CRISPR complex formation in eukaryotic cells and methods for utilizing the CRISPR-Cas system. In particular the present invention comprehends optimized functional CRISPR-Cas enzyme systems.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

DEAD GUIDES FOR CRISPR TRANSCRIPTION FACTORS

RELATED APPLICATIONS AND INCORPORATION BY REFERENCE

[0001] This application claims priority from US application Serial No. 62/091,462, filed December 12, 2014, US application Serial No. 62/096,324, filed December 23, 2014, US application Serial No. 62/180,681, filed June 17, 2015 and US application Serial No. 62/237,496, filed October 5, 2015.

[0002] The foregoing applications, and all documents cited therein or during their prosecution (“appln cited documents”) and all documents cited or referenced in the appln cited documents, and all documents cited or referenced herein (“herein cited documents”), and all documents cited or referenced in herein cited documents, together with any manufacturer’s instructions, descriptions, product specifications, and product sheets for any products mentioned herein or in any document incorporated by reference herein, are hereby incorporated herein by reference, and may be employed in the practice of the invention. More specifically, all referenced documents are incorporated by reference to the same extent as if each individual document was specifically and individually indicated to be incorporated by reference.

[0003] Mention is made of US applications 62/091,455, filed December 12, 2014, 62/096,708, filed December 24, 2014, 62/180,709, filed June 17, 2015, and PCT/US2015/____ (Broad Institute reference no. BI-2014/100.WO1, attorney docket 46783.99.2085) entitled PROTECTED GUIDE RNAS (PGRNAS). Mention is also made of US applications 62/091,456, filed December 12, 2014, 62/180,692, filed June 17, 2015, and PCT/US2015/____ entitled ESCORTED AND FUNCTIONALIZED GUIDES FOR CRISPR-CAS SYSTEMS.

STATEMENT AS TO FEDERALLY SPONSORED RESEARCH

[0004] This invention was made with government support under grant numbers MH100706 and MH110049 awarded by the National Institutes of Health. The government has certain rights in the invention.

FIELD OF THE INVENTION

[0005] The present invention generally relates to systems, methods and compositions used for the control of gene expression involving sequence targeting, such as perturbation of gene

transcripts or nucleic acid editing, that may use vector systems related to Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) and components thereof.

BACKGROUND OF THE INVENTION

[0006] Recent advances in genome sequencing techniques and analysis methods have significantly accelerated the ability to catalog and map genetic factors associated with a diverse range of biological functions and diseases. Precise genome targeting technologies are needed to enable systematic reverse engineering of causal genetic variations by allowing selective perturbation of individual genetic elements, as well as to advance synthetic biology, biotechnological, and medical applications. Although genome-editing techniques such as designer zinc fingers, transcription activator-like effectors (TALEs), or homing meganucleases are available for producing targeted genome perturbations, there remains a need for new genome engineering technologies that employ novel strategies and molecular mechanisms and are affordable, easy to set up, scalable, and amenable to targeting multiple positions within the eukaryotic genome. This would provide a major resource for new applications in genome engineering and biotechnology.

[0007] Citation or identification of any document in this application is not an admission that such document is available as prior art to the present invention.

SUMMARY OF THE INVENTION

[0008] There exists a pressing need for alternative and robust systems and techniques for sequence targeting with a wide array of applications. This invention addresses this need and provides related advantages. The CRISPR/Cas9 or the CRISPR-Cas9 system (both terms are used interchangeably throughout this application) does not require the generation of customized proteins to target specific sequences but rather a single Cas9 enzyme can be programmed by a short RNA molecule to recognize a specific DNA target, in other words the Cas9 enzyme can be recruited to a specific DNA target using said short RNA molecule. Adding the CRISPR-Cas9 system to the repertoire of genome sequencing techniques and analysis methods may significantly simplify the methodology and accelerate the ability to catalog and map genetic factors associated with a diverse range of biological functions and diseases. To utilize the CRISPR-Cas9 system effectively for genome editing without deleterious effects, it is critical to understand aspects of engineering and optimization of these genome engineering tools, which are

aspects of the claimed invention. The terms 'CRISPR-Cas9' or 'CRISPR-Cas9 system' and 'nucleic acid-targeting system' may be used interchangeably. The terms 'CRISPR complex' and 'nucleic acid-targeting complex' be used interchangeably. Where reference is made herein to a 'target locus,' for example a target locus of interest, then it will be appreciated that this may be used interchangeably with the phrase 'sequences associated with or at a target locus of interest.'

[0009] In one aspect, the invention provides a method for altering or modifying expression of a gene product. The said method may comprise introducing into a cell containing and expressing a DNA molecule encoding the gene product an engineered, non-naturally occurring CRISPR-Cas system comprising a Cas9 protein and guide RNA that targets the DNA molecule, whereby the guide RNA targets the DNA molecule encoding the gene product and the Cas9 protein cleaves the DNA molecule encoding the gene product, whereby expression of the gene product is altered; and, wherein the Cas9 protein and the guide RNA do not naturally occur together. The invention comprehends the guide RNA comprising a guide sequence fused to a tracr sequence. The invention further comprehends the Cas9 protein being codon optimized for expression in a Eukaryotic cell. In a preferred embodiment the Eukaryotic cell is a mammalian cell and in a more preferred embodiment the mammalian cell is a human cell. In a further embodiment of the invention, the expression of the gene product is decreased.

[0010] In particular, an object of the current invention is to further enhance the specificity of Cas9 given individual guide RNAs through thermodynamic tuning of the binding specificity of the guide RNA to target DNA.

[0011] In one aspect, the invention provides an engineered, non-naturally occurring CRISPR-Cas9 system comprising a Cas9 protein and a guide RNA that targets a DNA molecule encoding a gene product in a cell, whereby the guide RNA targets the DNA molecule encoding the gene product and the Cas9 protein cleaves the DNA molecule encoding the gene product, whereby expression of the gene product is altered; and, wherein the Cas9 protein and the guide RNA do not naturally occur together. The invention comprehends the guide RNA comprising a guide sequence fused to a tracr sequence. The invention further comprehends the Cas9 protein being codon optimized for expression in a eukaryotic cell. In a preferred embodiment the Eukaryotic cell is a mammalian cell and in a more preferred embodiment the mammalian cell is a human cell. In a further embodiment of the invention, the expression of the gene product is decreased.

[0012] In another aspect, the invention provides an engineered, non-naturally occurring vector system comprising one or more vectors comprising a first regulatory element operably linked to a CRISPR-Cas9 system guide RNA that targets a DNA molecule encoding a gene product and a second regulatory element operably linked to a Cas9 protein. Components (a) and (b) may be located on same or different vectors of the system. The guide RNA targets the DNA molecule encoding the gene product in a cell and the Cas9 protein cleaves the DNA molecule encoding the gene product, whereby expression of the gene product is altered; and, wherein the Cas9 protein and the guide RNA do not naturally occur together. The invention comprehends the guide RNA comprising a guide sequence fused to a tracr sequence. The invention further comprehends the Cas9 protein being codon optimized for expression in a Eukaryotic cell. In a preferred embodiment the eukaryotic cell is a mammalian cell and in a more preferred embodiment the mammalian cell is a human cell. In a further embodiment of the invention, the expression of the gene product is decreased.

[0013] In one aspect, the invention provides a vector system comprising one or more vectors. In some embodiments, the system comprises: (a) a first regulatory element operably linked to a tracr mate sequence and one or more insertion sites for inserting one or more guide sequences upstream of the tracr mate sequence, wherein when expressed, the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell, wherein the CRISPR complex comprises a CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence, and (2) the tracr mate sequence that is hybridized to the tracr sequence; and (b) a second regulatory element operably linked to an enzyme-coding sequence encoding said CRISPR enzyme comprising a nuclear localization sequence; wherein components (a) and (b) are located on the same or different vectors of the system. In some embodiments, component (a) further comprises the tracr sequence downstream of the tracr mate sequence under the control of the first regulatory element. In some embodiments, component (a) further comprises two or more guide sequences operably linked to the first regulatory element, wherein when expressed, each of the two or more guide sequences direct sequence specific binding of a CRISPR complex to a different target sequence in a eukaryotic cell. In some embodiments, the system comprises the tracr sequence under the control of a third regulatory element, such as a polymerase III promoter. In some embodiments, the tracr sequence exhibits at least 50%, 60%, 70%, 80%, 90%, 95%, or 99% of sequence

complementarity along the length of the tracr mate sequence when optimally aligned. Determining optimal alignment is within the purview of one of skill in the art. For example, there are publically and commercially available alignment algorithms and programs such as, but not limited to, ClustalW, Smith-Waterman in matlab, Bowtie, Geneious, Biopython and SeqMan. In some embodiments, the CRISPR complex comprises one or more nuclear localization sequences of sufficient strength to drive accumulation of said CRISPR complex in a detectable amount in the nucleus of a eukaryotic cell. Without wishing to be bound by theory, it is believed that a nuclear localization sequence is not necessary for CRISPR complex activity in eukaryotes, but that including such sequences enhances activity of the system, especially as to targeting nucleic acid molecules in the nucleus. In some embodiments, the CRISPR enzyme is a type II CRISPR system enzyme. In some embodiments, the CRISPR enzyme is a Cas9 enzyme. In some embodiments, the Cas9 enzyme is *S. pneumoniae*, *S. pyogenes*, or *S. thermophilus* Cas9, and may include mutated Cas9 derived from these organisms. The enzyme may be a Cas9 homolog or ortholog. In some embodiments, the CRISPR-Cas9 enzyme is codon-optimized for expression in a eukaryotic cell. In some embodiments, the CRISPR-Cas9 enzyme directs cleavage of one or two strands at the location of the target sequence. In some embodiments, the first regulatory element is a polymerase III promoter. In some embodiments, the second regulatory element is a polymerase II promoter. In some embodiments, the guide sequence is at least 15, 16, 17, 18, 19, 20, 25 nucleotides, or between 10-30, or between 15-25, or between 15-20 nucleotides in length.

[0014] In general, and throughout this specification, the term “vector” refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. Vectors include, but are not limited to, nucleic acid molecules that are single-stranded, double-stranded, or partially double-stranded; nucleic acid molecules that comprise one or more free ends, no free ends (e.g. circular); nucleic acid molecules that comprise DNA, RNA, or both; and other varieties of polynucleotides known in the art. One type of vector is a “plasmid,” which refers to a circular double stranded DNA loop into which additional DNA segments can be inserted, such as by standard molecular cloning techniques. Another type of vector is a viral vector, wherein virally-derived DNA or RNA sequences are present in the vector for packaging into a virus (e.g. retroviruses, replication defective retroviruses, adenoviruses, replication defective adenoviruses, and adeno-associated viruses). Viral vectors also include polynucleotides carried by a virus for

transfection into a host cell. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g. bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively-linked. Such vectors are referred to herein as “expression vectors.” Vectors for and that result in expression in a eukaryotic cell can be referred to herein as “eukaryotic expression vectors.” Common expression vectors of utility in recombinant DNA techniques are often in the form of plasmids.

[0015] Recombinant expression vectors can comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory elements, which may be selected on the basis of the host cells to be used for expression, that is operatively-linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, “operably linked” is intended to mean that the nucleotide sequence of interest is linked to the regulatory element(s) in a manner that allows for expression of the nucleotide sequence (e.g. in an in vitro transcription/translation system or in a host cell when the vector is introduced into the host cell).

[0016] The term “regulatory element” is intended to include promoters, enhancers, internal ribosomal entry sites (IRES), and other expression control elements (e.g. transcription termination signals, such as polyadenylation signals and poly-U sequences). Such regulatory elements are described, for example, in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Regulatory elements include those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). A tissue-specific promoter may direct expression primarily in a desired tissue of interest, such as muscle, neuron, bone, skin, blood, specific organs (e.g. liver, pancreas), or particular cell types (e.g. lymphocytes). Regulatory elements may also direct expression in a temporal-dependent manner, such as in a cell-cycle dependent or developmental stage-dependent manner, which may or may not also be tissue or cell-type specific. In some embodiments, a vector comprises one or more pol III promoter (e.g. 1, 2, 3, 4, 5, or more pol I promoters), one or more pol II promoters (e.g. 1, 2, 3, 4, 5, or more pol II

promoters), one or more pol I promoters (e.g. 1, 2, 3, 4, 5, or more pol I promoters), or combinations thereof. Examples of pol III promoters include, but are not limited to, U6 and H1 promoters. Examples of pol II promoters include, but are not limited to, the retroviral Rous sarcoma virus (RSV) LTR promoter (optionally with the RSV enhancer), the cytomegalovirus (CMV) promoter (optionally with the CMV enhancer) [see, e.g., Boshart et al, Cell, 41:521-530 (1985)], the SV40 promoter, the dihydrofolate reductase promoter, the β -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1 α promoter. Also encompassed by the term "regulatory element" are enhancer elements, such as WPRE; CMV enhancers; the R-U5' segment in LTR of HTLV-I (Mol. Cell. Biol., Vol. 8(1), p. 466-472, 1988); SV40 enhancer; and the intron sequence between exons 2 and 3 of rabbit β -globin (Proc. Natl. Acad. Sci. USA., Vol. 78(3), p. 1527-31, 1981). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression desired, etc. A vector can be introduced into host cells to thereby produce transcripts, proteins, or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (e.g., clustered regularly interspersed short palindromic repeats (CRISPR) transcripts, proteins, enzymes, mutant forms thereof, fusion proteins thereof, etc.). Advantageous vectors further include lentiviruses and adeno-associated viruses, and types of such vectors can also be selected for targeting particular types of cells.

[0017] In one aspect, the invention provides a eukaryotic host cell comprising (a) a first regulatory element operably linked to a tracr mate sequence and one or more insertion sites for inserting one or more guide sequences upstream of the tracr mate sequence, wherein when expressed, the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell, wherein the CRISPR complex comprises a CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence, and (2) the tracr mate sequence that is hybridized to the tracr sequence; and/or (b) a second regulatory element operably linked to an enzyme-coding sequence encoding said CRISPR enzyme comprising a nuclear localization sequence. In some embodiments, the host cell comprises components (a) and (b). In some embodiments, component (a), component (b), or components (a) and (b) are stably integrated into a genome of the host eukaryotic cell. In some embodiments, component (a) further comprises the tracr sequence downstream of the tracr mate sequence under the control of the first regulatory element. In some embodiments, component (a) further comprises two or

more guide sequences operably linked to the first regulatory element, wherein when expressed, each of the two or more guide sequences direct sequence specific binding of a CRISPR complex to a different target sequence in a eukaryotic cell. In some embodiments, the eukaryotic host cell further comprises a third regulatory element, such as a polymerase III promoter, operably linked to said tracr sequence. In some embodiments, the tracr sequence exhibits at least 50%, 60%, 70%, 80%, 90%, 95%, or 99% of sequence complementarity along the length of the tracr mate sequence when optimally aligned. The enzyme may be a Cas9 homolog or ortholog. In some embodiments, the CRISPR-Cas9 enzyme is codon-optimized for expression in a eukaryotic cell. In some embodiments, the CRISPR-Cas9 enzyme directs cleavage of one or two strands at the location of the target sequence. In some embodiments, the CRISPR-Cas9 enzyme lacks DNA strand cleavage activity. In some embodiments, the first regulatory element is a polymerase III promoter. In some embodiments, the second regulatory element is a polymerase II promoter. In some embodiments, the guide sequence is at least 15, 16, 17, 18, 19, 20, 25 nucleotides, or between 10-30, or between 15-25, or between 15-20 nucleotides in length. In an aspect, the invention provides a non-human eukaryotic organism; preferably a multicellular eukaryotic organism, comprising a eukaryotic host cell according to any of the described embodiments. In other aspects, the invention provides a eukaryotic organism; preferably a multicellular eukaryotic organism, comprising a eukaryotic host cell according to any of the described embodiments. The organism in some embodiments of these aspects may be an animal; for example a mammal. Also, the organism may be an arthropod such as an insect. The organism also may be a plant. Further, the organism may be a fungus.

[0018] With respect to use of the CRISPR-Cas9 system generally, mention is made of the documents, including patent applications, patents, and patent publications cited throughout this disclosure as embodiments of the invention can be used as in those documents. CRISPR-Cas9 system(s) (e.g., single or multiplexed) can be used in conjunction with recent advances in crop genomics. Such CRISPR-Cas9 system(s) can be used to perform efficient and cost effective plant gene or genome interrogation or editing or manipulation—for instance, for rapid investigation and/or selection and/or interrogations and/or comparison and/or manipulations and/or transformation of plant genes or genomes; e.g., to create, identify, develop, optimize, or confer trait(s) or characteristic(s) to plant(s) or to transform a plant genome. There can accordingly be improved production of plants, new plants with new combinations of traits or characteristics or

new plants with enhanced traits. Such CRISPR-Cas9 system(s) can be used with regard to plants in Site-Directed Integration (SDI) or Gene Editing (GE) or any Near Reverse Breeding (NRB) or Reverse Breeding (RB) techniques. With respect to use of the CRISPR-Cas9 system in plants, mention is made of the University of Arizona website "CRISPR-PLANT" (<http://www.genome.arizona.edu/crispr/>) (supported by Penn State and AGI). Embodiments of the invention can be used in genome editing in plants or where RNAi or similar genome editing techniques have been used previously; see, e.g., Nekrasov, "Plant genome editing made easy: targeted mutagenesis in model and crop plants using the CRISPR/Cas system," *Plant Methods* 2013, 9:39 (doi:10.1186/1746-4811-9-39); Brooks, "Efficient gene editing in tomato in the first generation using the CRISPR/Cas9 system," *Plant Physiology* September 2014 pp 114.247577; Shan, "Targeted genome modification of crop plants using a CRISPR-Cas system," *Nature Biotechnology* 31, 686-688 (2013); Feng, "Efficient genome editing in plants using a CRISPR/Cas system," *Cell Research* (2013) 23:1229-1232. doi:10.1038/cr.2013.114; published online 20 August 2013; Xie, "RNA-guided genome editing in plants using a CRISPR-Cas system," *Mol Plant*. 2013 Nov;6(6):1975-83. doi: 10.1093/mp/sst119. Epub 2013 Aug 17; Xu, "Gene targeting using the *Agrobacterium tumefaciens*-mediated CRISPR-Cas system in rice," *Rice* 2014, 7:5 (2014), Zhou et al., "Exploiting SNPs for biallelic CRISPR mutations in the outcrossing woody perennial *Populus* reveals 4-coumarate: CoA ligase specificity and Redundancy," *New Phytologist* (2015) (Forum) 1-4 (available online only at www.newphytologist.com); Caliando et al, "Targeted DNA degradation using a CRISPR device stably carried in the host genome, *NATURE COMMUNICATIONS* 6:6989, DOI: 10.1038/ncomms7989, www.nature.com/naturecommunications DOI: 10.1038/ncomms7989; US Patent No. 6,603,061 - *Agrobacterium*-Mediated Plant Transformation Method; US Patent No. 7,868,149 - Plant Genome Sequences and Uses Thereof and US 2009/0100536 - Transgenic Plants with Enhanced Agronomic Traits, all the contents and disclosure of each of which are herein incorporated by reference in their entirety. In the practice of the invention, the contents and disclosure of Morrell et al "Crop genomics: advances and applications," *Nat Rev Genet*. 2011 Dec 29;13(2):85-96; each of which is incorporated by reference herein including as to how herein embodiments may be used as to plants. Accordingly, reference herein to animal cells may also apply, *mutatis mutandis*, to plant cells unless otherwise apparent.

[0019] In one aspect, the invention provides guide sequences which are modified in a manner which allows for formation of the CRISPR complex and successful binding to the target, while at the same time, not allowing for successful nuclease activity (i.e. without nuclease activity / without indel activity). For matters of explanation such modified guide sequences are referred to as dead guides or dead guide sequences. These dead guides or dead guide sequences can be thought of as catalytically inactive or conformationally inactive with regard to nuclease activity. Nuclease activity may be measured using surveyor analysis or deep sequencing as commonly used in the art, preferably surveyor analysis. Similarly, dead guide sequences may not sufficiently engage in productive base pairing with respect to the ability to promote catalytic activity or to distinguish on-target and off-target binding activity. Briefly, the surveyor assay involves purifying and amplifying a CRISPR target site for a gene and forming heteroduplexes with primers amplifying the CRISPR target site. After re-anneal, the products are treated with SURVEYOR nuclease and SURVEYOR enhancer S (Transgenomics) following the manufacturer's recommended protocols, analyzed on gels, and quantified based upon relative band intensities.

[0020] As explained further herein, several structural parameters allow for a proper framework to arrive at such dead guides. For example, dead guides to be used for targeting Sp Cas9 are 10 – 16 nucleotides in length. Dead guides to be used for targeting Sa Cas9 are 15 - 19 nucleotides in length. Dead guide sequences are shorter than respective guide sequences which result in active Cas9-specific indel formation. Dead guides are 5%, 10%, 20%, 30%, 40%, 50%, shorter than respective guides directed to the same Cas9 leading to active Cas9-specific indel formation. More specifically, the guide sequences are 10 – 16 nucleotides in length for guides specific to Sp Cas9, more preferably 12 – 15 nucleotides in length, even more preferably 13-14 nucleotides in length and most preferably 13 nucleotides in length. Dead guide sequences of Sa Cas9 - specific sgRNAs may be 15-19 nucleotides in length, preferably 17-18 nucleotides in length, and most preferably 17 nucleotides in length.

[0021] As explained below and known in the art, one aspect of sgRNA – Cas9 specificity is the tracr sequence, which is to be appropriately linked to such guides. In particular, this implies that the tracr sequences are designed dependent on the origin of the Cas9. Thus, structural data available for validated dead guide sequences specific to Sp Cas9 may be used for designing Cas9 specific equivalents (e.g. guides specific to Sa Cas9). Structural similarity between, e.g., the

orthologous nuclease domains RuvC and HNH of Sp Cas9 and Sa Cas9 may be used to transfer design equivalent dead guides specific to Sa Cas9 (e.g. Cas9 specific equivalent). Thus, the dead guide herein may be appropriately modified in length and sequence to reflect such Cas9 specific equivalents, allowing for formation of the CRISPR complex and successful binding to the target, while at the same time, not allowing for successful nuclease activity. As one example, dead guide specific to Sp Cas9 with a nucleotide length of 13 may be used as a standard for determining structural similarity of Cas9 specific equivalents (e.g. formation of bulges, loops; as determined and accepted in the art).

[0022] The use of dead guides in the context herein as well as the state of the art provides a surprising and unexpected platform for network biology and/or systems biology in both in vitro, ex vivo, and in vivo applications, allowing for multiplex gene targeting, and in particular bidirectional multiplex gene targeting. Prior to the use of dead guides, addressing multiple targets, for example for activation, repression and/or silencing of gene activity, has been challenging and in some cases not possible. With the use of dead guides, multiple targets, and thus multiple activities, may be addressed, for example, in the same cell, in the same animal, or in the same patient. Such multiplexing may occur at the same time or staggered for a desired timeframe.

[0023] For example, the dead guides now allow for the first time to use sgRNA as a means for gene targeting, without the consequence of nuclease activity, while at the same time providing directed means for activation or repression. sgRNA comprising a dead guide may be modified to further include elements in a manner which allow for activation or repression of gene activity, in particular protein adaptors (e.g. aptamers) allowing for functional placement of gene effectors (e.g. activators or repressors of gene activity) (Konermann et al., "Genome-scale transcription activation by an engineered CRISPR-Cas9 complex," doi:10.1038/nature14136, incorporated herein by reference.). One example, is the incorporation of aptamers, as explained herein and in the state of the art. By engineering the sgRNA comprising a dead guide to incorporate protein-interacting aptamers (Konermann et al., "Genome-scale transcription activation by an engineered CRISPR-Cas9 complex," doi:10.1038/nature14136, incorporated herein by reference.), one may assemble a synthetic transcription activation complex consisting of multiple distinct effector domains. Such may be modeled after natural transcription activation processes. For example, an aptamer, which selectively binds an effector (e.g. an activator or

repressor; dimerized MS2 bacteriophage coat proteins as fusion proteins with an activator or repressor), or a protein which itself binds an effector (e.g. activator or repressor) may be appended to a sgRNA tetraloop and/or a stem-loop 2. In the case of MS2, the fusion protein MS2-VP64 binds to the tetraloop and/or stem-loop 2 and in turn mediates transcriptional up-regulation, for example for *Neurog2*. Other transcriptional activators are, for example, VP64, P65, HSF1, and MyoD1. By mere example of this concept, replacement of the MS2 stem-loops with PP7-interacting stem-loops may be used to recruit repressive elements.

[0024] Thus, one aspect is a sgRNA of the invention which comprises a dead guide, wherein the sgRNA further comprises modifications which provide for gene activation or repression. The sgRNA may comprise one or more aptamers. The aptamers may be specific to gene effectors, gene activators or gene repressors. Alternatively, the aptamers may be specific to a protein which in turn is specific to and recruits / binds a specific gene effector, gene activator or gene repressor. If there are multiple sites for activator or repressor recruitment, it is preferred that the sites are specific to either activators or repressors. If there are multiple sites for activator or repressor binding, the sites may be specific to the same activators or same repressors. The sites may also be specific to different activators or different repressors. The gene effectors, gene activators, gene repressors may be present in the form of fusion proteins.

[0025] One aspect of the invention is to take advantage of the modularity and customizability of the sgRNA scaffold to establish a series of sgRNA scaffolds with different binding sites (in particular aptamers) for recruiting distinct types of effectors in an orthogonal manner. Again, for matters of example and illustration of the broader concept, replacement of the MS2 stem-loops with PP7-interacting stem-loops may be used to bind / recruit repressive elements, enabling multiplexed bidirectional transcriptional control. Thus, in general, sgRNA comprising a dead guide may be employed to provide for multiplex transcriptional control and preferred bidirectional transcriptional control. This transcriptional control is most preferred of genes. For example, one or more sgRNA comprising dead guide(s) may employed in targeting the activation of one or more target genes. At the same time, one or more sgRNA comprising dead guide(s) may employed in targeting the repression of one or more target genes. Such a sequence may be applied in a variety of different combinations, for example the target genes are first repressed and then at an appropriate period other targets are activated, or select genes are repressed at the same time as select genes are activated, followed by further activation and/or

repression. As a result, multiple components of one or more biological systems may advantageously be addressed together.

[0026] In another aspect, structural analysis may also be used to study interactions between the dead Guide and the active Cas9 nuclease that enable DNA binding, but no DNA cutting. In this way amino acids important for nuclease activity of Cas9 are determined. Modification of such amino acids allows for improved Cas9 enzymes used for gene editing.

[0027] A further aspect is combining the use of dead guides as explained herein with other applications of CRISPR, as explained herein as well as known in the art. For example, sgRNA comprising dead guide(s) for targeted multiplex gene activation or repression or targeted multiplex bidirectional gene activation / repression may be combined with sgRNA comprising guides which maintain nuclease activity, as explained herein. Such sgRNA comprising guides which maintain nuclease activity may or may not further include modifications which allow for repression of gene activity (e.g. aptamers). Such sgRNA comprising guides which maintain nuclease activity may or may not further include modifications which allow for activation of gene activity (e.g. aptamers). In such a manner, a further means for multiplex gene control is introduced (e.g. multiplex gene targeted activation without nuclease activity / without indel activity may be provided at the same time or in combination with gene targeted repression with nuclease activity).

[0028] For example, 1) using one or more sgRNA (e.g. 1-50, 1-40, 1-30, 1-20, preferably 1-10, more preferably 1-5) comprising dead guide(s) targeted to one or more genes and further modified with appropriate aptamers for the recruitment of gene activators; 2) may be combined with one or more sgRNA (e.g. 1-50, 1-40, 1-30, 1-20, preferably 1-10, more preferably 1-5) comprising dead guide(s) targeted to one or more genes and further modified with appropriate aptamers for the recruitment of gene repressors. 1) and/or 2) may then be combined with 3) one or more sgRNA (e.g. 1-50, 1-40, 1-30, 1-20, preferably 1-10, more preferably 1-5) targeted to one or more genes. This combination can then be carried out in turn with 1) + 2) + 3) with 4) one or more sgRNA (e.g. 1-50, 1-40, 1-30, 1-20, preferably 1-10, more preferably 1-5) targeted to one or more genes and further modified with appropriate aptamers for the recruitment of gene activators. This combination can then be carried in turn with 1) + 2) + 3) + 4) with 5) one or more sgRNA (e.g. 1-50, 1-40, 1-30, 1-20, preferably 1-10, more preferably 1-5) targeted to one or more genes and further modified with appropriate aptamers for the recruitment of gene

repressors. As a result various uses and combinations are included in the invention. For example, combination 1) + 2); combination 1) + 3); combination 2) + 3); combination 1) + 2) + 3); combination 1) + 2) + 3) + 4); combination 1) + 3) + 4); combination 2) + 3) + 4); combination 1) + 2) + 4); combination 1) + 2) + 3) + 4) + 5); combination 1) + 3) + 4) + 5); combination 2) + 3) + 4) + 5); combination 1) + 2) + 4) + 5); combination 1) + 2) + 3) + 5); combination 1) + 3) + 5); combination 2) + 3) + 5); combination 1) + 2) + 5).

[0029] In an aspect, the invention provides an algorithm for designing, evaluating, or selecting a guide RNA targeting sequence for guiding a CRISPR-Cas9 system to a target gene locus.

[0030] In particular, it has been determined that guide RNA specificity relates to and can be optimized by varying i) GC content and ii) targeting sequence length. In an aspect, the invention provides an algorithm for designing or evaluating a guide RNA targeting sequence that minimizes off-target binding or interaction of the guide RNA. In an embodiment of the invention, the algorithm for selecting a guide RNA targeting sequence for directing a CRISPR system to a gene locus in an organism comprises a) locating one or more CRISPR motifs in the gene locus, analyzing the 20 nt sequence upstream of each CRISPR motif by i) determining the GC content of the sequence; and ii) determining whether there are off-target matches of the 15 upstream nucleotides nearest to the CRISPR motif in the genome of the organism, and c) selecting the 15 nucleotide sequence for use in a guide RNA if the GC content of the sequence is 70% or less and no off-target matches are identified. In an embodiment of the invention, the sequence is selected for a targeting sequence if the GC content is 60% or less. In certain embodiments of the invention, the sequence is selected for a targeting sequence if the GC content is 55% or less, 50% or less, 45% or less, 40% or less, 35% or less or 30% or less. Preferably, no off target matches are identified. In some embodiments, one or more off-target matches may be tolerated, depending on the location of the off-target sequence. For example, an off-target match in an intergenic locus or in a non-regulatory, untranscribed, or untranslated region of a gene may be tolerated. In an embodiment of the invention, no off-target matches are identified in transcribed sequences. In an embodiment of the invention, In an embodiment of the invention, no off-target matches are identified in translated sequences.

[0031] In an embodiment, two or more sequences of the gene locus are analyzed and the sequence having the lowest GC content, or the next lowest GC content, or the next lowest GC

content is selected. In an embodiment of the invention, the sequence is selected for a targeting sequence if no off-target matches are identified in the genome of the organism. In an embodiment of the invention, the targeting sequence is selected if no off-target matches are identified in regulatory sequences of the genome.

[0032] In an aspect, the invention provides a guide RNA for targeting a functionalized CRISPR system to a gene locus in an organism. In an embodiment of the invention, the guide RNA comprises a targeting sequence wherein the CG content of the target sequence is 70% or less, and the first 15 nt of the targeting sequence does not match an off-target sequence upstream from a CRISPR motif in the regulatory sequence of another gene locus in the organism. In certain embodiments, the GC content of the targeting sequence 60% or less, 55% or less, 50% or less, 45% or less, 40% or less, 35% or less or 30% or less. In certain embodiments, the GC content of the targeting sequence is from 70% to 60% or from 60% to 50% or from 50% to 40% or from 40% to 30%. In an embodiment, the targeting sequence has the lowest CG content among potential targeting sequences of the locus.

[0033] In an embodiment of the invention, the first 15 nt of the guide upstream from the CRISPR motif match the target sequence. In another embodiment, the first 14 nt of the guide match the target sequence. In another embodiment, the first 13 nt of the guide match the target sequence. In another embodiment first 12 nt of the guide match the target sequence. In another embodiment, first 11 nt of the guide match the target sequence. In another embodiment, the first 10 nt of the guide match the target sequence. In an embodiment of the invention the first 15 nt of the guide does not match an off-target sequence upstream from a CRISPR motif in the regulatory region of another gene locus. In other embodiments, the first 14 nt, or the first 13 nt of the guide, or the first 12 nt of the guide, or the first 11 nt of the guide, or the first 10 nt of the guide, does not match an off-target sequence upstream from a CRISPR motif in the regulatory region of another gene locus. In other embodiments, the first 15 nt, or 14 nt, or 13 nt, or 12 nt, or 11 nt of the guide do not match an off-target sequence upstream from a CRISPR motif in the genome.

[0034] In certain embodiments, the guide RNA includes additional nucleotides at the 5'-end that do not match the target sequence. Thus, a guide RNA that includes the first 15 nt, or 14 nt, or 13 nt, or 12 nt, or 11 nt upstream of a CRISPR motif can be extended in length at the 5' end to 12 nt, 13 nt, 14 nt, 15 nt, 16 nt, 17 nt, 18 nt, 19 nt, 20 nt, or longer.

[0035] The invention provides a method for directing a CRISPR-Cas9 system, including but not limited to a dead Cas9 (dCas9) or functionalized Cas9 system (which may comprise a functionalized Cas9 or functionalized guide) to a gene locus. In an aspect, the invention provides a method for selecting a guide RNA targeting sequence and directing a functionalized CRISPR system to a gene locus in an organism. In an aspect, the invention provides a method for selecting a guide RNA targeting sequence and effecting gene regulation of a target gene locus by a functionalized CRISPR-Cas9 system. In certain embodiments, the method is used to effect target gene regulation while minimizing off-target effects. In an aspect, the invention provides a method for selecting two or more guide RNA targeting sequences and effecting gene regulation of two or more target gene loci by a functionalized CRISPR-Cas9 system. In certain embodiments, the method is used to effect regulation of two or more target gene loci while minimizing off-target effects.

[0036] In an aspect, the invention provides for a single effector to be directed to one or more, or two or more gene loci. In certain embodiments, the effector is associated with a CRISPR protein or enzyme, and one or more, or two or more selected guide RNAs are used to direct the CRISPR-associated effector to one or more, or two or more selected target gene loci. In certain embodiments, the effector is associated with one or more, or two or more selected guide RNAs, each selected guide RNA, when complexed with a CRISPR protein or enzyme, causing its associated effector to be localized to the guide RNA target. One non-limiting example of such CRISPR systems modulates activity of one or more, or two or more gene loci subject to regulation by the same transcription factor.

[0037] In an aspect, the invention provides for two or more effectors to be directed to one or more gene loci. In certain embodiments, two or more guide RNAs are employed, each of the two or more effectors being associated with a selected guide RNA, with each of the two or more effectors being localized to the selected target of its guide RNA. One non-limiting example of such CRISPR systems modulates activity of one or more, or two or more gene loci subject to regulation by different transcription factors. Thus, in one non-limiting embodiment, two or more transcription factors are localized to different regulatory sequences of a single gene. In another non-limiting embodiment, two or more transcription factors are localized to different regulatory sequences of different genes. In certain embodiments, one transcription factor is an activator. In certain embodiments, one transcription factor is an inhibitor. In certain embodiments, one

transcription factor is an activator and another transcription factor is an inhibitor. In certain embodiments, gene loci expressing different components of the same regulatory pathway are regulated. In certain embodiments, gene loci expressing components of different regulatory pathways are regulated.

[0038] In certain of the above embodiments, a catalytically incompetent CRISPR protein is used. In certain of the above embodiments, an active CRISPR enzyme is used.

[0039] In an aspect, the invention also provides a method and algorithm for designing and selecting guide RNAs that are specific for target DNA cleavage or target binding and gene regulation mediated by an active CRISPR-Cas9 system. In certain embodiments, the CRISPR-Cas9 system provides orthogonal gene control using an active CRISPR enzyme which cleaves target DNA at one gene locus while at the same time binds to and promotes regulation of another gene locus.

[0040] In an aspect, the invention provides an method of selecting a guide RNA targeting sequence for directing a functionalized CRISPR enzyme to a gene locus in an organism, without cleavage, which comprises a) locating one or more CRISPR motifs in the gene locus; b) analyzing the sequence upstream of each CRISPR motif by i) selecting 10 to 15 nt adjacent to the CRISPR motif, ii) determining the GC content of the sequence, and c) selecting the 10 to 15 nt sequence as a targeting sequence for use in a guide RNA if the GC content of the sequence is 30% more, 40% or more. In certain embodiments, the GC content of the targeting sequence is 35% or more, 40% or more, 45% or more, 50% or more, 55% or more, 60% or more, 65% or more, or 70% or more. In certain embodiments, the GC content of the targeting sequence is from 30% to 40% or from 40% to 50% or from 50% to 60% or from 60% to 70%. In an embodiment of the invention, two or more sequences in a gene locus are analyzed and the sequence having the highest GC content is selected.

[0041] In an embodiment of the invention, the portion of the guide targeting sequence in which GC content is evaluated is 10 to 15 contiguous nucleotides of the 15 target nucleotides nearest to the PAM. In an embodiment of the invention, the portion of the guide in which GC content is considered is the 10 to 11 nucleotides or 11 to 12 nucleotides or 12 to 13 nucleotides or 13, or 14, or 15 contiguous nucleotides of the 15 nucleotides nearest to the PAM.

[0042] In an aspect, the invention further provides an algorithm for identifying guide RNAs which promote CRISPR system gene locus cleavage while avoiding functional activation or

inhibition. It is observed that increased GC content in guide RNAs of 16 to 20 nucleotides coincides with increased DNA cleavage and reduced functional activation.

[0043] It is also demonstrated herein that efficiency of functionalized CRISPR proteins and enzymes can be increased by addition of nucleotides to the 5' end of a guide RNA which do not match a target sequence upstream of the CRISPR motif. For example, of guide RNA 11 to 15 nt in length, shorter guides may be less likely to promote target cleavage, but are also less efficient at promoting CRISPR system binding and functional control. In certain embodiments, addition of nucleotides that don't match the target sequence to the 5' end of the guide RNA increase activation efficiency while not increasing undesired target cleavage. In an aspect, the invention also provides a method and algorithm for identifying improved guide RNAs that effectively promote CRISPR system function in DNA binding and gene regulation while not promoting DNA cleavage. Thus, in certain embodiments, the invention provides a guide RNA that includes the first 15 nt, or 14 nt, or 13 nt, or 12 nt, or 11 nt upstream of a CRISPR motif and is extended in length at the 5' end by nucleotides that mismatch the target to 12 nt, 13 nt, 14 nt, 15 nt, 16 nt, 17 nt, 18 nt, 19 nt, 20 nt, or longer.

[0044] In an aspect, the invention provides a method for effecting selective orthogonal gene control. As will be appreciated from the disclosure herein, guide selection according to the invention, taking into account guide length and GC content, provides effective and selective transcription control by a functional CRISPR-Cas system, for example to regulate transcription of a gene locus by activation or inhibition and minimize off-target effects. Accordingly, by providing effective regulation of individual target loci, the invention also provides effective orthogonal regulation of two or more target loci.

[0045] In certain embodiments, orthogonal gene control is by activation or inhibition of two or more target loci. In certain embodiments, orthogonal gene control is by activation or inhibition of one or more target locus and cleavage of one or more target locus.

[0046] In one aspect, the invention provides a cell comprising a non-naturally occurring CRISPR-Cas9 system comprising one or more guide RNAs disclosed or made according to a method or algorithm described herein wherein the expression of one or more gene products has been altered. In an embodiment of the invention, the expression in the cell of two or more gene products has been altered. The invention also provides a cell line from such a cell.

[0047] In one aspect, the invention provides a multicellular organism comprising one or more cells comprising a non-naturally occurring CRISPR-Cas9 system comprising one or more guide RNAs disclosed or made according to a method or algorithm described herein. In one aspect, the invention provides a product from a cell, cell line, or multicellular organism comprising a non-naturally occurring CRISPR-Cas9 system comprising one or more guide RNAs disclosed or made according to a method or algorithm described herein.

[0048] A further aspect of this invention is the use of sgRNA comprising dead guide(s) as described herein, optionally in combination with sgRNA comprising guide(s) as described herein or in the state of the art, in combination with systems (e.g. cells, transgenic animals, transgenic mice, inducible transgenic animals, inducible transgenic mice) which are engineered for either overexpression of Cas9 or preferably knockin Cas9, as explained, for example, in Platt et al., Cell 159, 440-455, Oct. 2014. As a result a single system (e.g. transgenic animal, cell) can serve as a basis for multiplex gene modifications in systems / network biology. On account of the dead guides, this is now possible in both in vitro, ex vivo, and in vivo.

[0049] For example, once the Cas9 is provided for (e.g. expression is knocked in; Platt et al., Cell 159, 440-455, Oct. 2014), one or more sgRNAs may be provided to direct multiplex gene regulation, and preferably multiplex bidirectional gene regulation. The one or more sgRNAs may be provided in a spatially and temporally appropriate manner if necessary or desired (for example tissue specific induction of Cas9 expression). On account that the transgenic / inducible Cas9 is provided for (e.g. expressed) in the cell, tissue, animal of interest, both sgRNAs comprising dead guides or sgRNAs comprising guides are equally effective. In the same manner, a further aspect of this invention is the use of sgRNA comprising dead guide(s) as described herein, optionally in combination with sgRNA comprising guide(s) as described herein or in the state of the art, in combination with systems (e.g. cells, transgenic animals, transgenic mice, inducible transgenic animals, inducible transgenic mice) which are engineered for knockout CRISPR-Cas9 as explained, for example, in Shalem et al., Science 12 December 2013, pp 1-7/10.1126/science.1247005.

[0050] As a result, the combination of dead guides as described herein with CRISPR applications described herein and CRISPR applications known in the art (e.g. inducible Cas9) results in a highly efficient and accurate means for multiplex screening of systems (e.g. network biology). Such screening allows, for example, identification of specific combinations of gene

activities for identifying genes responsible for diseases (e.g. on/off combinations), in particular gene related diseases. A preferred application of such screening is cancer. In the same manner, screening for treatment for such diseases is included in the invention. Cells or animals may be exposed to aberrant conditions resulting in disease or disease like effects. Candidate compositions may be provided and screened for an effect in the desired multiplex environment. For example a patient's cancer cells may be screened for which gene combinations will cause them to die, and then use this information to establish appropriate therapies.

[0051] In one aspect, the invention provides a kit comprising one or more of the components described herein. In some embodiments, the kit comprises a vector system and instructions for using the kit. In some embodiments, the vector system comprises (a) a first regulatory element operably linked to a *tracr* mate sequence and one or more insertion sites for inserting one or more guide sequences upstream of the *tracr* mate sequence, wherein when expressed, the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell, wherein the CRISPR complex comprises a CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence, and (2) the *tracr* mate sequence that is hybridized to the *tracr* sequence; and/or (b) a second regulatory element operably linked to an enzyme-coding sequence encoding said CRISPR enzyme comprising a nuclear localization sequence. In some embodiments, the kit comprises components (a) and (b) located on the same or different vectors of the system. In some embodiments, component (a) further comprises the *tracr* sequence downstream of the *tracr* mate sequence under the control of the first regulatory element. In some embodiments, component (a) further comprises two or more guide sequences operably linked to the first regulatory element, wherein when expressed, each of the two or more guide sequences direct sequence specific binding of a CRISPR complex to a different target sequence in a eukaryotic cell. In some embodiments, the system further comprises a third regulatory element, such as a polymerase III promoter, operably linked to said *tracr* sequence. In some embodiments, the *tracr* sequence exhibits at least 50%, 60%, 70%, 80%, 90%, 95%, or 99% of sequence complementarity along the length of the *tracr* mate sequence when optimally aligned. In some embodiments, the CRISPR enzyme comprises one or more nuclear localization sequences of sufficient strength to drive accumulation of said CRISPR enzyme in a detectable amount in the nucleus of a eukaryotic cell. In some embodiments, the CRISPR enzyme is a type II CRISPR system enzyme. In some embodiments, the CRISPR enzyme is a Cas9 enzyme. In

some embodiments, the Cas9 enzyme is *S. pneumoniae*, *S. pyogenes* or *S. thermophilus* Cas9, and may include mutated Cas9 derived from these organisms. The enzyme may be a Cas9 homolog or ortholog. In some embodiments, the CRISPR enzyme is codon-optimized for expression in a eukaryotic cell. In some embodiments, the CRISPR enzyme directs cleavage of one or two strands at the location of the target sequence. In some embodiments, the CRISPR enzyme lacks DNA strand cleavage activity. In some embodiments, the first regulatory element is a polymerase III promoter. In some embodiments, the second regulatory element is a polymerase II promoter. In some embodiments, the guide sequence is at least 15, 16, 17, 18, 19, 20, 25 nucleotides, or between 10-30, or between 15-25, or between 15-20 nucleotides in length. The kit may include dead guides as described herein with or without guides as described herein.

[0052] In one aspect, the invention provides a method of modifying a target polynucleotide in a eukaryotic cell. In some embodiments, the method comprises allowing a CRISPR complex to bind to the target polynucleotide to effect cleavage of said target polynucleotide thereby modifying the target polynucleotide, wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within said target polynucleotide, wherein said guide sequence is linked to a tracr mate sequence which in turn hybridizes to a tracr sequence. In some embodiments, said cleavage comprises cleaving one or two strands at the location of the target sequence by said CRISPR enzyme. In some embodiments, said cleavage results in decreased transcription of a target gene. In some embodiments, the method further comprises repairing said cleaved target polynucleotide by homologous recombination with an exogenous template polynucleotide, wherein said repair results in a mutation comprising an insertion, deletion, or substitution of one or more nucleotides of said target polynucleotide. In some embodiments, said mutation results in one or more amino acid changes in a protein expressed from a gene comprising the target sequence. In some embodiments, the method further comprises delivering one or more vectors to said eukaryotic cell, wherein the one or more vectors drive expression of one or more of: the CRISPR enzyme, the guide sequence linked to the tracr mate sequence, and the tracr sequence. In some embodiments, said vectors are delivered to the eukaryotic cell in a subject. In some embodiments, said modifying takes place in said eukaryotic cell in a cell culture. In some embodiments, the method further comprises isolating said eukaryotic cell from a subject prior to

said modifying. In some embodiments, the method further comprises returning said eukaryotic cell and/or cells derived therefrom to said subject.

[0053] In one aspect, the invention provides a method of modifying expression of a polynucleotide in a eukaryotic cell. In some embodiments, the method comprises allowing a CRISPR complex to bind to the polynucleotide such that said binding results in increased or decreased expression of said polynucleotide; wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within said polynucleotide, wherein said guide sequence is linked to a tracr mate sequence which in turn hybridizes to a tracr sequence. In some embodiments, the method further comprises delivering one or more vectors to said eukaryotic cells, wherein the one or more vectors drive expression of one or more of: the CRISPR enzyme, the guide sequence linked to the tracr mate sequence, and the tracr sequence.

[0054] In one aspect, the invention provides a method of generating a model eukaryotic cell comprising a mutated disease gene. In some embodiments, a disease gene is any gene associated an increase in the risk of having or developing a disease. In some embodiments, the method comprises (a) introducing one or more vectors into a eukaryotic cell, wherein the one or more vectors drive expression of one or more of: a CRISPR enzyme, a guide sequence linked to a tracr mate sequence, and a tracr sequence; and (b) allowing a CRISPR complex to bind to a target polynucleotide to effect cleavage of the target polynucleotide within said disease gene, wherein the CRISPR complex comprises the CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence within the target polynucleotide, and (2) the tracr mate sequence that is hybridized to the tracr sequence, thereby generating a model eukaryotic cell comprising a mutated disease gene. In some embodiments, said cleavage comprises cleaving one or two strands at the location of the target sequence by said CRISPR enzyme. In some embodiments, said cleavage results in decreased transcription of a target gene. In some embodiments, the method further comprises repairing said cleaved target polynucleotide by homologous recombination with an exogenous template polynucleotide, wherein said repair results in a mutation comprising an insertion, deletion, or substitution of one or more nucleotides of said target polynucleotide. In some embodiments, said mutation results in one or more amino acid changes in a protein expression from a gene comprising the target sequence.

[0055] In one aspect, the invention provides a method for developing a biologically active agent that modulates a cell signaling event associated with a disease gene. In some embodiments, a disease gene is any gene associated an increase in the risk of having or developing a disease. In some embodiments, the method comprises (a) contacting a test compound with a model cell of any one of the described embodiments; and (b) detecting a change in a readout that is indicative of a reduction or an augmentation of a cell signaling event associated with said mutation in said disease gene, thereby developing said biologically active agent that modulates said cell signaling event associated with said disease gene.

[0056] In one aspect, the invention provides a recombinant polynucleotide comprising a guide sequence upstream of a tracr mate sequence, wherein the guide sequence when expressed directs sequence-specific binding of a CRISPR complex to a corresponding target sequence present in a eukaryotic cell. In some embodiments, the target sequence is a viral sequence present in a eukaryotic cell. In some embodiments, the target sequence is a proto-oncogene or an oncogene.

[0057] In one aspect the invention provides for a method of selecting one or more cell(s) by introducing one or more mutations in a gene in the one or more cell (s), the method comprising: introducing one or more vectors into the cell (s), wherein the one or more vectors drive expression of one or more of: a CRISPR enzyme, a guide sequence linked to a tracr mate sequence, a tracr sequence, and an editing template; wherein the editing template comprises the one or more mutations that abolish CRISPR enzyme cleavage; allowing homologous recombination of the editing template with the target polynucleotide in the cell(s) to be selected; allowing a CRISPR complex to bind to a target polynucleotide to effect cleavage of the target polynucleotide within said gene, wherein the CRISPR complex comprises the CRISPR enzyme complexed with (1) the guide sequence that is hybridized to the target sequence within the target polynucleotide, and (2) the tracr mate sequence that is hybridized to the tracr sequence, wherein binding of the CRISPR complex to the target polynucleotide induces cell death, thereby allowing one or more cell(s) in which one or more mutations have been introduced to be selected. In a preferred embodiment, the CRISPR enzyme is Cas9. In another preferred embodiment of the invention the cell to be selected may be a eukaryotic cell. Aspects of the invention allow for selection of specific cells without requiring a selection marker or a two-step process that may include a counter-selection system.

[0058] With respect to mutations of the CRISPR enzyme, when the enzyme is not SpCas9, mutations may be made at any or all residues corresponding to positions 10, 762, 840, 854, 863 and/or 986 of SpCas9 (which may be ascertained for instance by standard sequence comparison tools). In particular, any or all of the following mutations are preferred in SpCas9: D10A, E762A, H840A, N854A, N863A and/or D986A; as well as conservative substitution for any of the replacement amino acids is also envisaged. In an aspect the invention provides as to any or each or all embodiments herein-discussed wherein the CRISPR enzyme comprises at least one or more, or at least two or more mutations, wherein the at least one or more mutation or the at least two or more mutations is as to D10, E762, H840, N854, N863, or D986 according to SpCas9 protein, e.g., D10A, E762A, H840A, N854A, N863A and/or D986A as to SpCas9, or N580 according to SaCas9, e.g., N580A as to SaCas9, or any corresponding mutation(s) in a Cas9 of an ortholog to Sp or Sa, or the CRISPR enzyme comprises at least one mutation wherein at least H840 or N863A as to Sp Cas9 or N580A as to Sa Cas9 is mutated; e.g., wherein the CRISPR enzyme comprises H840A, or D10A and H840A, or D10A and N863A, according to SpCas9 protein, or any corresponding mutation(s) in a Cas9 of an ortholog to Sp protein or Sa protein.

[0059] In a further aspect, the invention involves a computer-assisted method for identifying or designing potential compounds to fit within or bind to CRISPR-Cas9 system or a functional portion thereof or vice versa (a computer-assisted method for identifying or designing potential CRISPR-Cas9 systems or a functional portion thereof for binding to desired compounds) or a computer-assisted method for identifying or designing potential CRISPR-Cas9 systems (e.g., with regard to predicting areas of the CRISPR-Cas9 system to be able to be manipulated—for instance, based on crystal structure data or based on data of Cas9 orthologs, or with respect to where a functional group such as an activator or repressor can be attached to the CRISPR-Cas9 system, or as to Cas9 truncations or as to designing nickases), said method comprising:

using a computer system, e.g., a programmed computer comprising a processor, a data storage system, an input device, and an output device, the steps of:

(a) inputting into the programmed computer through said input device data comprising the three-dimensional co-ordinates of a subset of the atoms from or pertaining to the CRISPR-Cas9 crystal structure, e.g., in the CRISPR-Cas9 system binding domain or alternatively or additionally in domains that vary based on variance among Cas9 orthologs or as to Cas9s or as to

nickases or as to functional groups, optionally with structural information from CRISPR-Cas9 system complex(es), thereby generating a data set;

(b) comparing, using said processor, said data set to a computer database of structures stored in said computer data storage system, e.g., structures of compounds that bind or putatively bind or that are desired to bind to a CRISPR-Cas9 system or as to Cas9 orthologs (e.g., as Cas9s or as to domains or regions that vary amongst Cas9 orthologs) or as to the CRISPR-Cas9 crystal structure or as to nickases or as to functional groups;

(c) selecting from said database, using computer methods, structure(s)—e.g., CRISPR-Cas9 structures that may bind to desired structures, desired structures that may bind to certain CRISPR-Cas9 structures, portions of the CRISPR-Cas9 system that may be manipulated, e.g., based on data from other portions of the CRISPR-Cas9 crystal structure and/or from Cas9 orthologs, truncated Cas9s, novel nickases or particular functional groups, or positions for attaching functional groups or functional-group-CRISPR-Cas9 systems;

(d) constructing, using computer methods, a model of the selected structure(s); and

(e) outputting to said output device the selected structure(s);

and optionally synthesizing one or more of the selected structure(s);

and further optionally testing said synthesized selected structure(s) as or in a CRISPR-Cas9 system;

or, said method comprising: providing the co-ordinates of at least two atoms of the CRISPR-Cas9 crystal structure, e.g., at least two atoms of the herein Crystal Structure Table of the CRISPR-Cas9 crystal structure or co-ordinates of at least a sub-domain of the CRISPR-Cas9 crystal structure (“selected co-ordinates”), providing the structure of a candidate comprising a binding molecule or of portions of the CRISPR-Cas9 system that may be manipulated, e.g., based on data from other portions of the CRISPR-Cas9 crystal structure and/or from Cas9 orthologs, or the structure of functional groups, and fitting the structure of the candidate to the selected co-ordinates, to thereby obtain product data comprising CRISPR-Cas9 structures that may bind to desired structures, desired structures that may bind to certain CRISPR-Cas9 structures, portions of the CRISPR-Cas9 system that may be manipulated, truncated Cas9s, novel nickases, or particular functional groups, or positions for attaching functional groups or functional-group-CRISPR-Cas9 systems, with output thereof; and optionally synthesizing

compound(s) from said product data and further optionally comprising testing said synthesized compound(s) as or in a CRISPR-Cas9 system.

[0060] The testing can comprise analyzing the CRISPR-Cas9 system resulting from said synthesized selected structure(s), e.g., with respect to binding, or performing a desired function.

[0061] The output in the foregoing methods can comprise data transmission, e.g., transmission of information via telecommunication, telephone, video conference, mass communication, e.g., presentation such as a computer presentation (e.g. POWERPOINT), internet, email, documentary communication such as a computer program (e.g. WORD) document and the like. Accordingly, the invention also comprehends computer readable media containing: atomic co-ordinate data according to the herein-referenced Crystal Structure, said data defining the three dimensional structure of CRISPR-Cas9 or at least one sub-domain thereof, or structure factor data for CRISPR-Cas9, said structure factor data being derivable from the atomic co-ordinate data of herein-referenced Crystal Structure. The computer readable media can also contain any data of the foregoing methods. The invention further comprehends methods a computer system for generating or performing rational design as in the foregoing methods containing either: atomic co-ordinate data according to herein-referenced Crystal Structure, said data defining the three dimensional structure of CRISPR-Cas9 or at least one sub-domain thereof, or structure factor data for CRISPR-Cas9, said structure factor data being derivable from the atomic co-ordinate data of herein-referenced Crystal Structure. The invention further comprehends a method of doing business comprising providing to a user the computer system or the media or the three dimensional structure of CRISPR-Cas9 or at least one sub-domain thereof, or structure factor data for CRISPR-Cas9, said structure set forth in and said structure factor data being derivable from the atomic co-ordinate data of herein-referenced Crystal Structure, or the herein computer media or a herein data transmission.

[0062] A “binding site” or an “active site” comprises or consists essentially of or consists of a site (such as an atom, a functional group of an amino acid residue or a plurality of such atoms and/or groups) in a binding cavity or region, which may bind to a compound such as a nucleic acid molecule, which is/are involved in binding.

[0063] By “fitting”, is meant determining by automatic, or semi-automatic means, interactions between one or more atoms of a candidate molecule and at least one atom of a structure of the invention, and calculating the extent to which such interactions are stable.

Interactions include attraction and repulsion, brought about by charge, steric considerations and the like. Various computer-based methods for fitting are described further

[0064] By “root mean square (or rms) deviation,” Applicants mean the square root of the arithmetic mean of the squares of the deviations from the mean.

[0065] By a “computer system”, is meant the hardware means, software means and data storage means used to analyze atomic coordinate data. The minimum hardware means of the computer-based systems of the present invention typically comprises a central processing unit (CPU), input means, output means and data storage means. Desirably a display or monitor is provided to visualize structure data. The data storage means may be RAM or means for accessing computer readable media of the invention. Examples of such systems are computer and tablet devices running Unix, Windows or Apple operating systems.

[0066] By “computer readable media”, is meant any medium or media, which can be read and accessed directly or indirectly by a computer e.g. so that the media is suitable for use in the above-mentioned computer system. Such media include, but are not limited to: magnetic storage media such as floppy discs, hard disc storage medium and magnetic tape; optical storage media such as optical discs or CD-ROM; electrical storage media such as RAM and ROM; thumb drive devices; cloud storage devices and hybrids of these categories such as magnetic/optical storage media.

[0067] In particular embodiments of the invention, the conformational variations in the crystal structures of the CRISPR-Cas9 system or of components of the CRISPR-Cas9 provide important and critical information about the flexibility or movement of protein structure regions relative to nucleotide (RNA or DNA) structure regions that may be important for CRISPR-Cas9 system function. The structural information provided for Cas9 (e.g. *S. pyogenes* Cas9) as the CRISPR enzyme in the present application may be used to further engineer and optimize the CRISPR-Cas9 system and this may be extrapolated to interrogate structure-function relationships in other CRISPR enzyme systems as well, e.g. other Type II CRISPR enzyme systems.

[0068] The invention comprehends optimized functional CRISPR-Cas9 enzyme systems. In particular the CRISPR enzyme comprises one or more mutations that converts it to a DNA binding protein to which functional domains exhibiting a function of interest may be recruited or appended or inserted or attached. In certain embodiments, the CRISPR enzyme comprises one or more mutations which include but are not limited to D10A, E762A, H840A, N854A, N863A or

D986A (based on the amino acid position numbering of a *S. pyogenes* Cas9) and/or the one or more mutations is in a RuvC1 or HNH domain of the CRISPR enzyme or is a mutation as otherwise as discussed herein. In some embodiments, the CRISPR enzyme has one or more mutations in a catalytic domain, wherein when transcribed, the tracr mate sequence hybridizes to the tracr sequence and the guide sequence directs sequence-specific binding of a CRISPR complex to the target sequence, and wherein the enzyme further comprises a functional domain.

[0069] The structural information provided herein allows for interrogation of sgRNA (or chimeric RNA) interaction with the target DNA and the CRISPR enzyme (e.g. Cas9) permitting engineering or alteration of sgRNA structure to optimize functionality of the entire CRISPR-Cas9 system. For example, loops of the sgRNA may be extended, without colliding with the Cas9 protein by the insertion of adaptor proteins that can bind to RNA. These adaptor proteins can further recruit effector proteins or fusions which comprise one or more functional domains.

[0070] In some preferred embodiments, the functional domain is a transcriptional activation domain, preferably VP64. In some embodiments, the functional domain is a transcription repression domain, preferably KRAB. In some embodiments, the transcription repression domain is SID, or concatemers of SID (e.g. SID4X). In some embodiments, the functional domain is an epigenetic modifying domain, such that an epigenetic modifying enzyme is provided. In some embodiments, the functional domain is an activation domain, which may be the P65 activation domain.

[0071] Aspects of the invention encompass a non-naturally occurring or engineered composition that may comprise a guide RNA (sgRNA) comprising a guide sequence capable of hybridizing to a target sequence in a genomic locus of interest in a cell and a CRISPR enzyme that may comprise at least one or more nuclear localization sequences, wherein the CRISPR enzyme comprises two or more mutations, such that the enzyme has altered or diminished nuclease activity compared with the wild type enzyme, wherein at least one loop of the sgRNA is modified by the insertion of distinct RNA sequence(s) that bind to one or more adaptor proteins, and wherein the adaptor protein further recruits one or more heterologous functional domains. In an embodiment of the invention the CRISPR enzyme comprises two or more mutations in a residue selected from D10, E762, H840, N854, N863, or D986. In a further embodiment the CRISPR enzyme comprises two or more mutations selected from the group comprising D10A, E762A, H840A, N854A, N863A or D986A. In another embodiment, the functional domain is a

transcriptional activation domain, e.g. VP64. In another embodiment, the functional domain is a transcriptional repressor domain, e.g. KRAB domain, SID domain or a SID4X domain. In embodiments of the invention, the one or more heterologous functional domains have one or more activities selected from methylase activity, demethylase activity, transcription activation activity, transcription repression activity, transcription release factor activity, histone modification activity, RNA cleavage activity and nucleic acid binding activity. In further embodiments of the invention the cell is a eukaryotic cell or a mammalian cell or a human cell. In further embodiments, the adaptor protein is selected from MS2, PP7, Q β , F2, GA, fr, JP501, M12, R17, BZ13, JP34, JP500, KU1, M11, MX1, TW18, VK, SP, FI, ID2, NL95, TW19, AP205, ϕ Cb5, ϕ Cb8r, ϕ Cb12r, ϕ Cb23r, 7s, and PRR1. In another embodiment, the at least one loop of the sgRNA is tetraloop and/or loop2. An aspect of the invention encompasses methods of modifying a genomic locus of interest to change gene expression in a cell by introducing into the cell any of the compositions described herein.

[0072] An aspect of the invention is that the above elements are comprised in a single composition or comprised in individual compositions. These compositions may advantageously be applied to a host to elicit a functional effect on the genomic level.

[0073] In general, the sgRNA are modified in a manner that provides specific binding sites (e.g. aptamers) for adapter proteins comprising one or more functional domains (e.g. via fusion protein) to bind to. The modified sgRNA are modified such that once the sgRNA forms a CRISPR complex (i.e. CRISPR enzyme binding to sgRNA and target) the adapter proteins bind and, the functional domain on the adapter protein is positioned in a spatial orientation which is advantageous for the attributed function to be effective. For example, if the functional domain is a transcription activator (e.g. VP64 or p65), the transcription activator is placed in a spatial orientation which allows it to affect the transcription of the target. Likewise, a transcription repressor will be advantageously positioned to affect the transcription of the target and a nuclease (e.g. FokI) will be advantageously positioned to cleave or partially cleave the target.

[0074] The skilled person will understand that modifications to the sgRNA which allow for binding of the adapter + functional domain but not proper positioning of the adapter + functional domain (e.g. due to steric hindrance within the three dimensional structure of the CRISPR complex) are modifications which are not intended. The one or more modified sgRNA may be modified at the tetra loop, the stem loop 1, stem loop 2, or stem loop 3, as described herein,

preferably at either the tetra loop or stem loop 2, and most preferably at both the tetra loop and stem loop 2.

[0075] As explained herein the functional domains may be, for example, one or more domains comprising methylase activity, demethylase activity, transcription activation activity, transcription repression activity, transcription release factor activity, histone modification activity, RNA cleavage activity, DNA cleavage activity, nucleic acid binding activity, and molecular switches (e.g. light inducible). In some cases it is advantageous that additionally at least one NLS is provided. In some instances, it is advantageous to position the NLS at the N terminus. When more than one functional domain is included, the functional domains may be the same or different.

[0076] The sgRNA may be designed to include multiple binding recognition sites (e.g. aptamers) specific to the same or different adapter protein. The sgRNA may be designed to bind to the promoter region -1000 - +1 nucleic acids upstream of the transcription start site (i.e. TSS), preferably -200 nucleic acids. This positioning improves functional domains which affect gene activation (e.g. transcription activators) or gene inhibition (e.g. transcription repressors). The modified sgRNA may be one or more modified sgRNAs targeted to one or more target loci (e.g. at least 1 sgRNA, at least 2 sgRNA, at least 5 sgRNA, at least 10 sgRNA, at least 20 sgRNA, at least 30 sg RNA, at least 50 sgRNA) comprised in a composition.

[0077] Further, the CRISPR enzyme with diminished nuclease activity is most effective when the nuclease activity is inactivated (e.g. nuclease inactivation of at least 70%, at least 80%, at least 90%, at least 95%, at least 97%, or 100% as compared with the wild type enzyme; or to put in another way, a Cas9 enzyme or CRISPR enzyme having advantageously about 0% of the nuclease activity of the non-mutated or wild type Cas9 enzyme or CRISPR enzyme, or no more than about 3% or about 5% or about 10% of the nuclease activity of the non-mutated or wild type Cas9 enzyme or CRISPR enzyme). This is possible by introducing mutations into the RuvC and HNH nuclease domains of the SpCas9 and orthologs thereof. For example utilizing mutations in a residue selected from D10, E762, H840, N854, N863, or D986 and more preferably introducing one or more of the mutations selected from D10A, E762A, H840A, N854A, N863A or D986A. A preferable pair of mutations is D10A with H840A, more preferable is D10A with N863A of SpCas9 and orthologs thereof.

[0078] The inactivated CRISPR enzyme may have associated (e.g. via fusion protein) one or more functional domains, like for example as described herein for the modified sgRNA adaptor proteins, including for example, one or more domains from methylase activity, demethylase activity, transcription activation activity, transcription repression activity, transcription release factor activity, histone modification activity, RNA cleavage activity, DNA cleavage activity, nucleic acid binding activity, and molecular switches (e.g. light inducible). Preferred domains are Fok1, VP64, P65, HSF1, and MyoD1. In the event that Fok1 is provided, it is advantageous that multiple Fok1 functional domains are provided to allow for a functional dimer and that sgRNAs are designed to provide proper spacing for functional use (Fok1) as specifically described in Tsai et al. *Nature Biotechnology*, Vol. 32, Number 6, June 2014). The adaptor protein may utilize known linkers to attach such functional domains. In some cases it is advantageous that additionally at least one NLS is provided. In some instances, it is advantageous to position the NLS at the N terminus. When more than one functional domain is included, the functional domains may be the same or different.

[0079] In general, the positioning of the one or more functional domains on the inactivated CRISPR enzyme is one which allows for correct spatial orientation for the functional domain to affect the target with the attributed functional effect. For example, if the functional domain is a transcription activator (e.g. VP64 or p65), the transcription activator is placed in a spatial orientation which allows it to affect the transcription of the target. Likewise, a transcription repressor will be advantageously positioned to affect the transcription of the target, and a nuclease (e.g. Fok1) will be advantageously positioned to cleave or partially cleave the target. This may include positions other than the N- / C- terminus of the CRISPR enzyme.

[0080] Due to crystal structure experiments, the Applicant has identified that positioning the functional domain in the Rec1 domain, the Rec2 domain, the HNH domain, or the PI domain of the SpCas9 protein or any ortholog corresponding to these domains is advantageous. Positioning of the functional domains to the Rec1 domain or the Rec2 domain, of the SpCas9 protein or any ortholog corresponding to these domains, in some instances may be preferred. Positioning of the functional domains to the Rec1 domain at position 553, Rec1 domain at 575, the Rec2 domain at any position of 175-306 or replacement thereof, the HNH domain at any position of 715-901 or replacement thereof, or the PI domain at position 1153 of the SpCas9 protein or any ortholog corresponding to these domains, in some instances may be preferred. Fok1 functional domain

may be attached at the N terminus. When more than one functional domain is included, the functional domains may be the same or different.

[0081] The adaptor protein may be any number of proteins that binds to an aptamer or recognition site introduced into the modified sgRNA and which allows proper positioning of one or more functional domains, once the sgRNA has been incorporated into the CRISPR complex, to affect the target with the attributed function. As explained in detail in this application such may be coat proteins, preferably bacteriophage coat proteins. The functional domains associated with such adaptor proteins (e.g. in the form of fusion protein) may include, for example, one or more domains selected from methylase activity, demethylase activity, transcription activation activity, transcription repression activity, transcription release factor activity, histone modification activity, RNA cleavage activity, DNA cleavage activity, nucleic acid binding activity, and molecular switches (e.g. light inducible). Preferred domains are Fok1, VP64, P65, HSF1, and MyoD1. In the event that the functional domain is a transcription activator or transcription repressor it is advantageous that additionally at least an NLS is provided and preferably at the N terminus. When more than one functional domain is included, the functional domains may be the same or different. The adaptor protein may utilize known linkers to attach such functional domains.

[0082] Thus, the modified sgRNA, the inactivated CRISPR enzyme (with or without functional domains), and the binding protein with one or more functional domains, may each individually be comprised in a composition and administered to a host individually or collectively. Alternatively, these components may be provided in a single composition for administration to a host. Administration to a host may be performed via viral vectors known to the skilled person or described herein for delivery to a host (e.g. lentiviral vector, adenoviral vector, AAV vector). As explained herein, use of different selection markers (e.g. for lentiviral sgRNA selection) and concentration of sgRNA (e.g. dependent on whether multiple sgRNAs are used) may be advantageous for eliciting an improved effect.

[0083] On the basis of this concept, several variations are appropriate to elicit a genomic locus event, including DNA cleavage, gene activation, or gene deactivation. Using the provided compositions, the person skilled in the art can advantageously and specifically target single or multiple loci with the same or different functional domains to elicit one or more genomic locus events. The compositions may be applied in a wide variety of methods for screening in libraries

in cells and functional modeling in vivo (e.g. gene activation of lincRNA and identification of function; gain-of-function modeling; loss-of-function modeling; the use the compositions of the invention to establish cell lines and transgenic animals for optimization and screening purposes).

[0084] The current invention comprehends the use of the compositions of the current invention to establish and utilize conditional or inducible CRISPR transgenic cell /animals. (*See, e.g.,* Platt et al., 2014, Cell 159(2):440-55, <http://dx.doi.org/10.1016/j.cell.2014.09.014>, or PCT patent publications cited herein, such as WO 2014/093622 (PCT/US2013/074667), which are not believed prior to the present invention or application). For example, the target cell comprises CRISPR enzyme (e.g. Cas9) conditionally or inducibly (e.g. in the form of Cre dependent constructs) and/or the adapter protein conditionally or inducibly and, on expression of a vector introduced into the target cell, the vector expresses that which induces or gives rise to the condition of CRISPR enzyme (e.g. Cas9) expression and/or adaptor expression in the target cell. By applying the teaching and compositions of the current invention with the known method of creating a CRISPR complex, inducible genomic events affected by functional domains are also an aspect of the current invention. One mere example of this is the creation of a CRISPR knock-in / conditional transgenic animal (e.g. mouse comprising e.g. a Lox-Stop-polyA-Lox(LSL) cassette) and subsequent delivery of one or more compositions providing one or more modified sgRNA (e.g. -200 nucleotides to TSS of a target gene of interest for gene activation purposes) as described herein (e.g. modified sgRNA with one or more aptamers recognized by coat proteins, e.g. MS2), one or more adapter proteins as described herein (MS2 binding protein linked to one or more VP64) and means for inducing the conditional animal (e.g. Cre recombinase for rendering Cas9 expression inducible). Alternatively, the adaptor protein may be provided as a conditional or inducible element with a conditional or inducible CRISPR enzyme to provide an effective model for screening purposes, which advantageously only requires minimal design and administration of specific sgRNAs for a broad number of applications.

[0085] In one aspect Sa Cas9 is utilized in a single construct used to target genes for editing. The construction of a single Sa based vector, simultaneously containing an Sa Cas9 nuclease, a deadGuide, and an active guide may be incorporated into a viral vector. Sa Cas9 is smaller than sp Cas9 and will allow viral vectors with limited insertion sizes to be utilized. This vector can be used to simultaneously up and downregulate different genes using a single viral construct. The

vector can be used in the treatment of a patient in need thereof or to study the interaction of genes in a eukaryotic system.

[0086] In another aspect *in vivo* activation screens can be used in a mouse constitutively expressing nuclease active Cas9. Nuclease deficient Cas9 is not required based on the current invention. An *in vivo* orthogonal screen using a mouse constitutively expressing Cas9 may be performed. The current invention may be used, for example, to upregulate MYC in all cells, and then knockdown pairs of genes to see which genetic knockdown inhibits tumor growth and metastasis *in vivo*. In another example, p53 is deleted, and simultaneously different genes are upregulated to determine genes that can rescue this effect.

[0087] In another aspect the dead guides are further modified to improve specificity. Protected dead guides may be synthesized, whereby secondary structure is introduced into the 5' end of the dead guide to improve its specificity. A protected guide RNA (pgRNA) comprises a guide sequence capable of hybridizing to a target sequence in a genomic locus of interest in a cell and a protector strand, wherein the protector strand is optionally complementary to the guide sequence and wherein the guide sequence may in part be hybridizable to the protector strand. The pgRNA optionally includes an extension sequence. The thermodynamics of the pgRNA-target DNA hybridization is determined by the number of bases complementary between the guide RNA and target DNA. By employing 'thermodynamic protection', specificity of sgRNA can be improved by adding a protector sequence. For example, one method adds a complementary protector strand of varying lengths to the 5' end of the guide sequence within the sgRNA. As a result, the protector strand is bound to at least a portion of the sgRNA and provides for a protected sgRNA (pgRNA). In turn, the sgRNA references herein may be easily protected using the described embodiments, resulting in pgRNA. The protector strand can be either a separate RNA transcript or strand or a chimeric version joined to the 5' end of the sgRNA guide sequence.

[0088] Accordingly, it is an object of the invention not to encompass within the invention any previously known product, process of making the product, or method of using the product such that Applicants reserve the right and hereby disclose a disclaimer of any previously known product, process, or method. It is further noted that the invention does not intend to encompass within the scope of the invention any product, process, or making of the product or method of using the product, which does not meet the written description and enablement requirements of

the USPTO (35 U.S.C. §112, first paragraph) or the EPO (Article 83 of the EPC), such that Applicants reserve the right and hereby disclose a disclaimer of any previously described product, process of making the product, or method of using the product. Nothing herein is to be construed as a promise.

[0089] It is noted that in this disclosure and particularly in the claims and/or paragraphs, terms such as "comprises", "comprised", "comprising" and the like can have the meaning attributed to it in U.S. Patent law; e.g., they can mean "includes", "included", "including", and the like; and that terms such as "consisting essentially of" and "consists essentially of" have the meaning ascribed to them in U.S. Patent law, e.g., they allow for elements not explicitly recited, but exclude elements that are found in the prior art or that affect a basic or novel characteristic of the invention. These and other embodiments are disclosed or are obvious from and encompassed by, the following Detailed Description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0090] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[0091] **Figure 1** depicts an experimental setup wherein in a 96 well plate, HEK.293 cells were transfected with 100 ng Cas9, 100 ng sgRNA, and 100 ng MS2-p65-HSF1. 48 hours later, cells were removed, and taken for either indel analysis (surveyor) or qPCR to analyze gene activation. Applicants attempted to activate the gene IL1B. The sequences had a targeting sequence 20, 15, 14, 13, 12, or 11 bp long. The samples in the top table were treated with the sgRNA shown to the left, an active Cas9, and MS2-p65-HSF1. As a positive control for cutting, Applicants also tested this same sgRNA, which was previously shown to activate (Konermann et al., "Genome-scale transcription activation by an engineered CRISPR-Cas9 complex," doi:10.1038/nature14136, incorporated herein by reference) + dCas9 + MS2-p65-HSF1.

[0092] **Figure 2** illustrates that dead Guides activate, but do not cut target DNA. The bars are quantify the IL1B activation. GAPDH is a standard 'housekeeper' gene, which is used to normalize the data.

[0093] **Figure 3A-3D** is a phylogenetic tree of Cas genes.

[0094] **Figure 4A-4F** shows the phylogenetic analysis revealing five families of Cas9s, including three groups of large Cas9s (~1400 amino acids) and two of small Cas9s (~1100 amino acids).

[0095] **Figure 5A-5D** illustrates the use of sgRNA scaffolds to establish activation and repression in an orthogonal manner utilizing a single nuclease active Cas9 enzyme. Panel A shows the transfected sgRNA scaffolds used. Panel B shows cutting of the EMX1.3 gene by active Cas9. Panel C shows that recruitment of active Cas9 to the IL1B gene using the sgRNA scaffolds does not result in cutting. Panel D shows that recruitment of Cas9 to the IL1B gene using the sgRNA scaffolds results in activation of gene expression.

[0096] **Figure 6A-6D:** The schematic of Figure 6A illustrates aspects of bimodal gene control systems that make combined use of dead guide RNAs having modifications that facilitate recruitment of transcriptional activators (such as HSF1/P65), left side, in combination with 'live' sgRNAs, with the alternative dead and live sgRNAs working with the same Cas9 to mediate opposite bimodal gene control. The plots of Figure 6B show the activity achieved by guides of different lengths at the illustrated target sites upstream of HBG1, with robust activation shown for guides less than 16bp in length. The results for truncated guides illustrated in Figure 6B are illustrated independently, apart from the results for mismatched guides, in the bar graphs of Figure 6C. As illustrated in the graphs in the first row of Figure 6B, the length of the RNA targeting sequence was varied from 11nt to 20nt. HBG1 mRNA levels (normalized to GAPDH, and compared to cells transfected with GFP plasmid) were quantified along with HBG1 indel frequency. No indel formation was observed when sgRNAs had less than 16bp of homology to target DNA. In all cases, guides were designed with MS2 binding loops in the tetraloops and stemloop two, and were co-transfected with active Cas9 and the MPH transcriptional activation complex. The graphs of Figure 6D illustrate the results of using 14 and 15bp dead sgRNA constructs having MS2 loops, to target three different genes (IL1B, HBG1, and ZFP42). The data demonstrate that the activation effect using a dead sgRNA is reproducible at these different loci. To produce the data illustrated in Figure 6D, three dRNAs targeting the promoter regions of IL1B, HBG1, and ZFP42 were tested for activation and indel formation. dRNAs with 14bp or 15bp of homology to target DNA did not induce detectable indel formation. dRNAs co-transfected with Cas9 and MPH activated transcription to a similar extent as 20nt sgRNA-MS2

co-transfected with dCas9 and MPH. (In all cases, mean +/- S.E.M. is plotted. N=2-3 replicates / group).

[0097] **Figure 7A-7C:** The plots of Figure 7 illustrate embodiments in which dRNAs can specifically upregulate gene expression, and have a specificity profile similar to 20bp sgRNA activators. Sequences targeted to the HBG1/2 promoter were tested for off-target transcriptional activation using RNAseq. 20nt sgRNAs with MS2 binding loops were co-transfected with dCas9 and the MPH activation complex. These were compared to dRNAs co-delivered with active Cas9 and the MPH activation complex. Both systems showed similar offtarget profiles. (a) Zero significantly upregulated genes apart from HBG1/2 were observed for both the 20nt/dCas9 and dRNA/Cas9 treated cells. (b) A second guide showed 55 significantly upregulated genes apart from HBG1/2 for the 20nt/dCas9-treated cells, while 31 significantly upregulated genes were measured for dRNA-treated cells. (In all cases, N=3 replicates / group).. The plot of Figure 7c illustrates the results of differential gene expression analysis, and shows that the off target genes have minimal gene expression differences when compared to the on target HBG1/2.

[0098] **Figure 8A-8B:** These graphs illustrate results showing bimodal gene control that confers resistance to BRAF-mutant A375 cells. The bar graphs of Figure 8A illustrate relative upregulation of expression for 5 target genes: CUL3, MED12, LPAR5, ITGA9, and EGFR. Figure 8B is a set of line graphs showing that the bimodal gene perturbations can also cause phenotypic effects, in this case an increase in resistance conferred to A375 cells under PLX4720 BRAF inhibition. As illustrated, each perturbation individually increased the resistance of these cells to PLX4720 and the combinations shifted resistance even more, with some combinations exhibiting synergistic behaviour (e.g. MED12 and LPAR4, which exhibit a perturbation index (P.I.) > 1, indicating synergistic behaviour

[0099] **Figure 9A-9E:** Illustrates data evidencing orthogonal gene control using a single Cas9 nuclease. (a) Orthogonal gene control in melanoma A375 cells expressing an active Cas9 and the MS2-P65-HSF1 fusion protein. Cells were transduced with lentivirus containing a dRNA targeting one gene and an sgRNA targeting a second gene. Selected cells were subsequently treated with BRAFinhibitor PLX4720 and their survival was quantified. (b) Activation and indel % were measured for individually and orthogonally controlled genes. Left: LPAR5 transcriptional upregulation mediated by dRNA was robust in the presence and absence of sgRNAs targeting MED12 or TADA2B. Right: LPAR5 indel formation was undetectable at the

dRNA target site. (c) Robust indel formation was detected at DNA sites targeted by MED12 and TADA2B sgRNAs alone and when delivered together with a dRNA targeting LPAR5. (d) Survival curves for A375 cells expressing active Cas9 and MPH with different combinations of sgRNAs targeting TADA2B and MED12 for knockout and dRNAs targeting LPAR5 for transcriptional activation. (e) PLX-4720 doses resulting in 50% cell death (IC50 values) for different treatment conditions shown in (d). LPAR5/MED12 and LPAR5/TADA2B combination treatments significantly increased resistance relative to cells treated with LPAR5, MED12, or TADA2B alone. In all cases, average +/- SEM is plotted, N=3-4 replicates/ group. *p<0.05.

[00100] **Figure 10:** is a bar graph illustrating results that show the effect of different length sgRNAs when combined with Cas9 mutants, showing that Cas9 mutations that affect nuclease activity can also affect interactions with sgRNAs, to give rise to embodiments have dead guide RNAs of different lengths.

[00101] **Figure 11:** provides a schematic summary, with it understood that Applicant(s) / inventor(s) are not necessarily bound by any particular theory set forth herein or in any particular Figure, including Figure 11. The Figure discusses mutation of positively charged residues binding to the non-targeted gDNA strand whereby specificity is improved. Data in the Table of the schematic summary is as follows and is as to mutations of SpCas9:

Cas9 mutant	Indel %		
	ON Target (EMX1)	OFF Target 1(OT25)	OFF Target 2(OT46)
WT	24.8	10.5	8.8
R780	22.9	0.0	0.1
K810	23.3	0.1	0.1
K848	24.3	0.1	0.1
K855	25.1	0.2	0.3
R976	15.6	0.1	0.1
H982	20.9	0.5	0.4
K1003	24.6	4.1	2.8
R1060	20.4	1.3	1.8
GFP	0.1	0.0	0.1
untrans.	0.1	0.0	0.1

With reference to the numbering of SpCas9, the Figure illustrates alanine mutations that improve specificity, distributed along the non-targeting strand groove, e.g., Arg780, Lys810, Lys855, Lys848, Lys1003, Arg1060, Arg976, His982. Without wishing to be bound by any one particular theory, the mechanism proposal is that nuclease activity is inactive until the non-targeted DNA strand sterically triggers HNH conformation change; non-targeted strand binding to the groove

between HNH and RuvC depends on RNA:DNA pairing; mutating DNA binding residues in the groove places more energetic demand on proper RNA:DNA pairing. Using the information herein, including in Figure 11, the skilled person can readily prepare mutants of other Cas9s (e.g., other than SpCas9) that exhibit improved or reduced off-target effects. For instance, the documents cited herein provide information on numerous orthologs to SpCas9 and SaCas9 exemplified herein. From that information, including the sequence information of those other Cas9s, one skilled in the art can, from the information in this disclosure, readily prepare analogous mutants having reduced off-target effects in Cas9 orthologs in addition to SpCas9 and SaCas9 exemplified herein. Further, documents herein provide crystal structure information as to Cas9, e.g., SpCas9; and one can readily make structural comparisons between crystal structures, e.g., between the crystal structure of SpCas9 and the crystal structure of an ortholog thereto, to also readily, without undue experimentation, obtain analogous mutants having reduced off-target effects in Cas9 orthologs in addition to SpCas9. Accordingly, the invention is broadly applicable to modification(s) or mutation(s) in various Cas9 orthologs to reduce off-target effects, including but not limited to SpCas9 and SaCas9. As discussed further herein, additional or further modification of the above-described Cas9 enzymes can readily be achieved whereby the enzyme in the CRISPR complex has increased capability of modifying the one or more target loci as compared to an unmodified enzyme.

[00102] **Figure 12A-12F:** shows structural aspects of SpCas9 and improved specificity. Panel A is a model of target unwinding. The nt-groove between the RuvC (teal) and HNH (magenta) domains stabilize DNA unwinding through non-specific DNA interactions with the non-complementary strand. RNA:cDNA and Cas9:ncDNA interactions drive DNA unwinding (top arrow) in competition against cDNA:ncDNA rehybridization (bottom arrow). Panel B: The structure of SpCas9 (PDB ID 4UN3) showing the nt-groove situated between the HNH (magenta) and RuvC (teal) domains. The non-target DNA strand (red) was manually modeled into the nt-groove (inset). Panel C: Screen of alanine point mutants for improvement in specificity. Panel D: Assessment of top point mutants at additional off-target loci. The top five specificity conferring mutants are highlighted in red. Panel E: Combination mutants improve specificity compared to single point mutants. eSpCas9(1.0) and eSpCas9(1.1) are highlighted in red. Panel F: Screen of top point mutants and combination mutants at 10 target loci for on-target cleavage efficiency. SpCas9(K855A), eSpCas9(1.0), and eSpCas9(1.1) are highlighted in red.

[00103] **Figure 13A-13C:** shows maintenance of on-target efficiency by spCas9 mutants. Panel A shows an assessment of efficiency of on-target cutting of SpCas9 mutants as compared to SpCas9 for 24 sgRNAs targeted to 9 genomic loci. Panel B is a Tukey plot of normalized on-target indel formation for mutants SpCas9(K855A), eSpCas9(1.0) and eSpCas9(1.1). Panel C is a Western blot of SpCas9 and mutants using anti-SpCas9 antibody.

[00104] **Figure 14A-14C:** shows sensitivity of spCas9 and mutants K855A, eSpCas9(1.0), and eSpCas9(1.1) to single and double base mismatches between the guide RNA and target DNA. Panel A depicts mismatched guide sequences against a VEGFA target. Panel B provides heat maps for spCas9 and three mutants showing indel % with guide sequences having a single base mismatch. Panel C shows indel formation with guide sequences containing consecutive transversion mismatches. Compared to wild type: eSpCas9(1.0) comprises K810A, K1003A, R1060A; eSpCas9(1.1) comprises K848A, K1003A, R1060A.

[00105] **Figure 15:** is a series of bar graphs illustrating the results from using 20bp dRNAs with mismatches at the 5' end to activate transcription. Four dRNAs were designed to target the HBG1 promoter region with a series of 5' end mismatches (red). Target indel formation occurred consistently when sixteen or more nucleotides were matched to target DNA. Gene activation was observed with as few as 11 bp of homology to the target DNA. Average +/- SEM is plotted, N=2-3 replicates/ group.

[00106] **Figure 16A-16C:** illustrates a new activator off-target (OT) score and target DNA GC content are significantly correlated with activator specificity. (a) Transcriptome-wide mRNA profiles for ten different sgRNAs targeting HBG1/2, ranked by GC content and activator off-target score. (b) 20nt sgRNAs with MS2 binding loops were co-transfected with dCas9 and the MPH complex. Low GC content and a high activator OT score of the guide sequence are significantly correlated with the number of statistically significant off-targets. A previously published nuclease OT score did not significantly correlate with guide specificity. (c) Model parameters for a multivariate linear regression derived from data on twelve sequences targeting HBG1/2 (ten from Fig. 16, and two 20nt sequences from Fig. 15). (In all cases, N=3 replicates / group).

[00107] **Figure 17A-17B:** shows dRNAs activate target gene expression with active Cas9. Cells were transduced with lentivirus containing a dRNA. (a) Indel formation was measured at 0.6% and 0.05% for DNA sites targeted by ITGA9 and EGFR dRNAs, respectively. (b) ITGA9

and EGFR mRNA levels (normalized to GAPDH) were quantified. (average + / - SEM; N=3 replicates / group.)

[00108] The figures herein are for illustrative purposes only and are not necessarily drawn to scale.

DETAILED DESCRIPTION OF THE INVENTION

[00109] In general, the CRISPR-Cas, CRISPR-Cas9 or CRISPR system is as used in the foregoing documents, such as WO 2014/093622 (PCT/US2013/074667) and refers collectively to transcripts and other elements involved in the expression of or directing the activity of CRISPR-associated (“Cas”) genes, including sequences encoding a Cas9 gene, in particular a Cas9 gene in the case of CRISPR-Cas9, a tracr (trans-activating CRISPR) sequence (e.g. tracrRNA or an active partial tracrRNA), a tracr-mate sequence (encompassing a “direct repeat” and a tracrRNA-processed partial direct repeat in the context of an endogenous CRISPR system), a guide sequence (also referred to as a “spacer” in the context of an endogenous CRISPR system), or “RNA(s)” as that term is herein used (e.g., RNA(s) to guide Cas9, e.g. CRISPR RNA and transactivating (tracr) RNA or a single guide RNA (sgRNA) (chimeric RNA)) or other sequences and transcripts from a CRISPR locus. In general, a CRISPR system is characterized by elements that promote the formation of a CRISPR complex at the site of a target sequence (also referred to as a protospacer in the context of an endogenous CRISPR system). In the context of formation of a CRISPR complex, “target sequence” refers to a sequence to which a guide sequence is designed to have complementarity, where hybridization between a target sequence and a guide sequence promotes the formation of a CRISPR complex. A target sequence may comprise any polynucleotide, such as DNA or RNA polynucleotides. In some embodiments, a target sequence is located in the nucleus or cytoplasm of a cell, and may include nucleic acids in or from mitochondrial, organelles, vesicles, liposomes or particles present within the cell. In some embodiments, especially for non-nuclear uses, NLSs are not preferred. In some embodiments, direct repeats may be identified *in silico* by searching for repetitive motifs that fulfill any or all of the following criteria: 1. found in a 2Kb window of genomic sequence flanking the type II CRISPR locus; 2. span from 20 to 50 bp; and 3. interspaced by 20 to 50 bp. In some embodiments, 2 of these criteria may be used, for instance 1 and 2, 2 and 3, or 1 and 3. In some embodiments, all 3 criteria may be used.

[00110] In embodiments of the invention the terms guide sequence and guide RNA are used interchangeably as in foregoing cited documents such as WO 2014/093622 (PCT/US2013/074667). In general, a guide sequence is any polynucleotide sequence having sufficient complementarity with a target polynucleotide sequence to hybridize with the target sequence and direct sequence-specific binding of a CRISPR complex to the target sequence. In some embodiments, the degree of complementarity between a guide sequence and its corresponding target sequence, when optimally aligned using a suitable alignment algorithm, is about or more than about 50%, 60%, 75%, 80%, 85%, 90%, 95%, 97.5%, 99%, or more. Optimal alignment may be determined with the use of any suitable algorithm for aligning sequences, non-limiting example of which include the Smith-Waterman algorithm, the Needleman-Wunsch algorithm, algorithms based on the Burrows-Wheeler Transform (e.g. the Burrows Wheeler Aligner), ClustalW, Clustal X, BLAT, Novoalign (Novocraft Technologies; available at www.novocraft.com), ELAND (Illumina, San Diego, CA), SOAP (available at soap.genomics.org.cn), and Maq (available at maq.sourceforge.net). In some embodiments, a guide sequence is about or more than about 5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 75, or more nucleotides in length. In some embodiments, a guide sequence is less than about 75, 50, 45, 40, 35, 30, 25, 20, 15, 12, or fewer nucleotides in length. Preferably the guide sequence is 10 - 30 nucleotides long. The ability of a guide sequence to direct sequence-specific binding of a CRISPR complex to a target sequence may be assessed by any suitable assay. For example, the components of a CRISPR system sufficient to form a CRISPR complex, including the guide sequence to be tested, may be provided to a host cell having the corresponding target sequence, such as by transfection with vectors encoding the components of the CRISPR sequence, followed by an assessment of preferential cleavage within the target sequence, such as by Surveyor assay as described herein. Similarly, cleavage of a target polynucleotide sequence may be evaluated in a test tube by providing the target sequence, components of a CRISPR complex, including the guide sequence to be tested and a control guide sequence different from the test guide sequence, and comparing binding or rate of cleavage at the target sequence between the test and control guide sequence reactions. Other assays are possible, and will occur to those skilled in the art.

[00111] In a classic CRISPR-Cas system, the degree of complementarity between a guide sequence and its corresponding target sequence can be about or more than about 50%, 60%,

75%, 80%, 85%, 90%, 95%, 97.5%, 99%, or 100%; a guide or RNA or sgRNA can be about or more than about 5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 75, or more nucleotides in length; or guide or RNA or sgRNA can be less than about 75, 50, 45, 40, 35, 30, 25, 20, 15, 12, or fewer nucleotides in length; and advantageously tracr RNA is 30 or 50 nucleotides in length. However, an aspect of the invention is to reduce off-target interactions, e.g., reduce the guide interacting with a target sequence having low complementarity. Indeed, in the examples, it is shown that the invention involves mutations that result in the CRISPR-Cas9 system being able to distinguish between target and off-target sequences that have greater than 80% to about 95% complementarity, e.g., 83%-84% or 88-89% or 94-95% complementarity (for instance, distinguishing between a target having 18 nucleotides from an off-target of 18 nucleotides having 1, 2 or 3 mismatches). Accordingly, in the context of the present invention the degree of complementarity between a guide sequence and its corresponding target sequence is greater than 94.5% or 95% or 95.5% or 96% or 96.5% or 97% or 97.5% or 98% or 98.5% or 99% or 99.5% or 99.9%, or 100%. Off target is less than 100% or 99.9% or 99.5% or 99% or 99% or 98.5% or 98% or 97.5% or 97% or 96.5% or 96% or 95.5% or 95% or 94.5% or 94% or 93% or 92% or 91% or 90% or 89% or 88% or 87% or 86% or 85% or 84% or 83% or 82% or 81% or 80% complementarity between the sequence and the guide, with it advantageous that off target is 100% or 99.9% or 99.5% or 99% or 99% or 98.5% or 98% or 97.5% or 97% or 96.5% or 96% or 95.5% or 95% or 94.5% complementarity between the sequence and the guide.

[00112] In particularly preferred embodiments according to the invention, the guide RNA (capable of guiding Cas9 to a target locus) may comprise (1) a guide sequence capable of hybridizing to a genomic target locus in the eukaryotic cell; (2) a tracr sequence; and (3) a tracr mate sequence. All (1) to (3) may reside in a single RNA, i.e. an sgRNA (arranged in a 5' to 3' orientation), or the tracr RNA may be a different RNA than the RNA containing the guide and tracr sequence. The tracr hybridizes to the tracr mate sequence and directs the CRISPR/Cas9 complex to the target sequence.

[00113] The methods according to the invention as described herein comprehend inducing one or more mutations in a eukaryotic cell (in vitro, i.e. in an isolated eukaryotic cell) as herein discussed comprising delivering to cell a vector as herein discussed. The mutation(s) can include the introduction, deletion, or substitution of one or more nucleotides at each target sequence of

cell(s) via the guide(s) RNA(s) or sgRNA(s). The mutations can include the introduction, deletion, or substitution of 1-75 nucleotides at each target sequence of said cell(s) via the guide(s) RNA(s) or sgRNA(s). The mutations can include the introduction, deletion, or substitution of 1, 5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, or 75 nucleotides at each target sequence of said cell(s) via the guide(s) RNA(s) or sgRNA(s). The mutations can include the introduction, deletion, or substitution of 5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, or 75 nucleotides at each target sequence of said cell(s) via the guide(s) RNA(s) or sgRNA(s). The mutations include the introduction, deletion, or substitution of 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, or 75 nucleotides at each target sequence of said cell(s) via the guide(s) RNA(s) or sgRNA(s). The mutations can include the introduction, deletion, or substitution of 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, or 75 nucleotides at each target sequence of said cell(s) via the guide(s) RNA(s) or sgRNA(s). The mutations can include the introduction, deletion, or substitution of 40, 45, 50, 75, 100, 200, 300, 400 or 500 nucleotides at each target sequence of said cell(s) via the guide(s) RNA(s) or sgRNA(s).

[00114] For minimization of toxicity and off-target effect, it will be important to control the concentration of Cas9 mRNA and guide RNA delivered. Optimal concentrations of Cas9 mRNA and guide RNA can be determined by testing different concentrations in a cellular or non-human eukaryote animal model and using deep sequencing to analyze the extent of modification at potential off-target genomic loci. Alternatively, to minimize the level of toxicity and off-target effect, Cas9 nickase mRNA (for example *S. pyogenes* Cas9 with the D10A mutation) can be delivered with a pair of guide RNAs targeting a site of interest. Guide sequences and strategies to minimize toxicity and off-target effects can be as in WO 2014/093622 (PCT/US2013/074667); or, via mutation as herein.

[00115] Typically, in the context of an endogenous CRISPR system, formation of a CRISPR complex (comprising a guide sequence hybridized to a target sequence and complexed with one or more Cas9 proteins) results in cleavage of one or both strands in or near (e.g. within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, or more base pairs from) the target sequence. Without wishing to be bound by theory, the tracr sequence, which may comprise or consist of all or a portion of a wild-type tracr sequence (e.g. about or more than about 20, 26, 32, 45, 48, 54, 63, 67, 85, or more

nucleotides of a wild-type tracr sequence), may also form part of a CRISPR complex, such as by hybridization along at least a portion of the tracr sequence to all or a portion of a tracr mate sequence that is operably linked to the guide sequence.

[00116] The nucleic acid molecule encoding a Cas9 is advantageously codon optimized Cas9. An example of a codon optimized sequence, is in this instance a sequence optimized for expression in a eukaryote, e.g., humans (i.e. being optimized for expression in humans), or for another eukaryote, animal or mammal as herein discussed; see, e.g., SaCas9 human codon optimized sequence in WO 2014/093622 (PCT/US2013/074667). Whilst this is preferred, it will be appreciated that other examples are possible and codon optimization for a host species other than human, or for codon optimization for specific organs is known. In some embodiments, an enzyme coding sequence encoding a Cas9 is codon optimized for expression in particular cells, such as eukaryotic cells. The eukaryotic cells may be those of or derived from a particular organism, such as a mammal, including but not limited to human, or non-human eukaryote or animal or mammal as herein discussed, e.g., mouse, rat, rabbit, dog, livestock, or non-human mammal or primate. In some embodiments, processes for modifying the germ line genetic identity of human beings and/or processes for modifying the genetic identity of animals which are likely to cause them suffering without any substantial medical benefit to man or animal, and also animals resulting from such processes, may be excluded. In general, codon optimization refers to a process of modifying a nucleic acid sequence for enhanced expression in the host cells of interest by replacing at least one codon (e.g. about or more than about 1, 2, 3, 4, 5, 10, 15, 20, 25, 50, or more codons) of the native sequence with codons that are more frequently or most frequently used in the genes of that host cell while maintaining the native amino acid sequence. Various species exhibit particular bias for certain codons of a particular amino acid. Codon bias (differences in codon usage between organisms) often correlates with the efficiency of translation of messenger RNA (mRNA), which is in turn believed to be dependent on, among other things, the properties of the codons being translated and the availability of particular transfer RNA (tRNA) molecules. The predominance of selected tRNAs in a cell is generally a reflection of the codons used most frequently in peptide synthesis. Accordingly, genes can be tailored for optimal gene expression in a given organism based on codon optimization. Codon usage tables are readily available, for example, at the "Codon Usage Database" available at www.kazusa.or.jp/codon/ and these tables can be adapted in a number of ways. See Nakamura,

Y., et al. "Codon usage tabulated from the international DNA sequence databases: status for the year 2000" Nucl. Acids Res. 28:292 (2000). Computer algorithms for codon optimizing a particular sequence for expression in a particular host cell are also available, such as Gene Forge (Aptagen; Jacobus, PA), are also available. In some embodiments, one or more codons (e.g. 1, 2, 3, 4, 5, 10, 15, 20, 25, 50, or more, or all codons) in a sequence encoding a Cas9 correspond to the most frequently used codon for a particular amino acid.

[00117] In certain embodiments, the methods as described herein may comprise providing a Cas9 transgenic cell in which one or more nucleic acids encoding one or more guide RNAs are provided or introduced operably connected in the cell with a regulatory element comprising a promoter of one or more gene of interest. As used herein, the term "Cas9 transgenic cell" refers to a cell, such as a eukaryotic cell, in which a Cas9 gene has been genomically integrated. The nature, type, or origin of the cell are not particularly limiting according to the present invention. Also the way in which the Cas9 transgene is introduced in the cell may vary and can be any method as is known in the art. In certain embodiments, the Cas9 transgenic cell is obtained by introducing the Cas9 transgene in an isolated cell. In certain other embodiments, the Cas9 transgenic cell is obtained by isolating cells from a Cas9 transgenic organism. By means of example, and without limitation, the Cas9 transgenic cell as referred to herein may be derived from a Cas9 transgenic eukaryote, such as a Cas9 knock-in eukaryote. Reference is made to WO 2014/093622 (PCT/US13/74667), incorporated herein by reference. Methods of US Patent Publication Nos. 20120017290 and 20110265198 assigned to Sangamo BioSciences, Inc. directed to targeting the Rosa locus may be modified to utilize the CRISPR-Cas9 system of the present invention. Methods of US Patent Publication No. 20130236946 assigned to Collectis directed to targeting the Rosa locus may also be modified to utilize the CRISPR-Cas9 system of the present invention. By means of further example reference is made to Platt et. al. (Cell; 159(2):440-455 (2014)), describing a Cas9 knock-in mouse, which is incorporated herein by reference. The Cas9 transgene can further comprise a Lox-Stop-polyA-Lox(LSL) cassette thereby rendering Cas9 expression inducible by Cre recombinase. Alternatively, the Cas9 transgenic cell may be obtained by introducing the Cas9 transgene in an isolated cell. Delivery systems for transgenes are well known in the art. By means of example, the Cas9 transgene may be delivered in for instance eukaryotic cell by means of vector (e.g., AAV, adenovirus, lentivirus) and/or particle and/or nanoparticle delivery, as also described herein elsewhere.

[00118] It will be understood by the skilled person that the cell, such as the Cas9 transgenic cell, as referred to herein may comprise further genomic alterations besides having an integrated Cas9 gene or the mutations arising from the sequence specific action of Cas9 when complexed with RNA capable of guiding Cas9 to a target locus, such as for instance one or more oncogenic mutations, as for instance and without limitation described in Platt et al. (2014), Chen et al., (2014) or Kumar et al.. (2009).

[00119] In some embodiments, the Cas9 sequence is fused to one or more nuclear localization sequences (NLSs), such as about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more NLSs. In some embodiments, the Cas9 comprises about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more NLSs at or near the amino-terminus, about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more NLSs at or near the carboxy-terminus, or a combination of these (e.g. zero or at least one or more NLS at the amino-terminus and zero or at one or more NLS at the carboxy terminus). When more than one NLS is present, each may be selected independently of the others, such that a single NLS may be present in more than one copy and/or in combination with one or more other NLSs present in one or more copies. In a preferred embodiment of the invention, the Cas9 comprises at most 6 NLSs. In some embodiments, an NLS is considered near the N- or C-terminus when the nearest amino acid of the NLS is within about 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 40, 50, or more amino acids along the polypeptide chain from the N- or C-terminus. Non-limiting examples of NLSs include an NLS sequence derived from: the NLS of the SV40 virus large T-antigen, having the amino acid sequence PKKKRKV(SEQ ID NO: X); the NLS from nucleoplasmin (e.g. the nucleoplasmin bipartite NLS with the sequence KRPAATKKAGQAKKKK) (SEQ ID NO: X); the c-myc NLS having the amino acid sequence PAAKRVKLD (SEQ ID NO: X) or RQRRNELKRSP(SEQ ID NO: X); the hRNPA1 M9 NLS having the sequence NQSSNFGPMKGGNFGGRSSGPYGGGGQYFAKPRNQGGY(SEQ ID NO: X); the sequence RMRIZFKNKGKDTAELRRRRVEVSVELRKAKKDEQILKRRNV (SEQ ID NO: X) of the IBB domain from importin-alpha; the sequences VSRKRPRP (SEQ ID NO: X) and PPKKARED (SEQ ID NO: X) of the myoma T protein; the sequence POPKKKPL (SEQ ID NO: X) of human p53; the sequence SALIKKKKKMAP (SEQ ID NO: X) of mouse c-abl IV; the sequences DRLRR (SEQ ID NO: X) and PKQKKRK (SEQ ID NO: X) of the influenza virus NS1; the sequence RKLKKKIKKL (SEQ ID NO: X) of the Hepatitis virus delta antigen; the sequence REKKKFLKRR (SEQ ID NO: X) of the mouse Mx1 protein; the sequence

KRKGDEVVDGVDEVAKKSKK (SEQ ID NO: X) of the human poly(ADP-ribose) polymerase; and the sequence RKCLQAGMNLEARKTKK (SEQ ID NO: X) of the steroid hormone receptors (human) glucocorticoid. In general, the one or more NLSs are of sufficient strength to drive accumulation of the Cas9 in a detectable amount in the nucleus of a eukaryotic cell. In general, strength of nuclear localization activity may derive from the number of NLSs in the Cas, the particular NLS(s) used, or a combination of these factors. Detection of accumulation in the nucleus may be performed by any suitable technique. For example, a detectable marker may be fused to the Cas, such that location within a cell may be visualized, such as in combination with a means for detecting the location of the nucleus (e.g. a stain specific for the nucleus such as DAPI). Cell nuclei may also be isolated from cells, the contents of which may then be analyzed by any suitable process for detecting protein, such as immunohistochemistry, Western blot, or enzyme activity assay. Accumulation in the nucleus may also be determined indirectly, such as by an assay for the effect of CRISPR complex formation (e.g. assay for DNA cleavage or mutation at the target sequence, or assay for altered gene expression activity affected by CRISPR complex formation and/or Cas9 enzyme activity), as compared to a control not exposed to the Cas9 or complex, or exposed to a Cas9 lacking the one or more NLSs. In other embodiments, no NLS is required.

[00120] In certain aspects the invention involves vectors, e.g. for delivering or introducing in a cell Cas9 and/or RNA capable of guiding Cas9 to a target locus (i.e. guide RNA), but also for propagating these components (e.g. in prokaryotic cells). As used herein, a “vector” is a tool that allows or facilitates the transfer of an entity from one environment to another. It is a replicon, such as a plasmid, phage, or cosmid, into which another DNA segment may be inserted so as to bring about the replication of the inserted segment. Generally, a vector is capable of replication when associated with the proper control elements. In general, the term “vector” refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. Vectors include, but are not limited to, nucleic acid molecules that are single-stranded, double-stranded, or partially double-stranded; nucleic acid molecules that comprise one or more free ends, no free ends (e.g. circular); nucleic acid molecules that comprise DNA, RNA, or both; and other varieties of polynucleotides known in the art. One type of vector is a “plasmid,” which refers to a circular double stranded DNA loop into which additional DNA segments can be inserted, such as by standard molecular cloning techniques. Another type of vector is a viral

vector, wherein virally-derived DNA or RNA sequences are present in the vector for packaging into a virus (e.g. retroviruses, replication defective retroviruses, adenoviruses, replication defective adenoviruses, and adeno-associated viruses (AAVs)). Viral vectors also include polynucleotides carried by a virus for transfection into a host cell. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g. bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively-linked. Such vectors are referred to herein as “expression vectors.” Common expression vectors of utility in recombinant DNA techniques are often in the form of plasmids.

[00121] Recombinant expression vectors can comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory elements, which may be selected on the basis of the host cells to be used for expression, that is operatively-linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, “operably linked” is intended to mean that the nucleotide sequence of interest is linked to the regulatory element(s) in a manner that allows for expression of the nucleotide sequence (e.g. in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). With regards to recombination and cloning methods, mention is made of U.S. patent application 10/815,730, published September 2, 2004 as US 2004-0171156 A1, the contents of which are herein incorporated by reference in their entirety.

[00122] The vector(s) can include the regulatory element(s), e.g., promoter(s). The vector(s) can comprise Cas9 encoding sequences, and/or a single, but possibly also can comprise at least 3 or 8 or 16 or 32 or 48 or 50 guide RNA(s) (e.g., sgRNAs) encoding sequences, such as 1-2, 1-3, 1-4 1-5, 3-6, 3-7, 3-8, 3-9, 3-10, 3-8, 3-16, 3-30, 3-32, 3-48, 3-50 RNA(s) (e.g., sgRNAs). In a single vector there can be a promoter for each RNA (e.g., sgRNA), advantageously when there are up to about 16 RNA(s) (e.g., sgRNAs); and, when a single vector provides for more than 16 RNA(s) (e.g., sgRNAs), one or more promoter(s) can drive expression of more than one of the RNA(s) (e.g., sgRNAs), e.g., when there are 32 RNA(s) (e.g., sgRNAs), each promoter can drive expression of two RNA(s) (e.g., sgRNAs), and when there are 48 RNA(s) (e.g., sgRNAs), each

promoter can drive expression of three RNA(s) (e.g., sgRNAs). By simple arithmetic and well established cloning protocols and the teachings in this disclosure one skilled in the art can readily practice the invention as to the RNA(s) (e.g., sgRNA(s) for a suitable exemplary vector such as AAV, and a suitable promoter such as the U6 promoter, e.g., U6-sgRNAs. For example, the packaging limit of AAV is ~4.7 kb. The length of a single U6-sgRNA (plus restriction sites for cloning) is 361 bp. Therefore, the skilled person can readily fit about 12-16, e.g., 13 U6-sgRNA cassettes in a single vector. This can be assembled by any suitable means, such as a golden gate strategy used for TALE assembly (<http://www.genome-engineering.org/taleffectors/>). The skilled person can also use a tandem guide strategy to increase the number of U6-sgRNAs by approximately 1.5 times, e.g., to increase from 12-16, e.g., 13 to approximately 18-24, e.g., about 19 U6-sgRNAs. Therefore, one skilled in the art can readily reach approximately 18-24, e.g., about 19 promoter-RNAs, e.g., U6-sgRNAs in a single vector, e.g., an AAV vector. A further means for increasing the number of promoters and RNAs, e.g., sgRNA(s) in a vector is to use a single promoter (e.g., U6) to express an array of RNAs, e.g., sgRNAs separated by cleavable sequences. And an even further means for increasing the number of promoter-RNAs, e.g., sgRNAs in a vector, is to express an array of promoter-RNAs, e.g., sgRNAs separated by cleavable sequences in the intron of a coding sequence or gene; and, in this instance it is advantageous to use a polymerase II promoter, which can have increased expression and enable the transcription of long RNA in a tissue specific manner. (see, e.g., <http://nar.oxfordjournals.org/content/34/7/e53.short>, <http://www.nature.com/mt/journal/v16/n9/abs/mt2008144a.html>). In an advantageous embodiment, AAV may package U6 tandem sgRNA targeting up to about 50 genes. Accordingly, from the knowledge in the art and the teachings in this disclosure the skilled person can readily make and use vector(s), e.g., a single vector, expressing multiple RNAs or guides or sgRNAs under the control or operatively or functionally linked to one or more promoters—especially as to the numbers of RNAs or guides or sgRNAs discussed herein, without any undue experimentation.

[00123] The guide RNA(s), e.g., sgRNA(s) encoding sequences and/or Cas9 encoding sequences, can be functionally or operatively linked to regulatory element(s) and hence the regulatory element(s) drive expression. The promoter(s) can be constitutive promoter(s) and/or conditional promoter(s) and/or inducible promoter(s) and/or tissue specific promoter(s). The

promoter can be selected from the group consisting of RNA polymerases, pol I, pol II, pol III, T7, U6, H1, retroviral Rous sarcoma virus (RSV) LTR promoter, the cytomegalovirus (CMV) promoter, the SV40 promoter, the dihydrofolate reductase promoter, the β -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1 α promoter. An advantageous promoter is the promoter is U6.

[00124] As used herein, the term “crRNA” or “guide RNA” or “single guide RNA” or “sgRNA” or “one or more nucleic acid components” of a Type II CRISPR-Cas9 locus effector protein comprises any polynucleotide sequence having sufficient complementarity with a target nucleic acid sequence to hybridize with the target nucleic acid sequence and direct sequence-specific binding of a nucleic acid-targeting complex to the target nucleic acid sequence. In some embodiments, the degree of complementarity, when optimally aligned using a suitable alignment algorithm, is about or more than about 50%, 60%, 75%, 80%, 85%, 90%, 95%, 97.5%, 99%, or more. Optimal alignment may be determined with the use of any suitable algorithm for aligning sequences, non-limiting example of which include the Smith-Waterman algorithm, the Needleman-Wunsch algorithm, algorithms based on the Burrows-Wheeler Transform (e.g., the Burrows Wheeler Aligner), ClustalW, Clustal X, BLAT, Novoalign (Novocraft Technologies; available at www.novocraft.com), ELAND (Illumina, San Diego, CA), SOAP (available at soap.genomics.org.cn), and Maq (available at maq.sourceforge.net). The ability of a guide sequence (within a nucleic acid-targeting guide RNA) to direct sequence-specific binding of a nucleic acid -targeting complex to a target nucleic acid sequence may be assessed by any suitable assay. For example, the components of a nucleic acid-targeting CRISPR system sufficient to form a nucleic acid-targeting complex, including the guide sequence to be tested, may be provided to a host cell having the corresponding target nucleic acid sequence, such as by transfection with vectors encoding the components of the nucleic acid-targeting complex, followed by an assessment of preferential targeting (e.g., cleavage) within the target nucleic acid sequence, such as by Surveyor assay as described herein. Similarly, cleavage of a target nucleic acid sequence may be evaluated in a test tube by providing the target nucleic acid sequence, components of a nucleic acid -targeting complex, including the guide sequence to be tested and a control guide sequence different from the test guide sequence, and comparing binding or rate of cleavage at the target sequence between the test and control guide sequence reactions. Other assays are possible, and will occur to those skilled in the art. A guide sequence, and hence a

nucleic acid-targeting guide RNA may be selected to target any target nucleic acid sequence. The target sequence may be DNA. The target sequence may be any DNA that encodes an RNA sequence. In some embodiments, the target sequence may be a sequence that encodes an RNA molecule selected from messenger RNA (mRNA), pre-mRNA, ribosomal RNA (rRNA), transfer RNA (tRNA), micro-RNA (miRNA), small interfering RNA (siRNA), small nuclear RNA (snRNA), small nucleolar RNA (snoRNA), double stranded RNA (dsRNA), non coding RNA (ncRNA), long non-coding RNA (lncRNA), and small cytoplasmic RNA (scRNA). In some embodiments, the target sequence may be a DNA sequence encoding a sequence within an RNA molecule selected from mRNA, pre-mRNA, and rRNA. In some embodiments, the target sequence may encode a sequence within a RNA molecule selected from ncRNA, and lncRNA. In some embodiments, the target sequence may encode a sequence within an mRNA molecule or a pre-mRNA molecule.

[00125] In some embodiments, a nucleic acid-targeting guide RNA is selected to reduce the degree secondary structure within the DNA-targeting guide RNA. In some embodiments, about or less than about 75%, 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 1%, or fewer of the nucleotides of the nucleic acid-targeting guide RNA participate in self-complementary base pairing when optimally folded. Optimal folding may be determined by any suitable polynucleotide folding algorithm. Some programs are based on calculating the minimal Gibbs free energy. An example of one such algorithm is mFold, as described by Zuker and Stiegler (Nucleic Acids Res. 9 (1981), 133-148). Another example folding algorithm is the online webserver RNAfold, developed at Institute for Theoretical Chemistry at the University of Vienna, using the centroid structure prediction algorithm (see e.g., A.R. Gruber et al., 2008, Cell 106(1): 23-24; and PA Carr and GM Church, 2009, Nature Biotechnology 27(12): 1151-62).

[00126] In certain embodiments, a guide RNA or crRNA may comprise, consist essentially of, or consist of a direct repeat (DR) sequence and a guide sequence or spacer sequence. In certain embodiments, the guide RNA or crRNA may comprise, consist essentially of, or consist of a direct repeat sequence fused or linked to a guide sequence or spacer sequence. In certain embodiments, the direct repeat sequence may be located upstream (i.e., 5') from the guide sequence or spacer sequence. In other embodiments, the direct repeat sequence may be located downstream (i.e., 3') from the guide sequence or spacer sequence.

[00127] In certain embodiments, the crRNA comprises a stem loop, preferably a single stem loop. In certain embodiments, the direct repeat sequence forms a stem loop, preferably a single stem loop.

[00128] The “tracrRNA” sequence or analogous terms includes any polynucleotide sequence that has sufficient complementarity with a crRNA sequence to hybridize. In general, degree of complementarity is with reference to the optimal alignment of the tracr mate sequence and tracr sequence, along the length of the shorter of the two sequences. Optimal alignment may be determined by any suitable alignment algorithm, and may further account for secondary structures, such as self-complementarity within either the tracr sequence or tracr mate sequence. In some embodiments, the degree of complementarity between the tracr sequence and the tracr mate sequence along the length of the shorter of the two when optimally aligned is about or more than about 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 97.5%, 99%, or higher.

[00129] A guide sequence may be selected to target any target sequence. In some embodiments, the target sequence is a sequence within a genome of a cell. Exemplary target sequences include those that are unique in the target genome. For example, for the *S. pyogenes* Cas9, a unique target sequence in a genome may include a Cas9 target site of the form MMMMMMMMNNNNNNNNNNNNXGG where NNNNNNNNNNNXGG (N is A, G, T, or C; and X can be anything) has a single occurrence in the genome. A unique target sequence in a genome may include an *S. pyogenes* Cas9 target site of the form MMMMMMMMNNNNNNNNNNNNXGG where NNNNNNNNNNNXGG (N is A, G, T, or C; and X can be anything) has a single occurrence in the genome. For the *S. thermophilus* CRISPR1 Cas9, a unique target sequence in a genome may include a Cas9 target site of the form MMMMMMMMNNNNNNNNNNNNXXAGAAW where NNNNNNNNNNNXXAGAAW (N is A, G, T, or C; X can be anything; and W is A or T) has a single occurrence in the genome. A unique target sequence in a genome may include an *S. thermophilus* CRISPR1 Cas9 target site of the form MMMMMMMMNNNNNNNNNNNNXXAGAAW where NNNNNNNNNNNXXAGAAW (N is A, G, T, or C; X can be anything; and W is A or T) has a single occurrence in the genome. For the *S. pyogenes* Cas9, a unique target sequence in a genome may include a Cas9 target site of the form MMMMMMMMNNNNNNNNNNNNXGGXG where NNNNNNNNNNNXGGXG (N is A, G, T, or C; and X can be anything) has a single occurrence in the genome. A unique target

sequence in a genome may include an *S. pyogenes* Cas9 target site of the form MMMMMMMMNNNNNNNNNNNXGGXG where NNNNNNNNNNNNXGGXG (N is A, G, T, or C; and X can be anything) has a single occurrence in the genome. In each of these sequences “M” may be A, G, T, or C, and need not be considered in identifying a sequence as unique. In some embodiments, a guide sequence is selected to reduce the degree secondary structure within the guide sequence. In some embodiments, about or less than about 75%, 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 1%, or fewer of the nucleotides of the guide sequence participate in self-complementary base pairing when optimally folded. Optimal folding may be determined by any suitable polynucleotide folding algorithm. Some programs are based on calculating the minimal Gibbs free energy. An example of one such algorithm is mFold, as described by Zuker and Stiegler (Nucleic Acids Res. 9 (1981), 133-148). Another example folding algorithm is the online webserver RNAfold, developed at Institute for Theoretical Chemistry at the University of Vienna, using the centroid structure prediction algorithm (see e.g. A.R. Gruber et al., 2008, Cell 106(1): 23-24; and PA Carr and GM Church, 2009, Nature Biotechnology 27(12): 1151-62).

[00130] In general, a tracr mate sequence includes any sequence that has sufficient complementarity with a tracr sequence to promote one or more of: (1) excision of a guide sequence flanked by tracr mate sequences in a cell containing the corresponding tracr sequence; and (2) formation of a CRISPR complex at a target sequence, wherein the CRISPR complex comprises the tracr mate sequence hybridized to the tracr sequence. In general, degree of complementarity is with reference to the optimal alignment of the tracr mate sequence and tracr sequence, along the length of the shorter of the two sequences. Optimal alignment may be determined by any suitable alignment algorithm, and may further account for secondary structures, such as self-complementarity within either the tracr sequence or tracr mate sequence. In some embodiments, the degree of complementarity between the tracr sequence and tracr mate sequence along the length of the shorter of the two when optimally aligned is about or more than about 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 97.5%, 99%, or higher. In some embodiments, the tracr sequence is about or more than about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, or more nucleotides in length. In some embodiments, the tracr sequence and tracr mate sequence are contained within a single transcript, such that hybridization between the two produces a transcript having a secondary structure, such as a hairpin. In an

embodiment of the invention, the transcript or transcribed polynucleotide sequence has at least two or more hairpins. In preferred embodiments, the transcript has two, three, four or five hairpins. In a further embodiment of the invention, the transcript has at most five hairpins. In a hairpin structure the portion of the sequence 5' of the final "N" and upstream of the loop corresponds to the tracr mate sequence, and the portion of the sequence 3' of the loop corresponds to the tracr sequence. Further non-limiting examples of single polynucleotides comprising a guide sequence, a tracr mate sequence, and a tracr sequence are as follows (listed 5' to 3'), where "N" represents a base of a guide sequence, the first block of lower case letters represent the tracr mate sequence, and the second block of lower case letters represent the tracr sequence, and the final poly-T sequence represents the transcription terminator:

(1) NNNNNNNNNNNNNNNNNNNNNggttttgtagagctcaGAAAtaaatcttgcagaagctacaaagataa
ggcttcatgccgaaatcaacaccctgtcattttatggcagggtgttttcgtatttaaTTTTTT;

(2) NNNNNNNNNNNNNNNNNNNNNggttttgtagagctcaGAAAtgcagaagctacaaagataaggcttcatgccg
aaatcaacaccctgtcattttatggcagggtgttttcgtatttaaTTTTTT;

(3) NNNNNNNNNNNNNNNNNNNNNggttttgtagagctcaGAAAtgcagaagctacaaagataaggcttcatgccg
aaatcaacaccctgtcattttatggcagggtgtTTTTTT;

(4) NNNNNNNNNNNNNNNNNNNNNggttttagagctcaGAAAtagcaagttaaaataaggctagtcggttatcaact
gaaaagtggcaccgagtcggtgcTTTTTT;

(5) NNNNNNNNNNNNNNNNNNNNNggttttagagctcaGAAATAGcaagttaaaataaggctagtcggttatcaac
ttgaaaagtTTTTTTT; and

(6) NNNNNNNNNNNNNNNNNNNNNggttttagagctagAAATAGcaagttaaaataaggctagtcggttatcaTT
TTTTTT. In some embodiments, sequences (1) to (3) are used in combination with Cas9 from *S. thermophilus* CRISPR1. In some embodiments, sequences (4) to (6) are used in combination with Cas9 from *S. pyogenes*. In some embodiments, the tracr sequence is a separate transcript from a transcript comprising the tracr mate sequence.

[00131] In some embodiments, candidate tracrRNA may be subsequently predicted by sequences that fulfill any or all of the following criteria: 1. sequence homology to direct repeats (motif search in Geneious with up to 18-bp mismatches); 2. presence of a predicted Rho-independent transcriptional terminator in direction of transcription; and 3. stable hairpin secondary structure between tracrRNA and direct repeat. In some embodiments, 2 of these

criteria may be used, for instance 1 and 2, 2 and 3, or 1 and 3. In some embodiments, all 3 criteria may be used.

[00132] In some embodiments, chimeric synthetic guide RNAs (sgRNAs) designs may incorporate at least 12 bp of duplex structure between the direct repeat and tracrRNA.

[00133] For minimization of toxicity and off-target effects, it will be important to control the concentration of CRISPR enzyme mRNA and guide RNA delivered. Optimal concentrations of CRISPR enzyme mRNA and guide RNA can be determined by testing different concentrations in a cellular or non-human eukaryote animal model and using deep sequencing to analyze the extent of modification at potential off-target genomic loci. For example, for the guide sequence targeting 5'-GAGTCCGAGCAGAAGAAGAA-3' in the EMX1 gene of the human genome, deep sequencing can be used to assess the level of modification at the following two off-target loci, 1: 5'-GAGTCCTAGCAGGAGAAGAA-3' and 2: 5'-GAGTCTAAGCAGAAGAAGAA-3'. The concentration that gives the highest level of on-target modification while minimizing the level of off-target modification should be chosen for in vivo delivery. Alternatively, to minimize the level of toxicity and off-target effect, CRISPR enzyme nickase mRNA (for example *S. pyogenes* Cas9 with the D10A mutation) can be delivered with a pair of guide RNAs targeting a site of interest. The two guide RNAs need to be spaced as follows. Guide sequences and strategies to minimize toxicity and off-target effects can be as in WO 2014/093622 (PCT/US2013/074667).

[00134] In an aspect of the invention, novel DNA targeting systems also referred to as DNA-targeting CRISPR/Cas or the CRISPR-Cas DNA-targeting system of the present application are based on identified Type II Cas9 proteins which do not require the generation of customized proteins to target specific DNA sequences but rather a single effector protein or enzyme can be programmed by a RNA molecule to recognize a specific DNA target, in other words the enzyme can be recruited to a specific DNA target using said RNA molecule. Aspects of the invention particularly relate to DNA targeting RNA-guided Cas9 CRISPR systems.

[00135] The nucleic acids-targeting systems, the vector systems, the vectors and the compositions described herein may be used in various nucleic acids-targeting applications, altering or modifying synthesis of a gene product, such as a protein, nucleic acids cleavage, nucleic acids editing, nucleic acids splicing; trafficking of target nucleic acids, tracing of target nucleic acids, isolation of target nucleic acids, visualization of target nucleic acids, etc.

[00136] Aspects of the invention also encompass methods and uses of the compositions and systems described herein in genome engineering, e.g. for altering or manipulating the expression of one or more genes or the one or more gene products, in prokaryotic or eukaryotic cells, *in vitro*, *in vivo* or *ex vivo*.

[00137] The CRISPR system is derived advantageously from a type II CRISPR system. In some embodiments, one or more elements of a CRISPR system is derived from a particular organism comprising an endogenous CRISPR system, such as *Streptococcus pyogenes*. The CRISPR system is a type II CRISPR system and the Cas enzyme is Cas9, which catalyzes DNA cleavage. Other non-limiting examples of Cas proteins include Cas1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4, homologues thereof, or modified versions thereof.

[00138] In an embodiment, the Cas9 protein may be an ortholog of an organism of a genus which includes but is not limited to Corynebacter, Sutterella, Legionella, Treponema, Filifactor, Eubacterium, Streptococcus, Lactobacillus, Mycoplasma, Bacteroides, Flavivola, Flavobacterium, Sphaerochaeta, Azospirillum, Gluconacetobacter, Neisseria, Roseburia, Parvibaculum, Staphylococcus, Nitratifactor, Mycoplasma and Campylobacter. Species of an organism of such a genus can be as otherwise herein discussed.

[00139] Some methods of identifying orthologs of CRISPR-Cas9 system enzymes may involve identifying tracr sequences in genomes of interest. Identification of tracr sequences may relate to the following steps: Search for the direct repeats or tracr mate sequences in a database to identify a CRISPR region comprising a CRISPR enzyme. Search for homologous sequences in the CRISPR region flanking the CRISPR enzyme in both the sense and antisense directions. Look for transcriptional terminators and secondary structures. Identify any sequence that is not a direct repeat or a tracr mate sequence but has more than 50% identity to the direct repeat or tracr mate sequence as a potential tracr sequence. Take the potential tracr sequence and analyze for transcriptional terminator sequences associated therewith.

[00140] It will be appreciated that any of the functionalities described herein may be engineered into CRISPR enzymes from other orthologs, including chimeric enzymes comprising fragments from multiple orthologs. Examples of such orthologs are described elsewhere herein.

Thus, chimeric enzymes may comprise fragments of CRISPR enzyme orthologs of an organism which includes but is not limited to *Corynebacter*, *Sutterella*, *Legionella*, *Treponema*, *Filifactor*, *Eubacterium*, *Streptococcus*, *Lactobacillus*, *Mycoplasma*, *Bacteroides*, *Flaviivola*, *Flavobacterium*, *Sphaerochaeta*, *Azospirillum*, *Gluconacetobacter*, *Neisseria*, *Roseburia*, *Parvibaculum*, *Staphylococcus*, *Nitratifactor*, *Mycoplasma* and *Campylobacter*. A chimeric enzyme can comprise a first fragment and a second fragment, and the fragments can be of CRISPR enzyme orthologs of organisms of genres herein mentioned or of species herein mentioned; advantageously the fragments are from CRISPR enzyme orthologs of different species

[00141] In some embodiments, the unmodified CRISPR enzyme has DNA cleavage activity, such as Cas9. In some embodiments, the CRISPR enzyme directs cleavage of one or both strands at the location of a target sequence, such as within the target sequence and/or within the complement of the target sequence. In some embodiments, the CRISPR enzyme directs cleavage of one or both strands within about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 50, 100, 200, 500, or more base pairs from the first or last nucleotide of a target sequence. In some embodiments, a vector encodes a CRISPR enzyme that is mutated to with respect to a corresponding wild-type enzyme such that the mutated CRISPR enzyme lacks the ability to cleave one or both strands of a target polynucleotide containing a target sequence. For example, an aspartate-to-alanine substitution (D10A) in the RuvC I catalytic domain of Cas9 from *S. pyogenes* converts Cas9 from a nuclease that cleaves both strands to a nickase (cleaves a single strand). Other examples of mutations that render Cas9 a nickase include, without limitation, H840A, N854A, and N863A. As a further example, two or more catalytic domains of Cas9 (RuvC I, RuvC II, and RuvC III or the HNH domain) may be mutated to produce a mutated Cas9 substantially lacking all DNA cleavage activity. In some embodiments, a D10A mutation is combined with one or more of H840A, N854A, or N863A mutations to produce a Cas9 enzyme substantially lacking all DNA cleavage activity. In some embodiments, a CRISPR enzyme is considered to substantially lack all DNA cleavage activity when the DNA cleavage activity of the mutated enzyme is about no more than 25%, 10%, 5%, 1%, 0.1%, 0.01%, or less of the DNA cleavage activity of the non-mutated form of the enzyme; an example can be when the DNA cleavage activity of the mutated form is nil or negligible as compared with the non-mutated form. Where the enzyme is not SpCas9, mutations may be made at any or all residues corresponding to positions 10, 762, 840, 854, 863 and/or 986 of SpCas9 (which may be ascertained for instance by standard sequence

comparison tools). In particular, any or all of the following mutations are preferred in SpCas9: D10A, E762A, H840A, N854A, N863A and/or D986A; as well as conservative substitution for any of the replacement amino acids is also envisaged. The same (or conservative substitutions of these mutations) at corresponding positions in other Cas9s are also preferred. Particularly preferred are D10 and H840 in SpCas9. However, in other Cas9s, residues corresponding to SpCas9 D10 and H840 are also preferred. Orthologs of SpCas9 can be used in the practice of the invention. Cas9 refers to the general class of enzymes that share homology to the biggest nuclease with multiple nuclease domains from the type II CRISPR system. Most preferably, the Cas9 enzyme is from, or is derived from, spCas9 (*S. pyogenes* Cas9) or saCas9 (*S. aureus* Cas9). "StCas9" refers to wild type Cas9 from *S. thermophilus*, the protein sequence of which is given in the SwissProt database under accession number G3ECR1. Similarly, *S. pyogenes* Cas9 or spCas9 is included in SwissProt under accession number Q99ZW2. By derived, Applicants mean that the derived enzyme is largely based, in the sense of having a high degree of sequence homology with, a wildtype enzyme, but that it has been mutated (modified) in some way as described herein. It will be appreciated that the terms Cas and CRISPR enzyme are generally used herein interchangeably, unless otherwise apparent. As mentioned above, many of the residue numberings used herein refer to the Cas9 enzyme from the type II CRISPR-Cas9 locus in *Streptococcus pyogenes*. However, it will be appreciated that this invention includes many more Cas9s from other species of microbes, such as SpCas9, SaCa9, St1Cas9 and so forth. Enzymatic action by Cas9 derived from *Streptococcus pyogenes* or any closely related Cas9 generates double stranded breaks at target site sequences which hybridize to 20 nucleotides of the guide sequence and that have a protospacer-adjacent motif (PAM) sequence (examples include NGG/NRG or a PAM that can be determined as described herein) following the 20 nucleotides of the target sequence. CRISPR activity through Cas9 for site-specific DNA recognition and cleavage is defined by the guide sequence, the tracr sequence that hybridizes in part to the guide sequence and the PAM sequence. More aspects of the CRISPR system are described in Karginov and Hannon, The CRISPR system: small RNA-guided defense in bacteria and archaea, *Mole Cell* 2010, January 15; 37(1): 7. The type II CRISPR locus from *Streptococcus pyogenes* SF370, which contains a cluster of four genes Cas9, Cas1, Cas2, and Csn1, as well as two non-coding RNA elements, tracrRNA and a characteristic array of repetitive sequences (direct repeats) interspaced by short stretches of non-repetitive sequences (spacers, about 30bp each). In this

system, targeted DNA double-strand break (DSB) is generated in four sequential steps. First, two non-coding RNAs, the pre-crRNA array and tracrRNA, are transcribed from the CRISPR locus. Second, tracrRNA hybridizes to the direct repeats of pre-crRNA, which is then processed into mature crRNAs containing individual spacer sequences. Third, the mature crRNA:tracrRNA complex directs Cas9 to the DNA target comprising, consisting essentially of, or consisting of the protospacer and the corresponding PAM via heteroduplex formation between the spacer region of the crRNA and the protospacer DNA. Finally, Cas9 mediates cleavage of target DNA upstream of PAM to create a DSB within the protospacer. A pre-crRNA array comprising, consisting essentially of, or consisting of a single spacer flanked by two direct repeats (DRs) is also encompassed by the term “tracr-mate sequences”). In certain embodiments, Cas9 may be constitutively present or inducibly present or conditionally present or administered or delivered. Cas9 optimization may be used to enhance function or to develop new functions, one can generate chimeric Cas9 proteins. And Cas9 may be used as a generic DNA binding protein.

[00142] Typically, in the context of an endogenous CRISPR system, formation of a CRISPR complex (comprising a guide sequence hybridized to a target sequence and complexed with one or more Cas9 proteins) results in cleavage of one or both strands in or near (e.g. within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, or more base pairs from) the target sequence. Without wishing to be bound by theory, the tracr sequence, which may comprise, consist essentially of, or consist of all or a portion of a wild-type tracr sequence (e.g. about or more than about 20, 26, 32, 45, 48, 54, 63, 67, 85, or more nucleotides of a wild-type tracr sequence), may also form part of a CRISPR complex, such as by hybridization along at least a portion of the tracr sequence to all or a portion of a tracr mate sequence that is operably linked to the guide sequence.

[00143] An example of a codon optimized sequence, is in this instance a sequence optimized for expression in a eukaryote, e.g., humans (i.e. being optimized for expression in humans), or for another eukaryote, animal or mammal as herein discussed; see, e.g., SaCas9 human codon optimized sequence in WO 2014/093622 (PCT/US2013/074667). Whilst this is preferred, it will be appreciated that other examples are possible and codon optimization for a host species other than human, or for codon optimization for specific organs is known. In some embodiments, an enzyme coding sequence encoding a CRISPR enzyme is codon optimized for expression in particular cells, such as eukaryotic cells. The eukaryotic cells may be those of or derived from a

particular organism, such as a mammal, including but not limited to human, or non-human eukaryote or animal or mammal as herein discussed, e.g., mouse, rat, rabbit, dog, livestock, or non-human mammal or primate. In some embodiments, processes for modifying the germ line genetic identity of human beings and/or processes for modifying the genetic identity of animals which are likely to cause them suffering without any substantial medical benefit to man or animal, and also animals resulting from such processes, may be excluded. In general, codon optimization refers to a process of modifying a nucleic acid sequence for enhanced expression in the host cells of interest by replacing at least one codon (e.g. about or more than about 1, 2, 3, 4, 5, 10, 15, 20, 25, 50, or more codons) of the native sequence with codons that are more frequently or most frequently used in the genes of that host cell while maintaining the native amino acid sequence. Various species exhibit particular bias for certain codons of a particular amino acid. Codon bias (differences in codon usage between organisms) often correlates with the efficiency of translation of messenger RNA (mRNA), which is in turn believed to be dependent on, among other things, the properties of the codons being translated and the availability of particular transfer RNA (tRNA) molecules. The predominance of selected tRNAs in a cell is generally a reflection of the codons used most frequently in peptide synthesis. Accordingly, genes can be tailored for optimal gene expression in a given organism based on codon optimization. Codon usage tables are readily available, for example, at the "Codon Usage Database" available at www.kazusa.or.jp/codon/ and these tables can be adapted in a number of ways. See Nakamura, Y., et al. "Codon usage tabulated from the international DNA sequence databases: status for the year 2000" *Nucl. Acids Res.* 28:292 (2000). Computer algorithms for codon optimizing a particular sequence for expression in a particular host cell are also available, such as Gene Forge (Aptagen; Jacobus, PA), are also available. In some embodiments, one or more codons (e.g. 1, 2, 3, 4, 5, 10, 15, 20, 25, 50, or more, or all codons) in a sequence encoding a CRISPR enzyme correspond to the most frequently used codon for a particular amino acid.

[00144] In some embodiments, a vector encodes a CRISPR enzyme comprising one or more nuclear localization sequences (NLSs), such as about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more NLSs. In some embodiments, the CRISPR enzyme comprises about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more NLSs at or near the amino-terminus, about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more NLSs at or near the carboxy-terminus, or a combination of these (e.g. zero or at least one or more NLS at the amino-terminus and zero or at one or more

NLS at the carboxy terminus). When more than one NLS is present, each may be selected independently of the others, such that a single NLS may be present in more than one copy and/or in combination with one or more other NLSs present in one or more copies. In a preferred embodiment of the invention, the CRISPR enzyme comprises at most 6 NLSs. In some embodiments, an NLS is considered near the N- or C-terminus when the nearest amino acid of the NLS is within about 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 40, 50, or more amino acids along the polypeptide chain from the N- or C-terminus. Non-limiting examples of NLSs include an NLS sequence derived from: the NLS of the SV40 virus large T-antigen, having the amino acid sequence PKKKRKV; the NLS from nucleoplasmin (e.g. the nucleoplasmin bipartite NLS with the sequence KRPAATKKAGQAKKKK); the c-myc NLS having the amino acid sequence PAAKRVKLD or RQRRNELKRSP; the hRNPA1 M9 NLS having the sequence NQSSNFGPMKGGNFGGRSSGPYGGGGQYFAKPRNQGGY; the sequence RMRIZFKNKGKDTAELRRRRVEVSVELRKAKKDEQILKRRNV of the IBB domain from importin-alpha; the sequences VSRKRPRP and PPKKARED of the myoma T protein; the sequence POPKKKPL of human p53; the sequence SALIKKKKKMAP of mouse c-abl IV; the sequences DRLRR and PKQKKRK of the influenza virus NS1; the sequence RKLKKKIKKL of the Hepatitis virus delta antigen; the sequence REKKKFLKRR of the mouse Mx1 protein; the sequence KRKGDEVDGVDEVAKKKSKK of the human poly(ADP-ribose) polymerase; and the sequence RKCLQAGMNLEARKTKK of the steroid hormone receptors (human) glucocorticoid. In general, the one or more NLSs are of sufficient strength to drive accumulation of the CRISPR enzyme in a detectable amount in the nucleus of a eukaryotic cell. In general, strength of nuclear localization activity may derive from the number of NLSs in the CRISPR enzyme, the particular NLS(s) used, or a combination of these factors. Detection of accumulation in the nucleus may be performed by any suitable technique. For example, a detectable marker may be fused to the CRISPR enzyme, such that location within a cell may be visualized, such as in combination with a means for detecting the location of the nucleus (e.g. a stain specific for the nucleus such as DAPI). Cell nuclei may also be isolated from cells, the contents of which may then be analyzed by any suitable process for detecting protein, such as immunohistochemistry, Western blot, or enzyme activity assay. Accumulation in the nucleus may also be determined indirectly, such as by an assay for the effect of CRISPR complex formation (e.g. assay for DNA cleavage or mutation at the target sequence, or assay for altered

gene expression activity affected by CRISPR complex formation and/or CRISPR enzyme activity), as compared to a control not exposed to the CRISPR enzyme or complex, or exposed to a CRISPR enzyme lacking the one or more NLSs.

[00145] While it is preferred for a dead guide to lack detectable nuclease activity in a CRISPR complex, in certain embodiments, a dead guide complexed with an active Cas9 may comprise reduced or residual nuclease activity as compared to an active guide. Reduced or residual nuclease activity can comprise 20% or less, or 10% or less, or 8% or less, or 5% or less, or 3% or less, or 2% or less, or 1% or less, or 0.5% or less, or 0.2% or less, or 0.1% or less than that of an active guide complexed with an active Cas9. Nuclease activity can be measured by indel formation, for example by Surveyor or sequencing.

[00146] In an aspect, the invention provides multiplex regulation of a plurality of gene loci. For example, in certain embodiments, an active Cas9 enzyme is used with a first guide, which is a dead guide associated with a functional domain operable at one locus and a second guide which directs the Cas9 enzyme to cleave a second locus. In such embodiments, a template polynucleotide can be introduced into the DNA molecule at the cleaved locus or an intervening sequence excised for example by generating overhangs that reanneal and ligate. sgRNA pairs creating 5' overhangs with less than 8bp overlap between the guide sequences (offset greater than -8 bp) were able to efficiently mediate detectable indel formation. Accordingly, the activity or function of a gene product from the cleaved locus can be altered or the expression of the gene product be increased or decreased. In an embodiment of the invention, the gene product is a protein. In embodiments involving overhangs and or recombination templates, the Cas9 is preferably a nickase. In certain embodiments, nickases are used in pairs to generate overhangs at the cleaved locus. In certain embodiments, a nickase pair generates 5' overhangs at the cleavage sites. In other embodiments, a nickase pair generate 3' overhangs at the cleavage sites. In other embodiments, a nickase pair generates a 5' overhang at one cleavage site and a 3' overhang at the other cleavage site.

[00147] In embodiments in which a recombination template is provided, the recombination template may be a component of the same vector as provides another CRISPR-Cas9 system component, contained in a separate vector, or provided as a separate polynucleotide. In some embodiments, a recombination template is designed to serve as a template in homologous recombination, such as within or near a target sequence nicked or cleaved by a CRISPR enzyme

as a part of a CRISPR complex. A template polynucleotide may be of any suitable length, such as about or more than about 10, 15, 20, 25, 50, 75, 100, 150, 200, 500, 1000, or more nucleotides in length. In some embodiments, the template polynucleotide is complementary to a portion of a polynucleotide comprising the target sequence. When optimally aligned, a template polynucleotide might overlap with one or more nucleotides of a target sequences (e.g. about or more than about 1, 5, 10, 15, 20, or more nucleotides). In some embodiments, when a template sequence and a polynucleotide comprising a target sequence are optimally aligned, the nearest nucleotide of the template polynucleotide is within about 1, 5, 10, 15, 20, 25, 50, 75, 100, 200, 300, 400, 500, 1000, 5000, 10000, or more nucleotides from the target sequence.

[00148] In some embodiments, one or more vectors driving expression of one or more elements of a CRISPR system are introduced into a host cell such that expression of the elements of the CRISPR system direct formation of a CRISPR complex at one or more target sites. For example, a Cas9 enzyme, a guide sequence linked to a tracr-mate sequence, and a tracr sequence could each be operably linked to separate regulatory elements on separate vectors. Or, RNA(s) of the CRISPR System can be delivered to a transgenic Cas9 animal or mammal, e.g., an animal or mammal that constitutively or inducibly or conditionally expresses Cas9; or an animal or mammal that is otherwise expressing Cas9 or has cells containing Cas9, such as by way of prior administration thereto of a vector or vectors that code for and express *in vivo* Cas9. Alternatively, two or more of the elements expressed from the same or different regulatory elements, may be combined in a single vector, with one or more additional vectors providing any components of the CRISPR system not included in the first vector. CRISPR system elements that are combined in a single vector may be arranged in any suitable orientation, such as one element located 5' with respect to ("upstream" of) or 3' with respect to ("downstream" of) a second element. The coding sequence of one element may be located on the same or opposite strand of the coding sequence of a second element, and oriented in the same or opposite direction. In some embodiments, a single promoter drives expression of a transcript encoding a CRISPR enzyme and one or more of the guide sequence, tracr mate sequence (optionally operably linked to the guide sequence), and a tracr sequence embedded within one or more intron sequences (e.g. each in a different intron, two or more in at least one intron, or all in a single intron). In some embodiments, the CRISPR enzyme, guide sequence, tracr mate sequence, and tracr sequence are operably linked to and expressed from the same promoter. Delivery vehicles,

vectors, particles, nanoparticles, formulations and components thereof for expression of one or more elements of a CRISPR system are as used in the foregoing documents, such as WO 2014/093622 (PCT/US2013/074667). In some embodiments, a vector comprises one or more insertion sites, such as a restriction endonuclease recognition sequence (also referred to as a “cloning site”). In some embodiments, one or more insertion sites (e.g. about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more insertion sites) are located upstream and/or downstream of one or more sequence elements of one or more vectors. In some embodiments, a vector comprises an insertion site upstream of a tracr mate sequence, and optionally downstream of a regulatory element operably linked to the tracr mate sequence, such that following insertion of a guide sequence into the insertion site and upon expression the guide sequence directs sequence-specific binding of a CRISPR complex to a target sequence in a eukaryotic cell. In some embodiments, a vector comprises two or more insertion sites, each insertion site being located between two tracr mate sequences so as to allow insertion of a guide sequence at each site. In such an arrangement, the two or more guide sequences may comprise two or more copies of a single guide sequence, two or more different guide sequences, or combinations of these. When multiple different guide sequences are used, a single expression construct may be used to target CRISPR activity to multiple different, corresponding target sequences within a cell. For example, a single vector may comprise about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, or more guide sequences. In some embodiments, about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more such guide-sequence-containing vectors may be provided, and optionally delivered to a cell. In some embodiments, a vector comprises a regulatory element operably linked to an enzyme-coding sequence encoding a CRISPR enzyme, such as a Cas protein. CRISPR enzyme or CRISPR enzyme mRNA or CRISPR guide RNA or RNA(s) can be delivered separately; and advantageously at least one of these is delivered via a nanoparticle complex. CRISPR enzyme mRNA can be delivered prior to the guide RNA to give time for CRISPR enzyme to be expressed. CRISPR enzyme mRNA might be administered 1-12 hours (preferably around 2-6 hours) prior to the administration of guide RNA. Alternatively, CRISPR enzyme mRNA and guide RNA can be administered together. Advantageously, a second booster dose of guide RNA can be administered 1-12 hours (preferably around 2-6 hours) after the initial administration of CRISPR enzyme mRNA + guide RNA. Additional administrations of CRISPR enzyme mRNA and/or guide RNA might be useful to achieve the most efficient levels of genome modification.

[00149] In one aspect, the invention provides methods for using one or more elements of a CRISPR system. The CRISPR complex of the invention provides an effective means for modifying a target polynucleotide. The CRISPR complex of the invention has a wide variety of utility including modifying (e.g., deleting, inserting, translocating, inactivating, activating) a target polynucleotide in a multiplicity of cell types. As such the CRISPR complex of the invention has a broad spectrum of applications in, e.g., gene therapy, drug screening, disease diagnosis, and prognosis. An exemplary CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within the target polynucleotide. The guide sequence is linked to a tracr mate sequence, which in turn hybridizes to a tracr sequence. In one embodiment, this invention provides a method of cleaving a target polynucleotide. The method comprises modifying a target polynucleotide using a CRISPR complex that binds to the target polynucleotide and effect cleavage of said target polynucleotide. Typically, the CRISPR complex of the invention, when introduced into a cell, creates a break (e.g., a single or a double strand break) in the genome sequence. For example, the method can be used to cleave a disease gene in a cell. The break created by the CRISPR complex can be repaired by a repair processes such as the error prone non-homologous end joining (NHEJ) pathway or the high fidelity homology-directed repair (HDR). During these repair process, an exogenous polynucleotide template can be introduced into the genome sequence. In some methods, the HDR process is used modify genome sequence. For example, an exogenous polynucleotide template comprising a sequence to be integrated flanked by an upstream sequence and a downstream sequence is introduced into a cell. The upstream and downstream sequences share sequence similarity with either side of the site of integration in the chromosome. Where desired, a donor polynucleotide can be DNA, e.g., a DNA plasmid, a bacterial artificial chromosome (BAC), a yeast artificial chromosome (YAC), a viral vector, a linear piece of DNA, a PCR fragment, a naked nucleic acid, or a nucleic acid complexed with a delivery vehicle such as a liposome or poloxamer. The exogenous polynucleotide template comprises a sequence to be integrated (e.g., a mutated gene). The sequence for integration may be a sequence endogenous or exogenous to the cell. Examples of a sequence to be integrated include polynucleotides encoding a protein or a non-coding RNA (e.g., a microRNA). Thus, the sequence for integration may be operably linked to an appropriate control sequence or sequences. Alternatively, the sequence to be integrated may provide a regulatory function. The upstream and downstream

sequences in the exogenous polynucleotide template are selected to promote recombination between the chromosomal sequence of interest and the donor polynucleotide. The upstream sequence is a nucleic acid sequence that shares sequence similarity with the genome sequence upstream of the targeted site for integration. Similarly, the downstream sequence is a nucleic acid sequence that shares sequence similarity with the chromosomal sequence downstream of the targeted site of integration. The upstream and downstream sequences in the exogenous polynucleotide template can have 75%, 80%, 85%, 90%, 95%, or 100% sequence identity with the targeted genome sequence. Preferably, the upstream and downstream sequences in the exogenous polynucleotide template have about 95%, 96%, 97%, 98%, 99%, or 100% sequence identity with the targeted genome sequence. In some methods, the upstream and downstream sequences in the exogenous polynucleotide template have about 99% or 100% sequence identity with the targeted genome sequence. An upstream or downstream sequence may comprise from about 20 bp to about 2500 bp, for example, about 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, or 2500 bp. In some methods, the exemplary upstream or downstream sequence have about 200 bp to about 2000 bp, about 600 bp to about 1000 bp, or more particularly about 700 bp to about 1000 bp. In some methods, the exogenous polynucleotide template may further comprise a marker. Such a marker may make it easy to screen for targeted integrations. Examples of suitable markers include restriction sites, fluorescent proteins, or selectable markers. The exogenous polynucleotide template of the invention can be constructed using recombinant techniques (see, for example, Sambrook et al., 2001 and Ausubel et al., 1996). In a method for modifying a target polynucleotide by integrating an exogenous polynucleotide template, a double stranded break is introduced into the genome sequence by the CRISPR complex, the break is repaired via homologous recombination an exogenous polynucleotide template such that the template is integrated into the genome. The presence of a double-stranded break facilitates integration of the template. In other embodiments, this invention provides a method of modifying expression of a polynucleotide in a eukaryotic cell. The method comprises increasing or decreasing expression of a target polynucleotide by using a CRISPR complex that binds to the polynucleotide. In some methods, a target polynucleotide can be inactivated to effect the modification of the expression in a cell. For example, upon the binding of a CRISPR complex to a target sequence in a cell, the target polynucleotide is inactivated such that the

sequence is not transcribed, the coded protein is not produced, or the sequence does not function as the wild-type sequence does. For example, a protein or microRNA coding sequence may be inactivated such that the protein or microRNA or pre-microRNA transcript is not produced. In some methods, a control sequence can be inactivated such that it no longer functions as a control sequence. As used herein, "control sequence" refers to any nucleic acid sequence that effects the transcription, translation, or accessibility of a nucleic acid sequence. Examples of a control sequence include, a promoter, a transcription terminator, and an enhancer are control sequences. The target polynucleotide of a CRISPR complex can be any polynucleotide endogenous or exogenous to the eukaryotic cell. For example, the target polynucleotide can be a polynucleotide residing in the nucleus of the eukaryotic cell. The target polynucleotide can be a sequence coding a gene product (e.g., a protein) or a non-coding sequence (e.g., a regulatory polynucleotide or a junk DNA). Examples of target polynucleotides include a sequence associated with a signaling biochemical pathway, e.g., a signaling biochemical pathway-associated gene or polynucleotide. Examples of target polynucleotides include a disease associated gene or polynucleotide. A "disease-associated" gene or polynucleotide refers to any gene or polynucleotide which is yielding transcription or translation products at an abnormal level or in an abnormal form in cells derived from a disease-affected tissues compared with tissues or cells of a non disease control. It may be a gene that becomes expressed at an abnormally high level; it may be a gene that becomes expressed at an abnormally low level, where the altered expression correlates with the occurrence and/or progression of the disease. A disease-associated gene also refers to a gene possessing mutation(s) or genetic variation that is directly responsible or is in linkage disequilibrium with a gene(s) that is responsible for the etiology of a disease. The transcribed or translated products may be known or unknown, and may be at a normal or abnormal level. The target polynucleotide of a CRISPR complex can be any polynucleotide endogenous or exogenous to the eukaryotic cell. For example, the target polynucleotide can be a polynucleotide residing in the nucleus of the eukaryotic cell. The target polynucleotide can be a sequence coding a gene product (e.g., a protein) or a non-coding sequence (e.g., a regulatory polynucleotide or a junk DNA).

[00150] The target polynucleotide of a CRISPR complex can be any polynucleotide endogenous or exogenous to the eukaryotic cell. For example, the target polynucleotide can be a polynucleotide residing in the nucleus of the eukaryotic cell. The target polynucleotide can be a

sequence coding a gene product (e.g., a protein) or a non-coding sequence (e.g., a regulatory polynucleotide or a junk DNA). The target can be a control element or a regulatory element or a promoter or an enhancer or a silencer. The promoter may, in some embodiments, be in the region of +200bp or even +1000 bp from the TTS. In some embodiments, the regulatory region may be an enhancer. The enhancer is typically more than +1000 bp from the TTS. More in particular, expression of eukaryotic protein-coding genes generally is regulated through multiple cis-acting transcription-control regions. Some control elements are located close to the start site (promoter-proximal elements), whereas others lie more distant (enhancers and silencers) Promoters determine the site of transcription initiation and direct binding of RNA polymerase II. Three types of promoter sequences have been identified in eukaryotic DNA. The TATA box, the most common, is prevalent in rapidly transcribed genes. Initiator promoters infrequently are found in some genes, and CpG islands are characteristic of transcribed genes. Promoter-proximal elements occur within ≈ 200 base pairs of the start site. Several such elements, containing up to ≈ 20 base pairs, may help regulate a particular gene. Enhancers, which are usually ≈ 100 – 200 base pairs in length, contain multiple 8- to 20-bp control elements. They may be located from 200 base pairs to tens of kilobases upstream or downstream from a promoter, within an intron, or downstream from the final exon of a gene. Promoter-proximal elements and enhancers may be cell-type specific, functioning only in specific differentiated cell types. However, any of these regions can be the target sequence and are encompassed by the concept that the target can be a control element or a regulatory element or a promoter or an enhancer or a silencer.

[00151] Without wishing to be bound by theory, it is believed that the target sequence should be associated with a PAM (protospacer adjacent motif); that is, a short sequence recognized by the CRISPR complex. The precise sequence and length requirements for the PAM differ depending on the CRISPR enzyme used, but PAMs are typically 2-5 base pair sequences adjacent the protospacer (that is, the target sequence) Examples of PAM sequences are given in the examples section below, and the skilled person will be able to identify further PAM sequences for use with a given CRISPR enzyme. In some embodiments, the method comprises allowing a CRISPR complex to bind to the target polynucleotide to effect cleavage of said target polynucleotide thereby modifying the target polynucleotide, wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within said target polynucleotide, wherein said guide sequence is linked to a tracr mate sequence

which in turn hybridizes to a tracr sequence. In one aspect, the invention provides a method of modifying expression of a polynucleotide in a eukaryotic cell. In some embodiments, the method comprises allowing a CRISPR complex to bind to the polynucleotide such that said binding results in increased or decreased expression of said polynucleotide; wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence within said polynucleotide, wherein said guide sequence is linked to a tracr mate sequence which in turn hybridizes to a tracr sequence. Similar considerations and conditions apply as above for methods of modifying a target polynucleotide. In fact, these sampling, culturing and re-introduction options apply across the aspects of the present invention. In one aspect, the invention provides for methods of modifying a target polynucleotide in a eukaryotic cell, which may be *in vivo*, *ex vivo* or *in vitro*. In some embodiments, the method comprises sampling a cell or population of cells from a human or non-human animal, and modifying the cell or cells. Culturing may occur at any stage *ex vivo*. The cell or cells may even be re-introduced into the non-human animal or plant. For re-introduced cells it is particularly preferred that the cells are stem cells.

[00152] Indeed, in any aspect of the invention, the CRISPR complex may comprise a CRISPR enzyme complexed with a guide sequence hybridized to a target sequence, wherein said guide sequence may be linked to a tracr mate sequence which in turn may hybridize to a tracr sequence.

[00153] The invention relates to the engineering and optimization of systems, methods and compositions used for the control of gene expression involving sequence targeting, such as genome perturbation or gene-editing, that relate to the CRISPR-Cas9 system and components thereof. An advantage of the present methods is that the CRISPR system minimizes or avoids off-target binding and its resulting side effects. This is achieved using systems arranged to have a high degree of sequence specificity for the target DNA.

[00154] In relation to a CRISPR-Cas9 complex or system preferably, the tracr sequence has one or more hairpins and is 30 or more nucleotides in length, 40 or more nucleotides in length, or 50 or more nucleotides in length; the guide sequence is between 10 to 30 nucleotides in length, the CRISPR/Cas enzyme is a Type II Cas9 enzyme.

[00155] One guide with a first aptamer/RNA-binding protein pair can be linked or fused to an activator, whilst a second guide with a second aptamer/RNA-binding protein pair can be linked

or fused to a repressor. The guides are for different targets (loci), so this allows one gene to be activated and one repressed. For example, the following schematic shows such an approach:

Guide 1 – MS2 aptamer-----MS2 RNA-binding protein-----VP64 activator; and

Guide 2 – PP7 aptamer-----PP7 RNA-binding protein-----SID4x repressor.

[00156] The present invention also relates to orthogonal PP7/MS2 gene targeting. In this example, sgRNA targeting different loci are modified with distinct RNA loops in order to recruit MS2-VP64 or PP7-SID4X, which activate and repress their target loci, respectively. PP7 is the RNA-binding coat protein of the bacteriophage Pseudomonas. Like MS2, it binds a specific RNA sequence and secondary structure. The PP7 RNA-recognition motif is distinct from that of MS2. Consequently, PP7 and MS2 can be multiplexed to mediate distinct effects at different genomic loci simultaneously. For example, an sgRNA targeting locus A can be modified with MS2 loops, recruiting MS2-VP64 activators, while another sgRNA targeting locus B can be modified with PP7 loops, recruiting PP7-SID4X repressor domains. In the same cell, dCas9 can thus mediate orthogonal, locus-specific modifications. This principle can be extended to incorporate other orthogonal RNA-binding proteins such as Q-beta.

[00157] An alternative option for orthogonal repression includes incorporating non-coding RNA loops with transactive repressive function into the guide (either at similar positions to the MS2/PP7 loops integrated into the guide or at the 3' terminus of the guide). For instance, guides were designed with non-coding (but known to be repressive) RNA loops (e.g. using the Alu repressor (in RNA) that interferes with RNA polymerase II in mammalian cells). The Alu RNA sequence was located: in place of the MS2 RNA sequences as used herein (e.g. at tetraloop and/or stem loop 2); and/or at 3' terminus of the guide. This gives possible combinations of MS2, PP7 or Alu at the tetraloop and/or stemloop 2 positions, as well as, optionally, addition of Alu at the 3' end of the guide (with or without a linker).

[00158] The use of two different aptamers (each associated with a distinct RNA) allows an activator-adaptor protein fusion and a repressor-adaptor protein fusion to be used, with different guides, to activate expression of one gene, whilst repressing another. They, along with their different guides can be administered together, or substantially together, in a multiplexed approach. A large number of such modified guides can be used all at the same time, for example 10 or 20 or 30 and so forth, whilst only one (or at least a minimal number) of Cas9s to be delivered, as a comparatively small number of Cas9s can be used with a large number modified

guides. The adaptor protein may be associated (preferably linked or fused to) one or more activators or one or more repressors. For example, the adaptor protein may be associated with a first activator and a second activator. The first and second activators may be the same, but they are preferably different activators. For example, one might be VP64, whilst the other might be p65, although these are just examples and other transcriptional activators are envisaged. Three or more or even four or more activators (or repressors) may be used, but package size may limit the number being higher than 5 different functional domains. Linkers are preferably used, over a direct fusion to the adaptor protein, where two or more functional domains are associated with the adaptor protein. Suitable linkers might include the GlySer linker.

[00159] It is also envisaged that the enzyme-guide complex as a whole may be associated with two or more functional domains. For example, there may be two or more functional domains associated with the enzyme, or there may be two or more functional domains associated with the guide (via one or more adaptor proteins), or there may be one or more functional domains associated with the enzyme and one or more functional domains associated with the guide (via one or more adaptor proteins).

[00160] The fusion between the adaptor protein and the activator or repressor may include a linker. For example, GlySer linkers GGGGS can be used. They can be used in repeats of 3 ((GGGGS)₃) or 6, 9 or even 12 or more, to provide suitable lengths, as required. Linkers can be used between the RNA-binding protein and the functional domain (activator or repressor), or between the CRISPR Enzyme (Cas9) and the functional domain (activator or repressor). The linkers the user to engineer appropriate amounts of “mechanical flexibility”.

[00161] The invention comprehends a CRISPR Cas9 complex comprising a CRISPR enzyme and a guide RNA (sgRNA), wherein the CRISPR enzyme comprises at least one mutation, such that the CRISPR enzyme has no more than 5% of the nuclease activity of the CRISPR enzyme not having the at least one mutation and, optional, at least one or more nuclear localization sequences; the guide RNA (sgRNA) comprises a guide sequence capable of hybridizing to a target sequence in a genomic locus of interest in a cell; and wherein: the CRISPR enzyme is associated with two or more functional domains; or at least one loop of the sgRNA is modified by the insertion of distinct RNA sequence(s) that bind to one or more adaptor proteins, and wherein the adaptor protein is associated with two or more functional domains; or the CRISPR enzyme is associated with one or more functional domains and at least one loop of the sgRNA is

modified by the insertion of distinct RNA sequence(s) that bind to one or more adaptor proteins, and wherein the adaptor protein is associated with one or more functional domains.

[00162] In an embodiment, nucleic acid molecule(s) encoding a CRISPR-Cas9 or an ortholog or homolog thereof, may be codon-optimized for expression in a eukaryotic cell. A eukaryote can be as herein discussed. Nucleic acid molecule(s) can be engineered or non-naturally occurring.

[00163] In an embodiment, the CRISPR-Cas9 effector protein may comprise one or more mutations. The mutations may be artificially introduced mutations and may include but are not limited to one or more mutations in a catalytic domain, to provide a nickase, for example. Examples of catalytic domains with reference to a Cas9 enzyme may include but are not limited to RuvC I, RuvC II, RuvC III, and HNH domains.

[00164] In an embodiment, the CRISPR-Cas9 effector protein may be used as a generic nucleic acid binding protein with fusion to or being operably linked to a functional domain. Exemplary functional domains may include but are not limited to translational initiator, translational activator, translational repressor, nucleases, in particular ribonucleases, a spliceosome, beads, a light inducible/controllable domain or a chemically inducible/controllable domain.

[00165] In some embodiments, the CRISPR-Cas9 effector protein may have cleavage activity. In some embodiments, the Cas9 effector protein may direct cleavage of one or both nucleic acid strands at the location of or near a target sequence, such as within the target sequence and/or within the complement of the target sequence or at sequences associated with the target sequence. In some embodiments, the Cas9 effector protein may direct cleavage of one or both DNA or RNA strands within about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 50, 100, 200, 500, or more base pairs from the first or last nucleotide of a target sequence. In some embodiments, the cleavage may be blunt, i.e., generating blunt ends. In some embodiments, the cleavage may be staggered, i.e., generating sticky ends. In some embodiments, the cleavage may be a staggered cut with a 5' overhang, e.g., a 5' overhang of 1 to 5 nucleotides. In some embodiments, the cleavage may be a staggered cut with a 3' overhang, e.g., a 3' overhang of 1 to 5 nucleotides. In some embodiments, a vector encodes a nucleic acid-targeting Cas9 protein that may be mutated with respect to a corresponding wild-type enzyme such that the mutated nucleic acid-targeting Cas9 protein lacks the ability to cleave one or both DNA or RNA strands of a target

polynucleotide containing a target sequence. As a further example, two or more catalytic domains of Cas9 (RuvC I, RuvC II, and RuvC III or the HNH domain) may be mutated to produce a mutated Cas9 substantially lacking all RNA cleavage activity. As described herein, corresponding catalytic domains of a Cas9 effector protein may also be mutated to produce a mutated Cas9 lacking all DNA cleavage activity or having substantially reduced DNA cleavage activity. In some embodiments, a nucleic acid -targeting effector protein may be considered to substantially lack all RNA cleavage activity when the RNA cleavage activity of the mutated enzyme is about no more than 25%, 10%, 5%, 1%, 0.1%, 0.01%, or less of the nucleic acid cleavage activity of the non-mutated form of the enzyme; an example can be when the nucleic acid cleavage activity of the mutated form is nil or negligible as compared with the non-mutated form. An effector protein may be identified with reference to the general class of enzymes that share homology to the biggest nuclease with multiple nuclease domains from the Type II CRISPR system. Most preferably, the effector protein is a Type II protein such as Cas9. By derived, Applicants mean that the derived enzyme is largely based, in the sense of having a high degree of sequence homology with, a wildtype enzyme, but that it has been mutated (modified) in some way as known in the art or as described herein.

[00166] In certain embodiments, Cas9 may be constitutively present or inducibly present or conditionally present or administered or delivered. Cas9 optimization may be used to enhance function or to develop new functions, one can generate chimeric Cas9 proteins. And Cas9 may be used as a generic nucleic acid binding protein.

[00167] Typically, in the context of an endogenous nucleic acid-targeting system, formation of a nucleic acid-targeting complex (comprising a guide RNA hybridized to a target sequence and complexed with one or more nucleic acid-targeting effector proteins) results in cleavage of one or both DNA or RNA strands in or near (e.g., within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, or more base pairs from) the target sequence. As used herein the term “sequence(s) associated with a target locus of interest” refers to sequences near the vicinity of the target sequence (e.g. within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, or more base pairs from the target sequence, wherein the target sequence is comprised within a target locus of interest).

[00168] An example of a codon optimized sequence, is in this instance a sequence optimized for expression in a eukaryote, e.g., humans (i.e. being optimized for expression in humans), or for another eukaryote, animal or mammal as herein discussed; see, e.g., SaCas9 human codon

optimized sequence in WO 2014/093622 (PCT/US2013/074667) as an example of a codon optimized sequence (from knowledge in the art and this disclosure, codon optimizing coding nucleic acid molecule(s), especially as to effector protein (e.g., Cas9) is within the ambit of the skilled artisan). Whilst this is preferred, it will be appreciated that other examples are possible and codon optimization for a host species other than human, or for codon optimization for specific organs is known. In some embodiments, an enzyme coding sequence encoding a DNA-targeting Cas9 protein is codon optimized for expression in particular cells, such as eukaryotic cells. The eukaryotic cells may be those of or derived from a particular organism, such as a mammal, including but not limited to human, or non-human eukaryote or animal or mammal as herein discussed, e.g., mouse, rat, rabbit, dog, livestock, or non-human mammal or primate. In some embodiments, processes for modifying the germ line genetic identity of human beings and/or processes for modifying the genetic identity of animals which are likely to cause them suffering without any substantial medical benefit to man or animal, and also animals resulting from such processes, may be excluded. In general, codon optimization refers to a process of modifying a nucleic acid sequence for enhanced expression in the host cells of interest by replacing at least one codon (e.g., about or more than about 1, 2, 3, 4, 5, 10, 15, 20, 25, 50, or more codons) of the native sequence with codons that are more frequently or most frequently used in the genes of that host cell while maintaining the native amino acid sequence. Various species exhibit particular bias for certain codons of a particular amino acid. Codon bias (differences in codon usage between organisms) often correlates with the efficiency of translation of messenger RNA (mRNA), which is in turn believed to be dependent on, among other things, the properties of the codons being translated and the availability of particular transfer RNA (tRNA) molecules. The predominance of selected tRNAs in a cell is generally a reflection of the codons used most frequently in peptide synthesis. Accordingly, genes can be tailored for optimal gene expression in a given organism based on codon optimization. Codon usage tables are readily available, for example, at the "Codon Usage Database" available at www.kazusa.or.jp/codon/ and these tables can be adapted in a number of ways. See Nakamura, Y., et al. "Codon usage tabulated from the international DNA sequence databases: status for the year 2000" *Nucl. Acids Res.* 28:292 (2000). Computer algorithms for codon optimizing a particular sequence for expression in a particular host cell are also available, such as Gene Forge (Aptagen; Jacobus, PA), are also available. In some embodiments, one or more codons (e.g., 1,

2, 3, 4, 5, 10, 15, 20, 25, 50, or more, or all codons) in a sequence encoding a DNA -targeting Cas9 protein corresponds to the most frequently used codon for a particular amino acid.

[00169] In one aspect, the invention provides methods for using one or more elements of a nucleic acid-targeting system. The nucleic acid-targeting complex of the invention provides an effective means for modifying a target DNA (double stranded, linear or super-coiled). The nucleic acid-targeting complex of the invention has a wide variety of utility including modifying (e.g., deleting, inserting, translocating, inactivating, activating) a target DNA in a multiplicity of cell types. As such the nucleic acid-targeting complex of the invention has a broad spectrum of applications in, e.g., gene therapy, drug screening, disease diagnosis, and prognosis. An exemplary nucleic acid-targeting complex comprises a DNA targeting effector protein complexed with a guide RNA hybridized to a target sequence within the target locus of interest.

[00170] In some embodiments, the method may comprise allowing a nucleic acid-targeting complex to bind to the target DNA to effect cleavage of said target DNA thereby modifying the target DNA, wherein the nucleic acid-targeting complex comprises a nucleic acid-targeting effector protein complexed with a guide RNA hybridized to a target sequence within said target DNA. In one aspect, the invention provides a method of modifying expression of DNA in a eukaryotic cell. In some embodiments, the method comprises allowing a nucleic acid-targeting complex to bind to the DNA such that said binding results in increased or decreased expression of said DNA; wherein the nucleic acid-targeting complex comprises a nucleic acid-targeting effector protein complexed with a guide RNA. Similar considerations and conditions apply as above for methods of modifying a target DNA. In fact, these sampling, culturing and re-introduction options apply across the aspects of the present invention. In one aspect, the invention provides for methods of modifying a target DNA in a eukaryotic cell, which may be *in vivo*, *ex vivo* or *in vitro*. In some embodiments, the method comprises sampling a cell or population of cells from a human or non-human animal, and modifying the cell or cells. Culturing may occur at any stage *ex vivo*. The cell or cells may even be re-introduced into the non-human animal or plant. For re-introduced cells it is particularly preferred that the cells are stem cells.

[00171] Indeed, in any aspect of the invention, the nucleic acid-targeting complex may comprise a nucleic acid-targeting effector protein complexed with a guide RNA hybridized to a target sequence.

[00172] The invention relates to the engineering and optimization of systems, methods and compositions used for the control of gene expression involving DNA sequence targeting, that relate to the nucleic acid-targeting system and components thereof. An advantage of the present methods is that the CRISPR system minimizes or avoids off-target binding and its resulting side effects. This is achieved using systems arranged to have a high degree of sequence specificity for the target DNA.

[00173] In relation to a nucleic acid-targeting complex or system preferably, the tracr sequence has one or more hairpins and is 30 or more nucleotides in length, 40 or more nucleotides in length, or 50 or more nucleotides in length; the crRNA sequence is between 10 to 30 nucleotides in length, the nucleic acid-targeting effector protein is a Type II Cas9 effector protein.

Crystallization of CRISPR-Cas9 and Characterization of Crystal Structure

[00174] The crystals of the Cas9 can be obtained by techniques of protein crystallography, including batch, liquid bridge, dialysis, vapor diffusion and hanging drop methods. Generally, the crystals of the invention are grown by dissolving substantially pure CRISPR-Cas9 and a nucleic acid molecule to which it binds in an aqueous buffer containing a precipitant at a concentration just below that necessary to precipitate. Water is removed by controlled evaporation to produce precipitating conditions, which are maintained until crystal growth ceases. The crystal structure information is described in U.S. provisional applications 61/915,251 filed December 12, 2013, 61/930,214 filed on January 22, 2014, 61/980,012 filed April 15, 2014 and international application PCT/US2014/069925, filed December 12, 2014; and Nishimasu et al, "Crystal Structure of Cas9 in Complex with Guide RNA and Target DNA," Cell 156(5):935-949, DOI: <http://dx.doi.org/10.1016/j.cell.2014.02.001> (2014), each and all of which are incorporated herein by reference.

[00175] Uses of the Crystals, Crystal Structure and Atomic Structure Co-Ordinates: The crystals of the Cas9, and particularly the atomic structure co-ordinates obtained therefrom, have a wide variety of uses. The crystals and structure co-ordinates are particularly useful for identifying compounds (nucleic acid molecules) that bind to CRISPR-Cas9, and CRISPR-Cas9s that can bind to particular compounds (nucleic acid molecules). Thus, the structure co-ordinates described herein can be used as phasing models in determining the crystal structures of additional synthetic or mutated CRISPR-Cas9s, Cas9s, nickases, binding domains. The provision

of the crystal structure of CRISPR-Cas9 complexed with a nucleic acid molecule as applied in conjunction with the herein teachings provides the skilled artisan with a detailed insight into the mechanisms of action of CRISPR-Cas9. This insight provides a means to design modified CRISPR-Cas9s, such as by attaching thereto a functional group, such as a repressor or activator. While one can attach a functional group such as a repressor or activator to the N or C terminal of CRISPR-Cas9, the crystal structure demonstrates that the N terminal seems obscured or hidden, whereas the C terminal is more available for a functional group such as repressor or activator. Moreover, the crystal structure demonstrates that there is a flexible loop between approximately CRISPR-Cas9 (*S. pyogenes*) residues 534-676 which is suitable for attachment of a functional group such as an activator or repressor. Attachment can be via a linker, e.g., a flexible glycine-serine (GlyGlyGlySer) or (GGGS)₃ or a rigid alpha-helical linker such as (Ala(GluAlaAlaAlaLys)Ala). In addition to the flexible loop there is also a nuclease or H3 region, an H2 region and a helical region. By "helix" or "helical", is meant a helix as known in the art, including, but not limited to an alpha-helix. Additionally, the term helix or helical may also be used to indicate a c-terminal helical element with an N-terminal turn.

[00176] The provision of the crystal structure of CRISPR-Cas9 complexed with a nucleic acid molecule allows a novel approach for drug or compound discovery, identification, and design for compounds that can bind to CRISPR-Cas9 and thus the invention provides tools useful in diagnosis, treatment, or prevention of conditions or diseases of multicellular organisms, e.g., algae, plants, invertebrates, fish, amphibians, reptiles, avians, mammals; for example domesticated plants, animals (e.g., production animals such as swine, bovine, chicken; companion animal such as felines, canines, rodents (rabbit, gerbil, hamster); laboratory animals such as mouse, rat), and humans.

[00177] In any event, the determination of the three-dimensional structure of CRISPR-Cas9 (*S. pyogenes* Cas9) complex provides a basis for the design of new and specific nucleic acid molecules that bind to CRISPR-Cas9 (e.g., *S. pyogenes* Cas9), as well as the design of new CRISPR-Cas9 systems, such as by way of modification of the CRISPR-Cas9 system to bind to various nucleic acid molecules, by way of modification of the CRISPR-Cas9 system to have linked thereto to any one or more of various functional groups that may interact with each other, with the CRISPR-Cas9 (e.g., an inducible system that provides for self-activation and/or self-termination of function), with the nucleic acid molecule nucleic acid molecules (e.g., the

functional group may be a regulatory or functional domain which may be selected from the group consisting of a transcriptional repressor, a transcriptional activator, a nuclease domain, a DNA methyl transferase, a protein acetyltransferase, a protein deacetylase, a protein methyltransferase, a protein deaminase, a protein kinase, and a protein phosphatase; and, in some aspects, the functional domain is an epigenetic regulator; see, e.g., Zhang et al., US Patent No. 8,507,272, and it is again mentioned that it and all documents cited herein and all appln cited documents are hereby incorporated herein by reference), by way of modification of Cas9, by way of novel nickases). Indeed, the herewith CRISPR-Cas9 (*S. pyogenes* Cas9) crystal structure has a multitude of uses. For example, from knowing the three-dimensional structure of CRISPR-Cas9 (*S. pyogenes* Cas9) crystal structure, computer modelling programs may be used to design or identify different molecules expected to interact with possible or confirmed sites such as binding sites or other structural or functional features of the CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9). Compound that potentially bind (“binder”) can be examined through the use of computer modeling using a docking program. Docking programs are known; for example GRAM, DOCK or AUTODOCK (see Walters et al. Drug Discovery Today, vol. 3, no. 4 (1998), 160-178, and Dunbrack et al. Folding and Design 2 (1997), 27-42). This procedure can include computer fitting of potential binders ascertain how well the shape and the chemical structure of the potential binder will bind to a CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9). Computer-assisted, manual examination of the active site or binding site of a CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9) may be performed. Programs such as GRID (P. Goodford, J. Med. Chem, 1985, 28, 849-57)—a program that determines probable interaction sites between molecules with various functional groups—may also be used to analyze the active site or binding site to predict partial structures of binding compounds. Computer programs can be employed to estimate the attraction, repulsion or steric hindrance of the two binding partners, e.g., CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9) and a candidate nucleic acid molecule or a nucleic acid molecule and a candidate CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9), and the CRISPR-Cas9 crystal structure (*S. pyogenes* Cas9) herewith enables such methods. Generally, the tighter the fit, the fewer the steric hindrances, and the greater the attractive forces, the more potent the potential binder, since these properties are consistent with a tighter binding constant. Furthermore, the more specificity in the design of a candidate CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9), the more likely it is that it will not interact with off-target molecules as well. Also, “wet” methods are enabled by the

instant invention. For example, in an aspect, the invention provides for a method for determining the structure of a binder (e.g., target nucleic acid molecule) of a candidate CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9) bound to the candidate CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9), said method comprising, (a) providing a first crystal of a candidate CRISPR-Cas9 system (*S. pyogenes* Cas9) according to the invention or a second crystal of a candidate a candidate CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9), (b) contacting the first crystal or second crystal with said binder under conditions whereby a complex may form; and (c) determining the structure of said a candidate (e.g., CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9) or CRISPR-Cas9 system (*S. pyogenes* Cas9) complex. The second crystal may have essentially the same coordinates discussed herein, however due to minor alterations in CRISPR-Cas9 system (e.g., from the Cas9 of such a system being e.g., *S. pyogenes* Cas9 versus being *S. pyogenes* Cas9), wherein “e.g., *S. pyogenes* Cas9” indicates that the Cas9 is a Cas9 and can be of or derived from *S. pyogenes* or an ortholog thereof), the crystal may form in a different space group.

[00178] The invention further involves, in place of or in addition to “in silico” methods, other “wet” methods, including high throughput screening of a binder (e.g., target nucleic acid molecule) and a candidate CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9), or a candidate binder (e.g., target nucleic acid molecule) and a CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9), or a candidate binder (e.g., target nucleic acid molecule) and a candidate CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9) (the foregoing CRISPR-Cas9 system(s) with or without one or more functional group(s)), to select compounds with binding activity. Those pairs of binder and CRISPR-Cas9 system which show binding activity may be selected and further crystallized with the CRISPR-Cas9 crystal having a structure herein, e.g., by co-crystallization or by soaking, for X-ray analysis. The resulting X-ray structure may be compared with that of the Cas9 Crystal Structure for a variety of purposes, e.g., for areas of overlap. Having designed, identified, or selected possible pairs of binder and CRISPR-Cas9 system by determining those which have favorable fitting properties, e.g., predicted strong attraction based on the pairs of binder and CRISPR-Cas9 crystal structure data herein, these possible pairs can then be screened by “wet” methods for activity. Consequently, in an aspect the invention can involve: obtaining or synthesizing the possible pairs; and contacting a binder (e.g., target nucleic acid molecule) and a candidate CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9), or a candidate binder (e.g., target nucleic acid molecule) and a CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9), or a candidate

binder (e.g., target nucleic acid molecule) and a candidate CRISPR-Cas9 system (e.g., *S. pyogenes* Cas9) (the foregoing CRISPR-Cas9 system(s) with or without one or more functional group(s)) to determine ability to bind. In the latter step, the contacting is advantageously under conditions to determine function. Instead of, or in addition to, performing such an assay, the invention may comprise: obtaining or synthesizing complex(es) from said contacting and analyzing the complex(es), e.g., by X-ray diffraction or NMR or other means, to determine the ability to bind or interact. Detailed structural information can then be obtained about the binding, and in light of this information, adjustments can be made to the structure or functionality of a candidate CRISPR-Cas9 system or components thereof. These steps may be repeated and re-repeated as necessary. Alternatively or additionally, potential CRISPR-Cas9 systems from or in the foregoing methods can be with nucleic acid molecules in vivo, including without limitation by way of administration to an organism (including non-human animal and human) to ascertain or confirm function, including whether a desired outcome (e.g., reduction of symptoms, treatment) results therefrom.

[00179] The invention further involves a method of determining three dimensional structures of CRISPR-Cas9 systems or complex(es) of unknown structure by using the structural coordinates of the Cas9 Crystal Structure. For example, if X-ray crystallographic or NMR spectroscopic data are provided for a CRISPR-Cas9 system or complex of unknown crystal structure, the structure of a CRISPR-Cas9 complex may be used to interpret that data to provide a likely structure for the unknown system or complex by such techniques as by phase modeling in the case of X-ray crystallography. Thus, an inventive method can comprise: aligning a representation of the CRISPR-Cas9 system or complex having an unknown crystal structure with an analogous representation of the CRISPR-Cas9 system and complex of the crystal structure herein to match homologous or analogous regions (e.g., homologous or analogous sequences); modeling the structure of the matched homologous or analogous regions (e.g., sequences) of the CRISPR-Cas9 system or complex of unknown crystal structure based on the structure of the Cas9 Crystal Structure of the corresponding regions (e.g., sequences); and, determining a conformation (e.g. taking into consideration favorable interactions should be formed so that a low energy conformation is formed) for the unknown crystal structure which substantially preserves the structure of said matched homologous regions. "Homologous regions" describes, for example as to amino acids, amino acid residues in two sequences that are identical or have

similar, e.g., aliphatic, aromatic, polar, negatively charged, or positively charged, side-chain chemical groups. Homologous regions as to nucleic acid molecules can include at least 85% or 86% or 87% or 88% or 89% or 90% or 91% or 92% or 93% or 94% or 95% or 96% or 97% or 98% or 99% homology or identity. Identical and similar regions are sometimes described as being respectively “invariant” and “conserved” by those skilled in the art. Homology modeling is a technique that is well known to those skilled in the art (see, e.g., Greer, *Science* vol. 228 (1985) 1055, and Blundell et al. *Eur J Biochem* vol 172 (1988), 513). The computer representation of the conserved regions of the CRISPR-Cas9 crystal structure and those of a CRISPR-Cas9 system of unknown crystal structure aid in the prediction and determination of the crystal structure of the CRISPR-Cas9 system of unknown crystal structure.

[00180] Further still, the aspects of the invention which employ the CRISPR-Cas9 crystal structure in silico may be equally applied to new CRISPR-Cas9 crystal structures divined by using the herein-referenced CRISPR-Cas9 crystal structure. In this fashion, a library of CRISPR-Cas9 crystal structures can be obtained. Rational CRISPR-Cas9 system design is thus provided by the instant invention. For instance, having determined a conformation or crystal structure of a CRISPR-Cas9 system or complex, by the methods described herein, such a conformation may be used in a computer-based methods herein for determining the conformation or crystal structure of other CRISPR-Cas9 systems or complexes whose crystal structures are yet unknown. Data from all of these crystal structures can be in a database, and the herein methods can be more robust by having herein comparisons involving the herein crystal structure or portions thereof be with respect to one or more crystal structures in the library. The invention further provides systems, such as computer systems, intended to generate structures and/or perform rational design of a CRISPR-Cas9 system or complex. The system can contain: atomic co-ordinate data according to the herein-referenced Crystal Structure or be derived therefrom e.g., by modeling, said data defining the three-dimensional structure of a CRISPR-Cas9 system or complex or at least one domain or sub-domain thereof, or structure factor data therefor, said structure factor data being derivable from the atomic co-ordinate data of the herein-referenced Crystal Structure. The invention also involves computer readable media with: atomic co-ordinate data according to the herein-referenced Crystal Structure or derived therefrom e.g., by homology modeling, said data defining the three-dimensional structure of a CRISPR-Cas9 system or complex or at least one domain or sub-domain thereof, or structure factor data therefor, said

structure factor data being derivable from the atomic co-ordinate data of the herein-referenced Crystal Structure. "Computer readable media" refers to any media which can be read and accessed directly by a computer, and includes, but is not limited to: magnetic storage media; optical storage media; electrical storage media; cloud storage and hybrids of these categories. By providing such computer readable media, the atomic co-ordinate data can be routinely accessed for modeling or other "in silico" methods. The invention further comprehends methods of doing business by providing access to such computer readable media, for instance on a subscription basis, via the Internet or a global communication/computer network; or, the computer system can be available to a user, on a subscription basis. A "computer system" refers to the hardware means, software means and data storage means used to analyze the atomic co-ordinate data of the present invention. The minimum hardware means of computer-based systems of the invention may comprise a central processing unit (CPU), input means, output means, and data storage means. Desirably, a display or monitor is provided to visualize structure data. The invention further comprehends methods of transmitting information obtained in any method or step thereof described herein or any information described herein, e.g., via telecommunications, telephone, mass communications, mass media, presentations, internet, email, etc. The crystal structures of the invention can be analyzed to generate Fourier electron density map(s) of CRISPR-Cas9 systems or complexes; advantageously, the three-dimensional structure being as defined by the atomic co-ordinate data according to the herein-referenced Crystal Structure. Fourier electron density maps can be calculated based on X-ray diffraction patterns. These maps can then be used to determine aspects of binding or other interactions. Electron density maps can be calculated using known programs such as those from the CCP4 computer package (Collaborative Computing Project, No. 4. The CCP4 Suite: Programs for Protein Crystallography, Acta Crystallographica, D50, 1994, 760-763). For map visualization and model building programs such as "QUANTA" (1994, San Diego, Calif.: Molecular Simulations, Jones et al., Acta Crystallography A47 (1991), 110-119) can be used.

[00181] The herein-referenced Crystal Structure gives atomic co-ordinate data for a CRISPR-Cas9 (*S. pyogenes*), and lists each atom by a unique number; the chemical element and its position for each amino acid residue (as determined by electron density maps and antibody sequence comparisons), the amino acid residue in which the element is located, the chain identifier, the number of the residue, co-ordinates (e.g., X, Y, Z) which define with respect to the

crystallographic axes the atomic position (in angstroms) of the respective atom, the occupancy of the atom in the respective position, "B", isotropic displacement parameter (in angstroms²) which accounts for movement of the atom around its atomic center, and atomic number.

[00182] In particular embodiments of the invention, the conformational variations in the crystal structures of the CRISPR-Cas9 system or of components of the CRISPR-Cas9 provide important and critical information about the flexibility or movement of protein structure regions relative to nucleotide (RNA or DNA) structure regions that may be important for CRISPR-Cas9 system function. The structural information provided for Cas9 (e.g. *S. pyogenes* Cas9) as the CRISPR enzyme in the present application may be used to further engineer and optimize the CRISPR-Cas9 system and this may be extrapolated to interrogate structure-function relationships in other CRISPR enzyme systems as well. An aspect of the invention relates to the crystal structure of *S. pyogenes* Cas9 in complex with sgRNA and its target DNA at 2.4 Å resolution. The structure revealed a bilobed architecture composed of target recognition and nuclease lobes, accommodating a sgRNA:DNA duplex in a positively-charged groove at their interface. The recognition lobe is essential for sgRNA and DNA binding and the nuclease lobe contains the HNH and RuvC nuclease domains, which are properly positioned for the cleavage of complementary and non-complementary strands of the target DNA, respectively. This high-resolution structure and the functional analyses provided herein elucidate the molecular mechanism of RNA-guided DNA targeting by Cas9, and provides an abundance of information for generating optimized CRISPR-Cas9 systems and components thereof.

[00183] In particular embodiments of the invention, the crystal structure provides a critical step towards understanding the molecular mechanism of RNA-guided DNA targeting by Cas9. The structural and functional analyses herein provide a useful scaffold for rational engineering of Cas9-based genome modulating technologies and may provide guidance as to Cas9-mediated recognition of PAM sequences on the target DNA or mismatch tolerance between the sgRNA:DNA duplex. Aspects of the invention also relate to truncation mutants, e.g. an *S. pyogenes* Cas9 truncation mutant may facilitate packaging of Cas9 into size-constrained viral vectors for in vivo and therapeutic applications. Similarly, future engineering of the PAM Interacting (PI) domain may allow programing of PAM specificity, improve target site recognition fidelity, and increase the versatility of the Cas9 genome engineering platform.

[00184] The invention comprehends optimized functional CRISPR-Cas9 enzyme systems. In particular the CRISPR enzyme comprises one or more mutations that converts it to a DNA binding protein to which functional domains exhibiting a function of interest may be recruited or appended or inserted or attached. In certain embodiments, the CRISPR enzyme comprises one or more mutations which include but are not limited to D10A, E762A, H840A, N854A, N863A or D986A (based on the amino acid position numbering of a *S. pyogenes* Cas9) and/or the one or more mutations is in a RuvC1 or HNH domain of the CRISPR enzyme or is a mutation as otherwise as discussed herein. In some embodiments, the CRISPR enzyme has one or more mutations in a catalytic domain, wherein when transcribed, the tracr mate sequence hybridizes to the tracr sequence and the guide sequence directs sequence-specific binding of a CRISPR complex to the target sequence, and wherein the enzyme further comprises a functional domain.

[00185] The structural information provided herein allows for interrogation of sgRNA (or chimeric RNA) interaction with the target DNA and the CRISPR enzyme (e.g. Cas9) permitting engineering or alteration of sgRNA structure to optimize functionality of the entire CRISPR-Cas9 system. For example, loops of the sgRNA may be extended, without colliding with the Cas9 protein by the insertion of distinct RNA loop(s) or distinct sequence(s) that may recruit adaptor proteins that can bind to the distinct RNA loop(s) or distinct sequence(s). The adaptor proteins may include but are not limited to orthogonal RNA-binding protein / aptamer combinations that exist within the diversity of bacteriophage coat proteins. A list of such coat proteins includes, but is not limited to: Q β , F2, GA, fr, JP501, M12, R17, BZ13, JP34, JP500, KU1, M11, MX1, TW18, VK, SP, FI, ID2, NL95, TW19, AP205, ϕ Cb5, ϕ Cb8r, ϕ Cb12r, ϕ Cb23r, 7s and PRR1. These adaptor proteins or orthogonal RNA binding proteins can further recruit effector proteins or fusions which comprise one or more functional domains. In some embodiments, the functional domain may be selected from the group consisting of: transposase domain, integrase domain, recombinase domain, resolvase domain, invertase domain, protease domain, DNA methyltransferase domain, DNA hydroxymethylase domain, DNA demethylase domain, histone acetylase domain, histone deacetylases domain, nuclease domain, repressor domain, activator domain, nuclear-localization signal domains, transcription-regulatory protein (or transcription complex recruiting) domain, cellular uptake activity associated domain, nucleic acid binding domain, antibody presentation domain, histone modifying enzymes, recruiter of histone modifying enzymes; inhibitor of histone modifying enzymes, histone methyltransferase,

histone demethylase, histone kinase, histone phosphatase, histone ribosylase, histone deribosylase, histone ubiquitinase, histone deubiquitinase, histone biotinase and histone tail protease.

[00186] In some preferred embodiments, the functional domain is a transcriptional activation domain, preferably VP64. In some embodiments, the functional domain is a transcription repression domain, preferably KRAB. In some embodiments, the transcription repression domain is SID, or concatemers of SID (eg SID4X). In some embodiments, the functional domain is an epigenetic modifying domain, such that an epigenetic modifying enzyme is provided. In some embodiments, the functional domain is an activation domain, which may be the P65 activation domain.

[00187] In one aspect surveyor analysis is used for identification of indel activity / nuclease activity. In general survey analysis includes extraction of genomic DNA, PCR amplification of the genomic region flanking the CRISPR target site, purification of products, re-annealing to enable heteroduplex formation. After re-annealing, products are treated with SURVEYOR nuclease and SURVEYOR enhancer S (Transgenomics) following the manufacturer's recommended protocol. Analysis may be performed with poly-acrylamide gels according to known methods. Quantification may be based on relative band intensities.

Delivery generally

Gene Editing or Altering a Target Loci with Cas9

[00188] The double strand break or single strand break in one of the strands advantageously should be sufficiently close to target position such that correction occurs. In an embodiment, the distance is not more than 50, 100, 200, 300, 350 or 400 nucleotides. While not wishing to be bound by theory, it is believed that the break should be sufficiently close to target position such that the break is within the region that is subject to exonuclease-mediated removal during end resection. If the distance between the target position and a break is too great, the mutation may not be included in the end resection and, therefore, may not be corrected, as the template nucleic acid sequence may only be used to correct sequence within the end resection region.

[00189] In an embodiment, in which a guide RNA and a Type II molecule, in particular Cas9 or an ortholog or homolog thereof, preferably a Cas9 nuclease induce a double strand break for the purpose of inducing HDR-mediated correction, the cleavage site is between 0-200 bp (e.g., 0 to 175, 0 to 150, 0 to 125, 0 to 100, 0 to 75, 0 to 50, 0 to 25, 25 to 200, 25 to 175, 25 to 150, 25

to 125, 25 to 100, 25 to 75, 25 to 50, 50 to 200, 50 to 175, 50 to 150, 50 to 125, 50 to 100, 50 to 75, 75 to 200, 75 to 175, 75 to 150, 75 to 125, 75 to 100 bp) away from the target position. In an embodiment, the cleavage site is between 0- 100 bp (e.g., 0 to 75, 0 to 50, 0 to 25, 25 to 100, 25 to 75, 25 to 50, 50 to 100, 50 to 75 or 75 to 100 bp) away from the target position. In a further embodiment, two or more guide RNAs complexing with Cas9 or an ortholog or homolog thereof, may be used to induce multiplexed breaks for purpose of inducing HDR-mediated correction.

[00190] The homology arm should extend at least as far as the region in which end resection may occur, e.g., in order to allow the resected single stranded overhang to find a complementary region within the donor template. The overall length could be limited by parameters such as plasmid size or viral packaging limits. In an embodiment, a homology arm may not extend into repeated elements. Exemplary homology arm lengths include a least 50, 100, 250, 500, 750 or 1000 nucleotides.

[00191] Target position, as used herein, refers to a site on a target nucleic acid or target gene (e.g., the chromosome) that is modified by a Type II, in particular Cas9 or an ortholog or homolog thereof, preferably Cas9 molecule-dependent process. For example, the target position can be a modified Cas9 molecule cleavage of the target nucleic acid and template nucleic acid directed modification, e.g., correction, of the target position. In an embodiment, a target position can be a site between two nucleotides, e.g., adjacent nucleotides, on the target nucleic acid into which one or more nucleotides is added. The target position may comprise one or more nucleotides that are altered, e.g., corrected, by a template nucleic acid. In an embodiment, the target position is within a target sequence (e.g., the sequence to which the guide RNA binds). In an embodiment, a target position is upstream or downstream of a target sequence (e.g., the sequence to which the guide RNA binds).

[00192] A template nucleic acid, as that term is used herein, refers to a nucleic acid sequence which can be used in conjunction with a Type II molecule, in particular Cas9 or an ortholog or homolog thereof, preferably a Cas9 molecule and a guide RNA molecule to alter the structure of a target position. In an embodiment, the target nucleic acid is modified to have some or all of the sequence of the template nucleic acid, typically at or near cleavage site(s). In an embodiment, the template nucleic acid is single stranded. In an alternate embodiment, the template nucleic acid is double stranded. In an embodiment, the template nucleic acid is DNA, e.g., double stranded DNA. In an alternate embodiment, the template nucleic acid is single stranded DNA.

[00193] In an embodiment, the template nucleic acid alters the structure of the target position by participating in homologous recombination. In an embodiment, the template nucleic acid alters the sequence of the target position. In an embodiment, the template nucleic acid results in the incorporation of a modified, or non-naturally occurring base into the target nucleic acid.

[00194] The template sequence may undergo a breakage mediated or catalyzed recombination with the target sequence. In an embodiment, the template nucleic acid may include sequence that corresponds to a site on the target sequence that is cleaved by a Cas9 mediated cleavage event. In an embodiment, the template nucleic acid may include sequence that corresponds to both, a first site on the target sequence that is cleaved in a first Cas9 mediated event, and a second site on the target sequence that is cleaved in a second Cas9 mediated event.

[00195] In certain embodiments, the template nucleic acid can include sequence which results in an alteration in the coding sequence of a translated sequence, e.g., one which results in the substitution of one amino acid for another in a protein product, e.g., transforming a mutant allele into a wild type allele, transforming a wild type allele into a mutant allele, and/or introducing a stop codon, insertion of an amino acid residue, deletion of an amino acid residue, or a nonsense mutation. In certain embodiments, the template nucleic acid can include sequence which results in an alteration in a non-coding sequence, e.g., an alteration in an exon or in a 5' or 3' non-translated or non-transcribed region. Such alterations include an alteration in a control element, e.g., a promoter, enhancer, and an alteration in a cis-acting or trans-acting control element.

[00196] A template nucleic acid having homology with a target position in a target gene may be used to alter the structure of a target sequence. The template sequence may be used to alter an unwanted structure, e.g., an unwanted or mutant nucleotide. The template nucleic acid may include sequence which, when integrated, results in: decreasing the activity of a positive control element; increasing the activity of a positive control element; decreasing the activity of a negative control element; increasing the activity of a negative control element; decreasing the expression of a gene; increasing the expression of a gene; increasing resistance to a disorder or disease; increasing resistance to viral entry; correcting a mutation or altering an unwanted amino acid residue conferring, increasing, abolishing or decreasing a biological property of a gene product, e.g., increasing the enzymatic activity of an enzyme, or increasing the ability of a gene product to interact with another molecule.

[00197] The template nucleic acid may include sequence which results in: a change in sequence of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 or more nucleotides of the target sequence. In an embodiment, the template nucleic acid may be 20 \pm 10, 30 \pm 10, 40 \pm 10, 50 \pm 10, 60 \pm 10, 70 \pm 10, 80 \pm 10, 90 \pm 10, 100 \pm 10, 110 \pm 10, 120 \pm 10, 130 \pm 10, 140 \pm 10, 150 \pm 10, 160 \pm 10, 170 \pm 10, 180 \pm 10, 190 \pm 10, 200 \pm 10, 210 \pm 10, of 220 \pm 10 nucleotides in length. In an embodiment, the template nucleic acid may be 30 \pm 20, 40 \pm 20, 50 \pm 20, 60 \pm 20, 70 \pm 20, 80 \pm 20, 90 \pm 20, 100 \pm 20, 110 \pm 20, 120 \pm 20, 130 \pm 20, 140 \pm 20, 150 \pm 20, 160 \pm 20, 170 \pm 20, 180 \pm 20, 190 \pm 20, 200 \pm 20, 210 \pm 20, of 220 \pm 20 nucleotides in length. In an embodiment, the template nucleic acid is 10 to 1,000, 20 to 900, 30 to 800, 40 to 700, 50 to 600, 50 to 500, 50 to 400, 50 to 300, 50 to 200, or 50 to 100 nucleotides in length.

[00198] A template nucleic acid comprises the following components: [5' homology arm]-[replacement sequence]-[3' homology arm]. The homology arms provide for recombination into the chromosome, thus replacing the undesired element, e.g., a mutation or signature, with the replacement sequence. In an embodiment, the homology arms flank the most distal cleavage sites. In an embodiment, the 3' end of the 5' homology arm is the position next to the 5' end of the replacement sequence. In an embodiment, the 5' homology arm can extend at least 10, 20, 30, 40, 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, or 2000 nucleotides 5' from the 5' end of the replacement sequence. In an embodiment, the 5' end of the 3' homology arm is the position next to the 3' end of the replacement sequence. In an embodiment, the 3' homology arm can extend at least 10, 20, 30, 40, 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1500, or 2000 nucleotides 3' from the 3' end of the replacement sequence.

[00199] In certain embodiments, one or both homology arms may be shortened to avoid including certain sequence repeat elements. For example, a 5' homology arm may be shortened to avoid a sequence repeat element. In other embodiments, a 3' homology arm may be shortened to avoid a sequence repeat element. In some embodiments, both the 5' and the 3' homology arms may be shortened to avoid including certain sequence repeat elements.

[00200] In certain embodiments, a template nucleic acid for correcting a mutation may be designed for use as a single-stranded oligonucleotide. When using a single-stranded oligonucleotide, 5' and 3' homology arms may range up to about 200 base pairs (bp) in length, e.g., at least 25, 50, 75, 100, 125, 150, 175, or 200 bp in length.

Cas9 Effector Protein Complex System Promoted Non-Homologous End-Joining

[00201] In certain embodiments, nuclease-induced non-homologous end-joining (NHEJ) can be used to target gene-specific knockouts. Nuclease-induced NHEJ can also be used to remove (e.g., delete) sequence in a gene of interest. Generally, NHEJ repairs a double-strand break in the DNA by joining together the two ends; however, generally, the original sequence is restored only if two compatible ends, exactly as they were formed by the double-strand break, are perfectly ligated. The DNA ends of the double-strand break are frequently the subject of enzymatic processing, resulting in the addition or removal of nucleotides, at one or both strands, prior to rejoining of the ends. This results in the presence of insertion and/or deletion (indel) mutations in the DNA sequence at the site of the NHEJ repair. Two-thirds of these mutations typically alter the reading frame and, therefore, produce a non-functional protein. Additionally, mutations that maintain the reading frame, but which insert or delete a significant amount of sequence, can destroy functionality of the protein. This is locus dependent as mutations in critical functional domains are likely less tolerable than mutations in non-critical regions of the protein. The indel mutations generated by NHEJ are unpredictable in nature; however, at a given break site certain indel sequences are favored and are over represented in the population, likely due to small regions of microhomology. The lengths of deletions can vary widely; most commonly in the 1-50 bp range, but they can easily be greater than 50 bp, e.g., they can easily reach greater than about 100-200 bp. Insertions tend to be shorter and often include short duplications of the sequence immediately surrounding the break site. However, it is possible to obtain large insertions, and in these cases, the inserted sequence has often been traced to other regions of the genome or to plasmid DNA present in the cells.

[00202] Because NHEJ is a mutagenic process, it may also be used to delete small sequence motifs as long as the generation of a specific final sequence is not required. If a double-strand break is targeted near to a short target sequence, the deletion mutations caused by the NHEJ repair often span, and therefore remove, the unwanted nucleotides. For the deletion of larger DNA segments, introducing two double-strand breaks, one on each side of the sequence, can result in NHEJ between the ends with removal of the entire intervening sequence. Both of these approaches can be used to delete specific DNA sequences; however, the error-prone nature of NHEJ may still produce indel mutations at the site of repair.

[00203] Both double strand cleaving Type II molecule, in particular Cas9 or an ortholog or homolog thereof, preferably Cas9 molecules and single strand, or nickase, Type II molecule, in particular Cas9 or an ortholog or homolog thereof, preferably Cas9 molecules can be used in the methods and compositions described herein to generate NHEJ- mediated indels. NHEJ-mediated indels targeted to the gene, e.g., a coding region, e.g., an early coding region of a gene of interest can be used to knockout (i.e., eliminate expression of) a gene of interest. For example, early coding region of a gene of interest includes sequence immediately following a transcription start site, within a first exon of the coding sequence, or within 500 bp of the transcription start site (e.g., less than 500, 450, 400, 350, 300, 250, 200, 150, 100 or 50 bp).

[00204] In an embodiment, in which a guide RNA and Type II molecule, in particular Cas9 or an ortholog or homolog thereof, preferably Cas9 nuclease generate a double strand break for the purpose of inducing NHEJ-mediated indels, a guide RNA may be configured to position one double-strand break in close proximity to a nucleotide of the target position. In an embodiment, the cleavage site may be between 0-500 bp away from the target position (e.g., less than 500, 400, 300, 200, 100, 50, 40, 30, 25, 20, 15, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 bp from the target position).

[00205] In an embodiment, in which two guide RNAs complexing with Type II molecules, in particular Cas9 or an ortholog or homolog thereof, preferably Cas9 nickases induce two single strand breaks for the purpose of inducing NHEJ-mediated indels, two guide RNAs may be configured to position two single-strand breaks to provide for NHEJ repair a nucleotide of the target position.

Cas9 Effector Protein Complexes Can Deliver Functional Effectors

[00206] Unlike CRISPR-Cas-mediated gene knockout, which permanently eliminates expression by mutating the gene at the DNA level, CRISPR-Cas9 knockdown allows for temporary reduction of gene expression through the use of artificial transcription factors. Mutating key residues in both DNA cleavage domains of the Cas9 protein results in the generation of a catalytically inactive Cas9. A catalytically inactive Cas9 complexes with a guide RNA and localizes to the DNA sequence specified by that guide RNA's targeting domain, however, it does not cleave the target DNA. Fusion of the inactive Cas9 protein to an effector domain, e.g., a transcription repression domain, enables recruitment of the effector to any DNA site specified by the guide RNA. In certain embodiments, Cas9 may be fused to a transcriptional

repression domain and recruited to the promoter region of a gene. Especially for gene repression, it is contemplated herein that blocking the binding site of an endogenous transcription factor would aid in downregulating gene expression. In another embodiment, an inactive Cas9 can be fused to a chromatin modifying protein. Altering chromatin status can result in decreased expression of the target gene.

[00207] In an embodiment, a guide RNA molecule can be targeted to a known transcription response elements (e.g., promoters, enhancers, etc.), a known upstream activating sequences, and/or sequences of unknown or known function that are suspected of being able to control expression of the target DNA.

[00208] In some methods, a target polynucleotide can be inactivated to effect the modification of the expression in a cell. For example, upon the binding of a CRISPR complex to a target sequence in a cell, the target polynucleotide is inactivated such that the sequence is not transcribed, the coded protein is not produced, or the sequence does not function as the wild-type sequence does. For example, a protein or microRNA coding sequence may be inactivated such that the protein is not produced.

[00209] In certain embodiments, the CRISPR enzyme comprises one or more mutations selected from the group consisting of D917A, E1006A and D1225A and/or the one or more mutations is in a RuvC domain of the CRISPR enzyme or is a mutation as otherwise as discussed herein. In some embodiments, the CRISPR enzyme has one or more mutations in a catalytic domain, wherein when transcribed, the direct repeat sequence forms a single stem loop and the guide sequence directs sequence-specific binding of a CRISPR complex to the target sequence, and wherein the enzyme further comprises a functional domain. In some embodiments, the functional domain is a transcriptional activation domain, preferably VP64. In some embodiments, the functional domain is a transcription repression domain, preferably KRAB. In some embodiments, the transcription repression domain is SID, or concatemers of SID (eg SID4X). In some embodiments, the functional domain is an epigenetic modifying domain, such that an epigenetic modifying enzyme is provided. In some embodiments, the functional domain is an activation domain, which may be the P65 activation domain.

Delivery of the CRISPR-Cas9 Complex or Components Thereof

[00210] Through this disclosure and the knowledge in the art, TALEs, CRISPR-Cas9 system, specifically the novel CRISPR systems described herein, or components thereof or nucleic acid

molecules thereof (including, for instance HDR template) or nucleic acid molecules encoding or providing components thereof may be delivered by a delivery system herein described both generally and in detail.

[00211] Vector delivery, e.g., plasmid, viral delivery: The CRISPR enzyme, for instance a Cas9, and/or any of the present RNAs, for instance a guide RNA, can be delivered using any suitable vector, e.g., plasmid or viral vectors, such as adeno associated virus (AAV), lentivirus, adenovirus or other viral vector types, or combinations thereof. Cas9 and one or more guide RNAs can be packaged into one or more vectors, e.g., plasmid or viral vectors. In some embodiments, the vector, e.g., plasmid or viral vector is delivered to the tissue of interest by, for example, an intramuscular injection, while other times the delivery is via intravenous, transdermal, intranasal, oral, mucosal, or other delivery methods. Such delivery may be either via a single dose, or multiple doses. One skilled in the art understands that the actual dosage to be delivered herein may vary greatly depending upon a variety of factors, such as the vector choice, the target cell, organism, or tissue, the general condition of the subject to be treated, the degree of transformation/modification sought, the administration route, the administration mode, the type of transformation/modification sought, etc.

[00212] Such a dosage may further contain, for example, a carrier (water, saline, ethanol, glycerol, lactose, sucrose, calcium phosphate, gelatin, dextran, agar, pectin, peanut oil, sesame oil, etc.), a diluent, a pharmaceutically-acceptable carrier (e.g., phosphate-buffered saline), a pharmaceutically-acceptable excipient, and/or other compounds known in the art. The dosage may further contain one or more pharmaceutically acceptable salts such as, for example, a mineral acid salt such as a hydrochloride, a hydrobromide, a phosphate, a sulfate, etc.; and the salts of organic acids such as acetates, propionates, malonates, benzoates, etc. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, gels or gelling materials, flavorings, colorants, microspheres, polymers, suspension agents, etc. may also be present herein. In addition, one or more other conventional pharmaceutical ingredients, such as preservatives, humectants, suspending agents, surfactants, antioxidants, anticaking agents, fillers, chelating agents, coating agents, chemical stabilizers, etc. may also be present, especially if the dosage form is a reconstitutable form. Suitable exemplary ingredients include microcrystalline cellulose, carboxymethylcellulose sodium, polysorbate 80, phenylethyl alcohol, chlorobutanol, potassium sorbate, sorbic acid, sulfur dioxide, propyl gallate, the parabens, ethyl

vanillin, glycerin, phenol, parachlorophenol, gelatin, albumin and a combination thereof. A thorough discussion of pharmaceutically acceptable excipients is available in REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., N.J. 1991) which is incorporated by reference herein.

[00213] In an embodiment herein the delivery is via an adenovirus, which may be at a single booster dose containing at least 1×10^5 particles (also referred to as particle units, pu) of adenoviral vector. In an embodiment herein, the dose preferably is at least about 1×10^6 particles (for example, about 1×10^6 - 1×10^{12} particles), more preferably at least about 1×10^7 particles, more preferably at least about 1×10^8 particles (e.g., about 1×10^8 - 1×10^{11} particles or about 1×10^8 - 1×10^{12} particles), and most preferably at least about 1×10^9 particles (e.g., about 1×10^9 - 1×10^{10} particles or about 1×10^9 - 1×10^{12} particles), or even at least about 1×10^{10} particles (e.g., about 1×10^{10} - 1×10^{12} particles) of the adenoviral vector. Alternatively, the dose comprises no more than about 1×10^{14} particles, preferably no more than about 1×10^{13} particles, even more preferably no more than about 1×10^{12} particles, even more preferably no more than about 1×10^{11} particles, and most preferably no more than about 1×10^{10} particles (e.g., no more than about 1×10^9 articles). Thus, the dose may contain a single dose of adenoviral vector with, for example, about 1×10^6 particle units (pu), about 2×10^6 pu, about 4×10^6 pu, about 1×10^7 pu, about 2×10^7 pu, about 4×10^7 pu, about 1×10^8 pu, about 2×10^8 pu, about 4×10^8 pu, about 1×10^9 pu, about 2×10^9 pu, about 4×10^9 pu, about 1×10^{10} pu, about 2×10^{10} pu, about 4×10^{10} pu, about 1×10^{11} pu, about 2×10^{11} pu, about 4×10^{11} pu, about 1×10^{12} pu, about 2×10^{12} pu, or about 4×10^{12} pu of adenoviral vector. See, for example, the adenoviral vectors in U.S. Patent No. 8,454,972 B2 to Nabel, et. al., granted on June 4, 2013; incorporated by reference herein, and the dosages at col 29, lines 36-58 thereof. In an embodiment herein, the adenovirus is delivered via multiple doses.

[00214] In an embodiment herein, the delivery is via an AAV. A therapeutically effective dosage for in vivo delivery of the AAV to a human is believed to be in the range of from about 20 to about 50 ml of saline solution containing from about 1×10^{10} to about 1×10^{10} functional AAV/ml solution. The dosage may be adjusted to balance the therapeutic benefit against any side effects. In an embodiment herein, the AAV dose is generally in the range of concentrations of from about 1×10^5 to 1×10^{50} genomes AAV, from about 1×10^8 to 1×10^{20} genomes AAV, from about 1×10^{10} to about 1×10^{16} genomes, or about 1×10^{11} to about 1×10^{16} genomes

AAV. A human dosage may be about 1×10^{13} genomes AAV. Such concentrations may be delivered in from about 0.001 ml to about 100 ml, about 0.05 to about 50 ml, or about 10 to about 25 ml of a carrier solution. Other effective dosages can be readily established by one of ordinary skill in the art through routine trials establishing dose response curves. See, for example, U.S. Patent No. 8,404,658 B2 to Hajjar, et al., granted on March 26, 2013, at col. 27, lines 45-60.

[00215] In an embodiment herein the delivery is via a plasmid. In such plasmid compositions, the dosage should be a sufficient amount of plasmid to elicit a response. For instance, suitable quantities of plasmid DNA in plasmid compositions can be from about 0.1 to about 2 mg, or from about 1 μ g to about 10 μ g per 70 kg individual. Plasmids of the invention will generally comprise (i) a promoter; (ii) a sequence encoding a CRISPR enzyme, operably linked to said promoter; (iii) a selectable marker; (iv) an origin of replication; and (v) a transcription terminator downstream of and operably linked to (ii). The plasmid can also encode the RNA components of a CRISPR complex, but one or more of these may instead be encoded on a different vector.

[00216] The doses herein are based on an average 70 kg individual. The frequency of administration is within the ambit of the medical or veterinary practitioner (e.g., physician, veterinarian), or scientist skilled in the art. It is also noted that mice used in experiments are typically about 20g and from mice experiments one can scale up to a 70 kg individual.

[00217] In some embodiments the RNA molecules of the invention are delivered in liposome or lipofectin formulations and the like and can be prepared by methods well known to those skilled in the art. Such methods are described, for example, in U.S. Pat. Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference. Delivery systems aimed specifically at the enhanced and improved delivery of siRNA into mammalian cells have been developed, (see, for example, Shen et al FEBS Let. 2003, 539:111-114; Xia et al., Nat. Biotech. 2002, 20:1006-1010; Reich et al., Mol. Vision. 2003, 9: 210-216; Sorensen et al., J. Mol. Biol. 2003, 327: 761-766; Lewis et al., Nat. Gen. 2002, 32: 107-108 and Simeoni et al., NAR 2003, 31, 11: 2717-2724) and may be applied to the present invention. siRNA has recently been successfully used for inhibition of gene expression in primates (see for example. Tolentino et al., Retina 24(4):660 which may also be applied to the present invention.

[00218] Indeed, RNA delivery is a useful method of *in vivo* delivery. It is possible to deliver Cas9 and gRNA (and, for instance, HR repair template) into cells using liposomes or

particles/nanoparticles. Thus delivery of the CRISPR enzyme, such as a Cas9 and/or delivery of the RNAs of the invention may be in RNA form and via microvesicles, liposomes or particles/nanoparticles. For example, Cas9 mRNA and gRNA can be packaged into liposomal particles for delivery *in vivo*. Liposomal transfection reagents such as lipofectamine from Life Technologies and other reagents on the market can effectively deliver RNA molecules into the liver.

[00219] Means of delivery of RNA also preferred include delivery of RNA via particles/nanoparticles (Cho, S., Goldberg, M., Son, S., Xu, Q., Yang, F., Mei, Y., Bogatyrev, S., Langer, R. and Anderson, D., Lipid-like nanoparticles for small interfering RNA delivery to endothelial cells, *Advanced Functional Materials*, 19: 3112-3118, 2010) or exosomes (Schroeder, A., Levins, C., Cortez, C., Langer, R., and Anderson, D., Lipid-based nanotherapeutics for siRNA delivery, *Journal of Internal Medicine*, 267: 9-21, 2010, PMID: 20059641). Indeed, exosomes have been shown to be particularly useful in delivery siRNA, a system with some parallels to the CRISPR system. For instance, El-Andaloussi S, et al. ("Exosome-mediated delivery of siRNA *in vitro* and *in vivo*." *Nat Protoc.* 2012 Dec;7(12):2112-26. doi: 10.1038/nprot.2012.131. Epub 2012 Nov 15.) describe how exosomes are promising tools for drug delivery across different biological barriers and can be harnessed for delivery of siRNA *in vitro* and *in vivo*. Their approach is to generate targeted exosomes through transfection of an expression vector, comprising an exosomal protein fused with a peptide ligand. The exosomes are then purified and characterized from transfected cell supernatant, then RNA is loaded into the exosomes. Delivery or administration according to the invention can be performed with exosomes, in particular but not limited to the brain. Vitamin E (α -tocopherol) may be conjugated with CRISPR-Cas9 and delivered to the brain along with high density lipoprotein (HDL), for example in a similar manner as was done by Uno et al. (*HUMAN GENE THERAPY* 22:711–719 (June 2011)) for delivering short-interfering RNA (siRNA) to the brain. Mice were infused via Osmotic minipumps (model 1007D; Alzet, Cupertino, CA) filled with phosphate-buffered saline (PBS) or free TocsiBACE or Toc-siBACE/HDL and connected with Brain Infusion Kit 3 (Alzet). A brain-infusion cannula was placed about 0.5mm posterior to the bregma at midline for infusion into the dorsal third ventricle. Uno et al. found that as little as 3 nmol of Toc-siRNA with HDL could induce a target reduction in comparable degree by the same ICV infusion method. A similar dosage of CRISPR Cas9 conjugated to α -tocopherol and co-administered with

HDL targeted to the brain may be contemplated for humans in the present invention, for example, about 3 nmol to about 3 μ mol of CRISPR Cas9 targeted to the brain may be contemplated. Zou et al. ((HUMAN GENE THERAPY 22:465-475 (April 2011)) describes a method of lentiviral-mediated delivery of short-hairpin RNAs targeting PKC γ for in vivo gene silencing in the spinal cord of rats. Zou et al. administered about 10 μ l of a recombinant lentivirus having a titer of 1×10^9 transducing units (TU)/ml by an intrathecal catheter. A similar dosage of CRISPR Cas9 expressed in a lentiviral vector targeted to the brain may be contemplated for humans in the present invention, for example, about 10-50 ml of CRISPR Cas9 targeted to the brain in a lentivirus having a titer of 1×10^9 transducing units (TU)/ml may be contemplated.

[00220] In terms of local delivery to the brain, this can be achieved in various ways. For instance, material can be delivered intrastrially e.g. by injection. Injection can be performed stereotactically via a craniotomy.

[00221] Enhancing NHEJ or HR efficiency is also helpful for delivery. It is preferred that NHEJ efficiency is enhanced by co-expressing end-processing enzymes such as Trex2 (Dumitrache et al. Genetics. 2011 August; 188(4): 787-797). It is preferred that HR efficiency is increased by transiently inhibiting NHEJ machineries such as Ku70 and Ku86. HR efficiency can also be increased by co-expressing prokaryotic or eukaryotic homologous recombination enzymes such as RecBCD, RecA.

Packaging and Promoters generally

[00222] Ways to package Cas9 coding nucleic acid molecules, e.g., DNA, into vectors, e.g., viral vectors, to mediate genome modification *in vivo* include:

To achieve NHEJ-mediated gene knockout:

Single virus vector:

Vector containing two or more expression cassettes:

Promoter-Cas9 coding nucleic acid molecule -terminator

Promoter-guide RNA1-terminator

Promoter-guide RNA2-terminator

Promoter-guide RNA(N)-terminator (up to size limit of vector)

Double virus vector:

Vector 1 containing one expression cassette for driving the expression of Cas9

Promoter-Cas9 coding nucleic acid molecule-terminator

Vector 2 containing one more expression cassettes for driving the expression of one or more guide RNAs

Promoter-guide RNA1-terminator

Promoter-guide RNA(N)-terminator (up to size limit of vector)

To mediate homology-directed repair.

In addition to the single and double virus vector approaches described above, an additional vector is used to deliver a homology-direct repair template.

[00223] The promoter used to drive Cas9 coding nucleic acid molecule expression can include:

AAV ITR can serve as a promoter: this is advantageous for eliminating the need for an additional promoter element (which can take up space in the vector). The additional space freed up can be used to drive the expression of additional elements (gRNA, etc.). Also, ITR activity is relatively weaker, so can be used to reduce potential toxicity due to over expression of Cas9.

For ubiquitous expression, can use promoters: CMV, CAG, CBh, PGK, SV40, Ferritin heavy or light chains, etc.

For brain or other CNS expression, can use promoters: SynapsinI for all neurons, CaMKIIalpha for excitatory neurons, GAD67 or GAD65 or VGAT for GABAergic neurons, etc.

For liver expression, can use Albumin promoter.

For lung expression, can use SP-B.

For endothelial cells, can use ICAM.

For hematopoietic cells can use IFNbeta or CD45.

For Osteoblasts can use OG-2.

[00224] The promoter used to drive guide RNA can include:

Pol III promoters such as U6 or H1

Use of Pol II promoter and intronic cassettes to express guide RNA

Adeno associated virus (AAV)

[00225] Cas9 and one or more guide RNA can be delivered using adeno associated virus (AAV), lentivirus, adenovirus or other plasmid or viral vector types, in particular, using formulations and doses from, for example, US Patents Nos. 8,454,972 (formulations, doses for

adenovirus), 8,404,658 (formulations, doses for AAV) and 5,846,946 (formulations, doses for DNA plasmids) and from clinical trials and publications regarding the clinical trials involving lentivirus, AAV and adenovirus. For examples, for AAV, the route of administration, formulation and dose can be as in US Patent No. 8,454,972 and as in clinical trials involving AAV. For Adenovirus, the route of administration, formulation and dose can be as in US Patent No. 8,404,658 and as in clinical trials involving adenovirus. For plasmid delivery, the route of administration, formulation and dose can be as in US Patent No 5,846,946 and as in clinical studies involving plasmids. Doses may be based on or extrapolated to an average 70 kg individual (e.g. a male adult human), and can be adjusted for patients, subjects, mammals of different weight and species. Frequency of administration is within the ambit of the medical or veterinary practitioner (e.g., physician, veterinarian), depending on usual factors including the age, sex, general health, other conditions of the patient or subject and the particular condition or symptoms being addressed. The viral vectors can be injected into the tissue of interest. For cell-type specific genome modification, the expression of Cas9 can be driven by a cell-type specific promoter. For example, liver-specific expression might use the Albumin promoter and neuron-specific expression (e.g. for targeting CNS disorders) might use the Synapsin I promoter.

[00226] In terms of *in vivo* delivery, AAV is advantageous over other viral vectors for a couple of reasons:

- Low toxicity (this may be due to the purification method not requiring ultra centrifugation of cell particles that can activate the immune response)
- Low probability of causing insertional mutagenesis because it doesn't integrate into the host genome.

[00227] AAV has a packaging limit of 4.5 or 4.75 Kb. This means that Cas9 as well as a promoter and transcription terminator have to be all fit into the same viral vector. Constructs larger than 4.5 or 4.75 Kb will lead to significantly reduced virus production. SpCas9 is quite large, the gene itself is over 4.1 Kb, which makes it difficult for packing into AAV. Therefore embodiments of the invention include utilizing homologs of Cas9 that are shorter. For example:

Species	Cas9 Size
Corynebacter diphtheriae	3252
Eubacterium ventriosum	3321
Streptococcus pasteurianus	3390
Lactobacillus farciminis	3378

Sphaerochaeta globus	3537
Azospirillum B510	3504
Gluconacetobacter diazotrophicus	3150
Neisseria cinerea	3246
Roseburia intestinalis	3420
Parvibaculum lavamentivorans	3111
Staphylococcus aureus	3159
Nitratifactor salsuginis DSM 16511	3396
Campylobacter lari CF89-12	3009
Streptococcus thermophilus LMD-9	3396

[00228] These species are therefore, in general, preferred Cas9 species with respect to both AAV delivery and in general.

[00229] As to AAV, the AAV can be AAV1, AAV2, AAV5 or any combination thereof. One can select the AAV of the AAV with regard to the cells to be targeted; e.g., one can select AAV serotypes 1, 2, 5 or a hybrid capsid AAV1, AAV2, AAV5 or any combination thereof for targeting brain or neuronal cells; and one can select AAV4 for targeting cardiac tissue. AAV8 is useful for delivery to the liver. The herein promoters and vectors are preferred individually. A tabulation of certain AAV serotypes as to these cells (see Grimm, D. et al, J. Virol. 82: 5887-5911 (2008)) is as follows:

Cell Line	AAV-1	AAV-2	AAV-3	AAV-4	AAV-5	AAV-6	AAV-8	AAV-9
Huh-7	13	100	2.5	0.0	0.1	10	0.7	0.0
HEK293	25	100	2.5	0.1	0.1	5	0.7	0.1
HeLa	3	100	2.0	0.1	6.7	1	0.2	0.1
HepG2	3	100	16.7	0.3	1.7	5	0.3	ND
Hep1A	20	100	0.2	1.0	0.1	1	0.2	0.0
911	17	100	11	0.2	0.1	17	0.1	ND
CHO	100	100	14	1.4	333	50	10	1.0
COS	33	100	33	3.3	5.0	14	2.0	0.5
MeWo	10	100	20	0.3	6.7	10	1.0	0.2
NIH3T3	10	100	2.9	2.9	0.3	10	0.3	ND
A549	14	100	20	ND	0.5	10	0.5	0.1
HT1180	20	100	10	0.1	0.3	33	0.5	0.1
Monocytes	1111	100	ND	ND	125	1429	ND	ND
Immature DC	2500	100	ND	ND	222	2857	ND	ND
Mature DC	2222	100	ND	ND	333	3333	ND	ND

Lentivirus

[00230] Lentiviruses are complex retroviruses that have the ability to infect and express their genes in both mitotic and post-mitotic cells. The most commonly known lentivirus is the human immunodeficiency virus (HIV), which uses the envelope glycoproteins of other viruses to target a broad range of cell types.

[00231] Lentiviruses may be prepared as follows. After cloning pCasES10 (which contains a lentiviral transfer plasmid backbone), HEK293FT at low passage (p=5) were seeded in a T-75 flask to 50% confluence the day before transfection in DMEM with 10% fetal bovine serum and without antibiotics. After 20 hours, media was changed to OptiMEM (serum-free) media and transfection was done 4 hours later. Cells were transfected with 10 µg of lentiviral transfer plasmid (pCasES10) and the following packaging plasmids: 5 µg of pMD2.G (VSV-g pseudotype), and 7.5ug of psPAX2 (gag/pol/rev/tat). Transfection was done in 4mL OptiMEM with a cationic lipid delivery agent (50uL Lipofectamine 2000 and 100ul Plus reagent). After 6 hours, the media was changed to antibiotic-free DMEM with 10% fetal bovine serum. These methods use serum during cell culture, but serum-free methods are preferred.

[00232] Lentivirus may be purified as follows. Viral supernatants were harvested after 48 hours. Supernatants were first cleared of debris and filtered through a 0.45µm low protein binding (PVDF) filter. They were then spun in a ultracentrifuge for 2 hours at 24,000 rpm. Viral pellets were resuspended in 50ul of DMEM overnight at 4C. They were then aliquotted and immediately frozen at -80°C.

[00233] In another embodiment, minimal non-primate lentiviral vectors based on the equine infectious anemia virus (EIAV) are also contemplated, especially for ocular gene therapy (see, e.g., Balagaan, *J Gene Med* 2006; 8: 275 – 285). In another embodiment, RetinoStat®, an equine infectious anemia virus-based lentiviral gene therapy vector that expresses angiostatic proteins endostatin and angiostatin that is delivered via a subretinal injection for the treatment of the web form of age-related macular degeneration is also contemplated (see, e.g., Binley et al., *HUMAN GENE THERAPY* 23:980–991 (September 2012)) and this vector may be modified for the CRISPR-Cas9 system of the present invention.

[00234] In another embodiment, self-inactivating lentiviral vectors with an siRNA targeting a common exon shared by HIV tat/rev, a nucleolar-localizing TAR decoy, and an anti-CCR5-specific hammerhead ribozyme (see, e.g., DiGiusto et al. (2010) *Sci Transl Med* 2:36ra43) may

be used/and or adapted to the CRISPR-Cas9 system of the present invention. A minimum of 2.5×10^6 CD34+ cells per kilogram patient weight may be collected and prestimulated for 16 to 20 hours in X-VIVO 15 medium (Lonza) containing 2 $\mu\text{mol/L}$ -glutamine, stem cell factor (100 ng/ml), Flt-3 ligand (Flt-3L) (100 ng/ml), and thrombopoietin (10 ng/ml) (CellGenix) at a density of 2×10^6 cells/ml. Prestimulated cells may be transduced with lentiviral at a multiplicity of infection of 5 for 16 to 24 hours in 75-cm² tissue culture flasks coated with fibronectin (25 mg/cm²) (RetroNectin, Takara Bio Inc.).

[00235] Lentiviral vectors have been disclosed as in the treatment for Parkinson's Disease, see, e.g., US Patent Publication No. 20120295960 and US Patent Nos. 7303910 and 7351585. Lentiviral vectors have also been disclosed for the treatment of ocular diseases, see e.g., US Patent Publication Nos. 20060281180, 20090007284, US20110117189; US20090017543; US20070054961, US20100317109. Lentiviral vectors have also been disclosed for delivery to the brain, see, e.g., US Patent Publication Nos. US20110293571; US20110293571, US20040013648, US20070025970, US20090111106 and US Patent No. US7259015.

RNA delivery

[00236] RNA delivery: The CRISPR enzyme, for instance a Cas9, and/or any of the present RNAs, for instance a guide RNA, can also be delivered in the form of RNA. Cas9 mRNA can be generated using *in vitro* transcription. For example, Cas9 mRNA can be synthesized using a PCR cassette containing the following elements: T7_promoter-kozak sequence (GCCACC)-Cas9-3' UTR from beta globin-polyA tail (a string of 120 or more adenines). The cassette can be used for transcription by T7 polymerase. Guide RNAs can also be transcribed using *in vitro* transcription from a cassette containing T7_promoter-GG-guide RNA sequence.

[00237] To enhance expression and reduce possible toxicity, the CRISPR enzyme-coding sequence and/or the guide RNA can be modified to include one or more modified nucleoside e.g. using pseudo-U or 5-Methyl-C.

[00238] mRNA delivery methods are especially promising for liver delivery currently.

[00239] Much clinical work on RNA delivery has focused on RNAi or antisense, but these systems can be adapted for delivery of RNA for implementing the present invention. References below to RNAi *etc.* should be read accordingly.

Particle delivery systems and/or formulations:

[00240] Several types of particle delivery systems and/or formulations are known to be useful in a diverse spectrum of biomedical applications. In general, a particle is defined as a small object that behaves as a whole unit with respect to its transport and properties. Particles are further classified according to diameter. Coarse particles cover a range between 2,500 and 10,000 nanometers. Fine particles are sized between 100 and 2,500 nanometers. Ultrafine particles, or nanoparticles, are generally between 1 and 100 nanometers in size. The basis of the 100-nm limit is the fact that novel properties that differentiate particles from the bulk material typically develop at a critical length scale of under 100 nm.

[00241] As used herein, a particle delivery system/formulation is defined as any biological delivery system/formulation which includes a particle in accordance with the present invention. A particle in accordance with the present invention is any entity having a greatest dimension (e.g. diameter) of less than 100 microns (μm). In some embodiments, inventive particles have a greatest dimension of less than 10 μm . In some embodiments, inventive particles have a greatest dimension of less than 2000 nanometers (nm). In some embodiments, inventive particles have a greatest dimension of less than 1000 nanometers (nm). In some embodiments, inventive particles have a greatest dimension of less than 900 nm, 800 nm, 700 nm, 600 nm, 500 nm, 400 nm, 300 nm, 200 nm, or 100 nm. Typically, inventive particles have a greatest dimension (e.g., diameter) of 500 nm or less. In some embodiments, inventive particles have a greatest dimension (e.g., diameter) of 250 nm or less. In some embodiments, inventive particles have a greatest dimension (e.g., diameter) of 200 nm or less. In some embodiments, inventive particles have a greatest dimension (e.g., diameter) of 150 nm or less. In some embodiments, inventive particles have a greatest dimension (e.g., diameter) of 100 nm or less. Smaller particles, e.g., having a greatest dimension of 50 nm or less are used in some embodiments of the invention. In some embodiments, inventive particles have a greatest dimension ranging between 25 nm and 200 nm.

[00242] Particle characterization (including e.g., characterizing morphology, dimension, etc.) is done using a variety of different techniques. Common techniques are electron microscopy (TEM, SEM), atomic force microscopy (AFM), dynamic light scattering (DLS), X-ray photoelectron spectroscopy (XPS), powder X-ray diffraction (XRD), Fourier transform infrared spectroscopy (FTIR), matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF), ultraviolet-visible spectroscopy, dual polarisation interferometry and nuclear magnetic resonance (NMR). Characterization (dimension measurements) may be

made as to native particles (i.e., preloading) or after loading of the cargo (herein cargo refers to e.g., one or more components of CRISPR-Cas9 system e.g., CRISPR enzyme or mRNA or guide RNA, or any combination thereof, and may include additional carriers and/or excipients) to provide particles of an optimal size for delivery for any *in vitro*, *ex vivo* and/or *in vivo* application of the present invention. In certain preferred embodiments, particle dimension (e.g., diameter) characterization is based on measurements using dynamic laser scattering (DLS). Mention is made of US Patent No. 8,709,843; US Patent No. 6,007,845; US Patent No. 5,855,913; US Patent No. 5,985,309; US Patent No. 5,543,158; and the publication by James E. Dahlman and Carmen Barnes et al. Nature Nanotechnology (2014) published online 11 May 2014, doi:10.1038/nnano.2014.84, concerning particles, methods of making and using them and measurements thereof.

[00243] Particles delivery systems within the scope of the present invention may be provided in any form, including but not limited to solid, semi-solid, emulsion, or colloidal particles. As such any of the delivery systems described herein, including but not limited to, e.g., lipid-based systems, liposomes, micelles, microvesicles, exosomes, or gene gun may be provided as particle delivery systems within the scope of the present invention.

Particles

[00244] CRISPR enzyme mRNA and guide RNA may be delivered simultaneously using particles or lipid envelopes; for instance, CRISPR enzyme and RNA of the invention, e.g., as a complex, can be delivered via a particle as in Dahlman et al., WO2015089419 A2 and documents cited therein, such as 7C1 (see, e.g., James E. Dahlman and Carmen Barnes et al. Nature Nanotechnology (2014) published online 11 May 2014, doi:10.1038/nnano.2014.84), e.g., delivery particle comprising lipid or lipidoid and hydrophilic polymer, e.g., cationic lipid and hydrophilic polymer, for instance wherein the cationic lipid comprises 1,2-dioleoyl-3-trimethylammonium-propane (DOTAP) or 1,2-ditetradecanoyl-*sn*-glycero-3-phosphocholine (DMPC) and/or wherein the hydrophilic polymer comprises ethylene glycol or polyethylene glycol (PEG); and/or wherein the particle further comprises cholesterol (e.g., particle from formulation 1 = DOTAP 100, DMPC 0, PEG 0, Cholesterol 0; formulation number 2 = DOTAP 90, DMPC 0, PEG 10, Cholesterol 0; formulation number 3 = DOTAP 90, DMPC 0, PEG 5, Cholesterol 5), wherein particles are formed using an efficient, multistep process wherein first, effector protein and RNA are mixed together, e.g., at a 1:1 molar ratio, e.g., at room temperature,

e.g., for 30 minutes, e.g., in sterile, nuclease free 1X PBS; and separately, DOTAP, DMPC, PEG, and cholesterol as applicable for the formulation are dissolved in alcohol, e.g., 100% ethanol; and, the two solutions are mixed together to form particles containing the complexes).

[00245] Nucleic acid-targeting effector proteins (such as a Type II protein such as Cas9) mRNA and guide RNA may be delivered simultaneously using particles or lipid envelopes. For example, Su X, Fricke J, Kavanagh DG, Irvine DJ (“In vitro and in vivo mRNA delivery using lipid-enveloped pH-responsive polymer nanoparticles” *Mol Pharm*. 2011 Jun 6;8(3):774-87. doi: 10.1021/mp100390w. Epub 2011 Apr 1) describes biodegradable core-shell structured nanoparticles with a poly(β -amino ester) (PBAE) core enveloped by a phospholipid bilayer shell. These were developed for in vivo mRNA delivery. The pH-responsive PBAE component was chosen to promote endosome disruption, while the lipid surface layer was selected to minimize toxicity of the polycation core. Such are, therefore, preferred for delivering RNA of the present invention.

[00246] In one embodiment, particles based on self assembling bioadhesive polymers are contemplated, which may be applied to oral delivery of peptides, intravenous delivery of peptides and nasal delivery of peptides, all to the brain. Other embodiments, such as oral absorption and ocular delivery of hydrophobic drugs are also contemplated. The molecular envelope technology involves an engineered polymer envelope which is protected and delivered to the site of the disease (see, e.g., Mazza, M. et al. *ACSNano*, 2013. 7(2): 1016-1026; Siew, A., et al. *Mol Pharm*, 2012. 9(1):14-28; Lalatsa, A., et al. *J Contr Rel*, 2012. 161(2):523-36; Lalatsa, A., et al., *Mol Pharm*, 2012. 9(6):1665-80; Lalatsa, A., et al. *Mol Pharm*, 2012. 9(6):1764-74; Garrett, N.L., et al. *J Biophotonics*, 2012. 5(5-6):458-68; Garrett, N.L., et al. *J Raman Spect*, 2012. 43(5):681-688; Ahmad, S., et al. *J Royal Soc Interface* 2010. 7:S423-33; Uchegbu, I.F. *Expert Opin Drug Deliv*, 2006. 3(5):629-40; Qu, X., et al. *Biomacromolecules*, 2006. 7(12):3452-9 and Uchegbu, I.F., et al. *Int J Pharm*, 2001. 224:185-199). Doses of about 5 mg/kg are contemplated, with single or multiple doses, depending on the target tissue.

[00247] In one embodiment, particles that can deliver RNA to a cancer cell to stop tumor growth developed by Dan Anderson’s lab at MIT may be used/and or adapted to the CRISPR Cas9 system of the present invention. In particular, the Anderson lab developed fully automated, combinatorial systems for the synthesis, purification, characterization, and formulation of new biomaterials and nanoformulations. See, e.g., Alabi et al., *Proc Natl Acad Sci U S A*. 2013 Aug

6;110(32):12881-6; Zhang et al., *Adv Mater.* 2013 Sep 6;25(33):4641-5; Jiang et al., *Nano Lett.* 2013 Mar 13;13(3):1059-64; Karagiannis et al., *ACS Nano.* 2012 Oct 23;6(10):8484-7; Whitehead et al., *ACS Nano.* 2012 Aug 28;6(8):6922-9 and Lee et al., *Nat Nanotechnol.* 2012 Jun 3;7(6):389-93.

[00248] US patent application 20110293703 relates to lipidoid compounds are also particularly useful in the administration of polynucleotides, which may be applied to deliver the CRISPR-Cas9 system of the present invention. In one aspect, the aminoalcohol lipidoid compounds are combined with an agent to be delivered to a cell or a subject to form microparticles, nanoparticles, liposomes, or micelles. The agent to be delivered by the particles, liposomes, or micelles may be in the form of a gas, liquid, or solid, and the agent may be a polynucleotide, protein, peptide, or small molecule. The aminoalcohol lipidoid compounds may be combined with other aminoalcohol lipidoid compounds, polymers (synthetic or natural), surfactants, cholesterol, carbohydrates, proteins, lipids, etc. to form the particles. These particles may then optionally be combined with a pharmaceutical excipient to form a pharmaceutical composition.

[00249] US Patent Publication No. 20110293703 also provides methods of preparing the aminoalcohol lipidoid compounds. One or more equivalents of an amine are allowed to react with one or more equivalents of an epoxide-terminated compound under suitable conditions to form an aminoalcohol lipidoid compound of the present invention. In certain embodiments, all the amino groups of the amine are fully reacted with the epoxide-terminated compound to form tertiary amines. In other embodiments, all the amino groups of the amine are not fully reacted with the epoxide-terminated compound to form tertiary amines thereby resulting in primary or secondary amines in the aminoalcohol lipidoid compound. These primary or secondary amines are left as is or may be reacted with another electrophile such as a different epoxide-terminated compound. As will be appreciated by one skilled in the art, reacting an amine with less than excess of epoxide-terminated compound will result in a plurality of different aminoalcohol lipidoid compounds with various numbers of tails. Certain amines may be fully functionalized with two epoxide-derived compound tails while other molecules will not be completely functionalized with epoxide-derived compound tails. For example, a diamine or polyamine may include one, two, three, or four epoxide-derived compound tails off the various amino moieties of the molecule resulting in primary, secondary, and tertiary amines. In certain embodiments, all

the amino groups are not fully functionalized. In certain embodiments, two of the same types of epoxide-terminated compounds are used. In other embodiments, two or more different epoxide-terminated compounds are used. The synthesis of the aminoalcohol lipidoid compounds is performed with or without solvent, and the synthesis may be performed at higher temperatures ranging from 30-100 °C., preferably at approximately 50-90 °C. The prepared aminoalcohol lipidoid compounds may be optionally purified. For example, the mixture of aminoalcohol lipidoid compounds may be purified to yield an aminoalcohol lipidoid compound with a particular number of epoxide-derived compound tails. Or the mixture may be purified to yield a particular stereo- or regioisomer. The aminoalcohol lipidoid compounds may also be alkylated using an alkyl halide (e.g., methyl iodide) or other alkylating agent, and/or they may be acylated.

[00250] US Patent Publication No. 20110293703 also provides libraries of aminoalcohol lipidoid compounds prepared by the inventive methods. These aminoalcohol lipidoid compounds may be prepared and/or screened using high-throughput techniques involving liquid handlers, robots, microtiter plates, computers, etc. In certain embodiments, the aminoalcohol lipidoid compounds are screened for their ability to transfect polynucleotides or other agents (e.g., proteins, peptides, small molecules) into the cell.

[00251] US Patent Publication No. 20130302401 relates to a class of poly(beta-amino alcohols) (PBAAAs) has been prepared using combinatorial polymerization. The inventive PBAAAs may be used in biotechnology and biomedical applications as coatings (such as coatings of films or multilayer films for medical devices or implants), additives, materials, excipients, non-biofouling agents, micropatterning agents, and cellular encapsulation agents. When used as surface coatings, these PBAAAs elicited different levels of inflammation, both in vitro and in vivo, depending on their chemical structures. The large chemical diversity of this class of materials allowed us to identify polymer coatings that inhibit macrophage activation in vitro. Furthermore, these coatings reduce the recruitment of inflammatory cells, and reduce fibrosis, following the subcutaneous implantation of carboxylated polystyrene microparticles. These polymers may be used to form polyelectrolyte complex capsules for cell encapsulation. The invention may also have many other biological applications such as antimicrobial coatings, DNA or siRNA delivery, and stem cell tissue engineering. The teachings of US Patent Publication No. 20130302401 may be applied to the CRISPR Cas9 system of the present invention.

[00252] In another embodiment, lipid nanoparticles (LNPs) are contemplated. An antitransthyretin small interfering RNA has been encapsulated in lipid nanoparticles and delivered to humans (see, e.g., Coelho et al., *N Engl J Med* 2013;369:819-29), and such a system may be adapted and applied to the CRISPR Cas9 system of the present invention. Doses of about 0.01 to about 1 mg per kg of body weight administered intravenously are contemplated. Medications to reduce the risk of infusion-related reactions are contemplated, such as dexamethasone, acetaminophen, diphenhydramine or cetirizine, and ranitidine are contemplated. Multiple doses of about 0.3 mg per kilogram every 4 weeks for five doses are also contemplated.

[00253] LNPs have been shown to be highly effective in delivering siRNAs to the liver (see, e.g., Taberero et al., *Cancer Discovery*, April 2013, Vol. 3, No. 4, pages 363-470) and are therefore contemplated for delivering RNA encoding CRISPR Cas9 to the liver. A dosage of about four doses of 6 mg/kg of the LNP every two weeks may be contemplated. Taberero et al. demonstrated that tumor regression was observed after the first 2 cycles of LNPs dosed at 0.7 mg/kg, and by the end of 6 cycles the patient had achieved a partial response with complete regression of the lymph node metastasis and substantial shrinkage of the liver tumors. A complete response was obtained after 40 doses in this patient, who has remained in remission and completed treatment after receiving doses over 26 months. Two patients with RCC and extrahepatic sites of disease including kidney, lung, and lymph nodes that were progressing following prior therapy with VEGF pathway inhibitors had stable disease at all sites for approximately 8 to 12 months, and a patient with PNET and liver metastases continued on the extension study for 18 months (36 doses) with stable disease.

[00254] However, the charge of the LNP must be taken into consideration. As cationic lipids combined with negatively charged lipids to induce nonbilayer structures that facilitate intracellular delivery. Because charged LNPs are rapidly cleared from circulation following intravenous injection, ionizable cationic lipids with pKa values below 7 were developed (see, e.g., Rosin et al, *Molecular Therapy*, vol. 19, no. 12, pages 1286-2200, Dec. 2011). Negatively charged polymers such as RNA may be loaded into LNPs at low pH values (e.g., pH 4) where the ionizable lipids display a positive charge. However, at physiological pH values, the LNPs exhibit a low surface charge compatible with longer circulation times. Four species of ionizable cationic lipids have been focused upon, namely 1,2-dilinoyleoyl-3-dimethylammonium-propane (DLinDAP), 1,2-dilinoyleoxy-3-N,N-dimethylaminopropane (DLinDMA), 1,2-dilinoyleoxy-

keto-N,N-dimethyl-3-aminopropane (DLinKDMA), and 1,2-dilinoleyl-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLinKC2-DMA). It has been shown that LNP siRNA systems containing these lipids exhibit remarkably different gene silencing properties in hepatocytes *in vivo*, with potencies varying according to the series DLinKC2-DMA>DLinKDMA>DLinDMA>>DLinDAP employing a Factor VII gene silencing model (see, e.g., Rosin et al, *Molecular Therapy*, vol. 19, no. 12, pages 1286-2200, Dec. 2011). A dosage of 1 µg/ml of LNP or CRISPR-Cas9 RNA in or associated with the LNP may be contemplated, especially for a formulation containing DLinKC2-DMA.

[00255] Preparation of LNPs and CRISPR Cas9 encapsulation may be used/and or adapted from Rosin et al, *Molecular Therapy*, vol. 19, no. 12, pages 1286-2200, Dec. 2011). The cationic lipids 1,2-dilinoeyl-3-dimethylammonium-propane (DLinDAP), 1,2-dilinolexyloxy-3-N,N-dimethylaminopropane (DLinDMA), 1,2-dilinolexyloxyketo-N,N-dimethyl-3-aminopropane (DLinK-DMA), 1,2-dilinoleyl-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLinKC2-DMA), (3-*o*-[2''-(methoxypolyethyleneglycol 2000) succinoyl]-1,2-dimyristoyl-sn-glycol (PEG-S-DMG), and R-3-[(ω -methoxy-poly(ethylene glycol)2000) carbamoyl]-1,2-dimyristyloxylpropyl-3-amine (PEG-C-DOMG) may be provided by Tekmira Pharmaceuticals (Vancouver, Canada) or synthesized. Cholesterol may be purchased from Sigma (St Louis, MO). The specific CRISPR Cas9 RNA may be encapsulated in LNPs containing DLinDAP, DLinDMA, DLinK-DMA, and DLinKC2-DMA (cationic lipid:DSPC:CHOL: PEGS-DMG or PEG-C-DOMG at 40:10:40:10 molar ratios). When required, 0.2% SP-DiOC18 (Invitrogen, Burlington, Canada) may be incorporated to assess cellular uptake, intracellular delivery, and biodistribution. Encapsulation may be performed by dissolving lipid mixtures comprised of cationic lipid:DSPC:cholesterol:PEG-c-DOMG (40:10:40:10 molar ratio) in ethanol to a final lipid concentration of 10 mmol/l. This ethanol solution of lipid may be added drop-wise to 50 mmol/l citrate, pH 4.0 to form multilamellar vesicles to produce a final concentration of 30% ethanol vol/vol. Large unilamellar vesicles may be formed following extrusion of multilamellar vesicles through two stacked 80 nm Nuclepore polycarbonate filters using the Extruder (Northern Lipids, Vancouver, Canada). Encapsulation may be achieved by adding RNA dissolved at 2 mg/ml in 50 mmol/l citrate, pH 4.0 containing 30% ethanol vol/vol drop-wise to extruded preformed large unilamellar vesicles and incubation at 31 °C for 30 minutes with constant mixing to a final RNA/lipid weight ratio of 0.06/1 wt/wt. Removal of ethanol and neutralization of formulation

buffer were performed by dialysis against phosphate-buffered saline (PBS), pH 7.4 for 16 hours using Spectra/Por 2 regenerated cellulose dialysis membranes. Nanoparticle size distribution may be determined by dynamic light scattering using a NICOMP 370 particle sizer, the vesicle/intensity modes, and Gaussian fitting (Nicomp Particle Sizing, Santa Barbara, CA). The particle size for all three LNP systems may be ~70 nm in diameter. RNA encapsulation efficiency may be determined by removal of free RNA using VivaPureD MiniH columns (Sartorius Stedim Biotech) from samples collected before and after dialysis. The encapsulated RNA may be extracted from the eluted nanoparticles and quantified at 260 nm. RNA to lipid ratio was determined by measurement of cholesterol content in vesicles using the Cholesterol E enzymatic assay from Wako Chemicals USA (Richmond, VA). In conjunction with the herein discussion of LNPs and PEG lipids, PEGylated liposomes or LNPs are likewise suitable for delivery of a CRISPR-Cas9 system or components thereof.

[00256] Preparation of large LNPs may be used/and or adapted from Rosin et al, *Molecular Therapy*, vol. 19, no. 12, pages 1286-2200, Dec. 2011. A lipid premix solution (20.4 mg/ml total lipid concentration) may be prepared in ethanol containing DLinKC2-DMA, DSPC, and cholesterol at 50:10:38.5 molar ratios. Sodium acetate may be added to the lipid premix at a molar ratio of 0.75:1 (sodium acetate:DLinKC2-DMA). The lipids may be subsequently hydrated by combining the mixture with 1.85 volumes of citrate buffer (10 mmol/l, pH 3.0) with vigorous stirring, resulting in spontaneous liposome formation in aqueous buffer containing 35% ethanol. The liposome solution may be incubated at 37 °C to allow for time-dependent increase in particle size. Aliquots may be removed at various times during incubation to investigate changes in liposome size by dynamic light scattering (Zetasizer Nano ZS, Malvern Instruments, Worcestershire, UK). Once the desired particle size is achieved, an aqueous PEG lipid solution (stock = 10 mg/ml PEG-DMG in 35% (vol/vol) ethanol) may be added to the liposome mixture to yield a final PEG molar concentration of 3.5% of total lipid. Upon addition of PEG-lipids, the liposomes should their size, effectively quenching further growth. RNA may then be added to the empty liposomes at an RNA to total lipid ratio of approximately 1:10 (wt:wt), followed by incubation for 30 minutes at 37 °C to form loaded LNPs. The mixture may be subsequently dialyzed overnight in PBS and filtered with a 0.45- μ m syringe filter.

[00257] Spherical Nucleic Acid (SNATM) constructs and other nanoparticles (particularly gold nanoparticles) are also contemplated as a means to delivery CRISPR-Cas9 system to intended

targets. Significant data show that AuraSense Therapeutics' Spherical Nucleic Acid (SNA™) constructs, based upon nucleic acid-functionalized gold nanoparticles, are useful.

[00258] Literature that may be employed in conjunction with herein teachings include: Cutler et al., *J. Am. Chem. Soc.* 2011 133:9254-9257, Hao et al., *Small*. 2011 7:3158-3162, Zhang et al., *ACS Nano*. 2011 5:6962-6970, Cutler et al., *J. Am. Chem. Soc.* 2012 134:1376-1391, Young et al., *Nano Lett.* 2012 12:3867-71, Zheng et al., *Proc. Natl. Acad. Sci. USA*. 2012 109:11975-80, Mirkin, *Nanomedicine* 2012 7:635-638 Zhang et al., *J. Am. Chem. Soc.* 2012 134:16488-1691, Weintraub, *Nature* 2013 495:S14-S16, Choi et al., *Proc. Natl. Acad. Sci. USA*. 2013 110(19):7625-7630, Jensen et al., *Sci. Transl. Med.* 5, 209ra152 (2013) and Mirkin, et al., *Small*, 10:186-192.

[00259] Self-assembling nanoparticles with RNA may be constructed with polyethyleneimine (PEI) that is PEGylated with an Arg-Gly-Asp (RGD) peptide ligand attached at the distal end of the polyethylene glycol (PEG). This system has been used, for example, as a means to target tumor neovasculature expressing integrins and deliver siRNA inhibiting vascular endothelial growth factor receptor-2 (VEGF R2) expression and thereby achieve tumor angiogenesis (see, e.g., Schiffelers et al., *Nucleic Acids Research*, 2004, Vol. 32, No. 19). Nanoplexes may be prepared by mixing equal volumes of aqueous solutions of cationic polymer and nucleic acid to give a net molar excess of ionizable nitrogen (polymer) to phosphate (nucleic acid) over the range of 2 to 6. The electrostatic interactions between cationic polymers and nucleic acid resulted in the formation of polyplexes with average particle size distribution of about 100 nm, hence referred to here as nanoplexes. A dosage of about 100 to 200 mg of CRISPR Cas9 is envisioned for delivery in the self-assembling nanoparticles of Schiffelers et al.

[00260] The nanoplexes of Bartlett et al. (*PNAS*, September 25, 2007, vol. 104, no. 39) may also be applied to the present invention. The nanoplexes of Bartlett et al. are prepared by mixing equal volumes of aqueous solutions of cationic polymer and nucleic acid to give a net molar excess of ionizable nitrogen (polymer) to phosphate (nucleic acid) over the range of 2 to 6. The electrostatic interactions between cationic polymers and nucleic acid resulted in the formation of polyplexes with average particle size distribution of about 100 nm, hence referred to here as nanoplexes. The DOTA-siRNA of Bartlett et al. was synthesized as follows: 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid mono(N-hydroxysuccinimide ester) (DOTA-NHSester) was ordered from Macrocyclics (Dallas, TX). The amine modified RNA sense strand

with a 100-fold molar excess of DOTA-NHS-ester in carbonate buffer (pH 9) was added to a microcentrifuge tube. The contents were reacted by stirring for 4 h at room temperature. The DOTA-RNAsense conjugate was ethanol-precipitated, resuspended in water, and annealed to the unmodified antisense strand to yield DOTA-siRNA. All liquids were pretreated with Chelex-100 (Bio-Rad, Hercules, CA) to remove trace metal contaminants. Tf-targeted and nontargeted siRNA nanoparticles may be formed by using cyclodextrin-containing polycations. Typically, nanoparticles were formed in water at a charge ratio of 3 (+/-) and an siRNA concentration of 0.5 g/liter. One percent of the adamantane-PEG molecules on the surface of the targeted nanoparticles were modified with Tf (adamantane-PEG-Tf). The nanoparticles were suspended in a 5% (wt/vol) glucose carrier solution for injection.

[00261] Davis et al. (Nature, Vol 464, 15 April 2010) conducts a RNA clinical trial that uses a targeted nanoparticle-delivery system (clinical trial registration number NCT00689065). Patients with solid cancers refractory to standard-of-care therapies are administered doses of targeted particles on days 1, 3, 8 and 10 of a 21-day cycle by a 30-min intravenous infusion. The particles comprise, consist essentially of, or consist of a synthetic delivery system containing: (1) a linear, cyclodextrin-based polymer (CDP), (2) a human transferrin protein (TF) targeting ligand displayed on the exterior of the nanoparticle to engage TF receptors (TFR) on the surface of the cancer cells, (3) a hydrophilic polymer (polyethylene glycol (PEG) used to promote nanoparticle stability in biological fluids), and (4) siRNA designed to reduce the expression of the RRM2 (sequence used in the clinic was previously denoted siR2B+5). The TFR has long been known to be upregulated in malignant cells, and RRM2 is an established anti-cancer target. These nanoparticles (clinical version denoted as CALAA-01) have been shown to be well tolerated in multi-dosing studies in non-human primates. Although a single patient with chronic myeloid leukaemia has been administered siRNA by liposomal delivery, Davis et al.'s clinical trial is the initial human trial to systemically deliver siRNA with a targeted delivery system and to treat patients with solid cancer. To ascertain whether the targeted delivery system can provide effective delivery of functional siRNA to human tumors, Davis et al. investigated biopsies from three patients from three different dosing cohorts; patients A, B and C, all of whom had metastatic melanoma and received CALAA-01 doses of 18, 24 and 30 mg m⁻² siRNA, respectively. Similar doses may also be contemplated for the CRISPR Cas9 system of the present invention. The delivery of the invention may be achieved with nanoparticles containing a linear,

cyclodextrin-based polymer (CDP), a human transferrin protein (TF) targeting ligand displayed on the exterior of the nanoparticle to engage TF receptors (TFR) on the surface of the cancer cells and/or a hydrophilic polymer (for example, polyethylene glycol (PEG) used to promote nanoparticle stability in biological fluids).

Particles

[00262] In terms of this invention, it is preferred to have one or more components of CRISPR complex, e.g., CRISPR enzyme or mRNA or guide RNA delivered using nanoparticles or lipid envelopes. Other delivery systems or vectors are may be used in conjunction with the nanoparticle aspects of the invention.

[00263] In general, a "nanoparticle" refers to any particle having a diameter of less than 1000 nm. In certain preferred embodiments, nanoparticles of the invention have a greatest dimension (e.g., diameter) of 500 nm or less. In other preferred embodiments, nanoparticles of the invention have a greatest dimension ranging between 25 nm and 200 nm. In other preferred embodiments, nanoparticles of the invention have a greatest dimension of 100 nm or less. In other preferred embodiments, nanoparticles of the invention have a greatest dimension ranging between 35 nm and 60 nm.

[00264] Nanoparticles encompassed in the present invention may be provided in different forms, e.g., as solid nanoparticles (e.g., metal such as silver, gold, iron, titanium), non-metal, lipid-based solids, polymers), suspensions of nanoparticles, or combinations thereof. Metal, dielectric, and semiconductor nanoparticles may be prepared, as well as hybrid structures (e.g., core-shell nanoparticles). Nanoparticles made of semiconducting material may also be labeled quantum dots if they are small enough (typically sub 10 nm) that quantization of electronic energy levels occurs. Such nanoscale particles are used in biomedical applications as drug carriers or imaging agents and may be adapted for similar purposes in the present invention.

[00265] Semi-solid and soft nanoparticles have been manufactured, and are within the scope of the present invention. A prototype nanoparticle of semi-solid nature is the liposome. Various types of liposome nanoparticles are currently used clinically as delivery systems for anticancer drugs and vaccines. Nanoparticles with one half hydrophilic and the other half hydrophobic are termed Janus particles and are particularly effective for stabilizing emulsions. They can self-assemble at water/oil interfaces and act as solid surfactants.

[00266] US Patent No. 8,709,843, incorporated herein by reference, provides a drug delivery system for targeted delivery of therapeutic agent-containing particles to tissues, cells, and intracellular compartments. The invention provides targeted particles comprising comprising polymer conjugated to a surfactant, hydrophilic polymer or lipid.

[00267] US Patent No. 6,007,845, incorporated herein by reference, provides particles which have a core of a multiblock copolymer formed by covalently linking a multifunctional compound with one or more hydrophobic polymers and one or more hydrophilic polymers, and contain a biologically active material.

[00268] US Patent No. 5,855,913, incorporated herein by reference, provides a particulate composition having aerodynamically light particles having a tap density of less than 0.4 g/cm³ with a mean diameter of between 5 μm and 30 μm, incorporating a surfactant on the surface thereof for drug delivery to the pulmonary system.

[00269] US Patent No. 5,985,309, incorporated herein by reference, provides particles incorporating a surfactant and/or a hydrophilic or hydrophobic complex of a positively or negatively charged therapeutic or diagnostic agent and a charged molecule of opposite charge for delivery to the pulmonary system.

[00270] US Patent No. 5,543,158, incorporated herein by reference, provides biodegradable injectable nanoparticles having a biodegradable solid core containing a biologically active material and poly(alkylene glycol) moieties on the surface.

[00271] WO2012135025 (also published as US20120251560), incorporated herein by reference, describes conjugated polyethyleneimine (PEI) polymers and conjugated azamacrocycles (collectively referred to as “conjugated lipomer” or “lipomers”). In certain embodiments, it can envisioned that such conjugated lipomers can be used in the context of the CRISPR-Cas9 system to achieve *in vitro*, *ex vivo* and *in vivo* genomic perturbations to modify gene expression, including modulation of protein expression.

[00272] In one embodiment, the nanoparticle may be epoxide-modified lipid-polymer, advantageously 7C1 (see, e.g., James E. Dahlman and Carmen Barnes et al. *Nature Nanotechnology* (2014) published online 11 May 2014, doi:10.1038/nnano.2014.84). C71 was synthesized by reacting C15 epoxide-terminated lipids with PEI600 at a 14:1 molar ratio, and was formulated with C14PEG2000 to produce nanoparticles (diameter between 35 and 60 nm) that were stable in PBS solution for at least 40 days.

[00273] An epoxide-modified lipid-polymer may be utilized to deliver the CRISPR-Cas9 system of the present invention to pulmonary, cardiovascular or renal cells, however, one of skill in the art may adapt the system to deliver to other target organs. Dosage ranging from about 0.05 to about 0.6 mg/kg are envisioned. Dosages over several days or weeks are also envisioned, with a total dosage of about 2 mg/kg.

Exosomes

[00274] Exosomes are endogenous nano-vesicles that transport RNAs and proteins, and which can deliver RNA to the brain and other target organs. To reduce immunogenicity, Alvarez-Erviti et al. (2011, Nat Biotechnol 29: 341) used self-derived dendritic cells for exosome production. Targeting to the brain was achieved by engineering the dendritic cells to express Lamp2b, an exosomal membrane protein, fused to the neuron-specific RVG peptide. Purified exosomes were loaded with exogenous RNA by electroporation. Intravenously injected RVG-targeted exosomes delivered GAPDH siRNA specifically to neurons, microglia, oligodendrocytes in the brain, resulting in a specific gene knockdown. Pre-exposure to RVG exosomes did not attenuate knockdown, and non-specific uptake in other tissues was not observed. The therapeutic potential of exosome-mediated siRNA delivery was demonstrated by the strong mRNA (60%) and protein (62%) knockdown of BACE1, a therapeutic target in Alzheimer's disease.

[00275] To obtain a pool of immunologically inert exosomes, Alvarez-Erviti et al. harvested bone marrow from inbred C57BL/6 mice with a homogenous major histocompatibility complex (MHC) haplotype. As immature dendritic cells produce large quantities of exosomes devoid of T-cell activators such as MHC-II and CD86, Alvarez-Erviti et al. selected for dendritic cells with granulocyte/macrophage-colony stimulating factor (GM-CSF) for 7 d. Exosomes were purified from the culture supernatant the following day using well-established ultracentrifugation protocols. The exosomes produced were physically homogenous, with a size distribution peaking at 80 nm in diameter as determined by nanoparticle tracking analysis (NTA) and electron microscopy. Alvarez-Erviti et al. obtained 6-12 µg of exosomes (measured based on protein concentration) per 10⁶ cells.

[00276] Next, Alvarez-Erviti et al. investigated the possibility of loading modified exosomes with exogenous cargoes using electroporation protocols adapted for nanoscale applications. As electroporation for membrane particles at the nanometer scale is not well-characterized, nonspecific Cy5-labeled RNA was used for the empirical optimization of the electroporation

protocol. The amount of encapsulated RNA was assayed after ultracentrifugation and lysis of exosomes. Electroporation at 400 V and 125 μ F resulted in the greatest retention of RNA and was used for all subsequent experiments.

[00277] Alvarez-Erviti et al. administered 150 μ g of each BACE1 siRNA encapsulated in 150 μ g of RVG exosomes to normal C57BL/6 mice and compared the knockdown efficiency to four controls: untreated mice, mice injected with RVG exosomes only, mice injected with BACE1 siRNA complexed to an *in vivo* cationic liposome reagent and mice injected with BACE1 siRNA complexed to RVG-9R, the RVG peptide conjugated to 9 D-arginines that electrostatically binds to the siRNA. Cortical tissue samples were analyzed 3 d after administration and a significant protein knockdown (45%, $P < 0.05$, versus 62%, $P < 0.01$) in both siRNA-RVG-9R-treated and siRNA-RVG exosome-treated mice was observed, resulting from a significant decrease in BACE1 mRNA levels (66% [+ or -] 15%, $P < 0.001$ and 61% [+ or -] 13% respectively, $P < 0.01$). Moreover, Applicants demonstrated a significant decrease (55%, $P < 0.05$) in the total [beta]-amyloid 1-42 levels, a main component of the amyloid plaques in Alzheimer's pathology, in the RVG-exosome-treated animals. The decrease observed was greater than the β -amyloid 1-40 decrease demonstrated in normal mice after intraventricular injection of BACE1 inhibitors. Alvarez-Erviti et al. carried out 5'-rapid amplification of cDNA ends (RACE) on BACE1 cleavage product, which provided evidence of RNAi-mediated knockdown by the siRNA.

[00278] Finally, Alvarez-Erviti et al. investigated whether RNA-RVG exosomes induced immune responses *in vivo* by assessing IL-6, IP-10, TNF α and IFN- α serum concentrations. Following exosome treatment, nonsignificant changes in all cytokines were registered similar to siRNA-transfection reagent treatment in contrast to siRNA-RVG-9R, which potently stimulated IL-6 secretion, confirming the immunologically inert profile of the exosome treatment. Given that exosomes encapsulate only 20% of siRNA, delivery with RVG-exosome appears to be more efficient than RVG-9R delivery as comparable mRNA knockdown and greater protein knockdown was achieved with fivefold less siRNA without the corresponding level of immune stimulation. This experiment demonstrated the therapeutic potential of RVG-exosome technology, which is potentially suited for long-term silencing of genes related to neurodegenerative diseases. The exosome delivery system of Alvarez-Erviti et al. may be applied to deliver the CRISPR-Cas9 system of the present invention to therapeutic targets, especially

neurodegenerative diseases. A dosage of about 100 to 1000 mg of CRISPR Cas9 encapsulated in about 100 to 1000 mg of RVG exosomes may be contemplated for the present invention.

[00279] El-Andaloussi et al. (Nature Protocols 7,2112–2126(2012)) discloses how exosomes derived from cultured cells can be harnessed for delivery of RNA in vitro and in vivo. This protocol first describes the generation of targeted exosomes through transfection of an expression vector, comprising an exosomal protein fused with a peptide ligand. Next, El-Andaloussi et al. explain how to purify and characterize exosomes from transfected cell supernatant. Next, El-Andaloussi et al. detail crucial steps for loading RNA into exosomes. Finally, El-Andaloussi et al. outline how to use exosomes to efficiently deliver RNA in vitro and in vivo in mouse brain. Examples of anticipated results in which exosome-mediated RNA delivery is evaluated by functional assays and imaging are also provided. The entire protocol takes ~3 weeks. Delivery or administration according to the invention may be performed using exosomes produced from self-derived dendritic cells. From the herein teachings, this can be employed in the practice of the invention

[00280] In another embodiment, the plasma exosomes of Wahlgren et al. (Nucleic Acids Research, 2012, Vol. 40, No. 17 e130) are contemplated. Exosomes are nano-sized vesicles (30–90nm in size) produced by many cell types, including dendritic cells (DC), B cells, T cells, mast cells, epithelial cells and tumor cells. These vesicles are formed by inward budding of late endosomes and are then released to the extracellular environment upon fusion with the plasma membrane. Because exosomes naturally carry RNA between cells, this property may be useful in gene therapy, and from this disclosure can be employed in the practice of the instant invention.

[00281] Exosomes from plasma can be prepared by centrifugation of buffy coat at 900g for 20 min to isolate the plasma followed by harvesting cell supernatants, centrifuging at 300g for 10 min to eliminate cells and at 16 500g for 30 min followed by filtration through a 0.22 mm filter. Exosomes are pelleted by ultracentrifugation at 120 000g for 70 min. Chemical transfection of siRNA into exosomes is carried out according to the manufacturer's instructions in RNAi Human/Mouse Starter Kit (Quiagen, Hilden, Germany). siRNA is added to 100 ml PBS at a final concentration of 2 mmol/ml. After adding HiPerFect transfection reagent, the mixture is incubated for 10 min at RT. In order to remove the excess of micelles, the exosomes are re-isolated using aldehyde/sulfate latex beads. The chemical transfection of CRISPR Cas9 into exosomes may be conducted similarly to siRNA. The exosomes may be co-cultured with

monocytes and lymphocytes isolated from the peripheral blood of healthy donors. Therefore, it may be contemplated that exosomes containing CRISPR Cas9 may be introduced to monocytes and lymphocytes of and autologously reintroduced into a human. Accordingly, delivery or administration according to the invention may be performed using plasma exosomes.

Liposomes

[00282] Delivery or administration according to the invention can be performed with liposomes. Liposomes are spherical vesicle structures composed of a uni- or multilamellar lipid bilayer surrounding internal aqueous compartments and a relatively impermeable outer lipophilic phospholipid bilayer. Liposomes have gained considerable attention as drug delivery carriers because they are biocompatible, nontoxic, can deliver both hydrophilic and lipophilic drug molecules, protect their cargo from degradation by plasma enzymes, and transport their load across biological membranes and the blood brain barrier (BBB) (see, e.g., Spuch and Navarro, Journal of Drug Delivery, vol. 2011, Article ID 469679, 12 pages, 2011. doi:10.1155/2011/469679 for review).

[00283] Liposomes can be made from several different types of lipids; however, phospholipids are most commonly used to generate liposomes as drug carriers. Although liposome formation is spontaneous when a lipid film is mixed with an aqueous solution, it can also be expedited by applying force in the form of shaking by using a homogenizer, sonicator, or an extrusion apparatus (see, e.g., Spuch and Navarro, Journal of Drug Delivery, vol. 2011, Article ID 469679, 12 pages, 2011. doi:10.1155/2011/469679 for review).

[00284] Several other additives may be added to liposomes in order to modify their structure and properties. For instance, either cholesterol or sphingomyelin may be added to the liposomal mixture in order to help stabilize the liposomal structure and to prevent the leakage of the liposomal inner cargo. Further, liposomes are prepared from hydrogenated egg phosphatidylcholine or egg phosphatidylcholine, cholesterol, and dicetyl phosphate, and their mean vesicle sizes were adjusted to about 50 and 100 nm. (see, e.g., Spuch and Navarro, Journal of Drug Delivery, vol. 2011, Article ID 469679, 12 pages, 2011. doi:10.1155/2011/469679 for review).

[00285] A liposome formulation may be mainly comprised of natural phospholipids and lipids such as 1,2-distearoyl-sn-glycero-3-phosphatidyl choline (DSPC), sphingomyelin, egg phosphatidylcholines and monosialoganglioside. Since this formulation is made up of

phospholipids only, liposomal formulations have encountered many challenges, one of the ones being the instability in plasma. Several attempts to overcome these challenges have been made, specifically in the manipulation of the lipid membrane. One of these attempts focused on the manipulation of cholesterol. Addition of cholesterol to conventional formulations reduces rapid release of the encapsulated bioactive compound into the plasma or 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE) increases the stability (see, e.g., Spuch and Navarro, *Journal of Drug Delivery*, vol. 2011, Article ID 469679, 12 pages, 2011. doi:10.1155/2011/469679 for review).

[00286] In a particularly advantageous embodiment, Trojan Horse liposomes (also known as Molecular Trojan Horses) are desirable and protocols may be found at <http://cshprotocols.cshlp.org/content/2010/4/pdb.prot5407.long>. These particles allow delivery of a transgene to the entire brain after an intravascular injection. Without being bound by limitation, it is believed that neutral lipid particles with specific antibodies conjugated to surface allow crossing of the blood brain barrier via endocytosis. Applicant postulates utilizing Trojan Horse Liposomes to deliver the CRISPR family of nucleases to the brain via an intravascular injection, which would allow whole brain transgenic animals without the need for embryonic manipulation. About 1-5 g of DNA or RNA may be contemplated for in vivo administration in liposomes.

[00287] In another embodiment, the CRISPR Cas9 system or components thereof may be administered in liposomes, such as a stable nucleic-acid-lipid particle (SNALP) (see, e.g., Morrissey et al., *Nature Biotechnology*, Vol. 23, No. 8, August 2005). Daily intravenous injections of about 1, 3 or 5 mg/kg/day of a specific CRISPR Cas9 targeted in a SNALP are contemplated. The daily treatment may be over about three days and then weekly for about five weeks. In another embodiment, a specific CRISPR Cas9 encapsulated SNALP administered by intravenous injection to at doses of about 1 or 2.5 mg/kg are also contemplated (see, e.g., Zimmerman et al., *Nature Letters*, Vol. 441, 4 May 2006). The SNALP formulation may contain the lipids 3-N-[(w-methoxypoly(ethylene glycol) 2000) carbamoyl] -1,2-dimyristyloxypropylamine (PEG-C-DMA), 1,2-dilinoleyloxy-N,N-dimethyl-3-aminopropane (DLinDMA), 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC) and cholesterol, in a 2:40:10:48 molar per cent ratio (see, e.g., Zimmerman et al., *Nature Letters*, Vol. 441, 4 May 2006).

[00288] In another embodiment, stable nucleic-acid-lipid particles (SNALPs) have proven to be effective delivery molecules to highly vascularized HepG2-derived liver tumors but not in poorly vascularized HCT-116 derived liver tumors (see, e.g., Li, *Gene Therapy* (2012) 19, 775–780). The SNALP liposomes may be prepared by formulating D-Lin-DMA and PEG-C-DMA with distearoylphosphatidylcholine (DSPC), Cholesterol and siRNA using a 25:1 lipid/siRNA ratio and a 48/40/10/2 molar ratio of Cholesterol/D-Lin-DMA/DSPC/PEG-C-DMA. The resulted SNALP liposomes are about 80–100 nm in size.

[00289] In yet another embodiment, a SNALP may comprise synthetic cholesterol (Sigma-Aldrich, St Louis, MO, USA), dipalmitoylphosphatidylcholine (Avanti Polar Lipids, Alabaster, AL, USA), 3-N-[(w-methoxy poly(ethylene glycol)2000)carbamoyl]-1,2-dimyrestyloxypropylamine, and cationic 1,2-dilinoleyloxy-3-N,Ndimethylaminopropane (see, e.g., Geisbert et al., *Lancet* 2010; 375: 1896-905). A dosage of about 2 mg/kg total CRISPR Cas9 per dose administered as, for example, a bolus intravenous infusion may be contemplated.

[00290] In yet another embodiment, a SNALP may comprise synthetic cholesterol (Sigma-Aldrich), 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC; Avanti Polar Lipids Inc.), PEG-cDMA, and 1,2-dilinoleyloxy-3-(N;N-dimethyl)aminopropane (DLinDMA) (see, e.g., Judge, *J. Clin. Invest.* 119:661-673 (2009)). Formulations used for in vivo studies may comprise a final lipid/RNA mass ratio of about 9:1.

[00291] The safety profile of RNAi nanomedicines has been reviewed by Barros and Gollob of Alnylam Pharmaceuticals (see, e.g., *Advanced Drug Delivery Reviews* 64 (2012) 1730–1737). The stable nucleic acid lipid particle (SNALP) is comprised of four different lipids — an ionizable lipid (DLinDMA) that is cationic at low pH, a neutral helper lipid, cholesterol, and a diffusible polyethylene glycol (PEG)-lipid. The particle is approximately 80 nm in diameter and is charge-neutral at physiologic pH. During formulation, the ionizable lipid serves to condense lipid with the anionic RNA during particle formation. When positively charged under increasingly acidic endosomal conditions, the ionizable lipid also mediates the fusion of SNALP with the endosomal membrane enabling release of RNA into the cytoplasm. The PEG-lipid stabilizes the particle and reduces aggregation during formulation, and subsequently provides a neutral hydrophilic exterior that improves pharmacokinetic properties.

[00292] To date, two clinical programs have been initiated using SNALP formulations with RNA. Tekmira Pharmaceuticals recently completed a phase I single-dose study of SNALP-ApoB

in adult volunteers with elevated LDL cholesterol. ApoB is predominantly expressed in the liver and jejunum and is essential for the assembly and secretion of VLDL and LDL. Seventeen subjects received a single dose of SNALP-ApoB (dose escalation across 7 dose levels). There was no evidence of liver toxicity (anticipated as the potential dose-limiting toxicity based on preclinical studies). One (of two) subjects at the highest dose experienced flu-like symptoms consistent with immune system stimulation, and the decision was made to conclude the trial.

[00293] Alnylam Pharmaceuticals has similarly advanced ALN-TTR01, which employs the SNALP technology described above and targets hepatocyte production of both mutant and wild-type TTR to treat TTR amyloidosis (ATTR). Three ATTR syndromes have been described: familial amyloidotic polyneuropathy (FAP) and familial amyloidotic cardiomyopathy (FAC) — both caused by autosomal dominant mutations in TTR; and senile systemic amyloidosis (SSA) cause by wildtype TTR. A placebo-controlled, single dose-escalation phase I trial of ALN-TTR01 was recently completed in patients with ATTR. ALN-TTR01 was administered as a 15-minute IV infusion to 31 patients (23 with study drug and 8 with placebo) within a dose range of 0.01 to 1.0 mg/kg (based on siRNA). Treatment was well tolerated with no significant increases in liver function tests. Infusion-related reactions were noted in 3 of 23 patients at ≥ 0.4 mg/kg; all responded to slowing of the infusion rate and all continued on study. Minimal and transient elevations of serum cytokines IL-6, IP-10 and IL-1ra were noted in two patients at the highest dose of 1 mg/kg (as anticipated from preclinical and NHP studies). Lowering of serum TTR, the expected pharmacodynamics effect of ALN-TTR01, was observed at 1 mg/kg.

[00294] In yet another embodiment, a SNALP may be made by solubilizing a cationic lipid, DSPC, cholesterol and PEG-lipid e.g., in ethanol, e.g., at a molar ratio of 40:10:40:10, respectively (see, Semple et al., Nature Biotechnology, Volume 28 Number 2 February 2010, pp. 172-177). The lipid mixture was added to an aqueous buffer (50 mM citrate, pH 4) with mixing to a final ethanol and lipid concentration of 30% (vol/vol) and 6.1 mg/ml, respectively, and allowed to equilibrate at 22 °C for 2 min before extrusion. The hydrated lipids were extruded through two stacked 80 nm pore-sized filters (Nuclepore) at 22 °C using a Lipex Extruder (Northern Lipids) until a vesicle diameter of 70–90 nm, as determined by dynamic light scattering analysis, was obtained. This generally required 1–3 passes. The siRNA (solubilized in a 50 mM citrate, pH 4 aqueous solution containing 30% ethanol) was added to the pre-equilibrated (35 °C) vesicles at a rate of ~5 ml/min with mixing. After a final target siRNA/lipid

ratio of 0.06 (wt/wt) was reached, the mixture was incubated for a further 30 min at 35 °C to allow vesicle reorganization and encapsulation of the siRNA. The ethanol was then removed and the external buffer replaced with PBS (155 mM NaCl, 3 mM Na₂HPO₄, 1 mM KH₂PO₄, pH 7.5) by either dialysis or tangential flow diafiltration. siRNA were encapsulated in SNALP using a controlled step-wise dilution method process. The lipid constituents of KC2-SNALP were DLin-KC2-DMA (cationic lipid), dipalmitoylphosphatidylcholine (DPPC; Avanti Polar Lipids), synthetic cholesterol (Sigma) and PEG-C-DMA used at a molar ratio of 57.1:7.1:34.3:1.4. Upon formation of the loaded particles, SNALP were dialyzed against PBS and filter sterilized through a 0.2 µm filter before use. Mean particle sizes were 75–85 nm and 90–95% of the siRNA was encapsulated within the lipid particles. The final siRNA/lipid ratio in formulations used for in vivo testing was ~0.15 (wt/wt). LNP-siRNA systems containing Factor VII siRNA were diluted to the appropriate concentrations in sterile PBS immediately before use and the formulations were administered intravenously through the lateral tail vein in a total volume of 10 ml/kg. This method and these delivery systems may be extrapolated to the CRISPR Cas9 system of the present invention.

Other Lipids

[00295] Other cationic lipids, such as amino lipid 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA) may be utilized to encapsulate CRISPR Cas9 or components thereof or nucleic acid molecule(s) coding therefor e.g., similar to SiRNA (see, e.g., Jayaraman, *Angew. Chem. Int. Ed.* 2012, 51, 8529–8533), and hence may be employed in the practice of the invention. A preformed vesicle with the following lipid composition may be contemplated: amino lipid, distearoylphosphatidylcholine (DSPC), cholesterol and (R)-2,3-bis(octadecyloxy) propyl-1-(methoxy poly(ethylene glycol)2000)propylcarbamate (PEG-lipid) in the molar ratio 40/10/40/10, respectively, and a FVII siRNA/total lipid ratio of approximately 0.05 (w/w). To ensure a narrow particle size distribution in the range of 70–90 nm and a low polydispersity index of 0.11 ± 0.04 (n=56), the particles may be extruded up to three times through 80 nm membranes prior to adding the guide RNA. Particles containing the highly potent amino lipid 16 may be used, in which the molar ratio of the four lipid components 16, DSPC, cholesterol and PEG-lipid (50/10/38.5/1.5) which may be further optimized to enhance in vivo activity.

[00296] Michael S D Kormann et al. ("Expression of therapeutic proteins after delivery of chemically modified mRNA in mice: *Nature Biotechnology*, Volume:29, Pages: 154–157

(2011)) describes the use of lipid envelopes to deliver RNA. Use of lipid envelopes is also preferred in the present invention.

[00297] In another embodiment, lipids may be formulated with the CRISPR Cas9 system of the present invention or component(s) thereof or nucleic acid molecule(s) coding therefor to form lipid nanoparticles (LNPs). Lipids include, but are not limited to, DLin-KC2-DMA4, C12-200 and colipids distearylphosphatidyl choline, cholesterol, and PEG-DMG may be formulated with CRISPR Cas9 instead of siRNA (see, e.g., Novobrantseva, *Molecular Therapy—Nucleic Acids* (2012) 1, e4; doi:10.1038/mtna.2011.3) using a spontaneous vesicle formation procedure. The component molar ratio may be about 50/10/38.5/1.5 (DLin-KC2-DMA or C12-200/distearylphosphatidyl choline/cholesterol/PEG-DMG). The final lipid:siRNA weight ratio may be ~12:1 and 9:1 in the case of DLin-KC2-DMA and C12-200 lipid nanoparticles (LNPs), respectively. The formulations may have mean particle diameters of ~80 nm with >90% entrapment efficiency. A 3 mg/kg dose may be contemplated.

[00298] Tekmira has a portfolio of approximately 95 patent families, in the U.S. and abroad, that are directed to various aspects of LNPs and LNP formulations (see, e.g., U.S. Pat. Nos. 7,982,027; 7,799,565; 8,058,069; 8,283,333; 7,901,708; 7,745,651; 7,803,397; 8,101,741; 8,188,263; 7,915,399; 8,236,943 and 7,838,658 and European Pat. Nos 1766035; 1519714; 1781593 and 1664316), all of which may be used and/or adapted to the present invention.

[00299] The CRISPR Cas9 system or components thereof or nucleic acid molecule(s) coding therefor may be delivered encapsulated in PLGA Microspheres such as that further described in US published applications 20130252281 and 20130245107 and 20130244279 (assigned to Moderna Therapeutics) which relate to aspects of formulation of compositions comprising modified nucleic acid molecules which may encode a protein, a protein precursor, or a partially or fully processed form of the protein or a protein precursor. The formulation may have a molar ratio 50:10:38.5:1.5-3.0 (cationic lipid: fusogenic lipid: cholesterol: PEG lipid). The PEG lipid may be selected from, but is not limited to PEG-c-DMG, PEG-DMG. The fusogenic lipid may be DSPC. See also, Schrum et al., *Delivery and Formulation of Engineered Nucleic Acids*, US published application 20120251618.

[00300] Nanomerics' technology addresses bioavailability challenges for a broad range of therapeutics, including low molecular weight hydrophobic drugs, peptides, and nucleic acid based therapeutics (plasmid, siRNA, miRNA). Specific administration routes for which the

technology has demonstrated clear advantages include the oral route, transport across the blood-brain-barrier, delivery to solid tumors, as well as to the eye. See, e.g., Mazza et al., 2013, ACS Nano. 2013 Feb 26;7(2):1016-26; Uchegbu and Siew, 2013, J Pharm Sci. 102(2):305-10 and Lalatsa et al., 2012, J Control Release. 2012 Jul 20; 161(2):523-36.

[00301] US Patent Publication No. 20050019923 describes cationic dendrimers for delivering bioactive molecules, such as polynucleotide molecules, peptides and polypeptides and/or pharmaceutical agents, to a mammalian body. The dendrimers are suitable for targeting the delivery of the bioactive molecules to, for example, the liver, spleen, lung, kidney or heart (or even the brain). Dendrimers are synthetic 3-dimensional macromolecules that are prepared in a step-wise fashion from simple branched monomer units, the nature and functionality of which can be easily controlled and varied. Dendrimers are synthesized from the repeated addition of building blocks to a multifunctional core (divergent approach to synthesis), or towards a multifunctional core (convergent approach to synthesis) and each addition of a 3-dimensional shell of building blocks leads to the formation of a higher generation of the dendrimers. Polypropylenimine dendrimers start from a diaminobutane core to which is added twice the number of amino groups by a double Michael addition of acrylonitrile to the primary amines followed by the hydrogenation of the nitriles. This results in a doubling of the amino groups. Polypropylenimine dendrimers contain 100% protonable nitrogens and up to 64 terminal amino groups (generation 5, DAB 64). Protonable groups are usually amine groups which are able to accept protons at neutral pH. The use of dendrimers as gene delivery agents has largely focused on the use of the polyamidoamine, and phosphorous containing compounds with a mixture of amine/amide or N--P(O₂)S as the conjugating units respectively with no work being reported on the use of the lower generation polypropylenimine dendrimers for gene delivery. Polypropylenimine dendrimers have also been studied as pH sensitive controlled release systems for drug delivery and for their encapsulation of guest molecules when chemically modified by peripheral amino acid groups. The cytotoxicity and interaction of polypropylenimine dendrimers with DNA as well as the transfection efficacy of DAB 64 has also been studied.

[00302] US Patent Publication No. 20050019923 is based upon the observation that, contrary to earlier reports, cationic dendrimers, such as polypropylenimine dendrimers, display suitable properties, such as specific targeting and low toxicity, for use in the targeted delivery of bioactive molecules, such as genetic material. In addition, derivatives of the cationic dendrimer

also display suitable properties for the targeted delivery of bioactive molecules. See also, Bioactive Polymers, US published application 20080267903, which discloses "Various polymers, including cationic polyamine polymers and dendrimeric polymers, are shown to possess anti-proliferative activity, and may therefore be useful for treatment of disorders characterised by undesirable cellular proliferation such as neoplasms and tumors, inflammatory disorders (including autoimmune disorders), psoriasis and atherosclerosis. The polymers may be used alone as active agents, or as delivery vehicles for other therapeutic agents, such as drug molecules or nucleic acids for gene therapy. In such cases, the polymers' own intrinsic anti-tumor activity may complement the activity of the agent to be delivered." The disclosures of these patent publications may be employed in conjunction with herein teachings for delivery of CRISPR Cas9 system(s) or component(s) thereof or nucleic acid molecule(s) coding therefor.

Supercharged proteins

[00303] Supercharged proteins are a class of engineered or naturally occurring proteins with unusually high positive or negative net theoretical charge and may be employed in delivery of CRISPR Cas9 system(s) or component(s) thereof or nucleic acid molecule(s) coding therefor. Both supernegatively and superpositively charged proteins exhibit a remarkable ability to withstand thermally or chemically induced aggregation. Superpositively charged proteins are also able to penetrate mammalian cells. Associating cargo with these proteins, such as plasmid DNA, RNA, or other proteins, can enable the functional delivery of these macromolecules into mammalian cells both in vitro and in vivo. David Liu's lab reported the creation and characterization of supercharged proteins in 2007 (Lawrence et al., 2007, Journal of the American Chemical Society 129, 10110–10112).

[00304] The nonviral delivery of RNA and plasmid DNA into mammalian cells are valuable both for research and therapeutic applications (Akinc et al., 2010, Nat. Biotech. 26, 561–569). Purified +36 GFP protein (or other superpositively charged protein) is mixed with RNAs in the appropriate serum-free media and allowed to complex prior addition to cells. Inclusion of serum at this stage inhibits formation of the supercharged protein-RNA complexes and reduces the effectiveness of the treatment. The following protocol has been found to be effective for a variety of cell lines (McNaughton et al., 2009, Proc. Natl. Acad. Sci. USA 106, 6111–6116). However, pilot experiments varying the dose of protein and RNA should be performed to optimize the procedure for specific cell lines.

- (1) One day before treatment, plate 1×10^5 cells per well in a 48-well plate.
- (2) On the day of treatment, dilute purified +36 GFP protein in serum free media to a final concentration 200nM. Add RNA to a final concentration of 50nM. Vortex to mix and incubate at room temperature for 10min.
- (3) During incubation, aspirate media from cells and wash once with PBS.
- (4) Following incubation of +36 GFP and RNA, add the protein-RNA complexes to cells.
- (5) Incubate cells with complexes at 37 °C for 4h.
- (6) Following incubation, aspirate the media and wash three times with 20 U/mL heparin PBS. Incubate cells with serum-containing media for a further 48h or longer depending upon the assay for activity.
- (7) Analyze cells by immunoblot, qPCR, phenotypic assay, or other appropriate method.

[00305] David Liu's lab has further found +36 GFP to be an effective plasmid delivery reagent in a range of cells. As plasmid DNA is a larger cargo than siRNA, proportionately more +36 GFP protein is required to effectively complex plasmids. For effective plasmid delivery Applicants have developed a variant of +36 GFP bearing a C-terminal HA2 peptide tag, a known endosome-disrupting peptide derived from the influenza virus hemagglutinin protein. The following protocol has been effective in a variety of cells, but as above it is advised that plasmid DNA and supercharged protein doses be optimized for specific cell lines and delivery applications.

- (1) One day before treatment, plate 1×10^5 per well in a 48-well plate.
- (2) On the day of treatment, dilute purified p36 GFP protein in serumfree media to a final concentration 2 mM. Add 1mg of plasmid DNA. Vortex to mix and incubate at room temperature for 10min.
- (3) During incubation, aspirate media from cells and wash once with PBS.
- (4) Following incubation of p36 GFP and plasmid DNA, gently add the protein-DNA complexes to cells.
- (5) Incubate cells with complexes at 37 C for 4h.
- (6) Following incubation, aspirate the media and wash with PBS. Incubate cells in serum-containing media and incubate for a further 24–48h.

(7) Analyze plasmid delivery (e.g., by plasmid-driven gene expression) as appropriate.

[00306] See also, e.g., McNaughton et al., Proc. Natl. Acad. Sci. USA 106, 6111-6116 (2009); Cronican et al., ACS Chemical Biology 5, 747-752 (2010); Cronican et al., Chemistry & Biology 18, 833-838 (2011); Thompson et al., Methods in Enzymology 503, 293-319 (2012); Thompson, D.B., et al., Chemistry & Biology 19 (7), 831-843 (2012). The methods of the super charged proteins may be used and/or adapted for delivery of the CRISPR Cas9 system of the present invention. These systems of Dr. Lui and documents herein in conjunction with herein teachings can be employed in the delivery of CRISPR Cas9 system(s) or component(s) thereof or nucleic acid molecule(s) coding therefor.

Cell Penetrating Peptides (CPPs)

[00307] In yet another embodiment, cell penetrating peptides (CPPs) are contemplated for the delivery of the CRISPR Cas9 system. CPPs are short peptides that facilitate cellular uptake of various molecular cargo (from nanosize particles to small chemical molecules and large fragments of DNA). The term "cargo" as used herein includes but is not limited to the group consisting of therapeutic agents, diagnostic probes, peptides, nucleic acids, antisense oligonucleotides, plasmids, proteins, particles including nanoparticles, liposomes, chromophores, small molecules and radioactive materials. In aspects of the invention, the cargo may also comprise any component of the CRISPR Cas9 system or the entire functional CRISPR Cas9 system. Aspects of the present invention further provide methods for delivering a desired cargo into a subject comprising: (a) preparing a complex comprising the cell penetrating peptide of the present invention and a desired cargo, and (b) orally, intraarticularly, intraperitoneally, intrathecally, intrarterially, intranasally, intraparenchymally, subcutaneously, intramuscularly, intravenously, dermally, intrarectally, or topically administering the complex to a subject. The cargo is associated with the peptides either through chemical linkage via covalent bonds or through non-covalent interactions.

[00308] The function of the CPPs are to deliver the cargo into cells, a process that commonly occurs through endocytosis with the cargo delivered to the endosomes of living mammalian cells. Cell-penetrating peptides are of different sizes, amino acid sequences, and charges but all CPPs have one distinct characteristic, which is the ability to translocate the plasma membrane and facilitate the delivery of various molecular cargoes to the cytoplasm or an organelle. CPP

translocation may be classified into three main entry mechanisms: direct penetration in the membrane, endocytosis-mediated entry, and translocation through the formation of a transitory structure. CPPs have found numerous applications in medicine as drug delivery agents in the treatment of different diseases including cancer and virus inhibitors, as well as contrast agents for cell labeling. Examples of the latter include acting as a carrier for GFP, MRI contrast agents, or quantum dots. CPPs hold great potential as *in vitro* and *in vivo* delivery vectors for use in research and medicine. CPPs typically have an amino acid composition that either contains a high relative abundance of positively charged amino acids such as lysine or arginine or has sequences that contain an alternating pattern of polar/charged amino acids and non-polar, hydrophobic amino acids. These two types of structures are referred to as polycationic or amphipathic, respectively. A third class of CPPs are the hydrophobic peptides, containing only apolar residues, with low net charge or have hydrophobic amino acid groups that are crucial for cellular uptake. One of the initial CPPs discovered was the trans-activating transcriptional activator (Tat) from Human Immunodeficiency Virus 1 (HIV-1) which was found to be efficiently taken up from the surrounding media by numerous cell types in culture. Since then, the number of known CPPs has expanded considerably and small molecule synthetic analogues with more effective protein transduction properties have been generated. CPPs include but are not limited to Penetratin, Tat (48-60), Transportan, and (R-Ahx-R4) (Ahx=aminohexanoyl).

[00309] US Patent 8,372,951, provides a CPP derived from eosinophil cationic protein (ECP) which exhibits highly cell-penetrating efficiency and low toxicity. Aspects of delivering the CPP with its cargo into a vertebrate subject are also provided. Further aspects of CPPs and their delivery are described in U. S. patents 8,575,305; 8,614,194 and 8,044,019. CPPs can be used to deliver the CRISPR-Cas9 system or components thereof. That CPPs can be employed to deliver the CRISPR-Cas9 system or components thereof is also provided in the manuscript "Gene disruption by cell-penetrating peptide-mediated delivery of Cas9 protein and guide RNA", by Suresh Ramakrishna, Abu-Bonsrah Kwaku Dad, Jagadish Beloor, et al. *Genome Res.* 2014 Apr 2. [Epub ahead of print], incorporated by reference in its entirety, wherein it is demonstrated that treatment with CPP-conjugated recombinant Cas9 protein and CPP-complexed guide RNAs lead to endogenous gene disruptions in human cell lines. In the paper the Cas9 protein was conjugated to CPP via a thioether bond, whereas the guide RNA was complexed with CPP, forming condensed, positively charged particles. It was shown that simultaneous and sequential treatment

of human cells, including embryonic stem cells, dermal fibroblasts, HEK293T cells, HeLa cells, and embryonic carcinoma cells, with the modified Cas9 and guide RNA led to efficient gene disruptions with reduced off-target mutations relative to plasmid transfections.

Implantable devices

[00310] In another embodiment, implantable devices are also contemplated for delivery of the CRISPR Cas9 system or component(s) thereof or nucleic acid molecule(s) coding therefor. For example, US Patent Publication 20110195123 discloses an implantable medical device which elutes a drug locally and in prolonged period is provided, including several types of such a device, the treatment modes of implementation and methods of implantation. The device comprising of polymeric substrate, such as a matrix for example, that is used as the device body, and drugs, and in some cases additional scaffolding materials, such as metals or additional polymers, and materials to enhance visibility and imaging. An implantable delivery device can be advantageous in providing release locally and over a prolonged period, where drug is released directly to the extracellular matrix (ECM) of the diseased area such as tumor, inflammation, degeneration or for symptomatic objectives, or to injured smooth muscle cells, or for prevention. One kind of drug is RNA, as disclosed above, and this system may be used/and or adapted to the CRISPR Cas9 system of the present invention. The modes of implantation in some embodiments are existing implantation procedures that are developed and used today for other treatments, including brachytherapy and needle biopsy. In such cases the dimensions of the new implant described in this invention are similar to the original implant. Typically a few devices are implanted during the same treatment procedure.

[00311] US Patent Publication 20110195123, provides a drug delivery implantable or insertable system, including systems applicable to a cavity such as the abdominal cavity and/or any other type of administration in which the drug delivery system is not anchored or attached, comprising a biostable and/or degradable and/or bioabsorbable polymeric substrate, which may for example optionally be a matrix. It should be noted that the term "insertion" also includes implantation. The drug delivery system is preferably implemented as a "Loder" as described in US Patent Publication 20110195123.

[00312] The polymer or plurality of polymers are biocompatible, incorporating an agent and/or plurality of agents, enabling the release of agent at a controlled rate, wherein the total volume of the polymeric substrate, such as a matrix for example, in some embodiments is

optionally and preferably no greater than a maximum volume that permits a therapeutic level of the agent to be reached. As a non-limiting example, such a volume is preferably within the range of 0.1 m³ to 1000 mm³, as required by the volume for the agent load. The Loder may optionally be larger, for example when incorporated with a device whose size is determined by functionality, for example and without limitation, a knee joint, an intra-uterine or cervical ring and the like.

[00313] The drug delivery system (for delivering the composition) is designed in some embodiments to preferably employ degradable polymers, wherein the main release mechanism is bulk erosion; or in some embodiments, non degradable, or slowly degraded polymers are used, wherein the main release mechanism is diffusion rather than bulk erosion, so that the outer part functions as membrane, and its internal part functions as a drug reservoir, which practically is not affected by the surroundings for an extended period (for example from about a week to about a few months). Combinations of different polymers with different release mechanisms may also optionally be used. The concentration gradient at the surface is preferably maintained effectively constant during a significant period of the total drug releasing period, and therefore the diffusion rate is effectively constant (termed "zero mode" diffusion). By the term "constant" it is meant a diffusion rate that is preferably maintained above the lower threshold of therapeutic effectiveness, but which may still optionally feature an initial burst and/or may fluctuate, for example increasing and decreasing to a certain degree. The diffusion rate is preferably so maintained for a prolonged period, and it can be considered constant to a certain level to optimize the therapeutically effective period, for example the effective silencing period.

[00314] The drug delivery system optionally and preferably is designed to shield the nucleotide based therapeutic agent from degradation, whether chemical in nature or due to attack from enzymes and other factors in the body of the subject.

[00315] The drug delivery system of US Patent Publication 20110195123 is optionally associated with sensing and/or activation appliances that are operated at and/or after implantation of the device, by non and/or minimally invasive methods of activation and/or acceleration/deceleration, for example optionally including but not limited to thermal heating and cooling, laser beams, and ultrasonic, including focused ultrasound and/or RF (radiofrequency) methods or devices.

[00316] According to some embodiments of US Patent Publication 20110195123, the site for local delivery may optionally include target sites characterized by high abnormal proliferation of cells, and suppressed apoptosis, including tumors, active and or chronic inflammation and infection including autoimmune diseases states, degenerating tissue including muscle and nervous tissue, chronic pain, degenerative sites, and location of bone fractures and other wound locations for enhancement of regeneration of tissue, and injured cardiac, smooth and striated muscle.

[00317] The site for implantation of the composition, or target site, preferably features a radius, area and/or volume that is sufficiently small for targeted local delivery. For example, the target site optionally has a diameter in a range of from about 0.1 mm to about 5 cm.

[00318] The location of the target site is preferably selected for maximum therapeutic efficacy. For example, the composition of the drug delivery system (optionally with a device for implantation as described above) is optionally and preferably implanted within or in the proximity of a tumor environment, or the blood supply associated thereof.

[00319] For example the composition (optionally with the device) is optionally implanted within or in the proximity to pancreas, prostate, breast, liver, via the nipple, within the vascular system and so forth.

[00320] The target location is optionally selected from the group comprising, consisting essentially of, or consisting of (as non-limiting examples only, as optionally any site within the body may be suitable for implanting a Loder): 1. brain at degenerative sites like in Parkinson or Alzheimer disease at the basal ganglia, white and gray matter; 2. spine as in the case of amyotrophic lateral sclerosis (ALS); 3. uterine cervix to prevent HPV infection; 4. active and chronic inflammatory joints; 5. dermis as in the case of psoriasis; 6. sympathetic and sensoric nervous sites for analgesic effect; 7. Intra osseous implantation; 8. acute and chronic infection sites; 9. Intra vaginal; 10. Inner ear--auditory system, labyrinth of the inner ear, vestibular system; 11. Intra tracheal; 12. Intra-cardiac, coronary, epicardiac; 13. urinary bladder; 14. biliary system; 15. parenchymal tissue including and not limited to the kidney, liver, spleen; 16. lymph nodes; 17. salivary glands; 18. dental gums; 19. Intra-articular (into joints); 20. Intra-ocular; 21. Brain tissue; 22. Brain ventricles; 23. Cavities, including abdominal cavity (for example but without limitation, for ovary cancer); 24. Intra esophageal and 25. Intra rectal.

[00321] Optionally insertion of the system (for example a device containing the composition) is associated with injection of material to the ECM at the target site and the vicinity of that site to affect local pH and/or temperature and/or other biological factors affecting the diffusion of the drug and/or drug kinetics in the ECM, of the target site and the vicinity of such a site.

[00322] Optionally, according to some embodiments, the release of said agent could be associated with sensing and/or activation appliances that are operated prior and/or at and/or after insertion, by non and/or minimally invasive and/or else methods of activation and/or acceleration/deceleration, including laser beam, radiation, thermal heating and cooling, and ultrasonic, including focused ultrasound and/or RF (radiofrequency) methods or devices, and chemical activators.

[00323] According to other embodiments of US Patent Publication 20110195123, the drug preferably comprises a RNA, for example for localized cancer cases in breast, pancreas, brain, kidney, bladder, lung, and prostate as described below. Although exemplified with RNAi, many drugs are applicable to be encapsulated in Loder, and can be used in association with this invention, as long as such drugs can be encapsulated with the Loder substrate, such as a matrix for example, and this system may be used and/or adapted to deliver the CRISPR Cas9 system of the present invention.

[00324] As another example of a specific application, neuro and muscular degenerative diseases develop due to abnormal gene expression. Local delivery of RNAs may have therapeutic properties for interfering with such abnormal gene expression. Local delivery of anti apoptotic, anti inflammatory and anti degenerative drugs including small drugs and macromolecules may also optionally be therapeutic. In such cases the Loder is applied for prolonged release at constant rate and/or through a dedicated device that is implanted separately. All of this may be used and/or adapted to the CRISPR Cas9 system of the present invention.

[00325] As yet another example of a specific application, psychiatric and cognitive disorders are treated with gene modifiers. Gene knockdown is a treatment option. Loders locally delivering agents to central nervous system sites are therapeutic options for psychiatric and cognitive disorders including but not limited to psychosis, bi-polar diseases, neurotic disorders and behavioral maladies. The Loders could also deliver locally drugs including small drugs and macromolecules upon implantation at specific brain sites. All of this may be used and/or adapted to the CRISPR Cas9 system of the present invention.

[00326] As another example of a specific application, silencing of innate and/or adaptive immune mediators at local sites enables the prevention of organ transplant rejection. Local delivery of RNAs and immunomodulating reagents with the Loder implanted into the transplanted organ and/or the implanted site renders local immune suppression by repelling immune cells such as CD8 activated against the transplanted organ. All of this may be used/and or adapted to the CRISPR Cas9 system of the present invention.

[00327] As another example of a specific application, vascular growth factors including VEGFs and angiogenin and others are essential for neovascularization. Local delivery of the factors, peptides, peptidomimetics, or suppressing their repressors is an important therapeutic modality; silencing the repressors and local delivery of the factors, peptides, macromolecules and small drugs stimulating angiogenesis with the Loder is therapeutic for peripheral, systemic and cardiac vascular disease.

[00328] The method of insertion, such as implantation, may optionally already be used for other types of tissue implantation and/or for insertions and/or for sampling tissues, optionally without modifications, or alternatively optionally only with non-major modifications in such methods. Such methods optionally include but are not limited to brachytherapy methods, biopsy, endoscopy with and/or without ultrasound, such as ERCP, stereotactic methods into the brain tissue, Laparoscopy, including implantation with a laparoscope into joints, abdominal organs, the bladder wall and body cavities.

[00329] Implantable device technology herein discussed can be employed with herein teachings and hence by this disclosure and the knowledge in the art, CRISPR-Cas9 system or components thereof or nucleic acid molecules thereof or encoding or providing components may be delivered via an implantable device.

Patient-specific screening methods

[00330] A nucleic acid-targeting system that targets DNA, e.g., trinucleotide repeats can be used to screen patients or patient samples for the presence of such repeats. The repeats can be the target of the RNA of the nucleic acid-targeting system, and if there is binding thereto by the nucleic acid-targeting system, that binding can be detected, to thereby indicate that such a repeat is present. Thus, a nucleic acid-targeting system can be used to screen patients or patient samples for the presence of the repeat. The patient can then be administered suitable

compound(s) to address the condition; or, can be administered a nucleic acid-targeting system to bind to and cause insertion, deletion or mutation and alleviate the condition.

CRISPR effector protein mRNA and guide RNA

[00331] CRISPR enzyme mRNA and guide RNA might also be delivered separately. CRISPR enzyme mRNA can be delivered prior to the guide RNA to give time for CRISPR enzyme to be expressed. CRISPR enzyme mRNA might be administered 1-12 hours (preferably around 2-6 hours) prior to the administration of guide RNA.

[00332] Alternatively, CRISPR enzyme mRNA and guide RNA can be administered together. Advantageously, a second booster dose of guide RNA can be administered 1-12 hours (preferably around 2-6 hours) after the initial administration of CRISPR enzyme mRNA + guide RNA.

[00333] The CRISPR effector protein of the present invention, i.e. a Cas9 effector protein is sometimes referred to herein as a CRISPR Enzyme. It will be appreciated that the effector protein is based on or derived from an enzyme, so the term 'effector protein' certainly includes 'enzyme' in some embodiments. However, it will also be appreciated that the effector protein may, as required in some embodiments, have DNA or RNA binding, but not necessarily cutting or nicking, activity, including a dead-Cas9 effector protein function.

[00334] Additional administrations of CRISPR enzyme mRNA and/or guide RNA might be useful to achieve the most efficient levels of genome modification. In some embodiments, phenotypic alteration is preferably the result of genome modification when a genetic disease is targeted, especially in methods of therapy and preferably where a repair template is provided to correct or alter the phenotype.

[00335] In some embodiments diseases that may be targeted include those concerned with disease-causing splice defects.

[00336] In some embodiments, cellular targets include Hemopoietic Stem/Progenitor Cells (CD34+); Human T cells; and Eye (retinal cells) – for example photoreceptor precursor cells.

[00337] In some embodiments Gene targets include: Human Beta Globin – HBB (for treating Sickle Cell Anemia, including by stimulating gene-conversion (using closely related HBD gene as an endogenous template)); CD3 (T-Cells); and CEP920 - retina (eye).

[00338] In some embodiments disease targets also include: cancer; Sickle Cell Anemia (based on a point mutation); HIV; Beta-Thalassemia; and ophthalmic or ocular disease – for example Leber Congenital Amaurosis (LCA)-causing Splice Defect.

[00339] In some embodiments delivery methods include: Cationic Lipid Mediated “direct” delivery of Enzyme-Guide complex (RiboNucleoProtein) and electroporation of plasmid DNA.

[00340] Inventive methods can further comprise delivery of templates, such as repair templates, which may be dsODN or ssODN, see below. Delivery of templates may be via the contemporaneous or separate from delivery of any or all the CRISPR enzyme, guide, tracr mate or tracrRNA and via the same delivery mechanism or different. In some embodiments, it is preferred that the template is delivered together with the guide, tracr mate and/or tracrRNA and, preferably, also the CRISPR enzyme. An example may be an AAV vector where the CRISPR enzyme is SaCas9 (with the N580 mutation).

[00341] Inventive methods can further comprise: (a) delivering to the cell a double-stranded oligodeoxynucleotide (dsODN) comprising overhangs complimentary to the overhangs created by said double strand break, wherein said dsODN is integrated into the locus of interest; or –(b) delivering to the cell a single-stranded oligodeoxynucleotide (ssODN), wherein said ssODN acts as a template for homology directed repair of said double strand break. Inventive methods can be for the prevention or treatment of disease in an individual, optionally wherein said disease is caused by a defect in said locus of interest. Inventive methods can be conducted *in vivo* in the individual or *ex vivo* on a cell taken from the individual, optionally wherein said cell is returned to the individual.

[00342] For minimization of toxicity and off-target effect, it will be important to control the concentration of CRISPR enzyme mRNA and guide RNA delivered. Optimal concentrations of CRISPR enzyme mRNA and guide RNA can be determined by testing different concentrations in a cellular or animal model and using deep sequencing to analyze the extent of modification at potential off-target genomic loci. For example, for the guide sequence targeting 5'-GAGTCCGAGCAGAAGAAGAA-3' in the EMX1 gene of the human genome, deep sequencing can be used to assess the level of modification at the following two off-target loci, 1: 5'-GAGTCCTAGCAGGAGAAGAA-3' and 2: 5'-GAGTCTAAGCAGAAGAAGAA-3'. The concentration that gives the highest level of on-target modification while minimizing the level of off-target modification should be chosen for *in vivo* delivery.

Inducible Systems

[00343] In some embodiments, a CRISPR enzyme may form a component of an inducible system. The inducible nature of the system would allow for spatiotemporal control of gene editing or gene expression using a form of energy. The form of energy may include but is not limited to electromagnetic radiation, sound energy, chemical energy and thermal energy. Examples of inducible system include tetracycline inducible promoters (Tet-On or Tet-Off), small molecule two-hybrid transcription activations systems (FKBP, ABA, etc), or light inducible systems (Phytochrome, LOV domains, or cryptochrome). In one embodiment, the CRISPR enzyme may be a part of a Light Inducible Transcriptional Effector (LITE) to direct changes in transcriptional activity in a sequence-specific manner. The components of a light may include a CRISPR enzyme, a light-responsive cytochrome heterodimer (e.g. from *Arabidopsis thaliana*), and a transcriptional activation/repression domain. Further examples of inducible DNA binding proteins and methods for their use are provided in US 61/736,465, US 61/721,283 and WO 2014/018423, which is hereby incorporated by reference in its entirety.

Self-Inactivating Systems

[00344] Once all copies of a gene in the genome of a cell have been edited, continued CRISPR/Cas9 expression in that cell is no longer necessary. Indeed, sustained expression would be undesirable in case of off-target effects at unintended genomic sites, *etc.* Thus time-limited expression would be useful. Inducible expression offers one approach, but in addition Applicants have engineered a Self-Inactivating CRISPR-Cas9 system that relies on the use of a non-coding guide target sequence within the CRISPR vector itself. Thus, after expression begins, the CRISPR system will lead to its own destruction, but before destruction is complete it will have time to edit the genomic copies of the target gene (which, with a normal point mutation in a diploid cell, requires at most two edits). Simply, the self inactivating CRISPR-Cas9 system includes additional RNA (i.e., guide RNA) that targets the coding sequence for the CRISPR enzyme itself or that targets one or more non-coding guide target sequences complementary to unique sequences present in one or more of the following:

- (a) within the promoter driving expression of the non-coding RNA elements,
- (b) within the promoter driving expression of the Cas9 gene,
- (c) within 100bp of the ATG translational start codon in the Cas9 coding sequence,
- (d) within the inverted terminal repeat (iTR) of a viral delivery vector, e.g., in the AAV genome.

[00345] Furthermore, that RNA can be delivered via a vector, e.g., a separate vector or the same vector that is encoding the CRISPR complex. When provided by a separate vector, the CRISPR RNA that targets Cas9 expression can be administered sequentially or simultaneously. When administered sequentially, the CRISPR RNA that targets Cas9 expression is to be delivered after the CRISPR RNA that is intended for e.g. gene editing or gene engineering. This period may be a period of minutes (e.g. 5 minutes, 10 minutes, 20 minutes, 30 minutes, 45 minutes, 60 minutes). This period may be a period of hours (e.g. 2 hours, 4 hours, 6 hours, 8 hours, 12 hours, 24 hours). This period may be a period of days (e.g. 2 days, 3 days, 4 days, 7 days). This period may be a period of weeks (e.g. 2 weeks, 3 weeks, 4 weeks). This period may be a period of months (e.g. 2 months, 4 months, 8 months, 12 months). This period may be a period of years (2 years, 3 years, 4 years). In this fashion, the Cas9 enzyme associates with a first gRNA/chiRNA capable of hybridizing to a first target, such as a genomic locus or loci of interest and undertakes the function(s) desired of the CRISPR-Cas9 system (e.g., gene engineering); and subsequently the Cas9 enzyme may then associate with the second gRNA/chiRNA capable of hybridizing to the sequence comprising at least part of the Cas9 or CRISPR cassette. Where the gRNA/chiRNA targets the sequences encoding expression of the Cas9 protein, the enzyme becomes impeded and the system becomes self inactivating. In the same manner, CRISPR RNA that targets Cas9 expression applied via, for example liposome, lipofection, particles, microvesicles as explained herein, may be administered sequentially or simultaneously. Similarly, self-inactivation may be used for inactivation of one or more guide RNA used to target one or more targets.

[00346] In some aspects, a single gRNA is provided that is capable of hybridization to a sequence downstream of a CRISPR enzyme start codon, whereby after a period of time there is a loss of the CRISPR enzyme expression. In some aspects, one or more gRNA(s) are provided that are capable of hybridization to one or more coding or non-coding regions of the polynucleotide encoding the CRISPR-Cas9 system, whereby after a period of time there is an inactivation of one or more, or in some cases all, of the CRISPR-Cas9 system. In some aspects of the system, and not to be limited by theory, the cell may comprise a plurality of CRISPR-Cas9 complexes, wherein a first subset of CRISPR complexes comprise a first chiRNA capable of targeting a genomic locus or loci to be edited, and a second subset of CRISPR complexes comprise at least one second chiRNA capable of targeting the polynucleotide encoding the

CRISPR-Cas9 system, wherein the first subset of CRISPR-Cas9 complexes mediate editing of the targeted genomic locus or loci and the second subset of CRISPR complexes eventually inactivate the CRISPR-Cas9 system, thereby inactivating further CRISPR-Cas9 expression in the cell.

[00347] Thus the invention provides a CRISPR-Cas9 system comprising one or more vectors for delivery to a eukaryotic cell, wherein the vector(s) encode(s): (i) a CRISPR enzyme; (ii) a first guide RNA capable of hybridizing to a target sequence in the cell; (iii) a second guide RNA capable of hybridizing to one or more target sequence(s) in the vector which encodes the CRISPR enzyme; (iv) at least one tracr mate sequence; and (v) at least one tracr sequence. The first and second complexes can use the same tracr and tracr mate, thus differing only by the guide sequence, wherein, when expressed within the cell: the first guide RNA directs sequence-specific binding of a first CRISPR complex to the target sequence in the cell; the second guide RNA directs sequence-specific binding of a second CRISPR complex to the target sequence in the vector which encodes the CRISPR enzyme; the CRISPR complexes comprise (a) a tracr mate sequence hybridised to a tracr sequence and (b) a CRISPR enzyme bound to a guide RNA, such that a guide RNA can hybridize to its target sequence; and the second CRISPR complex inactivates the CRISPR-Cas9 system to prevent continued expression of the CRISPR enzyme by the cell.

[00348] Further characteristics of the vector(s), the encoded enzyme, the guide sequences, *etc.* are disclosed elsewhere herein. For instance, one or both of the guide sequence(s) can be part of a chiRNA sequence which provides the guide, tracr mate and tracr sequences within a single RNA, such that the system can encode (i) a CRISPR enzyme; (ii) a first chiRNA comprising a sequence capable of hybridizing to a first target sequence in the cell, a first tracr mate sequence, and a first tracr sequence; (iii) a second guide RNA capable of hybridizing to the vector which encodes the CRISPR enzyme, a second tracr mate sequence, and a second tracr sequence. Similarly, the enzyme can include one or more NLS, *etc.*

[00349] The various coding sequences (CRISPR enzyme, guide RNAs, tracr and tracr mate) can be included on a single vector or on multiple vectors. For instance, it is possible to encode the enzyme on one vector and the various RNA sequences on another vector, or to encode the enzyme and one chiRNA on one vector, and the remaining chiRNA on another vector, or any other permutation. In general, a system using a total of one or two different vectors is preferred.

[00350] Where multiple vectors are used, it is possible to deliver them in unequal numbers, and ideally with an excess of a vector which encodes the first guide RNA relative to the second guide RNA, thereby assisting in delaying final inactivation of the CRISPR system until genome editing has had a chance to occur.

[00351] The first guide RNA can target any target sequence of interest within a genome, as described elsewhere herein. The second guide RNA targets a sequence within the vector which encodes the CRISPR Cas9 enzyme, and thereby inactivates the enzyme's expression from that vector. Thus the target sequence in the vector must be capable of inactivating expression. Suitable target sequences can be, for instance, near to or within the translational start codon for the Cas9 coding sequence, in a non-coding sequence in the promoter driving expression of the non-coding RNA elements, within the promoter driving expression of the Cas9 gene, within 100bp of the ATG translational start codon in the Cas9 coding sequence, and/or within the inverted terminal repeat (iTR) of a viral delivery vector, e.g., in the AAV genome. A double stranded break near this region can induce a frame shift in the Cas9 coding sequence, causing a loss of protein expression. An alternative target sequence for the "self-inactivating" guide RNA would aim to edit/inactivate regulatory regions/sequences needed for the expression of the CRISPR-Cas9 system or for the stability of the vector. For instance, if the promoter for the Cas9 coding sequence is disrupted then transcription can be inhibited or prevented. Similarly, if a vector includes sequences for replication, maintenance or stability then it is possible to target these. For instance, in a AAV vector a useful target sequence is within the iTR. Other useful sequences to target can be promoter sequences, polyadenylation sites, etc.

[00352] Furthermore, if the guide RNAs are expressed in array format, the "self-inactivating" guide RNAs that target both promoters simultaneously will result in the excision of the intervening nucleotides from within the CRISPR-Cas9 expression construct, effectively leading to its complete inactivation. Similarly, excision of the intervening nucleotides will result where the guide RNAs target both ITRs, or targets two or more other CRISPR-Cas9 components simultaneously. Self-inactivation as explained herein is applicable, in general, with CRISPR-Cas9 systems in order to provide regulation of the CRISPR-Cas9. For example, self-inactivation as explained herein may be applied to the CRISPR repair of mutations, for example expansion disorders, as explained herein. As a result of this self-inactivation, CRISPR repair is only transiently active.

[00353] Addition of non-targeting nucleotides to the 5' end (e.g. 1 – 10 nucleotides, preferably 1 – 5 nucleotides) of the “self-inactivating” guide RNA can be used to delay its processing and/or modify its efficiency as a means of ensuring editing at the targeted genomic locus prior to CRISPR-Cas9 shutdown.

[00354] In one aspect of the self-inactivating AAV-CRISPR-Cas9 system, plasmids that co-express one or more sgRNA targeting genomic sequences of interest (e.g. 1-2, 1-5, 1-10, 1 -15, 1-20, 1-30) may be established with “self-inactivating” sgRNAs that target an SpCas9 sequence at or near the engineered ATG start site (e.g. within 5 nucleotides, within 15 nucleotides, within 30 nucleotides, within 50 nucleotides, within 100 nucleotides). A regulatory sequence in the U6 promoter region can also be targeted with an sgRNA. The U6-driven sgRNAs may be designed in an array format such that multiple sgRNA sequences can be simultaneously released. When first delivered into target tissue/cells (left cell) sgRNAs begin to accumulate while Cas9 levels rise in the nucleus. Cas9 complexes with all of the sgRNAs to mediate genome editing and self-inactivation of the CRISPR-Cas9 plasmids.

[00355] One aspect of a self-inactivating CRISPR-Cas9 system is expression of singly or in tandem array format from 1 up to 4 or more different guide sequences; e.g. up to about 20 or about 30 guides sequences. Each individual self inactivating guide sequence may target a different target. Such may be processed from, e.g. one chimeric pol3 transcript. Pol3 promoters such as U6 or H1 promoters may be used. Pol2 promoters such as those mentioned throughout herein. Inverted terminal repeat (iTR) sequences may flank the Pol3 promoter - sgRNA(s)-Pol2 promoter- Cas9.

[00356] One aspect of a chimeric, tandem array transcript is that one or more guide(s) edit the one or more target(s) while one or more self inactivating guides inactivate the CRISPR/Cas9 system. Thus, for example, the described CRISPR-Cas9 system for repairing expansion disorders may be directly combined with the self-inactivating CRISPR-Cas9 system described herein. Such a system may, for example, have two guides directed to the target region for repair as well as at least a third guide directed to self-inactivation of the CRISPR-Cas9. Reference is made to Application Ser. No. PCT/US2014/069897, entitled “Compositions And Methods Of Use Of Crispr-Cas9 Systems In Nucleotide Repeat Disorders,” published Dec. 12, 2014 as WO/2015/089351.

[00357] The guideRNA may be a control guide. For example it may be engineered to target a nucleic acid sequence encoding the CRISPR Enzyme itself, as described in US2015232881A1, the disclosure of which is hereby incorporated by reference. In some embodiments, a system or composition may be provided with just the guideRNA engineered to target the nucleic acid sequence encoding the CRISPR Enzyme. In addition, the system or composition may be provided with the guideRNA engineered to target the nucleic acid sequence encoding the CRISPR Enzyme, as well as nucleic acid sequence encoding the CRISPR Enzyme and, optionally a second guide RNA and, further optionally, a repair template. The second guideRNA may be the primary target of the CRISPR system or composition (such a therapeutic, diagnostic, knock out etc. as defined herein). In this way, the system or composition is self-inactivating. This is exemplified in relation to Cas9 in US2015232881A1 (also published as WO2015070083 (A1), referenced elsewhere herein).

Kits

[00358] In one aspect, the invention provides kits containing any one or more of the elements disclosed in the above methods and compositions. In some embodiments, the kit comprises a vector system as taught herein and instructions for using the kit. Elements may be provided individually or in combinations, and may be provided in any suitable container, such as a vial, a bottle, or a tube. The kits may include the sgRNA and the unbound protector strand as described herein. The kits may include the sgRNA with the protector strand bound to at least partially to the guide sequence (i.e. pgRNA). Thus the kits may include the pgRNA in the form of a partially double stranded nucleotide sequence as described here. In some embodiments, the kit includes instructions in one or more languages, for example in more than one language. The instructions may be specific to the applications and methods described herein.

[00359] In some embodiments, a kit comprises one or more reagents for use in a process utilizing one or more of the elements described herein. Reagents may be provided in any suitable container. For example, a kit may provide one or more reaction or storage buffers. Reagents may be provided in a form that is usable in a particular assay, or in a form that requires addition of one or more other components before use (e.g. in concentrate or lyophilized form). A buffer can be any buffer, including but not limited to a sodium carbonate buffer, a sodium bicarbonate buffer, a borate buffer, a Tris buffer, a MOPS buffer, a HEPES buffer, and combinations thereof. In some embodiments, the buffer is alkaline. In some embodiments, the

buffer has a pH from about 7 to about 10. In some embodiments, the kit comprises one or more oligonucleotides corresponding to a guide sequence for insertion into a vector so as to operably link the guide sequence and a regulatory element. In some embodiments, the kit comprises a homologous recombination template polynucleotide. In some embodiments, the kit comprises one or more of the vectors and/or one or more of the polynucleotides described herein. The kit may advantageously allow to provide all elements of the systems of the invention.

[00360] In one aspect, the invention provides methods for using one or more elements of a CRISPR system. The CRISPR complex of the invention provides an effective means for modifying a target polynucleotide. The CRISPR complex of the invention has a wide variety of utility including modifying (e.g., deleting, inserting, translocating, inactivating, activating) a target polynucleotide in a multiplicity of cell types. As such the CRISPR complex of the invention has a broad spectrum of applications in, e.g., gene therapy, drug screening, disease diagnosis, and prognosis. An exemplary CRISPR complex comprises a CRISPR effector protein complexed with a guide sequence hybridized to a target sequence within the target polynucleotide. In certain embodiments, a direct repeat sequence is linked to the guide sequence.

[00361] In one embodiment, this invention provides a method of cleaving a target polynucleotide. The method comprises modifying a target polynucleotide using a CRISPR complex that binds to the target polynucleotide and effect cleavage of said target polynucleotide. Typically, the CRISPR complex of the invention, when introduced into a cell, creates a break (e.g., a single or a double strand break) in the genome sequence. For example, the method can be used to cleave a disease gene in a cell.

[00362] The break created by the CRISPR complex can be repaired by a repair processes such as the error prone non-homologous end joining (NHEJ) pathway or the high fidelity homology directed repair (HDR). During these repair process, an exogenous polynucleotide template can be introduced into the genome sequence. In some methods, the HDR process is used to modify genome sequence. For example, an exogenous polynucleotide template comprising a sequence to be integrated flanked by an upstream sequence and a downstream sequence is introduced into a cell. The upstream and downstream sequences share sequence similarity with either side of the site of integration in the chromosome.

[00363] Where desired, a donor polynucleotide can be DNA, e.g., a DNA plasmid, a bacterial artificial chromosome (BAC), a yeast artificial chromosome (YAC), a viral vector, a linear piece

of DNA, a PCR fragment, a naked nucleic acid, or a nucleic acid complexed with a delivery vehicle such as a liposome or poloxamer.

[00364] The exogenous polynucleotide template comprises a sequence to be integrated (e.g., a mutated gene). The sequence for integration may be a sequence endogenous or exogenous to the cell. Examples of a sequence to be integrated include polynucleotides encoding a protein or a non-coding RNA (e.g., a microRNA). Thus, the sequence for integration may be operably linked to an appropriate control sequence or sequences. Alternatively, the sequence to be integrated may provide a regulatory function.

[00365] The upstream and downstream sequences in the exogenous polynucleotide template are selected to promote recombination between the chromosomal sequence of interest and the donor polynucleotide. The upstream sequence is a nucleic acid sequence that shares sequence similarity with the genome sequence upstream of the targeted site for integration. Similarly, the downstream sequence is a nucleic acid sequence that shares sequence similarity with the chromosomal sequence downstream of the targeted site of integration. The upstream and downstream sequences in the exogenous polynucleotide template can have 75%, 80%, 85%, 90%, 95%, or 100% sequence identity with the targeted genome sequence. Preferably, the upstream and downstream sequences in the exogenous polynucleotide template have about 95%, 96%, 97%, 98%, 99%, or 100% sequence identity with the targeted genome sequence. In some methods, the upstream and downstream sequences in the exogenous polynucleotide template have about 99% or 100% sequence identity with the targeted genome sequence.

[00366] An upstream or downstream sequence may comprise from about 20 bp to about 2500 bp, for example, about 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, or 2500 bp. In some methods, the exemplary upstream or downstream sequence have about 200 bp to about 2000 bp, about 600 bp to about 1000 bp, or more particularly about 700 bp to about 1000 bp.

[00367] In some methods, the exogenous polynucleotide template may further comprise a marker. Such a marker may make it easy to screen for targeted integrations. Examples of suitable markers include restriction sites, fluorescent proteins, or selectable markers. The exogenous polynucleotide template of the invention can be constructed using recombinant techniques (see, for example, Sambrook et al., 2001 and Ausubel et al., 1996).

[00368] In an exemplary method for modifying a target polynucleotide by integrating an exogenous polynucleotide template, a double stranded break is introduced into the genome sequence by the CRISPR complex, the break is repaired via homologous recombination an exogenous polynucleotide template such that the template is integrated into the genome. The presence of a double-stranded break facilitates integration of the template.

[00369] In other embodiments, this invention provides a method of modifying expression of a polynucleotide in a eukaryotic cell. The method comprises increasing or decreasing expression of a target polynucleotide by using a CRISPR complex that binds to the polynucleotide.

[00370] In some methods, a target polynucleotide can be inactivated to effect the modification of the expression in a cell. For example, upon the binding of a CRISPR complex to a target sequence in a cell, the target polynucleotide is inactivated such that the sequence is not transcribed, the coded protein is not produced, or the sequence does not function as the wild-type sequence does. For example, a protein or microRNA coding sequence may be inactivated such that the protein is not produced.

[00371] In some methods, a control sequence can be inactivated such that it no longer functions as a control sequence. As used herein, “control sequence” refers to any nucleic acid sequence that effects the transcription, translation, or accessibility of a nucleic acid sequence. Examples of a control sequence include, a promoter, a transcription terminator, and an enhancer are control sequences. The inactivated target sequence may include a deletion mutation (i.e., deletion of one or more nucleotides), an insertion mutation (i.e., insertion of one or more nucleotides), or a nonsense mutation (i.e., substitution of a single nucleotide for another nucleotide such that a stop codon is introduced). In some methods, the inactivation of a target sequence results in “knockout” of the target sequence.

Exemplary Methods of Using of CRISPR Cas9 System

[00372] The invention provides a non-naturally occurring or engineered composition, or one or more polynucleotides encoding components of said composition, or vector or delivery systems comprising one or more polynucleotides encoding components of said composition for use in a modifying a target cell *in vivo*, *ex vivo* or *in vitro* and, may be conducted in a manner alters the cell such that once modified the progeny or cell line of the CRISPR modified cell retains the altered phenotype. The modified cells and progeny may be part of a multi-cellular organism such as a plant or animal with *ex vivo* or *in vivo* application of CRISPR system to desired cell

types. The CRISPR invention may be a therapeutic method of treatment. The therapeutic method of treatment may comprise gene or genome editing, or gene therapy.

Modifying a Target with CRISPR-Cas9 System or Complex

[00373] In one aspect, the invention provides for methods of modifying a target polynucleotide in a eukaryotic cell, which may be *in vivo*, *ex vivo* or *in vitro*. In some embodiments, the method comprises sampling a cell or population of cells from a human or non-human animal, and modifying the cell or cells. Culturing may occur at any stage *ex vivo*. The cell or cells may even be re-introduced into the non-human animal or plant. For re-introduced cells it is particularly preferred that the cells are stem cells.

[00374] In some embodiments, the method comprises allowing a CRISPR complex to bind to the target polynucleotide to effect cleavage of said target polynucleotide thereby modifying the target polynucleotide, wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized or hybridizable to a target sequence within said target polynucleotide, wherein said guide sequence is linked to a tracr mate sequence which in turn hybridizes to a tracr sequence.

[00375] In one aspect, the invention provides a method of modifying expression of a polynucleotide in a eukaryotic cell. In some embodiments, the method comprises allowing a CRISPR complex to bind to the polynucleotide such that said binding results in increased or decreased expression of said polynucleotide; wherein the CRISPR complex comprises a CRISPR enzyme complexed with a guide sequence hybridized or hybridizable to a target sequence within said polynucleotide, wherein said guide sequence is linked to a tracr mate sequence which in turn hybridizes to a tracr sequence. Similar considerations and conditions apply as above for methods of modifying a target polynucleotide. In fact, these sampling, culturing and re-introduction options apply across the aspects of the present invention.

[00376] Indeed, in any aspect of the invention, the CRISPR complex may comprise a CRISPR enzyme complexed with a guide sequence hybridized or hybridizable to a target sequence, wherein said guide sequence may be linked to a tracr mate sequence which in turn may hybridize to a tracr sequence.

[00377] Similar considerations and conditions apply as above for methods of modifying a target polynucleotide. Thus in any of the non-naturally-occurring CRISPR enzymes described herein comprise at least one modification and whereby the enzyme has certain improved

capabilities. In particular, any of the enzymes are capable of forming a CRISPR complex with a guide RNA. When such a complex forms, the guide RNA is capable of binding to a target polynucleotide sequence and the enzyme is capable of modifying a target locus. In addition, the enzyme in the CRISPR complex has reduced capability of modifying one or more off-target loci as compared to an unmodified enzyme.

[00378] In addition, the modified CRISPR enzymes described herein encompass enzymes whereby in the CRISPR complex the enzyme has increased capability of modifying the one or more target loci as compared to an unmodified enzyme. Such function may be provided separate to or provided in combination with the above-described function of reduced capability of modifying one or more off-target loci. Any such enzymes may be provided with any of the further modifications to the CRISPR enzyme as described herein, such as in combination with any activity provided by one or more associated heterologous functional domains, any further mutations to reduce nuclease activity and the like.

[00379] In advantageous embodiments of the invention, the modified CRISPR enzyme is provided with reduced capability of modifying one or more off-target loci as compared to an unmodified enzyme and increased capability of modifying the one or more target loci as compared to an unmodified enzyme. In combination with further modifications to the enzyme, significantly enhanced specificity may be achieved. For example, combination of such advantageous embodiments with one or more additional mutations is provided wherein the one or more additional mutations are in one or more catalytically active domains. Such further catalytic mutations may confer nickase functionality as described in detail elsewhere herein. In such enzymes, enhanced specificity may be achieved due to an improved specificity in terms of enzyme activity.

[00380] Modifications to reduce off-target effects and/or enhance on-target effects as described above may be made to amino acid residues located in a positively-charged region/groove situated between the RuvC-III and HNH domains. It will be appreciated that any of the functional effects described above may be achieved by modification of amino acids within the aforementioned groove but also by modification of amino acids adjacent to or outside of that groove.

[00381] Additional functionalities which may be engineered into modified CRISPR enzymes as described herein include the following. 1. modified CRISPR enzymes that disrupt

DNA:protein interactions without affecting protein tertiary or secondary structure. This includes residues that contact any part of the RNA:DNA duplex. 2. modified CRISPR enzymes that weaken intra-protein interactions holding Cas9 in conformation essential for nuclease cutting in response to DNA binding (on or off target). For example: a modification that mildly inhibits, but still allows, the nuclease conformation of the HNH domain (positioned at the scissile phosphate). 3. modified CRISPR enzymes that strengthen intra-protein interactions holding Cas9 in a conformation inhibiting nuclease activity in response to DNA binding (on or off targets). For example: a modification that stabilizes the HNH domain in a conformation away from the scissile phosphate. Any such additional functional enhancement may be provided in combination with any other modification to the CRISPR enzyme as described in detail elsewhere herein.

[00382] Any of the herein described improved functionalities may be made to any CRISPR enzyme, such as a Cas9 enzyme. Cas9 enzymes described herein are derived from Cas9 enzymes from *S. pyogenes* and *S. aureus*. However, it will be appreciated that any of the functionalities described herein may be engineered into Cas9 enzymes from other orthologs, including chimeric enzymes comprising fragments from multiple orthologs.

Nucleic acids, amino acids and proteins, Regulatory sequences, Vectors, etc.

[00383] The invention uses nucleic acids to bind target DNA sequences. This is advantageous as nucleic acids are much easier and cheaper to produce than proteins, and the specificity can be varied according to the length of the stretch where homology is sought. Complex 3-D positioning of multiple fingers, for example is not required. The terms “polynucleotide”, “nucleotide”, “nucleotide sequence”, “nucleic acid” and “oligonucleotide” are used interchangeably. They refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Polynucleotides may have any three dimensional structure, and may perform any function, known or unknown. The following are non-limiting examples of polynucleotides: coding or non-coding regions of a gene or gene fragment, loci (locus) defined from linkage analysis, exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, short interfering RNA (siRNA), short-hairpin RNA (shRNA), micro-RNA (miRNA), ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers. The term also encompasses nucleic-acid-like

structures with synthetic backbones, see, e.g., Eckstein, 1991; Baserga et al., 1992; Milligan, 1993; WO 97/03211; WO 96/39154; Mata, 1997; Strauss-Soukup, 1997; and Samstag, 1996. A polynucleotide may comprise one or more modified nucleotides, such as methylated nucleotides and nucleotide analogs. If present, modifications to the nucleotide structure may be imparted before or after assembly of the polymer. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labeling component. As used herein the term "wild type" is a term of the art understood by skilled persons and means the typical form of an organism, strain, gene or characteristic as it occurs in nature as distinguished from mutant or variant forms. A "wild type" can be a base line. As used herein the term "variant" should be taken to mean the exhibition of qualities that have a pattern that deviates from what occurs in nature. The terms "non-naturally occurring" or "engineered" are used interchangeably and indicate the involvement of the hand of man. The terms, when referring to nucleic acid molecules or polypeptides mean that the nucleic acid molecule or the polypeptide is at least substantially free from at least one other component with which they are naturally associated in nature and as found in nature. "Complementarity" refers to the ability of a nucleic acid to form hydrogen bond(s) with another nucleic acid sequence by either traditional Watson-Crick base pairing or other non-traditional types. A percent complementarity indicates the percentage of residues in a nucleic acid molecule which can form hydrogen bonds (e.g., Watson-Crick base pairing) with a second nucleic acid sequence (e.g., 5, 6, 7, 8, 9, 10 out of 10 being 50%, 60%, 70%, 80%, 90%, and 100% complementary). "Perfectly complementary" means that all the contiguous residues of a nucleic acid sequence will hydrogen bond with the same number of contiguous residues in a second nucleic acid sequence. "Substantially complementary" as used herein refers to a degree of complementarity that is at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100% over a region of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, or more nucleotides, or refers to two nucleic acids that hybridize under stringent conditions. As used herein, "stringent conditions" for hybridization refer to conditions under which a nucleic acid having complementarity to a target sequence predominantly hybridizes with the target sequence, and substantially does not hybridize to non-target sequences. Stringent conditions are generally sequence-dependent, and vary depending on a number of factors. In general, the longer the sequence, the higher the temperature at which the sequence specifically

hybridizes to its target sequence. Non-limiting examples of stringent conditions are described in detail in Tijssen (1993), *Laboratory Techniques In Biochemistry And Molecular Biology-Hybridization With Nucleic Acid Probes Part I, Second Chapter "Overview of principles of hybridization and the strategy of nucleic acid probe assay"*, Elsevier, N.Y. Where reference is made to a polynucleotide sequence, then complementary or partially complementary sequences are also envisaged. These are preferably capable of hybridizing to the reference sequence under highly stringent conditions. Generally, in order to maximize the hybridization rate, relatively low-stringency hybridization conditions are selected: about 20 to 25° C lower than the thermal melting point (T_m). The T_m is the temperature at which 50% of specific target sequence hybridizes to a perfectly complementary probe in solution at a defined ionic strength and pH. Generally, in order to require at least about 85% nucleotide complementarity of hybridized sequences, highly stringent washing conditions are selected to be about 5 to 15° C lower than the T_m . In order to require at least about 70% nucleotide complementarity of hybridized sequences, moderately-stringent washing conditions are selected to be about 15 to 30° C lower than the T_m . Highly permissive (very low stringency) washing conditions may be as low as 50° C below the T_m , allowing a high level of mis-matching between hybridized sequences. Those skilled in the art will recognize that other physical and chemical parameters in the hybridization and wash stages can also be altered to affect the outcome of a detectable hybridization signal from a specific level of homology between target and probe sequences. Preferred highly stringent conditions comprise incubation in 50% formamide, 5×SSC, and 1% SDS at 42° C, or incubation in 5×SSC and 1% SDS at 65° C, with wash in 0.2×SSC and 0.1% SDS at 65° C. "Hybridization" refers to a reaction in which one or more polynucleotides react to form a complex that is stabilized via hydrogen bonding between the bases of the nucleotide residues. The hydrogen bonding may occur by Watson Crick base pairing, Hoogstein binding, or in any other sequence specific manner. The complex may comprise two strands forming a duplex structure, three or more strands forming a multi stranded complex, a single self-hybridizing strand, or any combination of these. A hybridization reaction may constitute a step in a more extensive process, such as the initiation of PCR, or the cleavage of a polynucleotide by an enzyme. A sequence capable of hybridizing with a given sequence is referred to as the "complement" of the given sequence. As used herein, the term "genomic locus" or "locus" (plural loci) is the specific location of a gene or DNA sequence on a chromosome. A "gene" refers to stretches of DNA or

RNA that encode a polypeptide or an RNA chain that has functional role to play in an organism and hence is the molecular unit of heredity in living organisms. For the purpose of this invention it may be considered that genes include regions which regulate the production of the gene product, whether or not such regulatory sequences are adjacent to coding and/or transcribed sequences. Accordingly, a gene includes, but is not necessarily limited to, promoter sequences, terminators, translational regulatory sequences such as ribosome binding sites and internal ribosome entry sites, enhancers, silencers, insulators, boundary elements, replication origins, matrix attachment sites and locus control regions. As used herein, "expression of a genomic locus" or "gene expression" is the process by which information from a gene is used in the synthesis of a functional gene product. The products of gene expression are often proteins, but in non-protein coding genes such as rRNA genes or tRNA genes, the product is functional RNA. The process of gene expression is used by all known life - eukaryotes (including multicellular organisms), prokaryotes (bacteria and archaea) and viruses to generate functional products to survive. As used herein "expression" of a gene or nucleic acid encompasses not only cellular gene expression, but also the transcription and translation of nucleic acid(s) in cloning systems and in any other context. As used herein, "expression" also refers to the process by which a polynucleotide is transcribed from a DNA template (such as into and mRNA or other RNA transcript) and/or the process by which a transcribed mRNA is subsequently translated into peptides, polypeptides, or proteins. Transcripts and encoded polypeptides may be collectively referred to as "gene product." If the polynucleotide is derived from genomic DNA, expression may include splicing of the mRNA in a eukaryotic cell. The terms "polypeptide", "peptide" and "protein" are used interchangeably herein to refer to polymers of amino acids of any length. The polymer may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non amino acids. The terms also encompass an amino acid polymer that has been modified; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation, such as conjugation with a labeling component. As used herein the term "amino acid" includes natural and/or unnatural or synthetic amino acids, including glycine and both the D or L optical isomers, and amino acid analogs and peptidomimetics. As used herein, the term "domain" or "protein domain" refers to a part of a protein sequence that may exist and function independently of the rest of the protein chain. As described in aspects of the invention, sequence identity is related to sequence homology.

Homology comparisons may be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs may calculate percent (%) homology between two or more sequences and may also calculate the sequence identity shared by two or more amino acid or nucleic acid sequences.

[00384] In aspects of the invention the term “guide RNA”, refers to the polynucleotide sequence comprising one or more of a putative or identified tracr sequence and a putative or identified crRNA sequence or guide sequence. In particular embodiments, the “guide RNA” comprises a putative or identified crRNA sequence or guide sequence. In further embodiments, the guide RNA does not comprise a putative or identified tracr sequence.

[00385] As used herein the term “wild type” is a term of the art understood by skilled persons and means the typical form of an organism, strain, gene or characteristic as it occurs in nature as distinguished from mutant or variant forms. A “wild type” can be a base line.

[00386] As used herein the term “variant” should be taken to mean the exhibition of qualities that have a pattern that deviates from what occurs in nature.

[00387] The terms “non-naturally occurring” or “engineered” are used interchangeably and indicate the involvement of the hand of man. The terms, when referring to nucleic acid molecules or polypeptides mean that the nucleic acid molecule or the polypeptide is at least substantially free from at least one other component with which they are naturally associated in nature and as found in nature. In all aspects and embodiments, whether they include these terms or not, it will be understood that, preferably, they may be optional and thus preferably included or not preferably not included. Furthermore, the terms “non-naturally occurring” and “engineered” may be used interchangeably and so can therefore be used alone or in combination and one or other may replace mention of both together. In particular, “engineered” is preferred in place of “non-naturally occurring” or “non-naturally occurring and/or engineered.”

[00388] Sequence homologies may be generated by any of a number of computer programs known in the art, for example BLAST or FASTA, etc. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A.; Devereux et al., 1984, *Nucleic Acids Research* 12:387). Examples of other software that may perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel et al., 1999 *ibid* – Chapter 18), FASTA (Atschul et al., 1990, *J. Mol. Biol.*, 403-410) and the GENWORKS suite of comparison tools. Both BLAST and FASTA are available for offline and

online searching (see Ausubel et al., 1999 *ibid*, pages 7-58 to 7-60). However it is preferred to use the GCG Bestfit program. Percentage (%) sequence homology may be calculated over contiguous sequences, i.e., one sequence is aligned with the other sequence and each amino acid or nucleotide in one sequence is directly compared with the corresponding amino acid or nucleotide in the other sequence, one residue at a time. This is called an “ungapped” alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues. Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion may cause the following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without unduly penalizing the overall homology or identity score. This is achieved by inserting “gaps” in the sequence alignment to try to maximize local homology or identity. However, these more complex methods assign “gap penalties” to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible - reflecting higher relatedness between the two compared sequences - may achieve a higher score than one with many gaps. “Affinity gap costs” are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties may, of course, produce optimized alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons. For example, when using the GCG Wisconsin Bestfit package the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension. Calculation of maximum % homology therefore first requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (Devereux et al., 1984 *Nuc. Acids Research* 12 p387). Examples of other software that may perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel et al., 1999 *Short Protocols in Molecular Biology*, 4th Ed. – Chapter 18), FASTA (Altschul et al., 1990 *J. Mol. Biol.* 403-410) and the GENWORKS suite of comparison tools. Both BLAST and FASTA are available for offline and online searching (see Ausubel et al., 1999, *Short Protocols in Molecular*

Biology, pages 7-58 to 7-60). However, for some applications, it is preferred to use the GCG Bestfit program. A new tool, called BLAST 2 Sequences is also available for comparing protein and nucleotide sequences (see *FEMS Microbiol Lett.* 1999 174(2): 247-50; *FEMS Microbiol Lett.* 1999 177(1): 187-8 and the website of the National Center for Biotechnology information at the website of the National Institutes for Health). Although the final % homology may be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pair-wise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of programs. GCG Wisconsin programs generally use either the public default values or a custom symbol comparison table, if supplied (see user manual for further details). For some applications, it is preferred to use the public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62. Alternatively, percentage homologies may be calculated using the multiple alignment feature in DNASIS™ (Hitachi Software), based on an algorithm, analogous to CLUSTAL (Higgins DG & Sharp PM (1988), *Gene* 73(1), 237-244). Once the software has produced an optimal alignment, it is possible to calculate % homology, preferably % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result. The sequences may also have deletions, insertions or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent substance. Deliberate amino acid substitutions may be made on the basis of similarity in amino acid properties (such as polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues) and it is therefore useful to group amino acids together in functional groups. Amino acids may be grouped together based on the properties of their side chains alone. However, it is more useful to include mutation data as well. The sets of amino acids thus derived are likely to be conserved for structural reasons. These sets may be described in the form of a Venn diagram (Livingstone C.D. and Barton G.J. (1993) "Protein sequence alignments: a strategy for the hierarchical analysis of residue conservation" *Comput. Appl. Biosci.* 9: 745-756) (Taylor W.R. (1986) "The classification of amino acid conservation" *J. Theor. Biol.* 119; 205-218). Conservative substitutions may be made, for example according to the table below which describes a generally accepted Venn diagram grouping of amino acids.

Set		Sub-set	
Hydrophobic	F W Y H K M I L V A G C	Aromatic	F W Y H
		Aliphatic	I L V
Polar	W Y H K R E D C S T N Q	Charged	H K R E D
		Positively charged	H K R
		Negatively charged	E D
Small	V C A G S P T N D	Tiny	A G S

[00389] The terms “subject,” “individual,” and “patient” are used interchangeably herein to refer to a vertebrate, preferably a mammal, more preferably a human. Mammals include, but are not limited to, murines, simians, humans, farm animals, sport animals, and pets. Tissues, cells and their progeny of a biological entity obtained in vivo or cultured in vitro are also encompassed.

[00390] The terms “therapeutic agent”, “therapeutic capable agent” or “treatment agent” are used interchangeably and refer to a molecule or compound that confers some beneficial effect upon administration to a subject. The beneficial effect includes enablement of diagnostic determinations; amelioration of a disease, symptom, disorder, or pathological condition; reducing or preventing the onset of a disease, symptom, disorder or condition; and generally counteracting a disease, symptom, disorder or pathological condition.

[00391] As used herein, “treatment” or “treating,” or “palliating” or “ameliorating” are used interchangeably. These terms refer to an approach for obtaining beneficial or desired results including but not limited to a therapeutic benefit and/or a prophylactic benefit. By therapeutic benefit is meant any therapeutically relevant improvement in or effect on one or more diseases, conditions, or symptoms under treatment. For prophylactic benefit, the compositions may be administered to a subject at risk of developing a particular disease, condition, or symptom, or to a subject reporting one or more of the physiological symptoms of a disease, even though the disease, condition, or symptom may not have yet been manifested.

[00392] The term “effective amount” or “therapeutically effective amount” refers to the amount of an agent that is sufficient to effect beneficial or desired results. The therapeutically

effective amount may vary depending upon one or more of: the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art. The term also applies to a dose that will provide an image for detection by any one of the imaging methods described herein. The specific dose may vary depending on one or more of: the particular agent chosen, the dosing regimen to be followed, whether it is administered in combination with other compounds, timing of administration, the tissue to be imaged, and the physical delivery system in which it is carried.

[00393] The practice of the present invention employs, unless otherwise indicated, conventional techniques of immunology, biochemistry, chemistry, molecular biology, microbiology, cell biology, genomics and recombinant DNA, which are within the skill of the art. See Sambrook, Fritsch and Maniatis, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd edition (1989); *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY* (F. M. Ausubel, et al. eds., (1987)); the series *METHODS IN ENZYMOLOGY* (Academic Press, Inc.): *PCR 2: A PRACTICAL APPROACH* (M.J. MacPherson, B.D. Hames and G.R. Taylor eds. (1995)), Harlow and Lane, eds. (1988) *ANTIBODIES, A LABORATORY MANUAL*, and *ANIMAL CELL CULTURE* (R.I. Freshney, ed. (1987)).

[00394] Several aspects of the invention relate to vector systems comprising one or more vectors, or vectors as such. Vectors can be designed for expression of CRISPR transcripts (e.g. nucleic acid transcripts, proteins, or enzymes) in prokaryotic or eukaryotic cells. For example, CRISPR transcripts can be expressed in bacterial cells such as *Escherichia coli*, insect cells (using baculovirus expression vectors), yeast cells, or mammalian cells. Suitable host cells are discussed further in Goeddel, *GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY* 185, Academic Press, San Diego, Calif. (1990). Alternatively, the recombinant expression vector can be transcribed and translated in vitro, for example using T7 promoter regulatory sequences and T7 polymerase.

[00395] Embodiments of the invention include sequences (both polynucleotide or polypeptide) which may comprise homologous substitution (substitution and replacement are both used herein to mean the interchange of an existing amino acid residue or nucleotide, with an alternative residue or nucleotide) that may occur i.e., like-for-like substitution in the case of amino acids such as basic for basic, acidic for acidic, polar for polar, etc. Non-homologous

substitution may also occur i.e., from one class of residue to another or alternatively involving the inclusion of unnatural amino acids such as ornithine (hereinafter referred to as Z), diaminobutyric acid ornithine (hereinafter referred to as B), norleucine ornithine (hereinafter referred to as O), pyriylalanine, thienylalanine, naphthylalanine and phenylglycine. Variant amino acid sequences may include suitable spacer groups that may be inserted between any two amino acid residues of the sequence including alkyl groups such as methyl, ethyl or propyl groups in addition to amino acid spacers such as glycine or β -alanine residues. A further form of variation, which involves the presence of one or more amino acid residues in peptoid form, may be well understood by those skilled in the art. For the avoidance of doubt, “the peptoid form” is used to refer to variant amino acid residues wherein the α -carbon substituent group is on the residue’s nitrogen atom rather than the α -carbon. Processes for preparing peptides in the peptoid form are known in the art, for example Simon RJ et al., *PNAS* (1992) 89(20), 9367-9371 and Horwell DC, *Trends Biotechnol.* (1995) 13(4), 132-134.

[00396] Homology modelling: Corresponding residues in other Cas9 orthologs can be identified by the methods of Zhang et al., 2012 (*Nature*; 490(7421): 556-60) and Chen et al., 2015 (*PLoS Comput Biol*; 11(5): e1004248)—a computational protein-protein interaction (PPI) method to predict interactions mediated by domain-motif interfaces. PrePPI (Predicting PPI), a structure based PPI prediction method, combines structural evidence with non-structural evidence using a Bayesian statistical framework. The method involves taking a pair a query proteins and using structural alignment to identify structural representatives that correspond to either their experimentally determined structures or homology models. Structural alignment is further used to identify both close and remote structural neighbors by considering global and local geometric relationships. Whenever two neighbors of the structural representatives form a complex reported in the Protein Data Bank, this defines a template for modelling the interaction between the two query proteins. Models of the complex are created by superimposing the representative structures on their corresponding structural neighbor in the template. This approach is further described in Dey et al., 2013 (*Prot Sci*; 22: 359-66).

[00397] For purpose of this invention, amplification means any method employing a primer and a polymerase capable of replicating a target sequence with reasonable fidelity. Amplification may be carried out by natural or recombinant DNA polymerases such as

TaqGold™, T7 DNA polymerase, Klenow fragment of E.coli DNA polymerase, and reverse transcriptase. A preferred amplification method is PCR.

[00398] In certain aspects the invention involves vectors. As used herein, a “vector” is a tool that allows or facilitates the transfer of an entity from one environment to another. It is a replicon, such as a plasmid, phage, or cosmid, into which another DNA segment may be inserted so as to bring about the replication of the inserted segment. Generally, a vector is capable of replication when associated with the proper control elements. In general, the term “vector” refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. Vectors include, but are not limited to, nucleic acid molecules that are single-stranded, double-stranded, or partially double-stranded; nucleic acid molecules that comprise one or more free ends, no free ends (e.g. circular); nucleic acid molecules that comprise DNA, RNA, or both; and other varieties of polynucleotides known in the art. One type of vector is a “plasmid,” which refers to a circular double stranded DNA loop into which additional DNA segments can be inserted, such as by standard molecular cloning techniques. Another type of vector is a viral vector, wherein virally-derived DNA or RNA sequences are present in the vector for packaging into a virus (e.g. retroviruses, replication defective retroviruses, adenoviruses, replication defective adenoviruses, and adeno-associated viruses (AAVs)). Viral vectors also include polynucleotides carried by a virus for transfection into a host cell. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g. bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively-linked. Such vectors are referred to herein as “expression vectors.” Common expression vectors of utility in recombinant DNA techniques are often in the form of plasmids.

[00399] Recombinant expression vectors can comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory elements, which may be selected on the basis of the host cells to be used for expression, that is operatively-linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, “operably linked” is intended to mean that the nucleotide sequence of interest is linked to the regulatory element(s) in a manner that

allows for expression of the nucleotide sequence (e.g. in an in vitro transcription/translation system or in a host cell when the vector is introduced into the host cell). With regards to recombination and cloning methods, mention is made of U.S. patent application 10/815,730, published September 2, 2004 as US 2004-0171156 A1, the contents of which are herein incorporated by reference in their entirety.

[00400] Aspects of the invention relate to bicistronic vectors for chimeric RNA and Cas9. Bicistronic expression vectors for chimeric RNA and Cas9 are preferred. In general and particularly in this embodiment Cas9 is preferably driven by the CBh promoter. The chimeric RNA may preferably be driven by a Pol III promoter, such as a U6 promoter. Ideally the two are combined. The chimeric guide RNA typically comprises, consists essentially of, or consists of a 20bp guide sequence (Ns) and this may be joined to the tracr sequence (running from the first “U” of the lower strand to the end of the transcript). The tracr sequence may be truncated at various positions as indicated. The guide and tracr sequences are separated by the tracr-mate sequence, which may be GUUUUAGAGCUA. This may be followed by the loop sequence GAAA as shown. Both of these are preferred examples. Applicants have demonstrated Cas9-mediated indels at the human *EMXI* and *PVALB* loci by SURVEYOR assays. ChiRNAs are indicated by their “+n” designation, and crRNA refers to a hybrid RNA where guide and tracr sequences are expressed as separate transcripts. Throughout this application, chimeric RNA may also be called single guide, or synthetic guide RNA (sgRNA).

[00401] In some embodiments, a loop in the guide RNA is provided. This may be a stem loop or a tetra loop. The loop is preferably GAAA, but it is not limited to this sequence or indeed to being only 4bp in length. Indeed, preferred loop forming sequences for use in hairpin structures are four nucleotides in length, and most preferably have the sequence GAAA. However, longer or shorter loop sequences may be used, as may alternative sequences. The sequences preferably include a nucleotide triplet (for example, AAA), and an additional nucleotide (for example C or G). Examples of loop forming sequences include CAAA and AAAG. In practicing any of the methods disclosed herein, a suitable vector can be introduced to a cell or an embryo via one or more methods known in the art, including without limitation, microinjection, electroporation, sonoporation, biolistics, calcium phosphate-mediated transfection, cationic transfection, liposome transfection, dendrimer transfection, heat shock transfection, nucleofection transfection, magnetofection, lipofection, impalefection, optical transfection, proprietary agent-

enhanced uptake of nucleic acids, and delivery via liposomes, immunoliposomes, virosomes, or artificial virions. In some methods, the vector is introduced into an embryo by microinjection. The vector or vectors may be microinjected into the nucleus or the cytoplasm of the embryo. In some methods, the vector or vectors may be introduced into a cell by nucleofection.

[00402] The term “regulatory element” is intended to include promoters, enhancers, internal ribosomal entry sites (IRES), and other expression control elements (e.g. transcription termination signals, such as polyadenylation signals and poly-U sequences). Such regulatory elements are described, for example, in Goeddel, *GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY* 185, Academic Press, San Diego, Calif. (1990). Regulatory elements include those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). A tissue-specific promoter may direct expression primarily in a desired tissue of interest, such as muscle, neuron, bone, skin, blood, specific organs (e.g. liver, pancreas), or particular cell types (e.g. lymphocytes). Regulatory elements may also direct expression in a temporal-dependent manner, such as in a cell-cycle dependent or developmental stage-dependent manner, which may or may not also be tissue or cell-type specific. In some embodiments, a vector comprises one or more pol III promoter (e.g. 1, 2, 3, 4, 5, or more pol III promoters), one or more pol II promoters (e.g. 1, 2, 3, 4, 5, or more pol II promoters), one or more pol I promoters (e.g. 1, 2, 3, 4, 5, or more pol I promoters), or combinations thereof. Examples of pol III promoters include, but are not limited to, U6 and H1 promoters. Examples of pol II promoters include, but are not limited to, the retroviral Rous sarcoma virus (RSV) LTR promoter (optionally with the RSV enhancer), the cytomegalovirus (CMV) promoter (optionally with the CMV enhancer) [see, e.g., Boshart et al, *Cell*, 41:521-530 (1985)], the SV40 promoter, the dihydrofolate reductase promoter, the β -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1 α promoter. Also encompassed by the term “regulatory element” are enhancer elements, such as WPRE; CMV enhancers; the R-US’ segment in LTR of HTLV-I (*Mol. Cell. Biol.*, Vol. 8(1), p. 466-472, 1988); SV40 enhancer; and the intron sequence between exons 2 and 3 of rabbit β -globin (*Proc. Natl. Acad. Sci. USA.*, Vol. 78(3), p. 1527-31, 1981). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression desired, etc. A vector can be introduced into host cells to thereby produce

transcripts, proteins, or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (e.g., clustered regularly interspersed short palindromic repeats (CRISPR) transcripts, proteins, enzymes, mutant forms thereof, fusion proteins thereof, etc.). With regards to regulatory sequences, mention is made of U.S. patent application 10/491,026, the contents of which are incorporated by reference herein in their entirety. With regards to promoters, mention is made of PCT publication WO 2011/028929 and U.S. application 12/511,940, the contents of which are incorporated by reference herein in their entirety.

[00403] Vectors can be designed for expression of CRISPR transcripts (e.g. nucleic acid transcripts, proteins, or enzymes) in prokaryotic or eukaryotic cells. For example, CRISPR transcripts can be expressed in bacterial cells such as *Escherichia coli*, insect cells (using baculovirus expression vectors), yeast cells, or mammalian cells. Suitable host cells are discussed further in Goeddel, *GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY* 185, Academic Press, San Diego, Calif. (1990). Alternatively, the recombinant expression vector can be transcribed and translated in vitro, for example using T7 promoter regulatory sequences and T7 polymerase.

[00404] Vectors may be introduced and propagated in a prokaryote or prokaryotic cell. In some embodiments, a prokaryote is used to amplify copies of a vector to be introduced into a eukaryotic cell or as an intermediate vector in the production of a vector to be introduced into a eukaryotic cell (e.g. amplifying a plasmid as part of a viral vector packaging system). In some embodiments, a prokaryote is used to amplify copies of a vector and express one or more nucleic acids, such as to provide a source of one or more proteins for delivery to a host cell or host organism. Expression of proteins in prokaryotes is most often carried out in *Escherichia coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, such as to the amino terminus of the recombinant protein. Such fusion vectors may serve one or more purposes, such as: (i) to increase expression of recombinant protein; (ii) to increase the solubility of the recombinant protein; and (iii) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition

sequences, include Factor Xa, thrombin and enterokinase. Example fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith and Johnson, 1988. *Gene* 67: 31-40), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.) that fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amrann et al., (1988) *Gene* 69:301-315) and pET 11d (Studier et al., *GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY* 185, Academic Press, San Diego, Calif. (1990) 60-89). In some embodiments, a vector is a yeast expression vector. Examples of vectors for expression in yeast *Saccharomyces cerevisiae* include pYepSec1 (Baldari, et al., 1987. *EMBO J.* 6: 229-234), pMFa (Kuijan and Herskowitz, 1982. *Cell* 30: 933-943), pJRY88 (Schultz et al., 1987. *Gene* 54: 113-123), pYES2 (Invitrogen Corporation, San Diego, Calif.), and picZ (Invitrogen Corp, San Diego, Calif.). In some embodiments, a vector drives protein expression in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., SF9 cells) include the pAc series (Smith, et al., 1983. *Mol. Cell. Biol.* 3: 2156-2165) and the pVL series (Lucklow and Summers, 1989. *Virology* 170: 31-39).

[00405] In some embodiments, a vector is capable of driving expression of one or more sequences in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, 1987. *Nature* 329: 840) and pMT2PC (Kaufman, et al., 1987. *EMBO J.* 6: 187-195). When used in mammalian cells, the expression vector's control functions are typically provided by one or more regulatory elements. For example, commonly used promoters are derived from polyoma, adenovirus 2, cytomegalovirus, simian virus 40, and others disclosed herein and known in the art. For other suitable expression systems for both prokaryotic and eukaryotic cells see, e.g., Chapters 16 and 17 of Sambrook, et al., *MOLECULAR CLONING: A LABORATORY MANUAL*. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.

[00406] In some embodiments, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert, et al., 1987. *Genes Dev.* 1: 268-277),

lymphoid-specific promoters (Calame and Eaton, 1988. *Adv. Immunol.* 43: 235-275), in particular promoters of T cell receptors (Winoto and Baltimore, 1989. *EMBO J.* 8: 729-733) and immunoglobulins (Baneiji, et al., 1983. *Cell* 33: 729-740; Queen and Baltimore, 1983. *Cell* 33: 741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle, 1989. *Proc. Natl. Acad. Sci. USA* 86: 5473-5477), pancreas-specific promoters (Edlund, et al., 1985. *Science* 230: 912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Pat. No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, e.g., the murine hox promoters (Kessel and Gruss, 1990. *Science* 249: 374-379) and the α -fetoprotein promoter (Campes and Tilghman, 1989. *Genes Dev.* 3: 537-546). With regards to these prokaryotic and eukaryotic vectors, mention is made of U.S. Patent 6,750,059, the contents of which are incorporated by reference herein in their entirety. Other embodiments of the invention may relate to the use of viral vectors, with regards to which mention is made of U.S. Patent application 13/092,085, the contents of which are incorporated by reference herein in their entirety. Tissue-specific regulatory elements are known in the art and in this regard, mention is made of U.S. Patent 7,776,321, the contents of which are incorporated by reference herein in their entirety. In some embodiments, a regulatory element is operably linked to one or more elements of a CRISPR system so as to drive expression of the one or more elements of the CRISPR system. In general, CRISPRs (Clustered Regularly Interspaced Short Palindromic Repeats), also known as SPIDRs (SPacer Interspersed Direct Repeats), constitute a family of DNA loci that are usually specific to a particular bacterial species. The CRISPR locus comprises a distinct class of interspersed short sequence repeats (SSRs) that were recognized in *E. coli* (Ishino et al., *J. Bacteriol.*, 169:5429-5433 [1987]; and Nakata et al., *J. Bacteriol.*, 171:3553-3556 [1989]), and associated genes. Similar interspersed SSRs have been identified in *Haloferax mediterranei*, *Streptococcus pyogenes*, *Anabaena*, and *Mycobacterium tuberculosis* (See, Groenen et al., *Mol. Microbiol.*, 10:1057-1065 [1993]; Hoe et al., *Emerg. Infect. Dis.*, 5:254-263 [1999]; Masepohl et al., *Biochim. Biophys. Acta* 1307:26-30 [1996]; and Mojica et al., *Mol. Microbiol.*, 17:85-93 [1995]). The CRISPR loci typically differ from other SSRs by the structure of the repeats, which have been termed short regularly spaced repeats (SRSRs) (Janssen et al., *OMICS J. Integ. Biol.*, 6:23-33 [2002]; and Mojica et al., *Mol. Microbiol.*, 36:244-246 [2000]). In general, the repeats are short elements that occur in clusters that are regularly spaced by unique intervening sequences with a substantially constant length

(Mojica et al., [2000], supra). Although the repeat sequences are highly conserved between strains, the number of interspersed repeats and the sequences of the spacer regions typically differ from strain to strain (van Embden et al., *J. Bacteriol.*, 182:2393-2401 [2000]). CRISPR loci have been identified in more than 40 prokaryotes (See e.g., Jansen et al., *Mol. Microbiol.*, 43:1565-1575 [2002]; and Mojica et al., [2005]) including, but not limited to *Aeropyrum*, *Pyrobaculum*, *Sulfolobus*, *Archaeoglobus*, *Halocarcula*, *Methanobacterium*, *Methanococcus*, *Methanosarcina*, *Methanopyrus*, *Pyrococcus*, *Picrophilus*, *Thermoplasma*, *Corynebacterium*, *Mycobacterium*, *Streptomyces*, *Aquifex*, *Porphyromonas*, *Chlorobium*, *Thermus*, *Bacillus*, *Listeria*, *Staphylococcus*, *Clostridium*, *Thermoanaerobacter*, *Mycoplasma*, *Fusobacterium*, *Azarcus*, *Chromobacterium*, *Neisseria*, *Nitrosomonas*, *Desulfovibrio*, *Geobacter*, *Myxococcus*, *Campylobacter*, *Wolinella*, *Acinetobacter*, *Erwinia*, *Escherichia*, *Legionella*, *Methylococcus*, *Pasteurella*, *Photobacterium*, *Salmonella*, *Xanthomonas*, *Yersinia*, *Treponema*, and *Thermotoga*.

[00407] In general, “nucleic acid-targeting system” as used in the present application refers collectively to transcripts and other elements involved in the expression of or directing the activity of nucleic acid-targeting CRISPR-associated (“Cas”) genes (also referred to herein as an effector protein), including sequences encoding a nucleic acid-targeting Cas9 (effector) protein and a guide RNA (comprising crRNA sequence and a trans-activating CRISPR/Cas9 system RNA (tracrRNA) sequence), or other sequences and transcripts from a nucleic acid-targeting CRISPR locus. In some embodiments, one or more elements of a nucleic acid-targeting system are derived from a Type II nucleic acid-targeting CRISPR system. In some embodiments, one or more elements of a nucleic acid-targeting system is derived from a particular organism comprising an endogenous nucleic acid-targeting CRISPR system. In general, a nucleic acid-targeting system is characterized by elements that promote the formation of a nucleic acid-targeting complex at the site of a target sequence. In the context of formation of a nucleic acid-targeting complex, “target sequence” refers to a sequence to which a guide sequence is designed to have complementarity, where hybridization between a target sequence and a guide RNA promotes the formation of a DNA or RNA-targeting complex. Full complementarity is not necessarily required, provided there is sufficient complementarity to cause hybridization and promote formation of a nucleic acid-targeting complex. A target sequence may comprise RNA polynucleotides. In some embodiments, a target sequence is located in the nucleus or cytoplasm of a cell. In some embodiments, the target sequence may be within an organelle of a eukaryotic

cell, for example, mitochondrion or chloroplast. A sequence or template that may be used for recombination into the targeted locus comprising the target sequences is referred to as an “editing template” or “editing RNA” or “editing sequence”. In aspects of the invention, an exogenous template RNA may be referred to as an editing template. In an aspect of the invention the recombination is homologous recombination.

[00408] Typically, in the context of an endogenous nucleic acid-targeting system, formation of a nucleic acid-targeting complex (comprising a guide RNA hybridized to a target sequence and complexed with one or more nucleic acid-targeting effector proteins) results in cleavage of one or both RNA strands in or near (e.g. within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, or more base pairs from) the target sequence. In some embodiments, one or more vectors driving expression of one or more elements of a nucleic acid-targeting system are introduced into a host cell such that expression of the elements of the nucleic acid-targeting system direct formation of a nucleic acid-targeting complex at one or more target sites. For example, a nucleic acid-targeting effector protein and a guide RNA could each be operably linked to separate regulatory elements on separate vectors. Alternatively, two or more of the elements expressed from the same or different regulatory elements, may be combined in a single vector, with one or more additional vectors providing any components of the nucleic acid-targeting system not included in the first vector. nucleic acid-targeting system elements that are combined in a single vector may be arranged in any suitable orientation, such as one element located 5' with respect to (“upstream” of) or 3' with respect to (“downstream” of) a second element. The coding sequence of one element may be located on the same or opposite strand of the coding sequence of a second element, and oriented in the same or opposite direction. In some embodiments, a single promoter drives expression of a transcript encoding a nucleic acid-targeting effector protein and a guide RNA embedded within one or more intron sequences (e.g. each in a different intron, two or more in at least one intron, or all in a single intron). In some embodiments, the nucleic acid-targeting effector protein and guide RNA are operably linked to and expressed from the same promoter.

[00409] In general, a guide sequence is any polynucleotide sequence having sufficient complementarity with a target polynucleotide sequence to hybridize with the target sequence and direct sequence-specific binding of a nucleic acid-targeting complex to the target sequence. In some embodiments, the degree of complementarity between a guide sequence and its

corresponding target sequence, when optimally aligned using a suitable alignment algorithm, is about or more than about 50%, 60%, 75%, 80%, 85%, 90%, 95%, 97.5%, 99%, or more. Optimal alignment may be determined with the use of any suitable algorithm for aligning sequences, non-limiting example of which include the Smith-Waterman algorithm, the Needleman-Wunsch algorithm, algorithms based on the Burrows-Wheeler Transform (e.g. the Burrows Wheeler Aligner), ClustalW, Clustal X, BLAT, Novoalign (Novocraft Technologies, ELAND (Illumina, San Diego, CA), SOAP (available at soap.genomics.org.cn), and Maq (available at maq.sourceforge.net). In some embodiments, a guide sequence is about or more than about 5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 75, or more nucleotides in length. In some embodiments, a guide sequence is less than about 75, 50, 45, 40, 35, 30, 25, 20, 15, 12, or fewer nucleotides in length. The ability of a guide sequence to direct sequence-specific binding of a nucleic acid-targeting complex to a target sequence may be assessed by any suitable assay. For example, the components of a nucleic acid-targeting system sufficient to form a nucleic acid-targeting complex, including the guide sequence to be tested, may be provided to a host cell having the corresponding target sequence, such as by transfection with vectors encoding the components of the nucleic acid-targeting CRISPR sequence, followed by an assessment of preferential cleavage within or in the vicinity of the target sequence, such as by Surveyor assay as described herein. Similarly, cleavage of a target polynucleotide sequence (or a sequence in the vicinity thereof) may be evaluated in a test tube by providing the target sequence, components of a nucleic acid-targeting complex, including the guide sequence to be tested and a control guide sequence different from the test guide sequence, and comparing binding or rate of cleavage at or in the vicinity of the target sequence between the test and control guide sequence reactions. Other assays are possible, and will occur to those skilled in the art.

[00410] A guide sequence may be selected to target any target sequence. In some embodiments, the target sequence is a sequence within a gene transcript or mRNA.

[00411] In some embodiments, the target sequence is a sequence within a genome of a cell.

[00412] In some embodiments, a guide sequence is selected to reduce the degree of secondary structure within the guide sequence. Secondary structure may be determined by any suitable polynucleotide folding algorithm. Some programs are based on calculating the minimal Gibbs free energy. An example of one such algorithm is mFold, as described by Zuker and Stiegler

(*Nucleic Acids Res.* 9 (1981), 133-148). Another example folding algorithm is the online webserver RNAfold, developed at Institute for Theoretical Chemistry at the University of Vienna, using the centroid structure prediction algorithm (see e.g. A.R. Gruber *et al.*, 2008, *Cell* 106(1): 23-24; and PA Carr and GM Church, 2009, *Nature Biotechnology* 27(12): 1151-62). Further algorithms may be found in U.S. application Serial No. TBA (attorney docket 44790.11.2022; Broad Reference BI-2013/004A); incorporated herein by reference.

[00413] In some embodiments, a recombination template is also provided. A recombination template may be a component of another vector as described herein, contained in a separate vector, or provided as a separate polynucleotide. In some embodiments, a recombination template is designed to serve as a template in homologous recombination, such as within or near a target sequence nicked or cleaved by a nucleic acid-targeting effector protein as a part of a nucleic acid-targeting complex. A template polynucleotide may be of any suitable length, such as about or more than about 10, 15, 20, 25, 50, 75, 100, 150, 200, 500, 1000, or more nucleotides in length. In some embodiments, the template polynucleotide is complementary to a portion of a polynucleotide comprising the target sequence. When optimally aligned, a template polynucleotide might overlap with one or more nucleotides of a target sequences (e.g. about or more than about 1, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100 or more nucleotides). In some embodiments, when a template sequence and a polynucleotide comprising a target sequence are optimally aligned, the nearest nucleotide of the template polynucleotide is within about 1, 5, 10, 15, 20, 25, 50, 75, 100, 200, 300, 400, 500, 1000, 5000, 10000, or more nucleotides from the target sequence.

[00414] In some embodiments, the nucleic acid-targeting effector protein is part of a fusion protein comprising one or more heterologous protein domains (e.g., about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more domains in addition to the nucleic acid-targeting effector protein). In some embodiments, the CRISPR enzyme is part of a fusion protein comprising one or more heterologous protein domains (e.g. about or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more domains in addition to the CRISPR enzyme). A CRISPR enzyme fusion protein may comprise any additional protein sequence, and optionally a linker sequence between any two domains. Examples of protein domains that may be fused to a CRISPR enzyme include, without limitation, epitope tags, reporter gene sequences, and protein domains having one or more of the following activities: methylase activity, demethylase activity, transcription activation activity,

transcription repression activity, transcription release factor activity, histone modification activity, RNA cleavage activity and nucleic acid binding activity. Non-limiting examples of epitope tags include histidine (His) tags, V5 tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Examples of reporter genes include, but are not limited to, glutathione-S-transferase (GST), horseradish peroxidase (HRP), chloramphenicol acetyltransferase (CAT) beta-galactosidase, beta-glucuronidase, luciferase, green fluorescent protein (GFP), HcRed, DsRed, cyan fluorescent protein (CFP), yellow fluorescent protein (YFP), and autofluorescent proteins including blue fluorescent protein (BFP). A CRISPR enzyme may be fused to a gene sequence encoding a protein or a fragment of a protein that bind DNA molecules or bind other cellular molecules, including but not limited to maltose binding protein (MBP), S-tag, Lex A DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. Additional domains that may form part of a fusion protein comprising a CRISPR enzyme are described in US20110059502, incorporated herein by reference. In some embodiments, a tagged CRISPR enzyme is used to identify the location of a target sequence.

[00415] In some embodiments, a CRISPR enzyme may form a component of an inducible system. The inducible nature of the system would allow for spatiotemporal control of gene editing or gene expression using a form of energy. The form of energy may include but is not limited to electromagnetic radiation, sound energy, chemical energy and thermal energy. Examples of inducible system include tetracycline inducible promoters (Tet-On or Tet-Off), small molecule two-hybrid transcription activations systems (FKBP, ABA, etc), or light inducible systems (Phytochrome, LOV domains, or cryptochrome). In one embodiment, the CRISPR enzyme may be a part of a Light Inducible Transcriptional Effector (LITE) to direct changes in transcriptional activity in a sequence-specific manner. The components of a light may include a CRISPR enzyme, a light-responsive cytochrome heterodimer (e.g. from *Arabidopsis thaliana*), and a transcriptional activation/repression domain. Further examples of inducible DNA binding proteins and methods for their use are provided in US 61/736465 and US 61/721,283 and WO 2014/018423 and US8889418, US8895308, US20140186919, US20140242700, US20140273234, US20140335620, WO2014093635, which is hereby incorporated by reference in its entirety.

[00416] In some aspects, the invention provides methods comprising delivering one or more polynucleotides, such as or one or more vectors as described herein, one or more transcripts thereof, and/or one or more proteins transcribed therefrom, to a host cell. In some aspects, the invention further provides cells produced by such methods, and organisms (such as animals, plants, or fungi) comprising or produced from such cells. In some embodiments, a nucleic acid-targeting effector protein in combination with (and optionally complexed with) a guide RNA is delivered to a cell. Conventional viral and non-viral based gene transfer methods can be used to introduce nucleic acids in mammalian cells or target tissues. Such methods can be used to administer nucleic acids encoding components of a nucleic acid-targeting system to cells in culture, or in a host organism. Non-viral vector delivery systems include DNA plasmids, RNA (e.g. a transcript of a vector described herein), naked nucleic acid, and nucleic acid complexed with a delivery vehicle, such as a liposome. Viral vector delivery systems include DNA and RNA viruses, which have either episomal or integrated genomes after delivery to the cell. For a review of gene therapy procedures, see Anderson, *Science* 256:808-813 (1992); Nabel & Felgner, *TIBTECH* 11:211-217 (1993); Mitani & Caskey, *TIBTECH* 11:162-166 (1993); Dillon, *TIBTECH* 11:167-175 (1993); Miller, *Nature* 357:455-460 (1992); Van Brunt, *Biotechnology* 6(10):1149-1154 (1988); Vigne, *Restorative Neurology and Neuroscience* 8:35-36 (1995); Kremer & Perricaudet, *British Medical Bulletin* 51(1):31-44 (1995); Haddada et al., in *Current Topics in Microbiology and Immunology*, Doerfler and Böhm (eds) (1995); and Yu et al., *Gene Therapy* 1:13-26 (1994).

[00417] Methods of non-viral delivery of nucleic acids include lipofection, nucleofection, microinjection, biolistics, virosomes, liposomes, immunoliposomes, polycation or lipid:nucleic acid conjugates, naked DNA, artificial virions, and agent-enhanced uptake of DNA. Lipofection is described in e.g., U.S. Pat. Nos. 5,049,386, 4,946,787; and 4,897,355) and lipofection reagents are sold commercially (e.g., Transfectam™ and Lipofectin™). Cationic and neutral lipids that are suitable for efficient receptor-recognition lipofection of polynucleotides include those of Felgner, WO 91/17424; WO 91/16024. Delivery can be to cells (e.g. in vitro or ex vivo administration) or target tissues (e.g. in vivo administration).

[00418] The preparation of lipid:nucleic acid complexes, including targeted liposomes such as immunolipid complexes, is well known to one of skill in the art (see, e.g., Crystal, *Science* 270:404-410 (1995); Blaese et al., *Cancer Gene Ther.* 2:291-297 (1995); Behr et al.,

Bioconjugate Chem. 5:382-389 (1994); Remy et al., Bioconjugate Chem. 5:647-654 (1994); Gao et al., Gene Therapy 2:710-722 (1995); Ahmad et al., Cancer Res. 52:4817-4820 (1992); U.S. Pat. Nos. 4,186,183, 4,217,344, 4,235,871, 4,261,975, 4,485,054, 4,501,728, 4,774,085, 4,837,028, and 4,946,787).

[00419] The use of RNA or DNA viral based systems for the delivery of nucleic acids takes advantage of highly evolved processes for targeting a virus to specific cells in the body and trafficking the viral payload to the nucleus. Viral vectors can be administered directly to patients (in vivo) or they can be used to treat cells in vitro, and the modified cells may optionally be administered to patients (ex vivo). Conventional viral based systems could include retroviral, lentivirus, adenoviral, adeno-associated and herpes simplex virus vectors for gene transfer. Integration in the host genome is possible with the retrovirus, lentivirus, and adeno-associated virus gene transfer methods, often resulting in long term expression of the inserted transgene. Additionally, high transduction efficiencies have been observed in many different cell types and target tissues.

[00420] The tropism of a retrovirus can be altered by incorporating foreign envelope proteins, expanding the potential target population of target cells. Lentiviral vectors are retroviral vectors that are able to transduce or infect non-dividing cells and typically produce high viral titers. Selection of a retroviral gene transfer system would therefore depend on the target tissue. Retroviral vectors are comprised of cis-acting long terminal repeats with packaging capacity for up to 6-10 kb of foreign sequence. The minimum cis-acting LTRs are sufficient for replication and packaging of the vectors, which are then used to integrate the therapeutic gene into the target cell to provide permanent transgene expression. Widely used retroviral vectors include those based upon murine leukemia virus (MuLV), gibbon ape leukemia virus (GaLV), Simian Immuno deficiency virus (SIV), human immuno deficiency virus (HIV), and combinations thereof (see, e.g., Buchscher et al., J. Virol. 66:2731-2739 (1992); Johann et al., J. Virol. 66:1635-1640 (1992); Somnerfelt et al., Virol. 176:58-59 (1990); Wilson et al., J. Virol. 63:2374-2378 (1989); Miller et al., J. Virol. 65:2220-2224 (1991); PCT/US94/05700). In applications where transient expression is preferred, adenoviral based systems may be used. Adenoviral based vectors are capable of very high transduction efficiency in many cell types and do not require cell division. With such vectors, high titer and levels of expression have been obtained. This vector can be produced in large quantities in a relatively simple system. Adeno-associated virus

("AAV") vectors may also be used to transduce cells with target nucleic acids, e.g., in the *in vitro* production of nucleic acids and peptides, and for *in vivo* and *ex vivo* gene therapy procedures (see, e.g., West et al., *Virology* 160:38-47 (1987); U.S. Pat. No. 4,797,368; WO 93/24641; Kotin, *Human Gene Therapy* 5:793-801 (1994); Muzyczka, *J. Clin. Invest.* 94:1351 (1994). Construction of recombinant AAV vectors are described in a number of publications, including U.S. Pat. No. 5,173,414; Tratschin et al., *Mol. Cell. Biol.* 5:3251-3260 (1985); Tratschin, et al., *Mol. Cell. Biol.* 4:2072-2081 (1984); Hermonat & Muzyczka, *PNAS* 81:6466-6470 (1984); and Samulski et al., *J. Virol.* 63:03822-3828 (1989).

Models of Genetic and Epigenetic Conditions

[00421] A method of the invention may be used to create a plant, an animal or cell that may be used to model and/or study genetic or epigenetic conditions of interest, such as a through a model of mutations of interest or a as a disease model. As used herein, "disease" refers to a disease, disorder, or indication in a subject. For example, a method of the invention may be used to create an animal or cell that comprises a modification in one or more nucleic acid sequences associated with a disease, or a plant, animal or cell in which the expression of one or more nucleic acid sequences associated with a disease are altered. Such a nucleic acid sequence may encode a disease associated protein sequence or may be a disease associated control sequence. Accordingly, it is understood that in embodiments of the invention, a plant, subject, patient, organism or cell can be a non-human subject, patient, organism or cell. Thus, the invention provides a plant, animal or cell, produced by the present methods, or a progeny thereof. The progeny may be a clone of the produced plant or animal, or may result from sexual reproduction by crossing with other individuals of the same species to introgress further desirable traits into their offspring. The cell may be *in vivo* or *ex vivo* in the cases of multicellular organisms, particularly animals or plants. In the instance where the cell is in cultured, a cell line may be established if appropriate culturing conditions are met and preferably if the cell is suitably adapted for this purpose (for instance a stem cell). Bacterial cell lines produced by the invention are also envisaged. Hence, cell lines are also envisaged.

[00422] In some methods, the disease model can be used to study the effects of mutations on the animal or cell and development and/or progression of the disease using measures commonly used in the study of the disease. Alternatively, such a disease model is useful for studying the effect of a pharmaceutically active compound on the disease.

[00423] In some methods, the disease model can be used to assess the efficacy of a potential gene therapy strategy. That is, a disease-associated gene or polynucleotide can be modified such that the disease development and/or progression is inhibited or reduced. In particular, the method comprises modifying a disease-associated gene or polynucleotide such that an altered protein is produced and, as a result, the animal or cell has an altered response. Accordingly, in some methods, a genetically modified animal may be compared with an animal predisposed to development of the disease such that the effect of the gene therapy event may be assessed.

[00424] In another embodiment, this invention provides a method of developing a biologically active agent that modulates a cell signaling event associated with a disease gene. The method comprises contacting a test compound with a cell comprising one or more vectors that drive expression of one or more of a CRISPR enzyme, a guide sequence linked to a tracr mate sequence, and a tracr sequence; and detecting a change in a readout that is indicative of a reduction or an augmentation of a cell signaling event associated with, e.g., a mutation in a disease gene contained in the cell.

[00425] A cell model or animal model can be constructed in combination with the method of the invention for screening a cellular function change. Such a model may be used to study the effects of a genome sequence modified by the CRISPR complex of the invention on a cellular function of interest. For example, a cellular function model may be used to study the effect of a modified genome sequence on intracellular signaling or extracellular signaling. Alternatively, a cellular function model may be used to study the effects of a modified genome sequence on sensory perception. In some such models, one or more genome sequences associated with a signaling biochemical pathway in the model are modified.

[00426] Several disease models have been specifically investigated. These include *de novo* autism risk genes CHD8, KATNAL2, and SCN2A; and the syndromic autism (Angelman Syndrome) gene UBE3A. These genes and resulting autism models are of course preferred, but serve to show the broad applicability of the invention across genes and corresponding models.

[00427] An altered expression of one or more genome sequences associated with a signaling biochemical pathway can be determined by assaying for a difference in the mRNA levels of the corresponding genes between the test model cell and a control cell, when they are contacted with a candidate agent. Alternatively, the differential expression of the sequences associated with a

signaling biochemical pathway is determined by detecting a difference in the level of the encoded polypeptide or gene product.

[00428] To assay for an agent-induced alteration in the level of mRNA transcripts or corresponding polynucleotides, nucleic acid contained in a sample is first extracted according to standard methods in the art. For instance, mRNA can be isolated using various lytic enzymes or chemical solutions according to the procedures set forth in Sambrook et al. (1989), or extracted by nucleic-acid-binding resins following the accompanying instructions provided by the manufacturers. The mRNA contained in the extracted nucleic acid sample is then detected by amplification procedures or conventional hybridization assays (e.g. Northern blot analysis) according to methods widely known in the art or based on the methods exemplified herein.

[00429] For purpose of this invention, amplification means any method employing a primer and a polymerase capable of replicating a target sequence with reasonable fidelity. Amplification may be carried out by natural or recombinant DNA polymerases such as TaqGold™, T7 DNA polymerase, Klenow fragment of E.coli DNA polymerase, and reverse transcriptase. A preferred amplification method is PCR. In particular, the isolated RNA can be subjected to a reverse transcription assay that is coupled with a quantitative polymerase chain reaction (RT-PCR) in order to quantify the expression level of a sequence associated with a signaling biochemical pathway.

[00430] Detection of the gene expression level can be conducted in real time in an amplification assay. In one aspect, the amplified products can be directly visualized with fluorescent DNA-binding agents including but not limited to DNA intercalators and DNA groove binders. Because the amount of the intercalators incorporated into the double-stranded DNA molecules is typically proportional to the amount of the amplified DNA products, one can conveniently determine the amount of the amplified products by quantifying the fluorescence of the intercalated dye using conventional optical systems in the art. DNA-binding dye suitable for this application include SYBR green, SYBR blue, DAPI, propidium iodine, Hoeste, SYBR gold, ethidium bromide, acridines, proflavine, acridine orange, acriflavine, fluorcoumanin, ellipticine, daunomycin, chloroquine, distamycin D, chromomycin, homidium, mithramycin, ruthenium polypyridyls, anthramycin, and the like.

[00431] In another aspect, other fluorescent labels such as sequence specific probes can be employed in the amplification reaction to facilitate the detection and quantification of the

amplified products. Probe-based quantitative amplification relies on the sequence-specific detection of a desired amplified product. It utilizes fluorescent, target-specific probes (e.g., TaqMan® probes) resulting in increased specificity and sensitivity. Methods for performing probe-based quantitative amplification are well established in the art and are taught in U.S. Patent No. 5,210,015.

[00432] In yet another aspect, conventional hybridization assays using hybridization probes that share sequence homology with sequences associated with a signaling biochemical pathway can be performed. Typically, probes are allowed to form stable complexes with the sequences associated with a signaling biochemical pathway contained within the biological sample derived from the test subject in a hybridization reaction. It will be appreciated by one of skill in the art that where antisense is used as the probe nucleic acid, the target polynucleotides provided in the sample are chosen to be complementary to sequences of the antisense nucleic acids. Conversely, where the nucleotide probe is a sense nucleic acid, the target polynucleotide is selected to be complementary to sequences of the sense nucleic acid.

[00433] Hybridization can be performed under conditions of various stringency. Suitable hybridization conditions for the practice of the present invention are such that the recognition interaction between the probe and sequences associated with a signaling biochemical pathway is both sufficiently specific and sufficiently stable. Conditions that increase the stringency of a hybridization reaction are widely known and published in the art. See, for example, (Sambrook, et al., (1989); Nonradioactive In Situ Hybridization Application Manual, Boehringer Mannheim, second edition). The hybridization assay can be formed using probes immobilized on any solid support, including but are not limited to nitrocellulose, glass, silicon, and a variety of gene arrays. A preferred hybridization assay is conducted on high-density gene chips as described in U.S. Patent No. 5,445,934.

[00434] For a convenient detection of the probe-target complexes formed during the hybridization assay, the nucleotide probes are conjugated to a detectable label. Detectable labels suitable for use in the present invention include any composition detectable by photochemical, biochemical, spectroscopic, immunochemical, electrical, optical or chemical means. A wide variety of appropriate detectable labels are known in the art, which include fluorescent or chemiluminescent labels, radioactive isotope labels, enzymatic or other ligands. In preferred

embodiments, one will likely desire to employ a fluorescent label or an enzyme tag, such as digoxigenin, β -galactosidase, urease, alkaline phosphatase or peroxidase, avidin/biotin complex.

[00435] The detection methods used to detect or quantify the hybridization intensity will typically depend upon the label selected above. For example, radiolabels may be detected using photographic film or a phosphoimager. Fluorescent markers may be detected and quantified using a photodetector to detect emitted light. Enzymatic labels are typically detected by providing the enzyme with a substrate and measuring the reaction product produced by the action of the enzyme on the substrate; and finally colorimetric labels are detected by simply visualizing the colored label.

[00436] An agent-induced change in expression of sequences associated with a signaling biochemical pathway can also be determined by examining the corresponding gene products. Determining the protein level typically involves a) contacting the protein contained in a biological sample with an agent that specifically bind to a protein associated with a signaling biochemical pathway; and (b) identifying any agent:protein complex so formed. In one aspect of this embodiment, the agent that specifically binds a protein associated with a signaling biochemical pathway is an antibody, preferably a monoclonal antibody.

[00437] The reaction is performed by contacting the agent with a sample of the proteins associated with a signaling biochemical pathway derived from the test samples under conditions that will allow a complex to form between the agent and the proteins associated with a signaling biochemical pathway. The formation of the complex can be detected directly or indirectly according to standard procedures in the art. In the direct detection method, the agents are supplied with a detectable label and unreacted agents may be removed from the complex; the amount of remaining label thereby indicating the amount of complex formed. For such method, it is preferable to select labels that remain attached to the agents even during stringent washing conditions. It is preferable that the label does not interfere with the binding reaction. In the alternative, an indirect detection procedure may use an agent that contains a label introduced either chemically or enzymatically. A desirable label generally does not interfere with binding or the stability of the resulting agent:polypeptide complex. However, the label is typically designed to be accessible to an antibody for an effective binding and hence generating a detectable signal.

[00438] A wide variety of labels suitable for detecting protein levels are known in the art. Non-limiting examples include radioisotopes, enzymes, colloidal metals, fluorescent compounds, bioluminescent compounds, and chemiluminescent compounds.

[00439] The amount of agent:polypeptide complexes formed during the binding reaction can be quantified by standard quantitative assays. As illustrated above, the formation of agent:polypeptide complex can be measured directly by the amount of label remained at the site of binding. In an alternative, the protein associated with a signaling biochemical pathway is tested for its ability to compete with a labeled analog for binding sites on the specific agent. In this competitive assay, the amount of label captured is inversely proportional to the amount of protein sequences associated with a signaling biochemical pathway present in a test sample.

[00440] A number of techniques for protein analysis based on the general principles outlined above are available in the art. They include but are not limited to radioimmunoassays, ELISA (enzyme linked immunoradiometric assays), "sandwich" immunoassays, immunoradiometric assays, in situ immunoassays (using e.g., colloidal gold, enzyme or radioisotope labels), western blot analysis, immunoprecipitation assays, immunofluorescent assays, and SDS-PAGE.

[00441] Antibodies that specifically recognize or bind to proteins associated with a signaling biochemical pathway are preferable for conducting the aforementioned protein analyses. Where desired, antibodies that recognize a specific type of post-translational modifications (e.g., signaling biochemical pathway inducible modifications) can be used. Post-translational modifications include but are not limited to glycosylation, lipidation, acetylation, and phosphorylation. These antibodies may be purchased from commercial vendors. For example, anti-phosphotyrosine antibodies that specifically recognize tyrosine-phosphorylated proteins are available from a number of vendors including Invitrogen and Perkin Elmer. Anti-phosphotyrosine antibodies are particularly useful in detecting proteins that are differentially phosphorylated on their tyrosine residues in response to an ER stress. Such proteins include but are not limited to eukaryotic translation initiation factor 2 alpha (eIF-2 α). Alternatively, these antibodies can be generated using conventional polyclonal or monoclonal antibody technologies by immunizing a host animal or an antibody-producing cell with a target protein that exhibits the desired post-translational modification.

[00442] In practicing the subject method, it may be desirable to discern the expression pattern of an protein associated with a signaling biochemical pathway in different bodily tissue, in

different cell types, and/or in different subcellular structures. These studies can be performed with the use of tissue-specific, cell-specific or subcellular structure specific antibodies capable of binding to protein markers that are preferentially expressed in certain tissues, cell types, or subcellular structures.

[00443] An altered expression of a gene associated with a signaling biochemical pathway can also be determined by examining a change in activity of the gene product relative to a control cell. The assay for an agent-induced change in the activity of a protein associated with a signaling biochemical pathway will depend on the biological activity and/or the signal transduction pathway that is under investigation. For example, where the protein is a kinase, a change in its ability to phosphorylate the downstream substrate(s) can be determined by a variety of assays known in the art. Representative assays include but are not limited to immunoblotting and immunoprecipitation with antibodies such as anti-phosphotyrosine antibodies that recognize phosphorylated proteins. In addition, kinase activity can be detected by high throughput chemiluminescent assays such as AlphaScreen™ (available from Perkin Elmer) and eTag™ assay (Chan-Hui, et al. (2003) *Clinical Immunology* 111: 162-174).

[00444] Where the protein associated with a signaling biochemical pathway is part of a signaling cascade leading to a fluctuation of intracellular pH condition, pH sensitive molecules such as fluorescent pH dyes can be used as the reporter molecules. In another example where the protein associated with a signaling biochemical pathway is an ion channel, fluctuations in membrane potential and/or intracellular ion concentration can be monitored. A number of commercial kits and high-throughput devices are particularly suited for a rapid and robust screening for modulators of ion channels. Representative instruments include FLIPR™ (Molecular Devices, Inc.) and VIPR (Aurora Biosciences). These instruments are capable of detecting reactions in over 1000 sample wells of a microplate simultaneously, and providing real-time measurement and functional data within a second or even a minisecond.

[00445] In practicing any of the methods disclosed herein, a suitable vector can be introduced to a cell or an embryo via one or more methods known in the art, including without limitation, microinjection, electroporation, sonoporation, biolistics, calcium phosphate-mediated transfection, cationic transfection, liposome transfection, dendrimer transfection, heat shock transfection, nucleofection transfection, magnetofection, lipofection, impalefection, optical transfection, proprietary agent-enhanced uptake of nucleic acids, and delivery via liposomes,

immunoliposomes, virosomes, or artificial virions. In some methods, the vector is introduced into an embryo by microinjection. The vector or vectors may be microinjected into the nucleus or the cytoplasm of the embryo. In some methods, the vector or vectors may be introduced into a cell by nucleofection.

[00446] The target polynucleotide of a CRISPR complex can be any polynucleotide endogenous or exogenous to the eukaryotic cell. For example, the target polynucleotide can be a polynucleotide residing in the nucleus of the eukaryotic cell. The target polynucleotide can be a sequence coding a gene product (e.g., a protein) or a non-coding sequence (e.g., a regulatory polynucleotide or a junk DNA).

[00447] Examples of target polynucleotides include a sequence associated with a signaling biochemical pathway, e.g., a signaling biochemical pathway-associated gene or polynucleotide. Examples of target polynucleotides include a disease associated gene or polynucleotide. A “disease-associated” gene or polynucleotide refers to any gene or polynucleotide which is yielding transcription or translation products at an abnormal level or in an abnormal form in cells derived from a disease-affected tissues compared with tissues or cells of a non disease control. It may be a gene that becomes expressed at an abnormally high level; it may be a gene that becomes expressed at an abnormally low level, where the altered expression correlates with the occurrence and/or progression of the disease. A disease-associated gene also refers to a gene possessing mutation(s) or genetic variation that is directly responsible or is in linkage disequilibrium with a gene(s) that is responsible for the etiology of a disease. The transcribed or translated products may be known or unknown, and may be at a normal or abnormal level.

[00448] The target polynucleotide of a CRISPR complex can be any polynucleotide endogenous or exogenous to the eukaryotic cell. For example, the target polynucleotide can be a polynucleotide residing in the nucleus of the eukaryotic cell. The target polynucleotide can be a sequence coding a gene product (e.g., a protein) or a non-coding sequence (e.g., a regulatory polynucleotide or a junk DNA). Without wishing to be bound by theory, it is believed that the target sequence should be associated with a PAM (protospacer adjacent motif); that is, a short sequence recognized by the CRISPR complex. The precise sequence and length requirements for the PAM differ depending on the CRISPR enzyme used, but PAMs are typically 2-5 base pair sequences adjacent the protospacer (that is, the target sequence) Examples of PAM

sequences are given in the examples section below, and the skilled person will be able to identify further PAM sequences for use with a given CRISPR enzyme.

[00449] The target polynucleotide of a CRISPR complex may include a number of disease-associated genes and polynucleotides as well as signaling biochemical pathway-associated genes and polynucleotides as listed in US provisional patent applications 61/736,527 and 61/748,427 both entitled SYSTEMS METHODS AND COMPOSITIONS FOR SEQUENCE MANIPULATION filed on December 12, 2012 and January 2, 2013, respectively, and PCT Application PCT/US2013/074667, entitled DELIVERY, ENGINEERING AND OPTIMIZATION OF SYSTEMS, METHODS AND COMPOSITIONS FOR SEQUENCE MANIPULATION AND THERAPEUTIC APPLICATIONS, filed December 12, 2013, the contents of all of which are herein incorporated by reference in their entirety.

[00450] Examples of target polynucleotides include a sequence associated with a signaling biochemical pathway, e.g., a signaling biochemical pathway-associated gene or polynucleotide. Examples of target polynucleotides include a disease associated gene or polynucleotide. A “disease-associated” gene or polynucleotide refers to any gene or polynucleotide which is yielding transcription or translation products at an abnormal level or in an abnormal form in cells derived from a disease-affected tissues compared with tissues or cells of a non disease control. It may be a gene that becomes expressed at an abnormally high level; it may be a gene that becomes expressed at an abnormally low level, where the altered expression correlates with the occurrence and/or progression of the disease. A disease-associated gene also refers to a gene possessing mutation(s) or genetic variation that is directly responsible or is in linkage disequilibrium with a gene(s) that is responsible for the etiology of a disease. The transcribed or translated products may be known or unknown, and may be at a normal or abnormal level.

Genome-Wide Knock-out Screening

[00451] The CRISPR-Cas9 proteins and systems described herein can be used to perform efficient and cost effective functional genomic screens. Such screens can utilize CRISPR-Cas9 genome wide libraries. Such screens and libraries can provide for determining the function of genes, cellular pathways genes are involved in, and how any alteration in gene expression can result in a particular biological process. An advantage of the present invention is that the CRISPR system avoids off-target binding and its resulting side effects. This is achieved using systems arranged to have a high degree of sequence specificity for the target DNA.

[00452] A genome wide library may comprise a plurality of CRISPR-Cas9 system guide RNAs, as described herein, comprising guide sequences that are capable of targeting a plurality of target sequences in a plurality of genomic loci in a population of eukaryotic cells. The population of cells may be a population of embryonic stem (ES) cells. The target sequence in the genomic locus may be a non-coding sequence. The non-coding sequence may be an intron, regulatory sequence, splice site, 3' UTR, 5' UTR, or polyadenylation signal. Gene function of one or more gene products may be altered by said targeting. The targeting may result in a knockout of gene function. The targeting of a gene product may comprise more than one guide RNA. A gene product may be targeted by 2, 3, 4, 5, 6, 7, 8, 9, or 10 guide RNAs, preferably 3 to 4 per gene. Off-target modifications may be minimized by exploiting the staggered double strand breaks generated by Cas9 effector protein complexes or by utilizing methods analogous to those used in CRISPR-Cas9 systems. (See, e.g., DNA targeting specificity of RNA-guided Cas9 nucleases. Hsu, P., Scott, D., Weinstein, J., Ran, FA., Konermann, S., Agarwala, V., Li, Y., Fine, E., Wu, X., Shalem, O., Cradick, TJ., Marraffini, LA., Bao, G., & Zhang, F. *Nat Biotechnol* doi:10.1038/nbt.2647 (2013)), incorporated herein by reference. The targeting may be of about 100 or more sequences. The targeting may be of about 1000 or more sequences. The targeting may be of about 20,000 or more sequences. The targeting may be of the entire genome. The targeting may be of a panel of target sequences focused on a relevant or desirable pathway. The pathway may be an immune pathway. The pathway may be a cell division pathway.

[00453] One aspect of the invention comprehends a genome wide library that may comprise a plurality of CRISPR-Cas9 system guide RNAs that may comprise guide sequences that are capable of targeting a plurality of target sequences in a plurality of genomic loci, wherein said targeting results in a knockout of gene function. This library may potentially comprise guide RNAs that target each and every gene in the genome of an organism.

[00454] In some embodiments of the invention the organism or subject is a eukaryote (including mammal including human) or a non-human eukaryote or a non-human animal or a non-human mammal. In some embodiments, the organism or subject is a non-human animal, and may be an arthropod, for example, an insect, or may be a nematode. In some methods of the invention the organism or subject is a plant. In some methods of the invention the organism or subject is a mammal or a non-human mammal. A non-human mammal may be for example a

rodent (preferably a mouse or a rat), an ungulate, or a primate. In some methods of the invention the organism or subject is algae, including microalgae, or is a fungus.

[00455] The knockout of gene function may comprise: introducing into each cell in the population of cells a vector system of one or more vectors comprising an engineered, non-naturally occurring CRISPR-Cas9 system comprising I. a Cas9 protein, and II. one or more guide RNAs, wherein components I and II may be same or on different vectors of the system, integrating components I and II into each cell, wherein the guide sequence targets a unique gene in each cell, wherein the Cas9 protein is operably linked to a regulatory element, wherein when transcribed, the guide RNA comprising the guide sequence directs sequence-specific binding of a CRISPR-Cas9 system to a target sequence in the genomic loci of the unique gene, inducing cleavage of the genomic loci by the Cas9 protein, and confirming different knockout mutations in a plurality of unique genes in each cell of the population of cells thereby generating a gene knockout cell library. The invention comprehends that the population of cells is a population of eukaryotic cells, and in a preferred embodiment, the population of cells is a population of embryonic stem (ES) cells.

[00456] The one or more vectors may be plasmid vectors. The vector may be a single vector comprising Cas9, a sgRNA, and optionally, a selection marker into target cells. Not being bound by a theory, the ability to simultaneously deliver Cas9 and sgRNA through a single vector enables application to any cell type of interest, without the need to first generate cell lines that express Cas9. The regulatory element may be an inducible promoter. The inducible promoter may be a doxycycline inducible promoter. In some methods of the invention the expression of the guide sequence is under the control of the T7 promoter and is driven by the expression of T7 polymerase. The confirming of different knockout mutations may be by whole exome sequencing. The knockout mutation may be achieved in 100 or more unique genes. The knockout mutation may be achieved in 1000 or more unique genes. The knockout mutation may be achieved in 20,000 or more unique genes. The knockout mutation may be achieved in the entire genome. The knockout of gene function may be achieved in a plurality of unique genes which function in a particular physiological pathway or condition. The pathway or condition may be an immune pathway or condition. The pathway or condition may be a cell division pathway or condition.

[00457] The invention also provides kits that comprise the genome wide libraries mentioned herein. The kit may comprise a single container comprising vectors or plasmids comprising the library of the invention. The kit may also comprise a panel comprising a selection of unique CRISPR-Cas9 system guide RNAs comprising guide sequences from the library of the invention, wherein the selection is indicative of a particular physiological condition. The invention comprehends that the targeting is of about 100 or more sequences, about 1000 or more sequences or about 20,000 or more sequences or the entire genome. Furthermore, a panel of target sequences may be focused on a relevant or desirable pathway, such as an immune pathway or cell division.

[00458] In an additional aspect of the invention, a Cas9 enzyme may comprise one or more mutations and may be used as a generic DNA binding protein with or without fusion to a functional domain. The mutations may be artificially introduced mutations or gain- or loss-of-function mutations. The mutations may include but are not limited to mutations in one of the catalytic domains (D10 and H840) in the RuvC and HNH catalytic domains, respectively. Further mutations have been characterized. In one aspect of the invention, the functional domain may be a transcriptional activation domain, which may be VP64. In other aspects of the invention, the functional domain may be a transcriptional repressor domain, which may be KRAB or SID4X. Other aspects of the invention relate to the mutated Cas9 enzyme being fused to domains which include but are not limited to a transcriptional activator, repressor, a recombinase, a transposase, a histone remodeler, a demethylase, a DNA methyltransferase, a cryptochrome, a light inducible/controllable domain or a chemically inducible/controllable domain. Some methods of the invention can include inducing expression of targeted genes. In one embodiment, inducing expression by targeting a plurality of target sequences in a plurality of genomic loci in a population of eukaryotic cells is by use of a functional domain.

[00459] Useful in the practice of the instant invention, reference is made to:

- Genome-Scale CRISPR-Cas9 Knockout Screening in Human Cells. Shalem, O., Sanjana, NE., Hartenian, E., Shi, X., Scott, DA., Mikkelsen, T., Heckl, D., Ebert, BL., Root, DE., Doench, JG., Zhang, F. *Science* Dec 12. (2013). [Epub ahead of print]; Published in final edited form as: *Science*. 2014 Jan 3; 343(6166): 84–87.
- Shalem *et al.* involves a new way to interrogate gene function on a genome-wide scale. Their studies showed that delivery of a genome-scale CRISPR-Cas9 knockout (GeCKO)

library targeted 18,080 genes with 64,751 unique guide sequences enabled both negative and positive selection screening in human cells. First, the authors showed use of the GeCKO library to identify genes essential for cell viability in cancer and pluripotent stem cells. Next, in a melanoma model, the authors screened for genes whose loss is involved in resistance to vemurafenib, a therapeutic that inhibits mutant protein kinase BRAF. Their studies showed that the highest-ranking candidates included previously validated genes NF1 and MED12 as well as novel hits NF2, CUL3, TADA2B, and TADA1. The authors observed a high level of consistency between independent guide RNAs targeting the same gene and a high rate of hit confirmation, and thus demonstrated the promise of genome-scale screening with Cas9.

[00460] Reference is also made to US patent publication number US20140357530; and PCT Patent Publication WO2014093701, hereby incorporated herein by reference. Reference is also made to NIH Press Release of Oct. 22, 2015 entitled, "Researchers identify potential alternative to CRISPR-Cas genome editing tools: New Cas enzymes shed light on evolution of CRISPR-Cas systems, which is incorporated by reference.

Functional Alteration and Screening

[00461] In another aspect, the present invention provides for a method of functional evaluation and screening of genes. The use of the CRISPR system of the present invention to precisely deliver functional domains, to activate or repress genes or to alter epigenetic state by precisely altering the methylation site on a specific locus of interest, can be with one or more guide RNAs applied to a single cell or population of cells or with a library applied to genome in a pool of cells ex vivo or in vivo comprising the administration or expression of a library comprising a plurality of guide RNAs (sgRNAs) and wherein the screening further comprises use of a Cas9 effector protein, wherein the CRISPR complex comprising the Cas9 effector protein is modified to comprise a heterologous functional domain. In an aspect the invention provides a method for screening a genome comprising the administration to a host or expression in a host in vivo of a library. In an aspect the invention provides a method as herein discussed further comprising an activator administered to the host or expressed in the host. In an aspect the invention provides a method as herein discussed wherein the activator is attached to a Cas9 effector protein. In an aspect the invention provides a method as herein discussed wherein the activator is attached to the N terminus or the C terminus of the Cas9 effector protein. In an aspect

the invention provides a method as herein discussed wherein the activator is attached to a sgRNA loop. In an aspect the invention provides a method as herein discussed further comprising a repressor administered to the host or expressed in the host. In an aspect the invention provides a method as herein discussed, wherein the screening comprises affecting and detecting gene activation, gene inhibition, or cleavage in the locus.

[00462] In an aspect, the invention provides efficient on-target activity and minimizes off target activity. In an aspect, the invention provides efficient on-target cleavage by Cas9 effector protein and minimizes off-target cleavage by the Cas9 effector protein. In an aspect, the invention provides guide specific binding of Cas9 effector protein at a gene locus without DNA cleavage. Accordingly, in an aspect, the invention provides target-specific gene regulation. In an aspect, the invention provides guide specific binding of Cas9 effector protein at a gene locus without DNA cleavage. Accordingly, in an aspect, the invention provides for cleavage at one gene locus and gene regulation at a different gene locus using a single Cas9 effector protein. In an aspect, the invention provides orthogonal activation and/or inhibition and/or cleavage of multiple targets using one or more Cas9 effector protein and/or enzyme.

[00463] In an aspect the invention provides a method as herein discussed, wherein the host is a eukaryotic cell. In an aspect the invention provides a method as herein discussed, wherein the host is a mammalian cell. In an aspect the invention provides a method as herein discussed, wherein the host is a non-human eukaryote. In an aspect the invention provides a method as herein discussed, wherein the non-human eukaryote is a non-human mammal. In an aspect the invention provides a method as herein discussed, wherein the non-human mammal is a mouse. An aspect the invention provides a method as herein discussed comprising the delivery of the Cas9 effector protein complexes or component(s) thereof or nucleic acid molecule(s) coding therefor, wherein said nucleic acid molecule(s) are operatively linked to regulatory sequence(s) and expressed in vivo. In an aspect the invention provides a method as herein discussed wherein the expressing in vivo is via a lentivirus, an adenovirus, or an AAV. In an aspect the invention provides a method as herein discussed wherein the delivery is via a particle, a nanoparticle, a lipid or a cell penetrating peptide (CPP).

[00464] In an aspect the invention provides a pair of CRISPR complexes comprising Cas9 effector protein, each comprising a guide RNA (sgRNA) comprising a guide sequence capable of hybridizing to a target sequence in a genomic locus of interest in a cell, wherein at least one loop

of each sgRNA is modified by the insertion of distinct RNA sequence(s) that bind to one or more adaptor proteins, and wherein the adaptor protein is associated with one or more functional domains, wherein each sgRNA of each Cas9 effector protein complex comprises a functional domain having a DNA cleavage activity. In an aspect the invention provides paired Cas9 effector protein complexes as herein-discussed, wherein the DNA cleavage activity is due to a FokI nuclease.

[00465] In an aspect the invention provides a method for cutting a target sequence in a genomic locus of interest comprising delivery to a cell of the Cas9 effector protein complexes or component(s) thereof or nucleic acid molecule(s) coding therefor, wherein said nucleic acid molecule(s) are operatively linked to regulatory sequence(s) and expressed *in vivo*. In an aspect the invention provides a method as herein-discussed wherein the delivery is via a lentivirus, an adenovirus, or an AAV. In an aspect the invention provides a method as herein-discussed or paired Cas9 effector protein complexes as herein-discussed wherein the target sequence for a first complex of the pair is on a first strand of double stranded DNA and the target sequence for a second complex of the pair is on a second strand of double stranded DNA. In an aspect the invention provides a method as herein-discussed or paired Cas9 effector protein complexes as herein-discussed wherein the target sequences of the first and second complexes are in proximity to each other such that the DNA is cut in a manner that facilitates homology directed repair. In an aspect a herein method can further include introducing into the cell template DNA. In an aspect a herein method or herein paired Cas9 effector protein complexes can involve wherein each Cas9 effector protein complex has a Cas9 effector enzyme that is mutated such that it has no more than about 5% of the nuclease activity of the Cas9 effector enzyme that is not mutated.

[00466] In an aspect the invention provides a library, method or complex as herein-discussed wherein the sgRNA is modified to have at least one non-coding functional loop, e.g., wherein the at least one non-coding functional loop is repressive; for instance, wherein the at least one non-coding functional loop comprises Alu.

[00467] In one aspect, the invention provides a method for altering or modifying expression of a gene product. The said method may comprise introducing into a cell containing and expressing a DNA molecule encoding the gene product an engineered, non-naturally occurring CRISPR system comprising a Cas9 effector protein and guide RNA that targets the DNA molecule, whereby the guide RNA targets the DNA molecule encoding the gene product and the Cas9

effector protein cleaves the DNA molecule encoding the gene product, whereby expression of the gene product is altered; and, wherein the Cas9 effector protein and the guide RNA do not naturally occur together. The invention comprehends the guide RNA comprising a guide sequence linked to a direct repeat sequence. The invention further comprehends the Cas9 effector protein being codon optimized for expression in a Eukaryotic cell. In a preferred embodiment the Eukaryotic cell is a mammalian cell and in a more preferred embodiment the mammalian cell is a human cell. In a further embodiment of the invention, the expression of the gene product is decreased.

[00468] In some embodiments, one or more functional domains are associated with the CRISPR enzyme, for example a Type II Cas9 enzyme.

[00469] In some embodiments, one or more functional domains are associated with an adaptor protein, for example as used with the modified guides of Konnerman *et al.* (Nature **517**, 583–588, 29 January 2015).

[00470] In some embodiments, one or more functional domains are associated with an dead sgRNA (dRNA). In some embodiments, a dRNA complex with active cas9 directs gene regulation by a functional domain at on gene locus while an sgRNA directs DNA cleavage by the active cas9 at another locus, for example as described by Dahlman *et al.*, ‘Orthogonal gene control with a catalytically active Cas9 nuclease’ (*in press*). In some embodiments, dRNAs are selected to maximize selectivity of regulation for a gene locus of interest compared to off-target regulation. In some embodiments, dRNAs are selected to maximize target gene regulation and minimize target cleavage

[00471] For the purposes of the following discussion, reference to a functional domain could be a functional domain associated with the CRISPR enzyme or a functional domain associated with the adaptor protein.

[00472] In the practice of the invention, loops of the sgRNA may be extended, without colliding with the Cas9 protein by the insertion of distinct RNA loop(s) or distinct sequence(s) that may recruit adaptor proteins that can bind to the distinct RNA loop(s) or distinct sequence(s). The adaptor proteins may include but are not limited to orthogonal RNA-binding protein / aptamer combinations that exist within the diversity of bacteriophage coat proteins. A list of such coat proteins includes, but is not limited to: Q β , F2, GA, fr, JP501, M12, R17, BZ13, JP34, JP500, KUI, M11, MX1, TW18, VK, SP, FI, ID2, NL95, TW19, AP205, ϕ Cb5, ϕ Cb8r,

ϕ Cb12r, ϕ Cb23r, 7s and PRR1. These adaptor proteins or orthogonal RNA binding proteins can further recruit effector proteins or fusions which comprise one or more functional domains. In some embodiments, the functional domain may be selected from the group consisting of: transposase domain, integrase domain, recombinase domain, resolvase domain, invertase domain, protease domain, DNA methyltransferase domain, DNA hydroxymethylase domain, DNA demethylase domain, histone acetylase domain, histone deacetylases domain, nuclease domain, repressor domain, activator domain, nuclear-localization signal domains, transcription-regulatory protein (or transcription complex recruiting) domain, cellular uptake activity associated domain, nucleic acid binding domain, antibody presentation domain, histone modifying enzymes, recruiter of histone modifying enzymes; inhibitor of histone modifying enzymes, histone methyltransferase, histone demethylase, histone kinase, histone phosphatase, histone ribosylase, histone deribosylase, histone ubiquitinase, histone deubiquitinase, histone biotinase and histone tail protease. In some preferred embodiments, the functional domain is a transcriptional activation domain, such as, without limitation, VP64, p65, MyoD1, HSF1, RTA, SET7/9 or a histone acetyltransferase. In some embodiments, the functional domain is a transcription repression domain, preferably KRAB. In some embodiments, the transcription repression domain is SID, or concatemers of SID (eg SID4X). In some embodiments, the functional domain is an epigenetic modifying domain, such that an epigenetic modifying enzyme is provided. In some embodiments, the functional domain is an activation domain, which may be the P65 activation domain.

[00473] In some embodiments, the one or more functional domains is an NLS (Nuclear Localization Sequence) or an NES (Nuclear Export Signal). In some embodiments, the one or more functional domains is a transcriptional activation domain comprises VP64, p65, MyoD1, HSF1, RTA, SET7/9 and a histone acetyltransferase. Other references herein to activation (or activator) domains in respect of those associated with the CRISPR enzyme include any known transcriptional activation domain and specifically VP64, p65, MyoD1, HSF1, RTA, SET7/9 or a histone acetyltransferase.

[00474] In some embodiments, the one or more functional domains is a transcriptional repressor domain. In some embodiments, the transcriptional repressor domain is a KRAB domain. In some embodiments, the transcriptional repressor domain is a NuE domain, NcoR domain, SID domain or a SID4X domain.

[00475] In some embodiments, the one or more functional domains have one or more activities comprising methylase activity, demethylase activity, transcription activation activity, transcription repression activity, transcription release factor activity, histone modification activity, RNA cleavage activity, DNA cleavage activity, DNA integration activity or nucleic acid binding activity.

[00476] Histone modifying domains are also preferred in some embodiments. Exemplary histone modifying domains are discussed below. Transposase domains, HR (Homologous Recombination) machinery domains, recombinase domains, and/or integrase domains are also preferred as the present functional domains. In some embodiments, DNA integration activity includes HR machinery domains, integrase domains, recombinase domains and/or transposase domains. Histone acetyltransferases are preferred in some embodiments.

[00477] In some embodiments, the DNA cleavage activity is due to a nuclease. In some embodiments, the nuclease comprises a FokI nuclease. See, "Dimeric CRISPR RNA-guided FokI nucleases for highly specific genome editing", Shengdar Q. Tsai, Nicolas Wyvekens, Cyd Khayter, Jennifer A. Foden, Vishal Thapar, Deepak Reyon, Mathew J. Goodwin, Martin J. Aryee, J. Keith Joung *Nature Biotechnology* 32(6): 569-77 (2014), relates to dimeric RNA-guided FokI Nucleases that recognize extended sequences and can edit endogenous genes with high efficiencies in human cells.

[00478] In some embodiments, the one or more functional domains is attached to the CRISPR enzyme so that upon binding to the sgRNA and target the functional domain is in a spatial orientation allowing for the functional domain to function in its attributed function.

[00479] In some embodiments, the one or more functional domains is attached to the adaptor protein so that upon binding of the CRISPR enzyme to the sgRNA and target, the functional domain is in a spatial orientation allowing for the functional domain to function in its attributed function.

[00480] In an aspect the invention provides a composition as herein discussed wherein the one or more functional domains is attached to the CRISPR enzyme or adaptor protein via a linker, optionally a GlySer linker, as discussed herein.

[00481] Endogenous transcriptional repression is often mediated by chromatin modifying enzymes such as histone methyltransferases (HMTs) and deacetylases (HDACs). Repressive histone effector domains are known and an exemplary list is provided below. In the exemplary

table, preference was given to proteins and functional truncations of small size to facilitate efficient viral packaging (for instance via AAV). In general, however, the domains may include HDACs, histone methyltransferases (HMTs), and histone acetyltransferase (HAT) inhibitors, as well as HDAC and HMT recruiting proteins. The functional domain may be or include, in some embodiments, HDAC Effector Domains, HDAC Recruiter Effector Domains, Histone Methyltransferase (HMT) Effector Domains, Histone Methyltransferase (HMT) Recruiter Effector Domains, or Histone Acetyltransferase Inhibitor Effector Domains.

[00482] HDAC Effector Domains

Subtype/Complex	Name	Substrate (if known)	Modification (if known)	Organism	Full size (aa)	Selected truncation (aa)	Final size (aa)	Catalytic domain
HDAC I	HDAC8	-	-	<i>X. laevis</i>	325	1-325	325	1-272: HDAC
HDAC I	RPD3	-	-	<i>S. cerevisiae</i>	433	19-340	322 (Vannier)	19-331: HDAC
HDAC IV	MesoLo4	-	-	<i>M. loli</i>	300	1-300 (Gregorette)	300	-
HDAC IV	HDAC11	-	-	<i>H. sapiens</i>	347	1-347 (Gao)	347	14-326: HDAC
HD2	HDT1	-	-	<i>A. thaliana</i>	245	1-211 (Wu)	211	-
SIRT I	SIRT3	H3K9Ac H4K16Ac H3K56Ac	-	<i>H. sapiens</i>	399	143-399 (Scher)	257	126-382: SIRT
SIRT I	HST2	-	-	<i>C. albicans</i>	331	1-331 (Hnisz)	331	-
SIRT I	CobB	-	-	<i>E. coli (K12)</i>	242	1-242 (Landry)	242	-
SIRT I	HST2	-	-	<i>S. cerevisiae</i>	357	8-298 (Wilson)	291	-
SIRT III	SIRT5	H4K8Ac H4K16Ac	-	<i>H. sapiens</i>	310	37-310 (Gertz)	274	41-309: SIRT
SIRT III	Sir2A	-	-	<i>P. falciparum</i>	273	1-273 (Zhu)	273	19-273: SIRT
SIRT IV	SIRT6	H3K9Ac H3K56Ac	-	<i>H. sapiens</i>	355	1-289 (Tennen)	289	35-274: SIRT

[00483] Accordingly, the repressor domains of the present invention may be selected from histone methyltransferases (HMTs), histone deacetylases (HDACs), histone acetyltransferase (HAT) inhibitors, as well as HDAC and HMT recruiting proteins.

[00484] The HDAC domain may be any of those in the table above, namely: HDAC8, RPD3, MesoLo4, HDAC11, HDT1, SIRT3, HST2, CobB, HST2, SIRT5, Sir2A, or SIRT6.

[00485] In some embodiment, the functional domain may be a HDAC Recruiter Effector Domain. Preferred examples include those in the Table below, namely MeCP2, MBD2b, Sin3a, NcoR, SALL1, RCOR1. NcoR is exemplified in the present Examples and, although preferred, it is envisaged that others in the class will also be useful.

[00486] Table of HDAC Recruiter Effector Domains

Subtype/ Complex	Name	Substrate (if known)	Modification (if known)	Organism	Full size (aa)	Selected truncation (aa)	Final size (aa)	Catalytic domain
Sin3a	MeCP2	-	-	<i>R. norvegicus</i>	492	207-492 (Nan)	286	-
Sin3a	MBD2b	-	-	<i>H. sapiens</i>	262	45-262 (Boeke)	218	-
Sin3a	Sin3a	-	-	<i>H. sapiens</i>	1273	524-851 (Laherty)	328	627-829: HDAC1 interaction
NcoR	NcoR	-	-	<i>H. sapiens</i>	2440	420-488 (Zhang)	69	-
NuRD	SALL1	-	-	<i>M. musculus</i>	1322	1-93 (Lauberth)	93	-
CoREST	RCOR1	-	-	<i>H. sapiens</i>	482	81-300 (Gu, Ouyang)	220	-

[00487] In some embodiment, the functional domain may be a Methyltransferase (HMT) Effector Domain. Preferred examples include those in the Table below, namely NUE, vSET, EHMT2/G9A, SUV39H1, dim-5, KYP, SUV4, SET4, SET1, SETD8, and TgSET8. NUE is exemplified in the present Examples and, although preferred, it is envisaged that others in the class will also be useful.

[00488] Table of Histone Methyltransferase (HMT) Effector Domains

Subtype/ Complex	Name	Substrate (if known)	Modification (if known)	Organism	Full size (aa)	Selected truncation (aa)	Final size (aa)	Catalytic domain
SET	NUE	H2B, H3, H4	-	<i>C. trachomatis</i>	219	1-219 (Pennini)	219	-
SET	vSET	-	H3K27me3	<i>P. bursaria chlorella virus</i>	119	1-119 (Mujtaba)	119	4-112: SET2
SUV39 family	EHMT2/G9A	H1.4K2, H3K9, H3K27	H3K9me1/2, H1K25me1	<i>M. musculus</i>	1263	969-1263 (Tachibana)	295	1025-1233: preSET, SET, postSET
SUV39	SUV39H1	-	H3K9me2/3	<i>H. sapiens</i>	412	79-412 (Snowden)	334	172-412:

								preSET, SET, postSE T
Suvar3-9	dim-5	-	H3K9me3	<i>N. crassa</i>	331	1-331 (Rathert)	331	77-331: preSET, SET, postSE T
Suvar3-9 (SUVH subfamil y)	KYP	-	H3K9me1/2	<i>A. thaliana</i>	624	335-601	267 (Jackson)	-
Suvar3-9 (SUVR subfamil y)	SUVR4	H3K9me 1	H3K9me2/3	<i>A. thaliana</i>	492	180-492	313 (Thorstense n)	192- 462: preSET, SET, postSE T
Suvar4- 20	SET4	-	H4K20me3	<i>C. elegans</i>	288	1-288 (Vielle)	288	-
SET8	SET1	-	H4K20me1	<i>C. elegans</i>	242	1-242 (Vielle)	242	-
SET8	SETD8	-	H4K20me1	<i>H. sapiens</i>	393	185-393	209 (Couture)	256- 382: SET
SET8	TgSET8	-	H4K20me1/2 /3	<i>T. gondii</i>	1893	1590-1893 (Sautel)	304	1749- 1884: SET

[00489] In some embodiment, the functional domain may be a Histone Methyltransferase (HMT) Recruiter Effector Domain. Preferred examples include those in the Table below, namely Hp1a, PHF19, and NIPP1.

[00490] Table of Histone Methyltransferase (HMT) Recruiter Effector Domains

Subtype/ Complex	Name	Substrate (if known)	Modification (if known)	Organism	Full size (aa)	Selected truncation (aa)	Final size (aa)	Catalytic domain
-	Hp1a	-	H3K9me3	<i>M. musculus</i>	191	73-191	119 (Hathaway)	121-179: chromoshadow
-	PHF19	-	H3K27me3	<i>H. sapiens</i>	580	(1-250) + GGSG linker + (500-580)	335 (Ballaré)	163-250: PHD2
-	NIPP1	-	H3K27me3	<i>H. sapiens</i>	351	1-329 (Jin)	329	310-329: EED

[00491] In some embodiment, the functional domain may be Histone Acetyltransferase Inhibitor Effector Domain. Preferred examples include SET/TAF-1 β listed in the Table below.

[00492] Table of Histone Acetyltransferase Inhibitor Effector Domains

Subtype/ Complex	Name	Substrate (if known)	Modification (if known)	Organism	Full size (aa)	Selected truncation (aa)	Final size (aa)	Catalytic domain
-	SET/TAF- 1 β	-	-	<i>M. musculus</i>	289	1-289 (Cervoni)	289	-

[00493] It is also preferred to target endogenous (regulatory) control elements (such as enhancers and silencers) in addition to a promoter or promoter-proximal elements. Thus, the invention can also be used to target endogenous control elements (including enhancers and silencers) in addition to targeting of the promoter. These control elements can be located upstream and downstream of the transcriptional start site (TSS), starting from 200bp from the TSS to 100kb away. Targeting of known control elements can be used to activate or repress the gene of interest. In some cases, a single control element can influence the transcription of multiple target genes. Targeting of a single control element could therefore be used to control the transcription of multiple genes simultaneously.

[00494] Targeting of putative control elements on the other hand (e.g. by tiling the region of the putative control element as well as 200bp up to 100kB around the element) can be used as a means to verify such elements (by measuring the transcription of the gene of interest) or to detect novel control elements (e.g. by tiling 100kb upstream and downstream of the TSS of the gene of interest). In addition, targeting of putative control elements can be useful in the context of understanding genetic causes of disease. Many mutations and common SNP variants associated with disease phenotypes are located outside coding regions. Targeting of such regions with either the activation or repression systems described herein can be followed by readout of transcription of either a) a set of putative targets (e.g. a set of genes located in closest proximity to the control element) or b) whole-transcriptome readout by e.g. RNAseq or microarray. This would allow for the identification of likely candidate genes involved in the disease phenotype. Such candidate genes could be useful as novel drug targets.

[00495] Histone acetyltransferase (HAT) inhibitors are mentioned herein. However, an alternative in some embodiments is for the one or more functional domains to comprise an acetyltransferase, preferably a histone acetyltransferase. These are useful in the field of epigenomics, for example in methods of interrogating the epigenome. Methods of interrogating the epigenome may include, for example, targeting epigenomic sequences. Targeting epigenomic sequences may include the guide being directed to an epigenomic target sequence.

Epigenomic target sequence may include, in some embodiments, include a promoter, silencer or an enhancer sequence.

[00496] Use of a functional domain linked to a CRISPR-Cas9 enzyme as described herein, preferably a dead-Cas9, to target epigenomic sequences can be used to activate or repress promoters, silencer or enhancers.

[00497] Examples of acetyltransferases are known but may include, in some embodiments, histone acetyltransferases. In some embodiments, the histone acetyltransferase may comprise the catalytic core of the human acetyltransferase p300 (Gerbasch & Reddy, Nature Biotech 6th April 2015).

[00498] In some preferred embodiments, the functional domain is linked to a dead-Cas9 enzyme to target and activate epigenomic sequences such as promoters or enhancers. One or more guides directed to such promoters or enhancers may also be provided to direct the binding of the CRISPR enzyme to such promoters or enhancers.

[00499] The term “associated with” is used here in relation to the association of the functional domain to the CRISPR enzyme or the adaptor protein. It is used in respect of how one molecule ‘associates’ with respect to another, for example between an adaptor protein and a functional domain, or between the CRISPR enzyme and a functional domain. In the case of such protein-protein interactions, this association may be viewed in terms of recognition in the way an antibody recognizes an epitope. Alternatively, one protein may be associated with another protein via a fusion of the two, for instance one subunit being fused to another subunit. Fusion typically occurs by addition of the amino acid sequence of one to that of the other, for instance via splicing together of the nucleotide sequences that encode each protein or subunit. Alternatively, this may essentially be viewed as binding between two molecules or direct linkage, such as a fusion protein. In any event, the fusion protein may include a linker between the two subunits of interest (i.e. between the enzyme and the functional domain or between the adaptor protein and the functional domain). Thus, in some embodiments, the CRISPR enzyme or adaptor protein is associated with a functional domain by binding thereto. In other embodiments, the CRISPR enzyme or adaptor protein is associated with a functional domain because the two are fused together, optionally via an intermediate linker.

[00500] Attachment of a functional domain or fusion protein can be via a linker, e.g., a flexible glycine-serine (GlyGlyGlySer) or (GGGS)₃ or a rigid alpha-helical linker such as

(Ala(GluAlaAlaAlaLys)Ala). Linkers such as (GGGGS)₃ are preferably used herein to separate protein or peptide domains. (GGGGS)₃ is preferable because it is a relatively long linker (15 amino acids). The glycine residues are the most flexible and the serine residues enhance the chance that the linker is on the outside of the protein. (GGGGS)₆, (GGGGS)₉, or (GGGGS)₁₂ may preferably be used as alternatives. Other preferred alternatives are (GGGGS)₁, (GGGGS)₂, (GGGGS)₄, (GGGGS)₅, (GGGGS)₇, (GGGGS)₈, (GGGGS)₁₀, or (GGGGS)₁₁. Alternative linkers are available, but highly flexible linkers are thought to work best to allow for maximum opportunity for the 2 parts of the Cas9 to come together and thus reconstitute Cas9 activity. One alternative is that the NLS of nucleoplasmin can be used as a linker. For example, a linker can also be used between the Cas9 and any functional domain. Again, a (GGGGS)₃ linker may be used here (or the 6, 9, or 12 repeat versions therefore) or the NLS of nucleoplasmin can be used as a linker between Cas9 and the functional domain.

Saturating Mutagenesis

[00501] CRISPR-Cas System(s) can be used to perform saturating or deep scanning mutagenesis of genomic loci in conjunction with a cellular phenotype—for instance, for determining critical minimal features and discrete vulnerabilities of functional elements required for gene expression, drug resistance, and reversal of disease. By saturating or deep scanning mutagenesis is meant that every or essentially every DNA base is cut within the genomic loci. A library of CRISPR-Cas guide RNAs may be introduced into a population of cells. The library may be introduced, such that each cell receives a single guide RNA (sgRNA). In the case where the library is introduced by transduction of a viral vector, as described herein, a low multiplicity of infection (MOI) is used. The library may include sgRNAs targeting every sequence upstream of a (protospacer adjacent motif) (PAM) sequence in a genomic locus. The library may include at least 100 non-overlapping genomic sequences upstream of a PAM sequence for every 1000 base pairs within the genomic locus. The library may include sgRNAs targeting sequences upstream of at least one different PAM sequence. The CRISPR-Cas System(s) may include more than one Cas protein. Any Cas protein as described herein, including orthologues or engineered Cas proteins that recognize different PAM sequences may be used. The frequency of off target sites for a sgRNA may be less than 500. Off target scores may be generated to select sgRNAs with the lowest off target sites. Any phenotype determined to be associated with cutting at a sgRNA target site may be confirmed by using sgRNA's targeting

the same site in a single experiment. Validation of a target site may also be performed by using a nickase Cas9, as described herein, and two sgRNAs targeting the genomic site of interest. Not being bound by a theory, a target site is a true hit if the change in phenotype is observed in validation experiments.

[00502] The genomic loci may include at least one continuous genomic region. The at least one continuous genomic region may comprise up to the entire genome. The at least one continuous genomic region may comprise a functional element of the genome. The functional element may be within a non-coding region, coding gene, intronic region, promoter, or enhancer. The at least one continuous genomic region may comprise at least 1 kb, preferably at least 50 kb of genomic DNA. The at least one continuous genomic region may comprise a transcription factor binding site. The at least one continuous genomic region may comprise a region of DNase I hypersensitivity. The at least one continuous genomic region may comprise a transcription enhancer or repressor element. The at least one continuous genomic region may comprise a site enriched for an epigenetic signature. The at least one continuous genomic DNA region may comprise an epigenetic insulator. The at least one continuous genomic region may comprise two or more continuous genomic regions that physically interact. Genomic regions that interact may be determined by '4C technology'. 4C technology allows the screening of the entire genome in an unbiased manner for DNA segments that physically interact with a DNA fragment of choice, as is described in Zhao et al. ((2006) *Nat Genet* 38, 1341-7) and in U.S. patent 8,642,295, both incorporated herein by reference in its entirety. The epigenetic signature may be histone acetylation, histone methylation, histone ubiquitination, histone phosphorylation, DNA methylation, or a lack thereof.

[00503] CRISPR-Cas System(s) for saturating or deep scanning mutagenesis can be used in a population of cells. The CRISPR-Cas System(s) can be used in eukaryotic cells, including but not limited to mammalian and plant cells. The population of cells may be prokaryotic cells. The population of eukaryotic cells may be a population of embryonic stem (ES) cells, neuronal cells, epithelial cells, immune cells, endocrine cells, muscle cells, erythrocytes, lymphocytes, plant cells, or yeast cells.

[00504] In one aspect, the present invention provides for a method of screening for functional elements associated with a change in a phenotype. The library may be introduced into a population of cells that are adapted to contain a Cas protein. The cells may be sorted into at least

two groups based on the phenotype. The phenotype may be expression of a gene, cell growth, or cell viability. The relative representation of the guide RNAs present in each group are determined, whereby genomic sites associated with the change in phenotype are determined by the representation of guide RNAs present in each group. The change in phenotype may be a change in expression of a gene of interest. The gene of interest may be upregulated, downregulated, or knocked out. The cells may be sorted into a high expression group and a low expression group. The population of cells may include a reporter construct that is used to determine the phenotype. The reporter construct may include a detectable marker. Cells may be sorted by use of the detectable marker.

[00505] In another aspect, the present invention provides for a method of screening for genomic sites associated with resistance to a chemical compound. The chemical compound may be a drug or pesticide. The library may be introduced into a population of cells that are adapted to contain a Cas protein, wherein each cell of the population contains no more than one guide RNA; the population of cells are treated with the chemical compound; and the representation of guide RNAs are determined after treatment with the chemical compound at a later time point as compared to an early time point, whereby genomic sites associated with resistance to the chemical compound are determined by enrichment of guide RNAs. Representation of sgRNAs may be determined by deep sequencing methods.

[00506] Useful in the practice of the instant invention, reference is made to the article entitled BCL11A enhancer dissection by Cas9-mediated *in situ* saturating mutagenesis. Canver, M.C., Smith, E.C., Sher, F., Pinello, L., Sanjana, N.E., Shalem, O., Chen, D.D., Schupp, P.G., Vinjamur, D.S., Garcia, S.P., Luc, S., Kurita, R., Nakamura, Y., Fujiwara, Y., Maeda, T., Yuan, G., Zhang, F., Orkin, S.H., & Bauer, D.E. DOI:10.1038/nature15521, published online September 16, 2015, the article is herein incorporated by reference and discussed briefly below:

- Canver *et al.* describes novel pooled CRISPR-Cas9 guide RNA libraries to perform *in situ* saturating mutagenesis of the human and mouse BCL11A erythroid enhancers previously identified as an enhancer associated with fetal hemoglobin (HbF) level and whose mouse ortholog is necessary for erythroid BCL11A expression. This approach revealed critical minimal features and discrete vulnerabilities of these enhancers. Through editing of primary human progenitors and mouse transgenesis, the authors

validated the BCL11A erythroid enhancer as a target for HbF reinduction. The authors generated a detailed enhancer map that informs therapeutic genome editing.

Method of Using CRISPR-Cas Systems to Modify a Cell or Organism

[00507] The invention in some embodiments comprehends a method of modifying an cell or organism. The cell may be a prokaryotic cell or a eukaryotic cell. The cell may be a mammalian cell. The mammalian cell may be a non-human primate, bovine, porcine, rodent or mouse cell. The cell may be a non-mammalian eukaryotic cell such as poultry, fish or shrimp. The cell may also be a plant cell. The plant cell may be of a crop plant such as cassava, corn, sorghum, wheat, or rice. The plant cell may also be of an algae, tree or vegetable. The modification introduced to the cell by the present invention may be such that the cell and progeny of the cell are altered for improved production of biologic products such as an antibody, starch, alcohol or other desired cellular output. The modification introduced to the cell by the present invention may be such that the cell and progeny of the cell include an alteration that changes the biologic product produced.

[00508] The system may comprise one or more different vectors. In an aspect of the invention, the Cas protein is codon optimized for expression the desired cell type, preferentially a eukaryotic cell, preferably a mammalian cell or a human cell.

[00509] Packaging cells are typically used to form virus particles that are capable of infecting a host cell. Such cells include 293 cells, which package adenovirus, and ψ 2 cells or PA317 cells, which package retrovirus. Viral vectors used in gene therapy are usually generated by producing a cell line that packages a nucleic acid vector into a viral particle. The vectors typically contain the minimal viral sequences required for packaging and subsequent integration into a host, other viral sequences being replaced by an expression cassette for the polynucleotide(s) to be expressed. The missing viral functions are typically supplied in trans by the packaging cell line. For example, AAV vectors used in gene therapy typically only possess ITR sequences from the AAV genome which are required for packaging and integration into the host genome. Viral DNA is packaged in a cell line, which contains a helper plasmid encoding the other AAV genes, namely rep and cap, but lacking ITR sequences. The cell line may also be infected with adenovirus as a helper. The helper virus promotes replication of the AAV vector and expression of AAV genes from the helper plasmid. The helper plasmid is not packaged in significant amounts due to a lack of ITR sequences. Contamination with adenovirus can be reduced by,

e.g., heat treatment to which adenovirus is more sensitive than AAV. Additional methods for the delivery of nucleic acids to cells are known to those skilled in the art. See, for example, US20030087817, incorporated herein by reference.

[00510] In some embodiments, a host cell is transiently or non-transiently transfected with one or more vectors described herein. In some embodiments, a cell is transfected as it naturally occurs in a subject. In some embodiments, a cell that is transfected is taken from a subject. In some embodiments, the cell is derived from cells taken from a subject, such as a cell line. A wide variety of cell lines for tissue culture are known in the art. Examples of cell lines include, but are not limited to, C8161, CCRF-CEM, MOLT, mIMCD-3, NHDF, HeLa-S3, Huh1, Huh4, Huh7, HUVEC, HASMC, HEK_n, HEK_a, MiaPaCell, Panc1, PC-3, TF1, CTLL-2, C1R, Rat6, CV1, RPTE, A10, T24, J82, A375, ARH-77, Calu1, SW480, SW620, SKOV3, SK-UT, CaCo2, P388D1, SEM-K2, WEHI-231, HB56, TIB55, Jurkat, J45.01, LRMB, Bcl-1, BC-3, IC21, DLD2, Raw264.7, NRK, NRK-52E, MRC5, MEF, Hep G2, HeLa B, HeLa T4, COS, COS-1, COS-6, COS-M6A, BS-C-1 monkey kidney epithelial, BALB/ 3T3 mouse embryo fibroblast, 3T3 Swiss, 3T3-L1, 132-d5 human fetal fibroblasts, 10.1 mouse fibroblasts, 293-T, 3T3, 721, 9L, A2780, A2780ADR, A2780cis, A172, A20, A253, A431, A-549, ALC, B16, B35, BCP-1 cells, BEAS-2B, bEnd.3, BHK-21, BR 293, BxPC3, C3H-10T1/2, C6/36, Cal-27, CHO, CHO-7, CHO-IR, CHO-K1, CHO-K2, CHO-T, CHO Dhfr ^{-/-}, COR-L23, COR-L23/CPR, COR-L23/5010, COR-L23/R23, COS-7, COV-434, CML T1, CMT, CT26, D17, DH82, DU145, DuCaP, EL4, EM2, EM3, EMT6/AR1, EMT6/AR10.0, FM3, H1299, H69, HB54, HB55, HCA2, HEK-293, HeLa, Hepa1c1c7, HL-60, HMEC, HT-29, Jurkat, JY cells, K562 cells, Ku812, KCL22, KG1, KYO1, LNCap, Ma-Mel 1-48, MC-38, MCF-7, MCF-10A, MDA-MB-231, MDA-MB-468, MDA-MB-435, MDCK II, MDCK II, MOR/0.2R, MONO-MAC 6, MTD-1A, MyEnd, NCI-H69/CPR, NCI-H69/LX10, NCI-H69/LX20, NCI-H69/LX4, NIH-3T3, NALM-1, NW-145, OPCN / OPCT cell lines, Peer, PNT-1A / PNT 2, RenCa, RIN-5F, RMA/RMAS, Saos-2 cells, Sf-9, SkBr3, T2, T-47D, T84, THP1 cell line, U373, U87, U937, VCaP, Vero cells, WM39, WT-49, X63, YAC-1, YAR, and transgenic varieties thereof. Cell lines are available from a variety of sources known to those with skill in the art (see, e.g., the American Type Culture Collection (ATCC) (Manassus, Va.)). In some embodiments, a cell transfected with one or more vectors described herein is used to establish a new cell line comprising one or more vector-derived sequences. In some embodiments, a cell transiently transfected with the components of a nucleic acid-targeting

system as described herein (such as by transient transfection of one or more vectors, or transfection with RNA), and modified through the activity of a nucleic acid-targeting complex, is used to establish a new cell line comprising cells containing the modification but lacking any other exogenous sequence. In some embodiments, cells transiently or non-transiently transfected with one or more vectors described herein, or cell lines derived from such cells are used in assessing one or more test compounds.

[00511] In some embodiments, one or more vectors described herein are used to produce a non-human transgenic animal or transgenic plant. In some embodiments, the transgenic animal is a mammal, such as a mouse, rat, or rabbit. In certain embodiments, the organism or subject is a plant. In certain embodiments, the organism or subject or plant is algae. Methods for producing transgenic plants and animals are known in the art, and generally begin with a method of cell transfection, such as described herein.

[00512] In one aspect, the invention provides for methods of modifying a target polynucleotide in a eukaryotic cell. In some embodiments, the method comprises allowing a nucleic acid-targeting complex to bind to the target polynucleotide to effect cleavage of said target polynucleotide thereby modifying the target polynucleotide, wherein the nucleic acid-targeting complex comprises a nucleic acid-targeting effector protein complexed with a guide RNA hybridized to a target sequence within said target polynucleotide.

[00513] In one aspect, the invention provides a method of modifying expression of a polynucleotide in a eukaryotic cell. In some embodiments, the method comprises allowing a nucleic acid-targeting complex to bind to the polynucleotide such that said binding results in increased or decreased expression of said polynucleotide; wherein the nucleic acid-targeting complex comprises a nucleic acid-targeting effector protein complexed with a guide RNA hybridized to a target sequence within said polynucleotide.

CRISPR Systems Can Be Used In Plants

[00514] CRISPR-Cas system(s) (e.g., single or multiplexed) can be used in conjunction with recent advances in crop genomics. Such CRISPR-Cas system(s) can be used to perform efficient and cost effective plant gene or genome interrogation or editing or manipulation—for instance, for rapid investigation and/or selection and/or interrogations and/or comparison and/or manipulations and/or transformation of plant genes or genomes; e.g., to create, identify, develop, optimize, or confer trait(s) or characteristic(s) to plant(s) or to transform a plant genome. There

can accordingly be improved production of plants, new plants with new combinations of traits or characteristics or new plants with enhanced traits. Such CRISPR-Cas system(s) can be used with regard to plants in Site-Directed Integration (SDI) or Gene Editing (GE) or any Near Reverse Breeding (NRB) or Reverse Breeding (RB) techniques. With respect to use of the CRISPR-Cas system in plants, mention is made of the University of Arizona website "CRISPR-PLANT" (<http://www.genome.arizona.edu/crispr/>) (supported by Penn State and AGI). Embodiments of the invention can be used in genome editing in plants or where RNAi or similar genome editing techniques have been used previously; see, e.g., Nekrasov, "Plant genome editing made easy: targeted mutagenesis in model and crop plants using the CRISPR/Cas system," *Plant Methods* 2013, 9:39 (doi:10.1186/1746-4811-9-39); Brooks, "Efficient gene editing in tomato in the first generation using the CRISPR/Cas9 system," *Plant Physiology* September 2014 pp 114.247577; Shan, "Targeted genome modification of crop plants using a CRISPR-Cas system," *Nature Biotechnology* 31, 686-688 (2013); Feng, "Efficient genome editing in plants using a CRISPR/Cas system," *Cell Research* (2013) 23:1229-1232. doi:10.1038/cr.2013.114; published online 20 August 2013; Xie, "RNA-guided genome editing in plants using a CRISPR-Cas system," *Mol Plant*. 2013 Nov;6(6):1975-83. doi: 10.1093/mp/sst119. Epub 2013 Aug 17; Xu, "Gene targeting using the *Agrobacterium tumefaciens*-mediated CRISPR-Cas system in rice," *Rice* 2014, 7:5 (2014), Zhou et al., "Exploiting SNPs for biallelic CRISPR mutations in the outcrossing woody perennial *Populus* reveals 4-coumarate: CoA ligase specificity and Redundancy," *New Phytologist* (2015) (Forum) 1-4 (available online only at www.newphytologist.com); Caliendo et al, "Targeted DNA degradation using a CRISPR device stably carried in the host genome, *NATURE COMMUNICATIONS* 6:6989, DOI: 10.1038/ncomms7989, www.nature.com/naturecommunications DOI: 10.1038/ncomms7989; US Patent No. 6,603,061 - *Agrobacterium*-Mediated Plant Transformation Method; US Patent No. 7,868,149 - Plant Genome Sequences and Uses Thereof and US 2009/0100536 - Transgenic Plants with Enhanced Agronomic Traits, all the contents and disclosure of each of which are herein incorporated by reference in their entirety. In the practice of the invention, the contents and disclosure of Morrell et al "Crop genomics: advances and applications," *Nat Rev Genet*. 2011 Dec 29;13(2):85-96; each of which is incorporated by reference herein including as to how herein embodiments may be used as to plants. Accordingly, reference herein to animal cells may also apply, mutatis mutandis, to plant cells unless otherwise apparent; and, the enzymes herein

having reduced off-target effects and systems employing such enzymes can be used in plant applications, including those mentioned herein.

[00515] Sugano et al. (Plant Cell Physiol. 2014 Mar;55(3):475-81. doi: 10.1093/pcp/pcu014. Epub 2014 Jan 18) reports the application of CRISPR/Cas9 to targeted mutagenesis in the liverwort *Marchantia polymorpha* L., which has emerged as a model species for studying land plant evolution. The U6 promoter of *M. polymorpha* was identified and cloned to express the gRNA. The target sequence of the gRNA was designed to disrupt the gene encoding auxin response factor 1 (ARF1) in *M. polymorpha*. Using *Agrobacterium*-mediated transformation, Sugano et al. isolated stable mutants in the gametophyte generation of *M. polymorpha*. CRISPR/Cas9-based site-directed mutagenesis in vivo was achieved using either the Cauliflower mosaic virus 35S or *M. polymorpha* EF1 α promoter to express Cas9. Isolated mutant individuals showing an auxin-resistant phenotype were not chimeric. Moreover, stable mutants were produced by asexual reproduction of T1 plants. Multiple *arf1* alleles were easily established using CRISPR/Cas9-based targeted mutagenesis. The methods of Sugano et al. may be applied to the CRISPR Cas system of the present invention.

[00516] Kabadi et al. (Nucleic Acids Res. 2014 Oct 29;42(19):e147. doi: 10.1093/nar/gku749. Epub 2014 Aug 13) developed a single lentiviral system to express a Cas9 variant, a reporter gene and up to four sgRNAs from independent RNA polymerase III promoters that are incorporated into the vector by a convenient Golden Gate cloning method. Each sgRNA was efficiently expressed and can mediate multiplex gene editing and sustained transcriptional activation in immortalized and primary human cells. The methods of Kabadi et al. may be applied to the CRISPR Cas system of the present invention.

[00517] Ling et al. (BMC Plant Biology 2014, 14:327) developed a CRISPR/Cas9 binary vector set based on the pGreen or pCAMBIA backbone, as well as a gRNA. This toolkit requires no restriction enzymes besides *Bsa*I to generate final constructs harboring maize-codon optimized Cas9 and one or more gRNAs with high efficiency in as little as one cloning step. The toolkit was validated using maize protoplasts, transgenic maize lines, and transgenic *Arabidopsis* lines and was shown to exhibit high efficiency and specificity. More importantly, using this toolkit, targeted mutations of three *Arabidopsis* genes were detected in transgenic seedlings of the T1 generation. Moreover, the multiple-gene mutations could be inherited by the next

generation. (guide RNA) module vector set, as a toolkit for multiplex genome editing in plants. The toolbox of Lin et al. may be applied to the CRISPR Cas system of the present invention.

[00518] Protocols for targeted plant genome editing via CRISPR/Cas9 are also available in volume 1284 of the series *Methods in Molecular Biology* pp 239-255 10 February 2015. A detailed procedure to design, construct, and evaluate dual gRNAs for plant codon optimized Cas9 (pcoCas9) mediated genome editing using *Arabidopsis thaliana* and *Nicotiana benthamiana* protoplasts s model cellular systems are described. Strategies to apply the CRISPR/Cas9 system to generating targeted genome modifications in whole plants are also discussed. The protocols described in the chapter may be applied to the CRISPR Cas system of the present invention.

[00519] Ma et al. (*Mol Plant*. 2015 Aug 3;8(8):1274-84. doi: 10.1016/j.molp.2015.04.007) reports robust CRISPR/Cas9 vector system, utilizing a plant codon optimized Cas9 gene, for convenient and high-efficiency multiplex genome editing in monocot and dicot plants. Ma et al. designed PCR-based procedures to rapidly generate multiple sgRNA expression cassettes, which can be assembled into the binary CRISPR/Cas9 vectors in one round of cloning by Golden Gate ligation or Gibson Assembly. With this system, Ma et al. edited 46 target sites in rice with an average 85.4% rate of mutation, mostly in biallelic and homozygous status. Ma et al. provide examples of loss-of-function gene mutations in T0 rice and T1 *Arabidopsis* plants by simultaneous targeting of multiple (up to eight) members of a gene family, multiple genes in a biosynthetic pathway, or multiple sites in a single gene. The methods of Ma et al. may be applied to the CRISPR Cas system of the present invention.

[00520] Lowder et al. (*Plant Physiol*. 2015 Aug 21. pii: pp.00636.2015) also developed a CRISPR/Cas9 toolbox enables multiplex genome editing and transcriptional regulation of expressed, silenced or non-coding genes in plants. This toolbox provides researchers with a protocol and reagents to quickly and efficiently assemble functional CRISPR/Cas9 T-DNA constructs for monocots and dicots using Golden Gate and Gateway cloning methods. It comes with a full suite of capabilities, including multiplexed gene editing and transcriptional activation or repression of plant endogenous genes. T-DNA based transformation technology is fundamental to modern plant biotechnology, genetics, molecular biology and physiology. As such, Applicants developed a method for the assembly of Cas9 (WT, nickase or dCas9) and gRNA(s) into a T-DNA destination-vector of interest. The assembly method is based on both Golden Gate assembly and MultiSite Gateway recombination. Three modules are required for

assembly. The first module is a Cas9 entry vector, which contains promoterless Cas9 or its derivative genes flanked by attL1 and attR5 sites. The second module is a gRNA entry vector which contains entry gRNA expression cassettes flanked by attL5 and attL2 sites. The third module includes attR1-attR2-containing destination T-DNA vectors that provide promoters of choice for Cas9 expression. The toolbox of Lowder et al. may be applied to the CRISPR Cas system of the present invention.

[00521] In an advantageous embodiment, the plant may be a tree. The present invention may also utilize the herein disclosed CRISPR Cas system for herbaceous systems (see, e.g., Belhaj et al., *Plant Methods* 9: 39 and Harrison et al., *Genes & Development* 28: 1859–1872). In a particularly advantageous embodiment, the CRISPR Cas system of the present invention may target single nucleotide polymorphisms (SNPs) in trees (see, e.g., Zhou et al., *New Phytologist*, Volume 208, Issue 2, pages 298–301, October 2015). In the Zhou et al. study, the authors applied a CRISPR Cas system in the woody perennial *Populus* using the 4-coumarate:CoA ligase (4CL) gene family as a case study and achieved 100% mutational efficiency for two 4CL genes targeted, with every transformant examined carrying biallelic modifications. In the Zhou et al., study, the CRISPR/Cas9 system was highly sensitive to single nucleotide polymorphisms (SNPs), as cleavage for a third 4CL gene was abolished due to SNPs in the target sequence.

[00522] The methods of Zhou et al. (*New Phytologist*, Volume 208, Issue 2, pages 298–301, October 2015) may be applied to the present invention as follows. Two 4CL genes, 4CL1 and 4CL2, associated with lignin and flavonoid biosynthesis, respectively are targeted for CRISPR/Cas9 editing. The *Populus tremula* × *alba* clone 717-1B4 routinely used for transformation is divergent from the genome-sequenced *Populus trichocarpa*. Therefore, the 4CL1 and 4CL2 gRNAs designed from the reference genome are interrogated with in-house 717 RNA-Seq data to ensure the absence of SNPs which could limit Cas efficiency. A third gRNA designed for 4CL5, a genome duplicate of 4CL1, is also included. The corresponding 717 sequence harbors one SNP in each allele near/within the PAM, both of which are expected to abolish targeting by the 4CL5-gRNA. All three gRNA target sites are located within the first exon. For 717 transformation, the gRNA is expressed from the *Medicago* U6.6 promoter, along with a human codon-optimized Cas under control of the CaMV 35S promoter in a binary vector. Transformation with the Cas-only vector can serve as a control. Randomly selected 4CL1 and

4CL2 lines are subjected to amplicon-sequencing. The data is then processed and biallelic mutations are confirmed in all cases.

[00523] In plants, pathogens are often host-specific. For example, *Fusarium oxysporum* f. sp. *lycopersici* causes tomato wilt but attacks only tomato, and *F. oxysporum* f. *dianthii* *Puccinia graminis* f. sp. *tritici* attacks only wheat. Plants have existing and induced defenses to resist most pathogens. Mutations and recombination events across plant generations lead to genetic variability that gives rise to susceptibility, especially as pathogens reproduce with more frequency than plants. In plants there can be non-host resistance, e.g., the host and pathogen are incompatible. There can also be Horizontal Resistance, e.g., partial resistance against all races of a pathogen, typically controlled by many genes and Vertical Resistance, e.g., complete resistance to some races of a pathogen but not to other races, typically controlled by a few genes. In a Gene-for-Gene level, plants and pathogens evolve together, and the genetic changes in one balance changes in other. Accordingly, using Natural Variability, breeders combine most useful genes for Yield, Quality, Uniformity, Hardiness, Resistance. The sources of resistance genes include native or foreign Varieties, Heirloom Varieties, Wild Plant Relatives, and Induced Mutations, e.g., treating plant material with mutagenic agents. Using the present invention, plant breeders are provided with a new tool to induce mutations. Accordingly, one skilled in the art can analyze the genome of sources of resistance genes, and in Varieties having desired characteristics or traits employ the present invention to induce the rise of resistance genes, with more precision than previous mutagenic agents and hence accelerate and improve plant breeding programs.

CRISPR Systems Can Be Used In Non-Human Organisms / Animals

[00524] The present application may also be extended to other agricultural applications such as, for example, farm and production animals. For example, pigs have many features that make them attractive as biomedical models, especially in regenerative medicine. In particular, pigs with severe combined immunodeficiency (SCID) may provide useful models for regenerative medicine, xenotransplantation, and tumor development and will aid in developing therapies for human SCID patients. Lee et al., (Proc Natl Acad Sci U S A. 2014 May 20;111(20):7260-5) utilized a reporter-guided transcription activator-like effector nuclease (TALEN) system to generate targeted modifications of recombination activating gene (RAG) 2 in somatic cells at

high efficiency, including some that affected both alleles. CRISPR Cas may be applied to a similar system.

[00525] The methods of Lee et al., (Proc Natl Acad Sci U S A. 2014 May 20;111(20):7260-5) may be applied to the present invention as follows. Mutated pigs are produced by targeted modification of RAG2 in fetal fibroblast cells followed by SCNT and embryo transfer. Constructs coding for CRISPR Cas and a reporter are electroporated into fetal-derived fibroblast cells. After 48 h, transfected cells expressing the green fluorescent protein are sorted into individual wells of a 96-well plate at an estimated dilution of a single cell per well. Targeted modification of RAG2 are screened by amplifying a genomic DNA fragment flanking any CRISPR Cas cutting sites followed by sequencing the PCR products. After screening and ensuring lack of off-site mutations, cells carrying targeted modification of RAG2 are used for SCNT. The polar body, along with a portion of the adjacent cytoplasm of oocyte, presumably containing the metaphase II plate, are removed, and a donor cell are placed in the perivitelline. The reconstructed embryos are then electrically porated to fuse the donor cell with the oocyte and then chemically activated. The activated embryos are incubated in Porcine Zygote Medium 3 (PZM3) with 0.5 μ M Scriptaid (S7817; Sigma-Aldrich) for 14–16 h. Embryos are then washed to remove the Scriptaid and cultured in PZM3 until they were transferred into the oviducts of surrogate pigs.

[00526] The present invention is also applicable to modifying SNPs of other animals, such as cows. Tan et al. (Proc Natl Acad Sci U S A. 2013 Oct 8; 110(41): 16526–16531) expanded the livestock gene editing toolbox to include transcription activator-like (TAL) effector nuclease (TALEN)- and clustered regularly interspaced short palindromic repeats (CRISPR)/Cas9-stimulated homology-directed repair (HDR) using plasmid, rAAV, and oligonucleotide templates. Gene specific gRNA sequences were cloned into the Church lab gRNA vector (Addgene ID: 41824) according to their methods (Mali P, et al. (2013) RNA-Guided Human Genome Engineering via Cas9. Science 339(6121):823-826). The Cas9 nuclease was provided either by co-transfection of the hCas9 plasmid (Addgene ID: 41815) or mRNA synthesized from RCIScript-hCas9. This RCIScript-hCas9 was constructed by sub-cloning the XbaI-AgeI fragment from the hCas9 plasmid (encompassing the hCas9 cDNA) into the RCIScript plasmid.

[00527] Heo et al. (Stem Cells Dev. 2015 Feb 1;24(3):393-402. doi: 10.1089/scd.2014.0278. Epub 2014 Nov 3) reported highly efficient gene targeting in the bovine genome using bovine

pluripotent cells and clustered regularly interspaced short palindromic repeat (CRISPR)/Cas9 nuclease. First, Heo et al. generate induced pluripotent stem cells (iPSCs) from bovine somatic fibroblasts by the ectopic expression of yamanaka factors and GSK3 β and MEK inhibitor (2i) treatment. Heo et al. observed that these bovine iPSCs are highly similar to naïve pluripotent stem cells with regard to gene expression and developmental potential in teratomas. Moreover, CRISPR/Cas9 nuclease, which was specific for the bovine NANOG locus, showed highly efficient editing of the bovine genome in bovine iPSCs and embryos.

[00528] Igenity® provides a profile analysis of animals, such as cows, to perform and transmit traits of economic traits of economic importance, such as carcass composition, carcass quality, maternal and reproductive traits and average daily gain. The analysis of a comprehensive Igenity® profile begins with the discovery of DNA markers (most often single nucleotide polymorphisms or SNPs). All the markers behind the Igenity® profile were discovered by independent scientists at research institutions, including universities, research organizations, and government entities such as USDA. Markers are then analyzed at Igenity® in validation populations. Igenity® uses multiple resource populations that represent various production environments and biological types, often working with industry partners from the seedstock, cow-calf, feedlot and/or packing segments of the beef industry to collect phenotypes that are not commonly available. Cattle genome databases are widely available, see, e.g., the NAGRP Cattle Genome Coordination Program (<http://www.animalgenome.org/cattle/maps/db.html>). Thus, the present invention maybe applied to target bovine SNPs. One of skill in the art may utilize the above protocols for targeting SNPs and apply them to bovine SNPs as described, for example, by Tan et al. or Heo et al.

Therapeutic Targeting with RNA-guided Effector Protein Complex

[00529] As will be apparent, it is envisaged that the present system can be used to target any polynucleotide sequence of interest. The invention provides a non-naturally occurring or engineered composition, or one or more polynucleotides encoding components of said composition, or vector or delivery systems comprising one or more polynucleotides encoding components of said composition for use in a modifying a target cell *in vivo*, *ex vivo* or *in vitro* and, may be conducted in a manner alters the cell such that once modified the progeny or cell line of the CRISPR modified cell retains the altered phenotype. The modified cells and progeny may be part of a multi-cellular organism such as a plant or animal with *ex vivo* or *in vivo*

application of CRISPR system to desired cell types. The CRISPR invention may be a therapeutic method of treatment. The therapeutic method of treatment may comprise gene or genome editing, or gene therapy.

Treating pathogens, like bacterial, fungal and parasitic pathogens

[00530] The present invention may also be applied to treat bacterial, fungal and parasitic pathogens. Most research efforts have focused on developing new antibiotics, which once developed, would nevertheless be subject to the same problems of drug resistance. The invention provides novel CRISPR-based alternatives which overcome those difficulties. Furthermore, unlike existing antibiotics, CRISPR-based treatments can be made pathogen specific, inducing bacterial cell death of a target pathogen while avoiding beneficial bacteria.

[00531] Jiang et al. ("RNA-guided editing of bacterial genomes using CRISPR-Cas systems," Nature Biotechnology vol. 31, p. 233-9, March 2013) used a CRISPR-Cas9 system to mutate or kill *S. pneumoniae* and *E. coli*. The work, which introduced precise mutations into the genomes, relied on dual-RNA:Cas9-directed cleavage at the targeted genomic site to kill unmutated cells and circumvented the need for selectable markers or counter-selection systems. CRISPR systems have been used to reverse antibiotic resistance and eliminate the transfer of resistance between strains. Bickard et al. showed that Cas9, reprogrammed to target virulence genes, kills virulent, but not avirulent, *S. aureus*. Reprogramming the nuclease to target antibiotic resistance genes destroyed staphylococcal plasmids that harbor antibiotic resistance genes and immunized against the spread of plasmid-borne resistance genes. (see, Bikard *et al.*, "Exploiting CRISPR-Cas nucleases to produce sequence-specific antimicrobials," Nature Biotechnology vol. 32, 1146–1150, doi:10.1038/nbt.3043, published online 05 October 2014.) Bikard showed that CRISPR-Cas9 antimicrobials function *in vivo* to kill *S. aureus* in a mouse skin colonization model. Similarly, Yosef et al used a CRISPR system to target genes encoding enzymes that confer resistance to β -lactam antibiotics (see Yousef et al., "Temperate and lytic bacteriophages programmed to sensitize and kill antibiotic-resistant bacteria," Proc. Natl. Acad. Sci. USA, vol. 112, p. 7267–7272, doi: 10.1073/pnas.1500107112 published online May 18, 2015).

[00532] CRISPR systems can be used to edit genomes of parasites that are resistant to other genetic approaches. For example, a CRISPR-Cas9 system was shown to introduce double-stranded breaks into the in the *Plasmodium yoelii* genome (see, Zhang et al., "Efficient Editing of Malaria Parasite Genome Using the CRISPR/Cas9 System," mBio. vol. 5, e01414-14, Jul-Aug

2014). Ghorbal *et al.* (“Genome editing in the human malaria parasite *Plasmodium falciparum* using the CRISPR-Cas9 system,” *Nature Biotechnology*, vol. 32, p. 819-821, doi: 10.1038/nbt.2925, published online June 1, 2014) modified the sequences of two genes, *orc1* and *kelch13*, which have putative roles in gene silencing and emerging resistance to artemisinin, respectively. Parasites that were altered at the appropriate sites were recovered with very high efficiency, despite there being no direct selection for the modification, indicating that neutral or even deleterious mutations can be generated using this system. CRISPR-Cas9 is also used to modify the genomes of other pathogenic parasites, including *Toxoplasma gondii* (see Shen *et al.*, “Efficient gene disruption in diverse strains of *Toxoplasma gondii* using CRISPR/CAS9,” *mBio* vol. 5:e01114-14, 2014; and Sidik *et al.*, “Efficient Genome Engineering of *Toxoplasma gondii* Using CRISPR/Cas9,” *PLoS One* vol. 9, e100450, doi: 10.1371/journal.pone.0100450, published online June 27, 2014).

[00533] Vyas *et al.* (“A *Candida albicans* CRISPR system permits genetic engineering of essential genes and gene families,” *Science Advances*, vol. 1, e1500248, DOI: 10.1126/sciadv.1500248, April 3, 2015) employed a CRISPR system to overcome long-standing obstacles to genetic engineering in *C. albicans* and efficiently mutate in a single experiment both copies of several different genes. In an organism where several mechanisms contribute to drug resistance, Vyas produced homozygous double mutants that no longer displayed the hyper-resistance to fluconazole or cycloheximide displayed by the parental clinical isolate Can90. Vyas also obtained homozygous loss-of-function mutations in essential genes of *C. albicans* by creating conditional alleles. Null alleles of *DCR1*, which is required for ribosomal RNA processing, are lethal at low temperature but viable at high temperature. Vyas used a repair template that introduced a nonsense mutation and isolated *der1/der1* mutants that failed to grow at 16°C.

[00534] The CRISPR system of the present invention for use in *P. falciparum* by disrupting chromosomal loci. Ghorbal *et al.* (“Genome editing in the human malaria parasite *Plasmodium falciparum* using the CRISPR-Cas9 system”, *Nature Biotechnology*, 32, 819-821 (2014), DOI: 10.1038/nbt.2925, June 1, 2014) employed a CRISPR system to introduce specific gene knockouts and single-nucleotide substitutions in the malaria genome. To adapt the CRISPR-Cas9 system to *P. falciparum*, Ghorbal *et al.* generated expression vectors for under the control of plasmoidal regulatory elements in the pUF1-Cas9 episome that also carries the drug-selectable

marker *ydhodh*, which gives resistance to DSM1, a *P. falciparum* dihydroorotate dehydrogenase (PfDHODH) inhibitor and for transcription of the sgRNA, used *P. falciparum* U6 small nuclear (sn)RNA regulatory elements placing the guide RNA and the donor DNA template for homologous recombination repair on the same plasmid, pL7. See also, Zhang C. *et al.* (“Efficient editing of malaria parasite genome using the CRISPR/Cas9 system”, *MBio*, 2014 Jul 1; 5(4):E01414-14, doi: 10.1128/MbIO.01414-14) and Wagner *et al.* (“Efficient CRISPR-Cas9-mediated genome editing in *Plasmodium falciparum*”, *Nature Methods* 11, 915-918 (2014), DOI: 10.1038/nmeth.3063).

Treating pathogens, like viral pathogens such as HIV

[00535] Cas-mediated genome editing might be used to introduce protective mutations in somatic tissues to combat nongenetic or complex diseases. For example, NHEJ-mediated inactivation of the CCR5 receptor in lymphocytes (Lombardo *et al.*, *Nat Biotechnol.* 2007 Nov; 25(11):1298-306) may be a viable strategy for circumventing HIV infection, whereas deletion of PCSK9 (Cohen *et al.*, *Nat Genet.* 2005 Feb; 37(2):161-5) orangioipoetin (Musunuru *et al.*, *N Engl J Med.* 2010 Dec 2; 363(23):2220-7) may provide therapeutic effects against statin-resistant hypercholesterolemia or hyperlipidemia. Although these targets may be also addressed using siRNA-mediated protein knockdown, a unique advantage of NHEJ-mediated gene inactivation is the ability to achieve permanent therapeutic benefit without the need for continuing treatment. As with all gene therapies, it will of course be important to establish that each proposed therapeutic use has a favorable benefit-risk ratio.

[00536] Hydrodynamic delivery of plasmid DNA encoding Cas9 and guide RNA along with a repair template into the liver of an adult mouse model of tyrosinemia was shown to be able to correct the mutant *Fah* gene and rescue expression of the wild-type *Fah* protein in ~1 out of 250 cells (*Nat Biotechnol.* 2014 Jun; 32(6):551-3). In addition, clinical trials successfully used ZFNucleases to combat HIV infection by *ex vivo* knockout of the CCR5 receptor. In all patients, HIV DNA levels decreased, and in one out of four patients, HIV RNA became undetectable (Tebas *et al.*, *N Engl J Med.* 2014 Mar 6; 370(10):901-10). Both of these results demonstrate the promise of programmable nucleases as a new therapeutic platform.

[00537] In another embodiment, self-inactivating lentiviral vectors with an siRNA targeting a common exon shared by HIV *tat/rev*, a nucleolar-localizing TAR decoy, and an anti-CCR5-specific hammerhead ribozyme (see, e.g., DiGiusto *et al.* (2010) *Sci Transl Med* 2:36ra43) may

be used/and or adapted to the CRISPR-Cas system of the present invention. A minimum of 2.5×10^6 CD34+ cells per kilogram patient weight may be collected and prestimulated for 16 to 20 hours in X-VIVO 15 medium (Lonza) containing 2 $\mu\text{mol/L}$ -glutamine, stem cell factor (100 ng/ml), Flt-3 ligand (Flt-3L) (100 ng/ml), and thrombopoietin (10 ng/ml) (CellGenix) at a density of 2×10^6 cells/ml. Prestimulated cells may be transduced with lentiviral at a multiplicity of infection of 5 for 16 to 24 hours in 75-cm² tissue culture flasks coated with fibronectin (25 mg/cm²) (RetroNectin, Takara Bio Inc.).

[00538] With the knowledge in the art and the teachings in this disclosure the skilled person can correct HSCs as to immunodeficiency condition such as HIV / AIDS comprising contacting an HSC with a CRISPR-Cas9 system that targets and knocks out CCR5. An guide RNA (and advantageously a dual guide approach, e.g., a pair of different guide RNAs; for instance, guide RNAs targeting of two clinically relevant genes, B2M and CCR5, in primary human CD4+ T cells and CD34+ hematopoietic stem and progenitor cells (HSPCs)) that targets and knocks out CCR5-and-Cas9 protein containing particle is contacted with HSCs. The so contacted cells can be administered; and optionally treated / expanded; cf. Cartier. *See also* Kiem, "Hematopoietic stem cell-based gene therapy for HIV disease," *Cell Stem Cell*. Feb 3, 2012; 10(2): 137–147; incorporated herein by reference along with the documents it cites; Mandal et al, "Efficient Ablation of Genes in Human Hematopoietic Stem and Effector Cells using CRISPR/Cas9," *Cell Stem Cell*, Volume 15, Issue 5, p643–652, 6 November 2014; incorporated herein by reference along with the documents it cites. Mention is also made of Ebina, "CRISPR/Cas9 system to suppress HIV-1 expression by editing HIV-1 integrated proviral DNA" *SCIENTIFIC REPORTS* | 3 : 2510 | DOI: 10.1038/srep02510, incorporated herein by reference along with the documents it cites, as another means for combatting HIV/AIDS using a CRISPR-Cas9 system.

[00539] The rationale for genome editing for HIV treatment originates from the observation that individuals homozygous for loss of function mutations in CCR5, a cellular co-receptor for the virus, are highly resistant to infection and otherwise healthy, suggesting that mimicking this mutation with genome editing could be a safe and effective therapeutic strategy [Liu, R., et al. *Cell* 86, 367-377 (1996)]. This idea was clinically validated when an HIV infected patient was given an allogeneic bone marrow transplant from a donor homozygous for a loss of function CCR5 mutation, resulting in undetectable levels of HIV and restoration of normal CD4 T-cell counts [Hutter, G., et al. *The New England journal of medicine* 360, 692-698 (2009)]. Although

bone marrow transplantation is not a realistic treatment strategy for most HIV patients, due to cost and potential graft vs. host disease, HIV therapies that convert a patient's own T-cells into CCR5 are desirable.

[00540] Early studies using ZFNs and NHEJ to knockout CCR5 in humanized mouse models of HIV showed that transplantation of CCR5 edited CD4 T cells improved viral load and CD4 T-cell counts [Perez, E.E., et al. *Nature biotechnology* 26, 808-816 (2008)]. Importantly, these models also showed that HIV infection resulted in selection for CCR5 null cells, suggesting that editing confers a fitness advantage and potentially allowing a small number of edited cells to create a therapeutic effect.

[00541] As a result of this and other promising preclinical studies, genome editing therapy that knocks out CCR5 in patient T cells has now been tested in humans [Holt, N., et al. *Nature biotechnology* 28, 839-847 (2010); Li, L., et al. *Molecular therapy : the journal of the American Society of Gene Therapy* 21, 1259-1269 (2013)]. In a recent phase I clinical trial, CD4+ T cells from patients with HIV were removed, edited with ZFNs designed to knockout the CCR5 gene, and autologously transplanted back into patients [Tebas, P., et al. *The New England journal of medicine* 370, 901-910 (2014)].

[00542] In another study (Mandal et al., *Cell Stem Cell*, Volume 15, Issue 5, p643–652, 6 November 2014), CRISPR-Cas9 has targeted two clinical relevant genes, B2M and CCR5, in human CD4+ T cells and CD34+ hematopoietic stem and progenitor cells (HSPCs). Use of single RNA guides led to highly efficient mutagenesis in HSPCs but not in T cells. A dual guide approach improved gene deletion efficacy in both cell types. HSPCs that had undergone genome editing with CRISPR-Cas9 retained multilineage potential. Predicted on- and off-target mutations were examined via target capture sequencing in HSPCs and low levels of off-target mutagenesis were observed at only one site. These results demonstrate that CRISPR-Cas9 can efficiently ablate genes in HSPCs with minimal off-target mutagenesis, which have broad applicability for hematopoietic cell-based therapy.

[00543] Wang et al. (*PLoS One*. 2014 Dec 26;9(12):e115987. doi: 10.1371/journal.pone.0115987) silenced CCR5 via CRISPR associated protein 9 (Cas9) and single guided RNAs (guide RNAs) with lentiviral vectors expressing Cas9 and CCR5 guide RNAs. Wang et al. showed that a single round transduction of lentiviral vectors expressing Cas9 and CCR5 guide RNAs into HIV-1 susceptible human CD4+ cells yields high frequencies of

CCR5 gene disruption. CCR5 gene-disrupted cells are not only resistant to R5-tropic HIV-1, including transmitted/founder (T/F) HIV-1 isolates, but also have selective advantage over CCR5 gene-undisrupted cells during R5-tropic HIV-1 infection. Genome mutations at potential off-target sites that are highly homologous to these CCR5 guide RNAs in stably transduced cells even at 84 days post transduction were not detected by a T7 endonuclease I assay.

[00544] Fine et al. (Sci Rep. 2015 Jul 1;5:10777. doi: 10.1038/srep10777) identified a two-cassette system expressing pieces of the *S. pyogenes* Cas9 (SpCas9) protein which splice together in cellula to form a functional protein capable of site-specific DNA cleavage. With specific CRISPR guide strands, Fine et al. demonstrated the efficacy of this system in cleaving the HBB and CCR5 genes in human HEK-293T cells as a single Cas9 and as a pair of Cas9 nickases. The trans-spliced SpCas9 (tsSpCas9) displayed ~35% of the nuclease activity compared with the wild-type SpCas9 (wtSpCas9) at standard transfection doses, but had substantially decreased activity at lower dosing levels. The greatly reduced open reading frame length of the tsSpCas9 relative to wtSpCas9 potentially allows for more complex and longer genetic elements to be packaged into an AAV vector including tissue-specific promoters, multiplexed guide RNA expression, and effector domain fusions to SpCas9.

[00545] Li et al. (J Gen Virol. 2015 Aug;96(8):2381-93. doi: 10.1099/vir.0.000139. Epub 2015 Apr 8) demonstrated that CRISPR-Cas9 can efficiently mediate the editing of the CCR5 locus in cell lines, resulting in the knockout of CCR5 expression on the cell surface. Next-generation sequencing revealed that various mutations were introduced around the predicted cleavage site of CCR5. For each of the three most effective guide RNAs that were analyzed, no significant off-target effects were detected at the 15 top-scoring potential sites. By constructing chimeric Ad5F35 adenoviruses carrying CRISPR-Cas9 components, Li et al. efficiently transduced primary CD4+ T-lymphocytes and disrupted CCR5 expression, and the positively transduced cells were conferred with HIV-1 resistance.

[00546] One of skill in the art may utilize the above studies of, for example, Holt, N., et al. Nature biotechnology 28, 839-847 (2010), Li, L., et al. Molecular therapy : the journal of the American Society of Gene Therapy 21, 1259-1269 (2013), Mandal et al., Cell Stem Cell, Volume 15, Issue 5, p643-652, 6 November 2014, Wang et al. (PLoS One. 2014 Dec 26;9(12):e115987. doi: 10.1371/journal.pone.0115987), Fine et al. (Sci Rep. 2015 Jul 1;5:10777. doi: 10.1038/srep10777) and Li et al. (J Gen Virol. 2015 Aug;96(8):2381-93. doi:

10.1099/vir.0.000139. Epub 2015 Apr 8) for targeting CCR5 with the CRISPR Cas system of the present invention.

Treating pathogens, like viral pathogens, such as HBV

[00547] The present invention may also be applied to treat hepatitis B virus (HBV). However, the CRISPR Cas system must be adapted to avoid the shortcomings of RNAi, such as the risk of oversatring endogenous small RNA pathways, by for example, optimizing dose and sequence (see, e.g., Grimm et al., Nature vol. 441, 26 May 2006). For example, low doses, such as about $1-10 \times 10^{14}$ particles per human are contemplated. In another embodiment, the CRISPR Cas system directed against HBV may be administered in liposomes, such as a stable nucleic-acid-lipid particle (SNALP) (see, e.g., Morrissey et al., Nature Biotechnology, Vol. 23, No. 8, August 2005). Daily intravenous injections of about 1, 3 or 5 mg/kg/day of CRISPR Cas targeted to HBV RNA in a SNALP are contemplated. The daily treatment may be over about three days and then weekly for about five weeks. In another embodiment, the system of Chen et al. (Gene Therapy (2007) 14, 11–19) may be used/and or adapted for the CRISPR Cas system of the present invention. Chen et al. use a double-stranded adenoassociated virus 8-pseudotyped vector (dsAAV2/8) to deliver shRNA. A single administration of dsAAV2/8 vector (1×10^{12} vector genomes per mouse), carrying HBV-specific shRNA, effectively suppressed the steady level of HBV protein, mRNA and replicative DNA in liver of HBV transgenic mice, leading to up to 2–3 \log_{10} decrease in HBV load in the circulation. Significant HBV suppression sustained for at least 120 days after vector administration. The therapeutic effect of shRNA was target sequence dependent and did not involve activation of interferon. For the present invention, a CRISPR Cas system directed to HBV may be cloned into an AAV vector, such as a dsAAV2/8 vector and administered to a human, for example, at a dosage of about 1×10^{15} vector genomes to about 1×10^{16} vector genomes per human. In another embodiment, the method of Wooddell et al. (Molecular Therapy vol. 21 no. 5, 973–985 May 2013) may be used/and or adapted to the CRISPR Cas system of the present invention. Wooddell et al. show that simple coinjection of a hepatocyte-targeted, N-acetylgalactosamine-conjugated melittin-like peptide (NAG-MLP) with a liver-tropic cholesterol-conjugated siRNA (chol-siRNA) targeting coagulation factor VII (F7) results in efficient F7 knockdown in mice and nonhuman primates without changes in clinical chemistry or induction of cytokines. Using transient and transgenic mouse models of HBV infection, Wooddell et al. show that a single coinjection of NAG-MLP with potent chol-siRNAs

targeting conserved HBV sequences resulted in multilog repression of viral RNA, proteins, and viral DNA with long duration of effect. Intravenous coinjections, for example, of about 6 mg/kg of NAG-MLP and 6 mg/kg of HBV specific CRISPR Cas may be envisioned for the present invention. In the alternative, about 3 mg/kg of NAG-MLP and 3 mg/kg of HBV specific CRISPR Cas may be delivered on day one, followed by administration of about 2-3 mg/kg of NAG-MLP and 2-3 mg/kg of HBV specific CRISPR Cas two weeks later.

[00548] Lin et al. (Mol Ther Nucleic Acids. 2014 Aug 19;3:e186. doi: 10.1038/mtna.2014.38) designed eight gRNAs against HBV of genotype A. With the HBV-specific gRNAs, the CRISPR-Cas9 system significantly reduced the production of HBV core and surface proteins in Huh-7 cells transfected with an HBV-expression vector. Among eight screened gRNAs, two effective ones were identified. One gRNA targeting the conserved HBV sequence acted against different genotypes. Using a hydrodynamics-HBV persistence mouse model, Lin et al. further demonstrated that this system could cleave the intrahepatic HBV genome-containing plasmid and facilitate its clearance *in vivo*, resulting in reduction of serum surface antigen levels. These data suggest that the CRISPR-Cas9 system could disrupt the HBV-expressing templates both *in vitro* and *in vivo*, indicating its potential in eradicating persistent HBV infection.

[00549] Dong et al. (Antiviral Res. 2015 Jun;118:110-7. doi: 10.1016/j.antiviral.2015.03.015. Epub 2015 Apr 3) used the CRISPR-Cas9 system to target the HBV genome and efficiently inhibit HBV infection. Dong et al. synthesized four single-guide RNAs (guide RNAs) targeting the conserved regions of HBV. The expression of these guide RNAs with Cas9 reduced the viral production in Huh7 cells as well as in HBV-replication cell HepG2.2.15. Dong et al. further demonstrated that CRISPR-Cas9 direct cleavage and cleavage-mediated mutagenesis occurred in HBV cccDNA of transfected cells. In the mouse model carrying HBV cccDNA, injection of guide RNA-Cas9 plasmids via rapid tail vein resulted in the low level of cccDNA and HBV protein.

[00550] Liu et al. (J Gen Virol. 2015 Aug;96(8):2252-61. doi: 10.1099/vir.0.000159. Epub 2015 Apr 22) designed eight guide RNAs (gRNAs) that targeted the conserved regions of different HBV genotypes, which could significantly inhibit HBV replication both *in vitro* and *in vivo* to investigate the possibility of using the CRISPR-Cas9 system to disrupt the HBV DNA templates. The HBV-specific gRNA/Cas9 system could inhibit the replication of HBV of

different genotypes in cells, and the viral DNA was significantly reduced by a single gRNA/Cas9 system and cleared by a combination of different gRNA/Cas9 systems.

[00551] Wang et al. (World J Gastroenterol. 2015 Aug 28;21(32):9554-65. doi: 10.3748/wjg.v21.i32.9554) designed 15 gRNAs against HBV of genotypes A-D. Eleven combinations of two above gRNAs (dual-gRNAs) covering the regulatory region of HBV were chosen. The efficiency of each gRNA and 11 dual-gRNAs on the suppression of HBV (genotypes A-D) replication was examined by the measurement of HBV surface antigen (HBsAg) or e antigen (HBeAg) in the culture supernatant. The destruction of HBV-expressing vector was examined in HuH7 cells co-transfected with dual-gRNAs and HBV-expressing vector using polymerase chain reaction (PCR) and sequencing method, and the destruction of cccDNA was examined in HepAD38 cells using KCl precipitation, plasmid-safe ATP-dependent DNase (PSAD) digestion, rolling circle amplification and quantitative PCR combined method. The cytotoxicity of these gRNAs was assessed by a mitochondrial tetrazolium assay. All of gRNAs could significantly reduce HBsAg or HBeAg production in the culture supernatant, which was dependent on the region in which gRNA against. All of dual gRNAs could efficiently suppress HBsAg and/or HBeAg production for HBV of genotypes A-D, and the efficacy of dual gRNAs in suppressing HBsAg and/or HBeAg production was significantly increased when compared to the single gRNA used alone. Furthermore, by PCR direct sequencing Applicants confirmed that these dual gRNAs could specifically destroy HBV expressing template by removing the fragment between the cleavage sites of the two used gRNAs. Most importantly, gRNA-5 and gRNA-12 combination not only could efficiently suppressing HBsAg and/or HBeAg production, but also destroy the cccDNA reservoirs in HepAD38 cells.

[00552] Karimova et al. (Sci Rep. 2015 Sep 3;5:13734. doi: 10.1038/srep13734) identified cross-genotype conserved HBV sequences in the S and X region of the HBV genome that were targeted for specific and effective cleavage by a Cas9 nickase. This approach disrupted not only episomal cccDNA and chromosomally integrated HBV target sites in reporter cell lines, but also HBV replication in chronically and de novo infected hepatoma cell lines.

[00553] One of skill in the art may utilize the above studies of, for example, Lin et al. (Mol Ther Nucleic Acids. 2014 Aug 19;3:e186. doi: 10.1038/mtna.2014.38), Dong et al. (Antiviral Res. 2015 Jun;118:110-7. doi: 10.1016/j.antiviral.2015.03.015. Epub 2015 Apr 3), Liu et al. (J Gen Virol. 2015 Aug;96(8):2252-61. doi: 10.1099/vir.0.000159. Epub 2015 Apr 22), Wang et al.

(World J Gastroenterol. 2015 Aug 28;21(32):9554-65. doi: 10.3748/wjg.v21.i32.9554) and Karimova et al. (Sci Rep. 2015 Sep 3;5:13734. doi: 10.1038/srep13734) for targeting HBV with the CRISPR Cas system of the present invention.

[00554] The present invention may also be applied to treat pathogens, e.g. bacterial, fungal and parasitic pathogens. Most research efforts have focused on developing new antibiotics, which once developed, would nevertheless be subject to the same problems of drug resistance. The invention provides novel CRISPR-based alternatives which overcome those difficulties. Furthermore, unlike existing antibiotics, CRISPR-based treatments can be made pathogen specific, inducing bacterial cell death of a target pathogen while avoiding beneficial bacteria.

[00555] Jiang et al. (“RNA-guided editing of bacterial genomes using CRISPR-Cas systems,” Nature Biotechnology vol. 31, p. 233-9, March 2013) used a CRISPR-Cas9 system to mutate or kill *S. pneumoniae* and *E. coli*. The work, which introduced precise mutations into the genomes, relied on dual-RNA:Cas9-directed cleavage at the targeted genomic site to kill unmutated cells and circumvented the need for selectable markers or counter-selection systems. CRISPR systems have been used to reverse antibiotic resistance and eliminate the transfer of resistance between strains. Bickard et al. showed that Cas9, reprogrammed to target virulence genes, kills virulent, but not avirulent, *S. aureus*. Reprogramming the nuclease to target antibiotic resistance genes destroyed staphylococcal plasmids that harbor antibiotic resistance genes and immunized against the spread of plasmid-borne resistance genes. (see, Bikard *et al.*, “Exploiting CRISPR-Cas nucleases to produce sequence-specific antimicrobials,” Nature Biotechnology vol. 32, 1146–1150, doi:10.1038/nbt.3043, published online 05 October 2014.) Bikard showed that CRISPR-Cas9 antimicrobials function *in vivo* to kill *S. aureus* in a mouse skin colonization model. Similarly, Yosef et al used a CRISPR system to target genes encoding enzymes that confer resistance to β -lactam antibiotics (see Yousef et al., “Temperate and lytic bacteriophages programmed to sensitize and kill antibiotic-resistant bacteria,” Proc. Natl. Acad. Sci. USA, vol. 112, p. 7267–7272, doi: 10.1073/pnas.1500107112 published online May 18, 2015).

[00556] CRISPR systems can be used to edit genomes of parasites that are resistant to other genetic approaches. For example, a CRISPR-Cas9 system was shown to introduce double-stranded breaks into the in the *Plasmodium yoelii* genome (see, Zhang et al., “Efficient Editing of Malaria Parasite Genome Using the CRISPR/Cas9 System,” mBio. vol. 5, e01414-14, Jul-Aug 2014). Ghorbal *et al.* (“Genome editing in the human malaria parasite *Plasmodium*

falciparum using the CRISPR-Cas9 system,” Nature Biotechnology, vol. 32, p. 819-821, doi: 10.1038/nbt.2925, published online June 1, 2014) modified the sequences of two genes, *orc1* and *kelch13*, which have putative roles in gene silencing and emerging resistance to artemisinin, respectively. Parasites that were altered at the appropriate sites were recovered with very high efficiency, despite there being no direct selection for the modification, indicating that neutral or even deleterious mutations can be generated using this system. CRISPR-Cas9 is also used to modify the genomes of other pathogenic parasites, including *Toxoplasma gondii* (see Shen *et al.*, “Efficient gene disruption in diverse strains of *Toxoplasma gondii* using CRISPR/CAS9,” mBio vol. 5:e01114-14, 2014; and Sidik *et al.*, “Efficient Genome Engineering of *Toxoplasma gondii* Using CRISPR/Cas9,” PLoS One vol. 9, e100450, doi: 10.1371/journal.pone.0100450, published online June 27, 2014).

[00557] Vyas *et al.* (“A *Candida albicans* CRISPR system permits genetic engineering of essential genes and gene families,” Science Advances, vol. 1, e1500248, DOI: 10.1126/sciadv.1500248, April 3, 2015) employed a CRISPR system to overcome long-standing obstacles to genetic engineering in *C. albicans* and efficiently mutate in a single experiment both copies of several different genes. In an organism where several mechanisms contribute to drug resistance, Vyas produced homozygous double mutants that no longer displayed the hyper-resistance to fluconazole or cycloheximide displayed by the parental clinical isolate Can90. Vyas also obtained homozygous loss-of-function mutations in essential genes of *C. albicans* by creating conditional alleles. Null alleles of *DCR1*, which is required for ribosomal RNA processing, are lethal at low temperature but viable at high temperature. Vyas used a repair template that introduced a nonsense mutation and isolated *dcr1/dcr1* mutants that failed to grow at 16°C.

Patient-specific screening methods

[00558] A CRISPR-Cas system that targets nucleotide, e.g., trinucleotide repeats can be used to screen patients or patient samples for the presence of such repeats. The repeats can be the target of the RNA of the CRISPR-Cas system, and if there is binding thereto by the CRISPR-Cas system, that binding can be detected, to thereby indicate that such a repeat is present. Thus, a CRISPR-Cas system can be used to screen patients or patient samples for the presence of the repeat. The patient can then be administered suitable compound(s) to address the condition; or,

can be administered a CRISPR-Cas system to bind to and cause insertion, deletion or mutation and alleviate the condition.

Treating Diseases with Genetic or Epigenetic Aspects

[00559] The CRISPR-Cas systems of the present invention can be used to correct genetic mutations that were previously attempted with limited success using TALEN and ZFN and have been identified as potential targets for Cas9 systems, including as in published applications of Editas Medicine describing methods to use Cas9 systems to target loci to therapeutically address diseases with gene therapy, including, WO 2015/048577 CRISPR-RELATED METHODS AND COMPOSITIONS of Glucksmann et al.; WO 2015/070083 CRISPR-RELATED METHODS AND COMPOSITIONS WITH GOVERNING gRNAs of Glucksmann et al.; WO 2015/134812 CRISPR/CAS-RELATED METHODS AND COMPOSITIONS FOR TREATING USHER SYNDROME AND RETINITIS PIGMENTOSA of Maeder et al.; and WO 2015/138510 CRISPR/CAS-RELATED METHODS AND COMPOSITIONS FOR TREATING LEBER'S CONGENITAL AMAUROSIS 10 (LCA10) of Maeder et al.

[00560] Researchers are contemplating whether gene therapies could be employed to treat a wide range of diseases. The CRISPR systems of the present invention based on Cas9 effector protein are envisioned for such therapeutic uses, including, but not limited to further exemplified targeted areas and with delivery methods as below. Some examples of conditions or diseases that might be usefully treated using the present system are included in the examples of genes and references included herein and are currently associated with those conditions are also provided there. The genes and conditions exemplified are not exhaustive.

Treating Diseases of the Circulatory System

[00561] The present invention also contemplates delivering the CRISPR-Cas system, specifically the novel CRISPR effector protein systems described herein, to the blood or hematopoietic stem cells. The plasma exosomes of Wahlgren et al. (Nucleic Acids Research, 2012, Vol. 40, No. 17 e130) were previously described and may be utilized to deliver the CRISPR Cas system to the blood. The nucleic acid-targeting system of the present invention is also contemplated to treat hemoglobinopathies, such as thalassemias and sickle cell disease. See, e.g., International Patent Publication No. WO 2013/126794 for potential targets that may be targeted by the CRISPR Cas system of the present invention.

[00562] Drakopoulou, “Review Article, The Ongoing Challenge of Hematopoietic Stem Cell-Based Gene Therapy for β -Thalassemia,” *Stem Cells International*, Volume 2011, Article ID 987980, 10 pages, doi:10.4061/2011/987980, incorporated herein by reference along with the documents it cites, as if set out in full, discuss modifying HSCs using a lentivirus that delivers a gene for β -globin or γ -globin. In contrast to using lentivirus, with the knowledge in the art and the teachings in this disclosure, the skilled person can correct HSCs as to β -Thalassemia using a CRISPR-Cas system that targets and corrects the mutation (e.g., with a suitable HDR template that delivers a coding sequence for β -globin or γ -globin, advantageously non-sickling β -globin or γ -globin); specifically, the guide RNA can target mutation that give rise to β -Thalassemia, and the HDR can provide coding for proper expression of β -globin or γ -globin. A guide RNA that targets the mutation-and-Cas protein containing particle is contacted with HSCs carrying the mutation. The particle also can contain a suitable HDR template to correct the mutation for proper expression of β -globin or γ -globin; or the HSC can be contacted with a second particle or a vector that contains or delivers the HDR template. The so contacted cells can be administered; and optionally treated / expanded; cf. Cartier. In this regard mention is made of: Cavazzana, “Outcomes of Gene Therapy for β -Thalassemia Major via Transplantation of Autologous Hematopoietic Stem Cells Transduced Ex Vivo with a Lentiviral β^{A-T87Q} -Globin Vector.” tif2014.org/abstractFiles/Jean%20Antoine%20Ribeil_Abtract.pdf; Cavazzana-Calvo, “Transfusion independence and HMGA2 activation after gene therapy of human β -thalassaemia”, *Nature* 467, 318–322 (16 September 2010) doi:10.1038/nature09328; Nienhuis, “Development of Gene Therapy for Thalassemia, *Cold Spring Harbor Perspectives in Medicine*, doi: 10.1101/cshperspect.a011833 (2012), LentiGlobin BB305, a lentiviral vector containing an engineered β -globin gene ($\beta A-T87Q$); and Xie et al., “Seamless gene correction of β -thalassaemia mutations in patient-specific iPSCs using CRISPR/Cas9 and piggyback” *Genome Research* gr.173427.114 (2014) <http://www.genome.org/cgi/doi/10.1101/gr.173427.114> (*Cold Spring Harbor Laboratory Press*); that is the subject of Cavazzana work involving human β -thalassaemia and the subject of the Xie work, are all incorporated herein by reference, together with all documents cited therein or associated therewith. In the instant invention, the HDR template can provide for the HSC to express an engineered β -globin gene (e.g., $\beta A-T87Q$), or β -globin as in Xie.

[00563] Xu et al. (Sci Rep. 2015 Jul 9;5:12065. doi: 10.1038/srep12065) have designed TALENs and CRISPR-Cas9 to directly target the intron2 mutation site IVS2-654 in the globin gene. Xu et al. observed different frequencies of double-strand breaks (DSBs) at IVS2-654 loci using TALENs and CRISPR-Cas9, and TALENs mediated a higher homologous gene targeting efficiency compared to CRISPR-Cas9 when combined with the piggyBac transposon donor. In addition, more obvious off-target events were observed for CRISPR-Cas9 compared to TALENs. Finally, TALENs-corrected iPSC clones were selected for erythroblast differentiation using the OP9 co-culture system and detected relatively higher transcription of HBB than the uncorrected cells.

[00564] Song et al. (Stem Cells Dev. 2015 May 1;24(9):1053-65. doi: 10.1089/scd.2014.0347. Epub 2015 Feb 5) used CRISPR/ Cas9 to correct β -Thal iPSCs; gene-corrected cells exhibit normal karyotypes and full pluripotency as human embryonic stem cells (hESCs) showed no off-targeting effects. Then, Song et al. evaluated the differentiation efficiency of the gene-corrected β -Thal iPSCs. Song et al. found that during hematopoietic differentiation, gene-corrected β -Thal iPSCs showed an increased embryoid body ratio and various hematopoietic progenitor cell percentages. More importantly, the gene-corrected β -Thal iPSC lines restored HBB expression and reduced reactive oxygen species production compared with the uncorrected group. Song et al.'s study suggested that hematopoietic differentiation efficiency of β -Thal iPSCs was greatly improved once corrected by the CRISPR-Cas9 system. Similar methods may be performed utilizing the CRISPR-Cas systems described herein, e.g. systems comprising Cas9 effector proteins.

[00565] Sickle cell anemia is an autosomal recessive genetic disease in which red blood cells become sickle-shaped. It is caused by a single base substitution in the β -globin gene, which is located on the short arm of chromosome 11. As a result, valine is produced instead of glutamic acid causing the production of sickle hemoglobin (HbS). This results in the formation of a distorted shape of the erythrocytes. Due to this abnormal shape, small blood vessels can be blocked, causing serious damage to the bone, spleen and skin tissues. This may lead to episodes of pain, frequent infections, hand-foot syndrome or even multiple organ failure. The distorted erythrocytes are also more susceptible to hemolysis, which leads to serious anemia. As in the case of β -thalassaemia, sickle cell anemia can be corrected by modifying HSCs with the CRISPR-Cas system. The system allows the specific editing of the cell's genome by cutting its

DNA and then letting it repair itself. The Cas protein is inserted and directed by a RNA guide to the mutated point and then it cuts the DNA at that point. Simultaneously, a healthy version of the sequence is inserted. This sequence is used by the cell's own repair system to fix the induced cut. In this way, the CRISPR-Cas allows the correction of the mutation in the previously obtained stem cells. With the knowledge in the art and the teachings in this disclosure, the skilled person can correct HSCs as to sickle cell anemia using a CRISPR-Cas system that targets and corrects the mutation (e.g., with a suitable HDR template that delivers a coding sequence for β -globin, advantageously non-sickling β -globin); specifically, the guide RNA can target mutation that give rise to sickle cell anemia, and the HDR can provide coding for proper expression of β -globin. An guide RNA that targets the mutation-and-Cas protein containing particle is contacted with HSCs carrying the mutation. The particle also can contain a suitable HDR template to correct the mutation for proper expression of β -globin; or the HSC can be contacted with a second particle or a vector that contains or delivers the HDR template. The so contacted cells can be administered; and optionally treated / expanded; cf. Cartier. The HDR template can provide for the HSC to express an engineered β -globin gene (e.g., β A-T87Q), or β -globin as in Xie.

[00566] Williams, "Broadening the Indications for Hematopoietic Stem Cell Genetic Therapies," *Cell Stem Cell* 13:263-264 (2013), incorporated herein by reference along with the documents it cites, as if set out in full, report lentivirus-mediated gene transfer into HSC/P cells from patients with the lysosomal storage disease metachromatic leukodystrophy disease (MLD), a genetic disease caused by deficiency of arylsulfatase A (ARSA), resulting in nerve demyelination; and lentivirus-mediated gene transfer into HSCs of patients with Wiskott-Aldrich syndrome (WAS) (patients with defective WAS protein, an effector of the small GTPase CDC42 that regulates cytoskeletal function in blood cell lineages and thus suffer from immune deficiency with recurrent infections, autoimmune symptoms, and thrombocytopenia with abnormally small and dysfunctional platelets leading to excessive bleeding and an increased risk of leukemia and lymphoma). In contrast to using lentivirus, with the knowledge in the art and the teachings in this disclosure, the skilled person can correct HSCs as to MLD (deficiency of arylsulfatase A (ARSA)) using a CRISPR-Cas system that targets and corrects the mutation (deficiency of arylsulfatase A (ARSA)) (e.g., with a suitable HDR template that delivers a coding sequence for ARSA); specifically, the guide RNA can target mutation that gives rise to MLD (deficient ARSA), and the HDR can provide coding for proper expression of ARSA. A

guide RNA that targets the mutation-and-Cas protein containing particle is contacted with HSCs carrying the mutation. The particle also can contain a suitable HDR template to correct the mutation for proper expression of ARSA; or the HSC can be contacted with a second particle or a vector that contains or delivers the HDR template. The so contacted cells can be administered; and optionally treated / expanded; cf. Cartier. In contrast to using lentivirus, with the knowledge in the art and the teachings in this disclosure, the skilled person can correct HSCs as to WAS using a CRISPR-Cas system that targets and corrects the mutation (deficiency of WAS protein) (e.g., with a suitable HDR template that delivers a coding sequence for WAS protein); specifically, the guide RNA can target mutation that gives rise to WAS (deficient WAS protein), and the HDR can provide coding for proper expression of WAS protein. A guide RNA that targets the mutation-and-Cas9 protein containing particle is contacted with HSCs carrying the mutation. The particle also can contain a suitable HDR template to correct the mutation for proper expression of WAS protein; or the HSC can be contacted with a second particle or a vector that contains or delivers the HDR template. The so contacted cells can be administered; and optionally treated / expanded; cf. Cartier.

[00567] Watts, "Hematopoietic Stem Cell Expansion and Gene Therapy" *Cytherapy* 13(10):1164–1171. doi:10.3109/14653249.2011.620748 (2011), incorporated herein by reference along with the documents it cites, as if set out in full, discusses hematopoietic stem cell (HSC) gene therapy, e.g., virus-mediated HSC gene therapy, as an highly attractive treatment option for many disorders including hematologic conditions, immunodeficiencies including HIV/AIDS, and other genetic disorders like lysosomal storage diseases, including SCID-X1, ADA-SCID, β -thalassemia, X-linked CGD, Wiskott-Aldrich syndrome, Fanconi anemia, adrenoleukodystrophy (ALD), and metachromatic leukodystrophy (MLD).

[00568] US Patent Publication Nos. 20110225664, 20110091441, 20100229252, 20090271881 and 20090222937 assigned to Cellectis, relates to CREI variants, wherein at least one of the two I-CreI monomers has at least two substitutions, one in each of the two functional subdomains of the LAGLIDADG core domain situated respectively from positions 26 to 40 and 44 to 77 of I-CreI, said variant being able to cleave a DNA target sequence from the human interleukin-2 receptor gamma chain (IL2RG) gene also named common cytokine receptor gamma chain gene or gamma C gene. The target sequences identified in US Patent Publication

Nos. 20110225664, 20110091441, 20100229252, 20090271881 and 20090222937 may be utilized for the nucleic acid-targeting system of the present invention.

[00569] Severe Combined Immune Deficiency (SCID) results from a defect in lymphocytes T maturation, always associated with a functional defect in lymphocytes B (Cavazzana-Calvo et al., *Annu. Rev. Med.*, 2005, 56, 585-602; Fischer et al., *Immunol. Rev.*, 2005, 203, 98-109). Overall incidence is estimated to 1 in 75 000 births. Patients with untreated SCID are subject to multiple opportunist micro-organism infections, and do generally not live beyond one year. SCID can be treated by allogenic hematopoietic stem cell transfer, from a familial donor. Histocompatibility with the donor can vary widely. In the case of Adenosine Deaminase (ADA) deficiency, one of the SCID forms, patients can be treated by injection of recombinant Adenosine Deaminase enzyme.

[00570] Since the ADA gene has been shown to be mutated in SCID patients (Giblett et al., *Lancet*, 1972, 2, 1067-1069), several other genes involved in SCID have been identified (Cavazzana-Calvo et al., *Annu. Rev. Med.*, 2005, 56, 585-602; Fischer et al., *Immunol. Rev.*, 2005, 203, 98-109). There are four major causes for SCID: (i) the most frequent form of SCID, SCID-X1 (X-linked SCID or X-SCID), is caused by mutation in the IL2RG gene, resulting in the absence of mature T lymphocytes and NK cells. IL2RG encodes the gamma C protein (Noguchi, et al., *Cell*, 1993, 73, 147-157), a common component of at least five interleukin receptor complexes. These receptors activate several targets through the JAK3 kinase (Macchi et al., *Nature*, 1995, 377, 65-68), which inactivation results in the same syndrome as gamma C inactivation; (ii) mutation in the ADA gene results in a defect in purine metabolism that is lethal for lymphocyte precursors, which in turn results in the quasi absence of B, T and NK cells; (iii) V(D)J recombination is an essential step in the maturation of immunoglobulins and T lymphocytes receptors (TCRs). Mutations in Recombination Activating Gene 1 and 2 (RAG1 and RAG2) and Artemis, three genes involved in this process, result in the absence of mature T and B lymphocytes; and (iv) Mutations in other genes such as CD45, involved in T cell specific signaling have also been reported, although they represent a minority of cases (Cavazzana-Calvo et al., *Annu. Rev. Med.*, 2005, 56, 585-602; Fischer et al., *Immunol. Rev.*, 2005, 203, 98-109). Since when their genetic bases have been identified, the different SCID forms have become a paradigm for gene therapy approaches (Fischer et al., *Immunol. Rev.*, 2005, 203, 98-109) for two major reasons. First, as in all blood diseases, an ex vivo treatment can be envisioned.

Hematopoietic Stem Cells (HSCs) can be recovered from bone marrow, and keep their pluripotent properties for a few cell divisions. Therefore, they can be treated in vitro, and then reinjected into the patient, where they repopulate the bone marrow. Second, since the maturation of lymphocytes is impaired in SCID patients, corrected cells have a selective advantage. Therefore, a small number of corrected cells can restore a functional immune system. This hypothesis was validated several times by (i) the partial restoration of immune functions associated with the reversion of mutations in SCID patients (Hirschhorn et al., *Nat. Genet.*, 1996, 13, 290-295; Stephan et al., *N. Engl. J. Med.*, 1996, 335, 1563-1567; Bousso et al., *Proc. Natl. Acad. Sci. USA*, 2000, 97, 274-278; Wada et al., *Proc. Natl. Acad. Sci. USA*, 2001, 98, 8697-8702; Nishikomori et al., *Blood*, 2004, 103, 4565-4572), (ii) the correction of SCID-X1 deficiencies in vitro in hematopoietic cells (Candotti et al., *Blood*, 1996, 87, 3097-3102; Cavazzana-Calvo et al., *Blood*, 1996, *Blood*, 88, 3901-3909; Taylor et al., *Blood*, 1996, 87, 3103-3107; Hacein-Bey et al., *Blood*, 1998, 92, 4090-4097), (iii) the correction of SCID-X1 (Soudais et al., *Blood*, 2000, 95, 3071-3077; Tsai et al., *Blood*, 2002, 100, 72-79), JAK-3 (Bunting et al., *Nat. Med.*, 1998, 4, 58-64; Bunting et al., *Hum. Gene Ther.*, 2000, 11, 2353-2364) and RAG2 (Yates et al., *Blood*, 2002, 100, 3942-3949) deficiencies in vivo in animal models and (iv) by the result of gene therapy clinical trials (Cavazzana-Calvo et al., *Science*, 2000, 288, 669-672; Aiuti et al., *Nat. Med.*, 2002; 8, 423-425; Gaspar et al., *Lancet*, 2004, 364, 2181-2187).

[00571] US Patent Publication No. 20110182867 assigned to the Children's Medical Center Corporation and the President and Fellows of Harvard College relates to methods and uses of modulating fetal hemoglobin expression (HbF) in a hematopoietic progenitor cells via inhibitors of BCL11A expression or activity, such as RNAi and antibodies. The targets disclosed in US Patent Publication No. 20110182867, such as BCL11A, may be targeted by the CRISPR Cas system of the present invention for modulating fetal hemoglobin expression. See also Bauer et al. (*Science* 11 October 2013: Vol. 342 no. 6155 pp. 253-257) and Xu et al. (*Science* 18 November 2011: Vol. 334 no. 6058 pp. 993-996) for additional BCL11A targets.

[00572] With the knowledge in the art and the teachings in this disclosure, the skilled person can correct HSCs as to a genetic hematologic disorder, e.g., β -Thalassemia, Hemophilia, or a genetic lysosomal storage disease.

Treating Disease of the Brain, Central Nervous and Immune Systems

[00573] The present invention also contemplates delivering the CRISPR-Cas system to the brain or neurons. For example, RNA interference (RNAi) offers therapeutic potential for this disorder by reducing the expression of *HTT*, the disease-causing gene of Huntington's disease (see, e.g., McBride et al., *Molecular Therapy* vol. 19 no. 12 Dec. 2011, pp. 2152-2162), therefore Applicant postulates that it may be used/and or adapted to the CRISPR-Cas system. The CRISPR-Cas system may be generated using an algorithm to reduce the off-targeting potential of antisense sequences. The CRISPR-Cas sequences may target either a sequence in exon 52 of mouse, rhesus or human huntingtin and expressed in a viral vector, such as AAV. Animals, including humans, may be injected with about three microinjections per hemisphere (six injections total): the first 1 mm rostral to the anterior commissure (12 μ l) and the two remaining injections (12 μ l and 10 μ l, respectively) spaced 3 and 6 mm caudal to the first injection with 1×10^{12} vg/ml of AAV at a rate of about 1 μ l/minute, and the needle was left in place for an additional 5 minutes to allow the injectate to diffuse from the needle tip.

[00574] DiFiglia et al. (*PNAS*, October 23, 2007, vol. 104, no. 43, 17204–17209) observed that single administration into the adult striatum of an siRNA targeting Htt can silence mutant Htt, attenuate neuronal pathology, and delay the abnormal behavioral phenotype observed in a rapid-onset, viral transgenic mouse model of HD. DiFiglia injected mice intrastrially with 2 μ l of Cy3-labeled cc-siRNA-Htt or unconjugated siRNA-Htt at 10 μ M. A similar dosage of CRISPR Cas targeted to Htt may be contemplated for humans in the present invention, for example, about 5-10 ml of 10 μ M CRISPR Cas targeted to Htt may be injected intrastrially.

[00575] In another example, Boudreau et al. (*Molecular Therapy* vol. 17 no. 6 June 2009) injects 5 μ l of recombinant AAV serotype 2/1 vectors expressing htt-specific RNAi virus (at 4×10^{12} viral genomes/ml) into the striatum. A similar dosage of CRISPR Cas targeted to Htt may be contemplated for humans in the present invention, for example, about 10-20 ml of 4×10^{12} viral genomes/ml CRISPR Cas9 targeted to Htt may be injected intrastrially.

[00576] In another example, a CRISPR Cas targeted to HTT may be administered continuously (see, e.g., Yu et al., *Cell* 150, 895–908, August 31, 2012). Yu et al. utilizes osmotic pumps delivering 0.25 ml/hr (Model 2004) to deliver 300 mg/day of ss-siRNA or phosphate-buffered saline (PBS) (Sigma Aldrich) for 28 days, and pumps designed to deliver 0.5 μ l/hr (Model 2002) were used to deliver 75 mg/day of the positive control MOE ASO for 14 days.

Pumps (Durect Corporation) were filled with ss-siRNA or MOE diluted in sterile PBS and then incubated at 37 C for 24 or 48 (Model 2004) hours prior to implantation. Mice were anesthetized with 2.5% isoflurane, and a midline incision was made at the base of the skull. Using stereotaxic guides, a cannula was implanted into the right lateral ventricle and secured with Loctite adhesive. A catheter attached to an Alzet osmotic mini pump was attached to the cannula, and the pump was placed subcutaneously in the midscapular area. The incision was closed with 5.0 nylon sutures. A similar dosage of CRISPR Cas targeted to Htt may be contemplated for humans in the present invention, for example, about 500 to 1000 g/day CRISPR Cas targeted to Htt may be administered.

[00577] In another example of continuous infusion, Stiles et al. (*Experimental Neurology* 233 (2012) 463–471) implanted an intraparenchymal catheter with a titanium needle tip into the right putamen. The catheter was connected to a SynchroMed® II Pump (Medtronic Neurological, Minneapolis, MN) subcutaneously implanted in the abdomen. After a 7 day infusion of phosphate buffered saline at 6 µL/day, pumps were re-filled with test article and programmed for continuous delivery for 7 days. About 2.3 to 11.52 mg/d of siRNA were infused at varying infusion rates of about 0.1 to 0.5 µL/min. A similar dosage of CRISPR Cas targeted to Htt may be contemplated for humans in the present invention, for example, about 20 to 200 mg/day CRISPR Cas targeted to Htt may be administered. In another example, the methods of US Patent Publication No. 20130253040 assigned to Sangamo may also be adapted from TALEs to the nucleic acid-targeting system of the present invention for treating Huntington's Disease.

[00578] A further aspect of the invention relates to utilizing the CRISPR-Cas system for correcting defects in the EMP2A and EMP2B genes that have been identified to be associated with Lafora disease. Lafora disease is an autosomal recessive condition which is characterized by progressive myoclonus epilepsy which may start as epileptic seizures in adolescence. A few cases of the disease may be caused by mutations in genes yet to be identified. The disease causes seizures, muscle spasms, difficulty walking, dementia, and eventually death. There is currently no therapy that has proven effective against disease progression. Other genetic abnormalities associated with epilepsy may also be targeted by the CRISPR-Cas system and the underlying genetics is further described in *Genetics of Epilepsy and Genetic Epilepsies*, edited by Giuliano Avanzini, Jeffrey L. Noebels, Mariani Foundation Paediatric Neurology:20; 2009).

[00579] The methods of US Patent Publication No. 20110158957 assigned to Sangamo BioSciences, Inc. involved in inactivating T cell receptor (TCR) genes may also be modified to the CRISPR Cas system of the present invention. In another example, the methods of US Patent Publication No. 20100311124 assigned to Sangamo BioSciences, Inc. and US Patent Publication No. 20110225664 assigned to Collectis, which are both involved in inactivating glutamine synthetase gene expression genes may also be modified to the CRISPR Cas system of the present invention.

Treating Hearing Diseases

[00580] The present invention also contemplates delivering the CRISPR-Cas system to one or both ears.

[00581] Researchers are looking into whether gene therapy could be used to aid current deafness treatments - namely, cochlear implants. Deafness is often caused by lost or damaged hair cells that cannot relay signals to auditory neurons. In such cases, cochlear implants may be used to respond to sound and transmit electrical signals to the nerve cells. But these neurons often degenerate and retract from the cochlea as fewer growth factors are released by impaired hair cells.

[00582] US patent application 20120328580 describes injection of a pharmaceutical composition into the ear (e.g., auricular administration), such as into the luminae of the cochlea (e.g., the Scala media, Sc vestibulae, and Sc tympani), e.g., using a syringe, e.g., a single-dose syringe. For example, one or more of the compounds described herein can be administered by intratympanic injection (e.g., into the middle ear), and/or injections into the outer, middle, and/or inner ear. Such methods are routinely used in the art, for example, for the administration of steroids and antibiotics into human ears. Injection can be, for example, through the round window of the ear or through the cochlear capsule. Other inner ear administration methods are known in the art (see, e.g., Salt and Plontke, *Drug Discovery Today*, 10:1299-1306, 2005).

[00583] In another mode of administration, the pharmaceutical composition can be administered in situ, via a catheter or pump. A catheter or pump can, for example, direct a pharmaceutical composition into the cochlear luminae or the round window of the ear and/or the lumen of the colon. Exemplary drug delivery apparatus and methods suitable for administering one or more of the compounds described herein into an ear, e.g., a human ear, are described by McKenna et al., (U.S. Publication No. 2006/0030837) and Jacobsen et al., (U.S. Pat. No.

7,206,639). In some embodiments, a catheter or pump can be positioned, e.g., in the ear (e.g., the outer, middle, and/or inner ear) of a patient during a surgical procedure. In some embodiments, a catheter or pump can be positioned, e.g., in the ear (e.g., the outer, middle, and/or inner ear) of a patient without the need for a surgical procedure.

[00584] Alternatively or in addition, one or more of the compounds described herein can be administered in combination with a mechanical device such as a cochlear implant or a hearing aid, which is worn in the outer ear. An exemplary cochlear implant that is suitable for use with the present invention is described by Edge et al., (U.S. Publication No. 2007/0093878).

[00585] In some embodiments, the modes of administration described above may be combined in any order and can be simultaneous or interspersed.

[00586] Alternatively or in addition, the present invention may be administered according to any of the Food and Drug Administration approved methods, for example, as described in CDER Data Standards Manual, version number 004 (which is available at fda.give/cder/dsm/DRG/drg00301.htm).

[00587] In general, the cell therapy methods described in US patent application 20120328580 can be used to promote complete or partial differentiation of a cell to or towards a mature cell type of the inner ear (e.g., a hair cell) *in vitro*. Cells resulting from such methods can then be transplanted or implanted into a patient in need of such treatment. The cell culture methods required to practice these methods, including methods for identifying and selecting suitable cell types, methods for promoting complete or partial differentiation of selected cells, methods for identifying complete or partially differentiated cell types, and methods for implanting complete or partially differentiated cells are described below.

[00588] Cells suitable for use in the present invention include, but are not limited to, cells that are capable of differentiating completely or partially into a mature cell of the inner ear, e.g., a hair cell (e.g., an inner and/or outer hair cell), when contacted, e.g., *in vitro*, with one or more of the compounds described herein. Exemplary cells that are capable of differentiating into a hair cell include, but are not limited to stem cells (e.g., inner ear stem cells, adult stem cells, bone marrow derived stem cells, embryonic stem cells, mesenchymal stem cells, skin stem cells, iPS cells, and fat derived stem cells), progenitor cells (e.g., inner ear progenitor cells), support cells (e.g., Deiters' cells, pillar cells, inner phalangeal cells, tectal cells and Hensen's cells), and/or germ cells. The use of stem cells for the replacement of inner ear sensory cells is described in Li

et al., (U.S. Publication No. 2005/0287127) and Li et al., (U.S. patent Ser. No. 11/953,797). The use of bone marrow derived stem cells for the replacement of inner ear sensory cells is described in Edge et al., PCT/US2007/084654. iPS cells are described, e.g., at Takahashi et al., *Cell*, Volume 131, Issue 5, Pages 861-872 (2007); Takahashi and Yamanaka, *Cell* 126, 663-76 (2006); Okita et al., *Nature* 448, 260-262 (2007); Yu, J. et al., *Science* 318(5858):1917-1920 (2007); Nakagawa et al., *Nat. Biotechnol.* 26:101-106 (2008); and Zaehres and Scholer, *Cell* 131(5):834-835 (2007). Such suitable cells can be identified by analyzing (e.g., qualitatively or quantitatively) the presence of one or more tissue specific genes. For example, gene expression can be detected by detecting the protein product of one or more tissue-specific genes. Protein detection techniques involve staining proteins (e.g., using cell extracts or whole cells) using antibodies against the appropriate antigen. In this case, the appropriate antigen is the protein product of the tissue-specific gene expression. Although, in principle, a first antibody (i.e., the antibody that binds the antigen) can be labeled, it is more common (and improves the visualization) to use a second antibody directed against the first (e.g., an anti-IgG). This second antibody is conjugated either with fluorochromes, or appropriate enzymes for colorimetric reactions, or gold beads (for electron microscopy), or with the biotin-avidin system, so that the location of the primary antibody, and thus the antigen, can be recognized.

[00589] The CRISPR Cas molecules of the present invention may be delivered to the ear by direct application of pharmaceutical composition to the outer ear, with compositions modified from US Published application, 20110142917. In some embodiments the pharmaceutical composition is applied to the ear canal. Delivery to the ear may also be referred to as aural or otic delivery.

[00590] In some embodiments the RNA molecules of the invention are delivered in liposome or lipofectin formulations and the like and can be prepared by methods well known to those skilled in the art. Such methods are described, for example, in U.S. Pat. Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference.

[00591] Delivery systems aimed specifically at the enhanced and improved delivery of siRNA into mammalian cells have been developed, (see, for example, Shen et al *FEBS Let.* 2003, 539:111-114; Xia et al., *Nat. Biotech.* 2002, 20:1006-1010; Reich et al., *Mol. Vision.* 2003, 9: 210-216; Sorensen et al., *J. Mol. Biol.* 2003, 327: 761-766; Lewis et al., *Nat. Gen.* 2002, 32: 107-108 and Simeoni et al., *NAR* 2003, 31, 11: 2717-2724) and may be applied to the

present invention. siRNA has recently been successfully used for inhibition of gene expression in primates (see for example, Tolentino et al., *Retina* 24(4):660 which may also be applied to the present invention.

[00592] Qi et al. discloses methods for efficient siRNA transfection to the inner ear through the intact round window by a novel proteidic delivery technology which may be applied to the nucleic acid-targeting system of the present invention (see, e.g., Qi et al., *Gene Therapy* (2013), 1-9). In particular, a TAT double stranded RNA-binding domains (TAT-DRBDs), which can transfect Cy3-labeled siRNA into cells of the inner ear, including the inner and outer hair cells, crista ampullaris, macula utriculi and macula sacculi, through intact round-window permeation was successful for delivering double stranded siRNAs in vivo for treating various inner ear ailments and preservation of hearing function. About 40 μ l of 10mM RNA may be contemplated as the dosage for administration to the ear.

[00593] According to Rejali et al. (*Hear Res.* 2007 Jun;228(1-2):180-7), cochlear implant function can be improved by good preservation of the spiral ganglion neurons, which are the target of electrical stimulation by the implant and brain derived neurotrophic factor (BDNF) has previously been shown to enhance spiral ganglion survival in experimentally deafened ears. Rejali et al. tested a modified design of the cochlear implant electrode that includes a coating of fibroblast cells transduced by a viral vector with a BDNF gene insert. To accomplish this type of ex vivo gene transfer, Rejali et al. transduced guinea pig fibroblasts with an adenovirus with a BDNF gene cassette insert, and determined that these cells secreted BDNF and then attached BDNF-secreting cells to the cochlear implant electrode via an agarose gel, and implanted the electrode in the scala tympani. Rejali et al. determined that the BDNF expressing electrodes were able to preserve significantly more spiral ganglion neurons in the basal turns of the cochlea after 48 days of implantation when compared to control electrodes and demonstrated the feasibility of combining cochlear implant therapy with ex vivo gene transfer for enhancing spiral ganglion neuron survival. Such a system may be applied to the nucleic acid-targeting system of the present invention for delivery to the ear.

[00594] Mukherjea et al. (*Antioxidants & Redox Signaling*, Volume 13, Number 5, 2010) document that knockdown of NOX3 using short interfering (si) RNA abrogated cisplatin ototoxicity, as evidenced by protection of OHCs from damage and reduced threshold shifts in auditory brainstem responses (ABRs). Different doses of siNOX3 (0.3, 0.6, and 0.9 μ g) were

administered to rats and NOX3 expression was evaluated by real time RT-PCR. The lowest dose of NOX3 siRNA used (0.3 μg) did not show any inhibition of NOX3 mRNA when compared to transtympanic administration of scrambled siRNA or untreated cochleae. However, administration of the higher doses of NOX3 siRNA (0.6 and 0.9 μg) reduced NOX3 expression compared to control scrambled siRNA. Such a system may be applied to the CRISPR Cas system of the present invention for transtympanic administration with a dosage of about 2 mg to about 4 mg of CRISPR Cas for administration to a human.

[00595] Jung et al. (Molecular Therapy, vol. 21 no. 4, 834–841 apr. 2013) demonstrate that Hes5 levels in the utricle decreased after the application of siRNA and that the number of hair cells in these utricles was significantly larger than following control treatment. The data suggest that siRNA technology may be useful for inducing repair and regeneration in the inner ear and that the Notch signaling pathway is a potentially useful target for specific gene expression inhibition. Jung et al. injected 8 μg of Hes5 siRNA in 2 μl volume, prepared by adding sterile normal saline to the lyophilized siRNA to a vestibular epithelium of the ear. Such a system may be applied to the nucleic acid-targeting system of the present invention for administration to the vestibular epithelium of the ear with a dosage of about 1 to about 30 mg of CRISPR Cas for administration to a human.

Treating Diseases of the Eye

[00596] The present invention also contemplates delivering the CRISPR-Cas system to one or both eyes.

[00597] In yet another aspect of the invention, the CRISPR-Cas system may be used to correct ocular defects that arise from several genetic mutations further described in Genetic Diseases of the Eye, Second Edition, edited by Elias I. Traboulsi, Oxford University Press, 2012.

[00598] For administration to the eye, lentiviral vectors, in particular equine infectious anemia viruses (EIAV) are particularly preferred.

[00599] In another embodiment, minimal non-primate lentiviral vectors based on the equine infectious anemia virus (EIAV) are also contemplated, especially for ocular gene therapy (see, e.g., Balagaan, J Gene Med 2006; 8: 275 – 285, Published online 21 November 2005 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/jgm.845). The vectors are contemplated to have cytomegalovirus (CMV) promoter driving expression of the target gene. Intracameral, subretinal, intraocular and intravitreal injections are all contemplated (see, e.g.,

Balagaan, J Gene Med 2006; 8: 275 – 285, Published online 21 November 2005 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/jgm.845). Intraocular injections may be performed with the aid of an operating microscope. For subretinal and intravitreal injections, eyes may be prolapsed by gentle digital pressure and fundi visualised using a contact lens system consisting of a drop of a coupling medium solution on the cornea covered with a glass microscope slide coverslip. For subretinal injections, the tip of a 10-mm 34-gauge needle, mounted on a 5- μ l Hamilton syringe may be advanced under direct visualisation through the superior equatorial sclera tangentially towards the posterior pole until the aperture of the needle was visible in the subretinal space. Then, 2 μ l of vector suspension may be injected to produce a superior bullous retinal detachment, thus confirming subretinal vector administration. This approach creates a self-sealing sclerotomy allowing the vector suspension to be retained in the subretinal space until it is absorbed by the RPE, usually within 48 h of the procedure. This procedure may be repeated in the inferior hemisphere to produce an inferior retinal detachment. This technique results in the exposure of approximately 70% of neurosensory retina and RPE to the vector suspension. For intravitreal injections, the needle tip may be advanced through the sclera 1 mm posterior to the corneoscleral limbus and 2 μ l of vector suspension injected into the vitreous cavity. For intracameral injections, the needle tip may be advanced through a corneoscleral limbal paracentesis, directed towards the central cornea, and 2 μ l of vector suspension may be injected. For intracameral injections, the needle tip may be advanced through a corneoscleral limbal paracentesis, directed towards the central cornea, and 2 μ l of vector suspension may be injected. These vectors may be injected at titres of either $1.0\text{--}1.4 \times 10^{10}$ or $1.0\text{--}1.4 \times 10^9$ transducing units (TU)/ml.

[00600] In another embodiment, RetinoStat®, an equine infectious anemia virus-based lentiviral gene therapy vector that expresses angiostatic proteins endostatin and angiostatin that is delivered via a subretinal injection for the treatment of the web form of age-related macular degeneration is also contemplated (see, e.g., Binley et al., HUMAN GENE THERAPY 23:980–991 (September 2012)). Such a vector may be modified for the CRISPR-Cas system of the present invention. Each eye may be treated with either RetinoStat® at a dose of 1.1×10^5 transducing units per eye (TU/eye) in a total volume of 100 μ l.

[00601] In another embodiment, an E1-, partial E3-, E4-deleted adenoviral vector may be contemplated for delivery to the eye. Twenty-eight patients with advanced neovascular

agerelated macular degeneration (AMD) were given a single intravitreal injection of an E1-, partial E3-, E4-deleted adenoviral vector expressing human pigment epithelium-derived factor (AdPEDF.II) (see, e.g., Campochiaro et al., *Human Gene Therapy* 17:167-176 (February 2006)). Doses ranging from 10^6 to $10^{9.5}$ particle units (PU) were investigated and there were no serious adverse events related to AdPEDF.II and no dose-limiting toxicities (see, e.g., Campochiaro et al., *Human Gene Therapy* 17:167-176 (February 2006)). Adenoviral vectormediated ocular gene transfer appears to be a viable approach for the treatment of ocular disorders and could be applied to the CRISPR Cas system.

[00602] In another embodiment, the sd-rxRNA® system of RXi Pharmaceuticals may be used/and or adapted for delivering CRISPR Cas to the eye. In this system, a single intravitreal administration of 3 µg of sd-rxRNA results in sequence-specific reduction of PPIB mRNA levels for 14 days. The the sd-rxRNA® system may be applied to the nucleic acid-targeting system of the present invention, contemplating a dose of about 3 to 20 mg of CRISPR administered to a human.

[00603] Millington-Ward et al. (*Molecular Therapy*, vol. 19 no. 4, 642–649 apr. 2011) describes adeno-associated virus (AAV) vectors to deliver an RNA interference (RNAi)-based rhodopsin suppressor and a codon-modified rhodopsin replacement gene resistant to suppression due to nucleotide alterations at degenerate positions over the RNAi target site. An injection of either 6.0×10^8 vp or 1.8×10^{10} vp AAV were subretinally injected into the eyes by Millington-Ward et al. The AAV vectors of Millington-Ward et al. may be applied to the CRISPR Cas system of the present invention, contemplating a dose of about 2×10^{11} to about 6×10^{13} vp administered to a human.

[00604] Dalkara et al. (*Sci Transl Med* 5, 189ra76 (2013)) also relates to in vivo directed evolution to fashion an AAV vector that delivers wild-type versions of defective genes throughout the retina after noninjurious injection into the eyes' vitreous humor. Dalkara describes a 7mer peptide display library and an AAV library constructed by DNA shuffling of cap genes from AAV1, 2, 4, 5, 6, 8, and 9. The rcAAV libraries and rAAV vectors expressing GFP under a CAG or Rho promoter were packaged and deoxyribonuclease-resistant genomic titers were obtained through quantitative PCR. The libraries were pooled, and two rounds of evolution were performed, each consisting of initial library diversification followed by three in vivo selection steps. In each such step, P30 rho-GFP mice were intravitreally injected

with 2 ml of iodixanol-purified, phosphate-buffered saline (PBS)-dialyzed library with a genomic titer of about 1×10^{12} vg/ml. The AAV vectors of Dalkara et al. may be applied to the nucleic acid-targeting system of the present invention, contemplating a dose of about 1×10^{15} to about 1×10^{16} vg/ml administered to a human.

[00605] In another embodiment, the rhodopsin gene may be targeted for the treatment of retinitis pigmentosa (RP), wherein the system of US Patent Publication No. 20120204282 assigned to Sangamo BioSciences, Inc. may be modified in accordance of the CRISPR Cas system of the present invention.

[00606] In another embodiment, the methods of US Patent Publication No. 20130183282 assigned to Collectis, which is directed to methods of cleaving a target sequence from the human rhodopsin gene, may also be modified to the nucleic acid-targeting system of the present invention.

[00607] US Patent Publication No. 20130202678 assigned to Academia Sinica relates to methods for treating retinopathies and sight-threatening ophthalmologic disorders relating to delivering of the Puf-A gene (which is expressed in retinal ganglion and pigmented cells of eye tissues and displays a unique anti-apoptotic activity) to the sub-retinal or intravitreal space in the eye. In particular, desirable targets are *zgc:193933*, *prdm1a*, *spata2*, *tex10*, *rbb4*, *ddx3*, *zp2.2*, *Blimp-1* and *Htra2*, all of which may be targeted by the nucleic acid-targeting system of the present invention.

[00608] Wu (Cell Stem Cell,13:659–62, 2013) designed a guide RNA that led Cas9 to a single base pair mutation that causes cataracts in mice, where it induced DNA cleavage. Then using either the other wild-type allele or oligos given to the zygotes repair mechanisms corrected the sequence of the broken allele and corrected the cataract-causing genetic defect in mutant mouse.

[00609] US Patent Publication No. 20120159653, describes use of zinc finger nucleases to genetically modify cells, animals and proteins associated with macular degeneration (MD). Macular degeneration (MD) is the primary cause of visual impairment in the elderly, but is also a hallmark symptom of childhood diseases such as Stargardt disease, Sorsby fundus, and fatal childhood neurodegenerative diseases, with an age of onset as young as infancy. Macular degeneration results in a loss of vision in the center of the visual field (the macula) because of damage to the retina. Currently existing animal models do not recapitulate major hallmarks of the disease as it is observed in humans. The available animal models comprising mutant genes

encoding proteins associated with MD also produce highly variable phenotypes, making translations to human disease and therapy development problematic.

[00610] One aspect of US Patent Publication No. 20120159653 relates to editing of any chromosomal sequences that encode proteins associated with MD which may be applied to the nucleic acid-targeting system of the present invention. The proteins associated with MD are typically selected based on an experimental association of the protein associated with MD to an MD disorder. For example, the production rate or circulating concentration of a protein associated with MD may be elevated or depressed in a population having an MD disorder relative to a population lacking the MD disorder. Differences in protein levels may be assessed using proteomic techniques including but not limited to Western blot, immunohistochemical staining, enzyme linked immunosorbent assay (ELISA), and mass spectrometry. Alternatively, the proteins associated with MD may be identified by obtaining gene expression profiles of the genes encoding the proteins using genomic techniques including but not limited to DNA microarray analysis, serial analysis of gene expression (SAGE), and quantitative real-time polymerase chain reaction (Q-PCR).

[00611] By way of non-limiting example, proteins associated with MD include but are not limited to the following proteins: (ABCA4) ATP-binding cassette, sub-family A (ABC1), member 4 ACHM1 achromatopsia (rod monochromacy) 1 ApoE Apolipoprotein E (ApoE) C1QTNF5 (CTRP5) C1q and tumor necrosis factor related protein 5 (C1QTNF5) C2 Complement component 2 (C2) C3 Complement components (C3) CCL2 Chemokine (C-C motif) Ligand 2 (CCL2) CCR2 Chemokine (C-C motif) receptor 2 (CCR2) CD36 Cluster of Differentiation 36 CFB Complement factor B CFH Complement factor CFH H CFHR1 complement factor H-related 1 CFHR3 complement factor H-related 3 CNGB3 cyclic nucleotide gated channel beta 3 CP ceruloplasmin (CP) CRP C reactive protein (CRP) CST3 cystatin C or cystatin 3 (CST3) CTSD Cathepsin D (CTSD) CX3CR1 chemokine (C-X3-C motif) receptor 1 ELOVL4 Elongation of very long chain fatty acids 4 ERCC6 excision repair crosscomplementing rodent repair deficiency, complementation group 6 FBLN5 Fibulin-5 FBLN5 Fibulin 5 FBLN6 Fibulin 6 FSCN2 fascin (FSCN2) HMCN1 Hemicentrin 1 HMCN1 hemicentrin 1 HTRA1 HtrA serine peptidase 1 (HTRA1) HTRA1 HtrA serine peptidase 1 IL-6 Interleukin 6 IL-8 Interleukin 8 LOC387715 Hypothetical protein PLEKHA1 Pleckstrin homology domain containing family A member 1 (PLEKHA1) PROM1 Prominin 1 (PROM1) or

CD133) PRPH2 Peripherin-2 RPGR retinitis pigmentosa GTPase regulator SERPING1 serpin peptidase inhibitor, clade G, member 1 (C1- inhibitor) TCOF1 Treacle TIMP3 Metalloproteinase inhibitor 3 (TIMP3) TLR3 Toll-like receptor 3.

[00612] The identity of the protein associated with MD whose chromosomal sequence is edited can and will vary. In preferred embodiments, the proteins associated with MD whose chromosomal sequence is edited may be the ATP-binding cassette, sub-family A (ABC1) member 4 protein (ABCA4) encoded by the ABCR gene, the apolipoprotein E protein (APOE) encoded by the APOE gene, the chemokine (C-C motif) Ligand 2 protein (CCL2) encoded by the CCL2 gene, the chemokine (C-C motif) receptor 2 protein (CCR2) encoded by the CCR2 gene, the ceruloplasmin protein (CP) encoded by the CP gene, the cathepsin D protein (CTSD) encoded by the CTSD gene, or the metalloproteinase inhibitor 3 protein (TIMP3) encoded by the TIMP3 gene. In an exemplary embodiment, the genetically modified animal is a rat, and the edited chromosomal sequence encoding the protein associated with MD may be: (ABCA4) ATPbinding cassette, NM_000350 sub-family A (ABC1), member 4 APOE Apolipoprotein E NM_138828 (APOE) CCL2 Chemokine (C-C NM_031530 motif) Ligand 2 (CCL2) CCR2 Chemokine (C-C NM_021866 motif) receptor 2 (CCR2) CP ceruloplasmin (CP) NM_012532 CTSD Cathepsin D (CTSD) NM_134334 TIMP3 Metalloproteinase NM_012886 inhibitor 3 (TIMP3) The animal or cell may comprise 1, 2, 3, 4, 5, 6, 7 or more disrupted chromosomal sequences encoding a protein associated with MD and zero, 1, 2, 3, 4, 5, 6, 7 or more chromosomally integrated sequences encoding the disrupted protein associated with MD.

[00613] The edited or integrated chromosomal sequence may be modified to encode an altered protein associated with MD. Several mutations in MD-related chromosomal sequences have been associated with MD. Non-limiting examples of mutations in chromosomal sequences associated with MD include those that may cause MD including in the ABCR protein, E471K (i.e. glutamate at position 471 is changed to lysine), R1129L (i.e. arginine at position 1129 is changed to leucine), T1428M (i.e. threonine at position 1428 is changed to methionine), R1517S (i.e. arginine at position 1517 is changed to serine), I1562T (i.e. isoleucine at position 1562 is changed to threonine), and G1578R (i.e. glycine at position 1578 is changed to arginine); in the CCR2 protein, V64I (i.e. valine at position 192 is changed to isoleucine); in CP protein, G969B (i.e. glycine at position 969 is changed to asparagine or aspartate); in TIMP3 protein, S156C (i.e. serine at position 156 is changed to cysteine), G166C (i.e. glycine at position 166 is changed to

cysteine), G167C (i.e. glycine at position 167 is changed to cysteine), Y168C (i.e. tyrosine at position 168 is changed to cysteine), S170C (i.e. serine at position 170 is changed to cysteine), Y172C (i.e. tyrosine at position 172 is changed to cysteine) and S181C (i.e. serine at position 181 is changed to cysteine). Other associations of genetic variants in MD-associated genes and disease are known in the art.

Treating Circulatory and Muscular Diseases

[00614] The present invention also contemplates delivering the CRISPR-Cas system described herein, e.g. Cas9 effector protein systems, to the heart. For the heart, a myocardium tropic adena-associated virus (AAVM) is preferred, in particular AAVM41 which showed preferential gene transfer in the heart (see, e.g., Lin-Yanga et al., PNAS, March 10, 2009, vol. 106, no. 10). Administration may be systemic or local. A dosage of about $1-10 \times 10^{14}$ vector genomes are contemplated for systemic administration. See also, e.g., Eulalio et al. (2012) Nature 492: 376 and Somasuntharam et al. (2013) Biomaterials 34: 7790.

[00615] For example, US Patent Publication No. 20110023139, describes use of zinc finger nucleases to genetically modify cells, animals and proteins associated with cardiovascular disease. Cardiovascular diseases generally include high blood pressure, heart attacks, heart failure, and stroke and TIA. Any chromosomal sequence involved in cardiovascular disease or the protein encoded by any chromosomal sequence involved in cardiovascular disease may be utilized in the methods described in this disclosure. The cardiovascular-related proteins are typically selected based on an experimental association of the cardiovascular-related protein to the development of cardiovascular disease. For example, the production rate or circulating concentration of a cardiovascular-related protein may be elevated or depressed in a population having a cardiovascular disorder relative to a population lacking the cardiovascular disorder. Differences in protein levels may be assessed using proteomic techniques including but not limited to Western blot, immunohistochemical staining, enzyme linked immunosorbent assay (ELISA), and mass spectrometry. Alternatively, the cardiovascular-related proteins may be identified by obtaining gene expression profiles of the genes encoding the proteins using genomic techniques including but not limited to DNA microarray analysis, serial analysis of gene expression (SAGE), and quantitative real-time polymerase chain reaction (Q-PCR).

Treating Diseases of the Liver and Kidney

[00616] The present invention also contemplates delivering the CRISPR-Cas system described herein, e.g. Cas9 effector protein systems, to the liver and/or kidney. Delivery strategies to induce cellular uptake of the therapeutic nucleic acid include physical force or vector systems such as viral-, lipid- or complex- based delivery, or nanocarriers. From the initial applications with less possible clinical relevance, when nucleic acids were addressed to renal cells with hydrodynamic high pressure injection systemically, a wide range of gene therapeutic viral and non-viral carriers have been applied already to target posttranscriptional events in different animal kidney disease models in vivo (Csaba Révész and Péter Hamar (2011). Delivery Methods to Target RNAs in the Kidney, Gene Therapy Applications, Prof. Chunsheng Kang (Ed.), ISBN: 978-953-307-541-9, InTech, Available from: <http://www.intechopen.com/books/gene-therapy-applications/delivery-methods-to-target-rnas-inthe-kidney>). Delivery methods to the kidney may include those in Yuan et al. (Am J Physiol Renal Physiol 295: F605–F617, 2008) investigated whether in vivo delivery of small interfering RNAs (siRNAs) targeting the 12/15-lipoxygenase (12/15-LO) pathway of arachidonate acid metabolism can ameliorate renal injury and diabetic nephropathy (DN) in a streptozotocin-injected mouse model of type 1 diabetes. To achieve greater in vivo access and siRNA expression in the kidney, Yuan et al. used double-stranded 12/15-LO siRNA oligonucleotides conjugated with cholesterol. About 400 µg of siRNA was injected subcutaneously into mice. The method of Yuan et al. may be applied to the CRISPR-Cas system of the present invention contemplating a 1-2 g subcutaneous injection of CRISPR Cas conjugated with cholesterol to a human for delivery to the kidneys.

[00617] Molitoris et al. (J Am Soc Nephrol 20: 1754–1764, 2009) exploited proximal tubule cells (PTCs), as the site of oligonucleotide reabsorption within the kidney to test the efficacy of siRNA targeted to p53, a pivotal protein in the apoptotic pathway, to prevent kidney injury. Naked synthetic siRNA to p53 injected intravenously 4 h after ischemic injury maximally protected both PTCs and kidney function. Molitoris et al.'s data indicates that rapid delivery of siRNA to proximal tubule cells follows intravenous administration. For dose-response analysis, rats were injected with doses of siP53, 0.33; 1, 3, or 5mg/kg, given at the same four time points, resulting in cumulative doses of 1.32; 4, 12, and 20 mg/kg, respectively. All siRNA doses tested produced a SCr reducing effect on day one with higher doses being effective over approximately

five days compared with PBS-treated ischemic control rats. The 12 and 20 mg/kg cumulative doses provided the best protective effect. The method of Molitoris et al. may be applied to the nucleic acid-targeting system of the present invention contemplating 12 and 20 mg/kg cumulative doses to a human for delivery to the kidneys.

[00618] Thompson et al. (Nucleic Acid Therapeutics, Volume 22, Number 4, 2012) reports the toxicological and pharmacokinetic properties of the synthetic, small interfering RNA I5NP following intravenous administration in rodents and nonhuman primates. I5NP is designed to act via the RNA interference (RNAi) pathway to temporarily inhibit expression of the pro-apoptotic protein p53 and is being developed to protect cells from acute ischemia/reperfusion injuries such as acute kidney injury that can occur during major cardiac surgery and delayed graft function that can occur following renal transplantation. Doses of 800mg/kg I5NP in rodents, and 1,000 mg/kg I5NP in nonhuman primates, were required to elicit adverse effects, which in the monkey were isolated to direct effects on the blood that included a sub-clinical activation of complement and slightly increased clotting times. In the rat, no additional adverse effects were observed with a rat analogue of I5NP, indicating that the effects likely represent class effects of synthetic RNA duplexes rather than toxicity related to the intended pharmacologic activity of I5NP. Taken together, these data support clinical testing of intravenous administration of I5NP for the preservation of renal function following acute ischemia/reperfusion injury. The no observed adverse effect level (NOAEL) in the monkey was 500 mg/kg. No effects on cardiovascular, respiratory, and neurologic parameters were observed in monkeys following i.v. administration at dose levels up to 25 mg/kg. Therefore, a similar dosage may be contemplated for intravenous administration of CRISPR Cas to the kidneys of a human.

[00619] Shimizu et al. (J Am Soc Nephrol 21: 622–633, 2010) developed a system to target delivery of siRNAs to glomeruli via poly(ethylene glycol)-poly(L-lysine)-based vehicles. The siRNA/nanocarrier complex was approximately 10 to 20 nm in diameter, a size that would allow it to move across the fenestrated endothelium to access to the mesangium. After intraperitoneal injection of fluorescence-labeled siRNA/nanocarrier complexes, Shimizu et al. detected siRNAs in the blood circulation for a prolonged time. Repeated intraperitoneal administration of a mitogen-activated protein kinase 1 (MAPK1) siRNA/nanocarrier complex suppressed glomerular MAPK1 mRNA and protein expression in a mouse model of glomerulonephritis. For the investigation of siRNA accumulation, Cy5-labeled siRNAs complexed with PIC nanocarriers

(0.5 ml, 5 nmol of siRNA content), naked Cy5-labeled siRNAs (0.5 ml, 5 nmol), or Cy5-labeled siRNAs encapsulated in HVJ-E (0.5 ml, 5 nmol of siRNA content) were administered to BALBc mice. The method of Shimizu et al. may be applied to the nucleic acid-targeting system of the present invention contemplating a dose of about of 10-20 μ mol CRISPR Cas complexed with nanocarriers in about 1-2 liters to a human for intraperitoneal administration and delivery to the kidneys.

Treating Epithelial and Lung Diseases

[00620] The present invention also contemplates delivering the CRISPR-Cas system described herein, e.g. Cas9 systems, to one or both lungs.

[00621] Although AAV-2-based vectors were originally proposed for CFTR delivery to CF airways, other serotypes such as AAV-1, AAV-5, AAV-6, and AAV-9 exhibit improved gene transfer efficiency in a variety of models of the lung epithelium (see, e.g., Li et al., *Molecular Therapy*, vol. 17 no. 12, 2067-2077 Dec 2009). AAV-1 was demonstrated to be ~100-fold more efficient than AAV-2 and AAV-5 at transducing human airway epithelial cells in vitro,⁵ although AAV-1 transduced murine tracheal airway epithelia in vivo with an efficiency equal to that of AAV-5. Other studies have shown that AAV-5 is 50-fold more efficient than AAV-2 at gene delivery to human airway epithelium (HAE) in vitro and significantly more efficient in the mouse lung airway epithelium in vivo. AAV-6 has also been shown to be more efficient than AAV-2 in human airway epithelial cells in vitro and murine airways in vivo.⁸ The more recent isolate, AAV-9, was shown to display greater gene transfer efficiency than AAV-5 in murine nasal and alveolar epithelia in vivo with gene expression detected for over 9 months suggesting AAV may enable long-term gene expression in vivo, a desirable property for a CFTR gene delivery vector. Furthermore, it was demonstrated that AAV-9 could be readministered to the murine lung with no loss of CFTR expression and minimal immune consequences. CF and non-CF HAE cultures may be inoculated on the apical surface with 100 μ l of AAV vectors for hours (see, e.g., Li et al., *Molecular Therapy*, vol. 17 no. 12, 2067-2077 Dec 2009). The MOI may vary from 1×10^3 to 4×10^5 vector genomes/cell, depending on virus concentration and purposes of the experiments. The above cited vectors are contemplated for the delivery and/or administration of the invention.

[00622] Zamora et al. (*Am J Respir Crit Care Med* Vol 183. pp 531–538, 2011) reported an example of the application of an RNA interference therapeutic to the treatment of human

infectious disease and also a randomized trial of an antiviral drug in respiratory syncytial virus (RSV)-infected lung transplant recipients. Zamora et al. performed a randomized, double-blind, placebocontrolled trial in LTX recipients with RSV respiratory tract infection. Patients were permitted to receive standard of care for RSV. Aerosolized ALN-RSV01 (0.6 mg/kg) or placebo was administered daily for 3 days. This study demonstrates that an RNAi therapeutic targeting RSV can be safely administered to LTX recipients with RSV infection. Three daily doses of ALN-RSV01 did not result in any exacerbation of respiratory tract symptoms or impairment of lung function and did not exhibit any systemic proinflammatory effects, such as induction of cytokines or CRP. Pharmacokinetics showed only low, transient systemic exposure after inhalation, consistent with preclinical animal data showing that ALN-RSV01, administered intravenously or by inhalation, is rapidly cleared from the circulation through exonucleasemediated digestion and renal excretion. The method of Zamora et al. may be applied to the nucleic acid-targeting system of the present invention and an aerosolized CRISPR Cas, for example with a dosage of 0.6 mg/kg, may be contemplated for the present invention.

[00623] Schwank et al. (*Cell Stem Cell*, 13:653–58, 2013) used CRISPR-Cas9 to correct a defect associated with cystic fibrosis in human stem cells. The team's target was the gene for an ion channel, cystic fibrosis transmembrane conductor receptor (CFTR). A deletion in CFTR causes the protein to misfold in cystic fibrosis patients. Using cultured intestinal stem cells developed from cell samples from two children with cystic fibrosis, Schwank et al. were able to correct the defect using CRISPR along with a donor plasmid containing the reparative sequence to be inserted. The researchers then grew the cells into intestinal "organoids," or miniature guts, and showed that they functioned normally. In this case, about half of clonal organoids underwent the proper genetic correction.

Treating Diseases of the Muscular System

[00624] The present invention also contemplates delivering the CRISPR-Cas system described herein, e.g. Cas9 systems, to muscle(s).

[00625] Bortolanza et al. (*Molecular Therapy* vol. 19 no. 11, 2055–2064 Nov. 2011) shows that systemic delivery of RNA interference expression cassettes in the FRG1 mouse, after the onset of facioscapulohumeral muscular dystrophy (FSHD), led to a dose-dependent long-term FRG1 knockdown without signs of toxicity. Bortolanza et al. found that a single intravenous injection of 5×10^{12} vg of rAAV6-sh1FRG1 rescues muscle histopathology and muscle function

of FRG1 mice. In detail, 200 μ l containing 2×10^{12} or 5×10^{12} vg of vector in physiological solution were injected into the tail vein using a 25-gauge Terumo syringe. The method of Bortolanza et al. may be applied to an AAV expressing CRISPR Cas and injected into humans at a dosage of about 2×10^{15} or 2×10^{16} vg of vector.

[00626] Dumonceaux et al. (Molecular Therapy vol. 18 no. 5, 881–887 May 2010) inhibit the myostatin pathway using the technique of RNA interference directed against the myostatin receptor AcvrIib mRNA (sh-AcvrIib). The restoration of a quasi-dystrophin was mediated by the vectorized U7 exon-skipping technique (U7-DYS). Adeno-associated vectors carrying either the sh-AcvrIib construct alone, the U7-DYS construct alone, or a combination of both constructs were injected in the tibialis anterior (TA) muscle of dystrophic mdx mice. The injections were performed with 10^{11} AAV viral genomes. The method of Dumonceaux et al. may be applied to an AAV expressing CRISPR Cas and injected into humans, for example, at a dosage of about 10^{14} to about 10^{15} vg of vector.

[00627] Kinouchi et al. (Gene Therapy (2008) 15, 1126–1130) report the effectiveness of in vivo siRNA delivery into skeletal muscles of normal or diseased mice through nanoparticle formation of chemically unmodified siRNAs with atelocollagen (ATCOL). ATCOL-mediated local application of siRNA targeting myostatin, a negative regulator of skeletal muscle growth, in mouse skeletal muscles or intravenously, caused a marked increase in the muscle mass within a few weeks after application. These results imply that ATCOL-mediated application of siRNAs is a powerful tool for future therapeutic use for diseases including muscular atrophy. MstsiRNAs (final concentration, 10 mM) were mixed with ATCOL (final concentration for local administration, 0.5%) (AteloGene, Kohken, Tokyo, Japan) according to the manufacturer's instructions. After anesthesia of mice (20-week-old male C57BL/6) by Nembutal (25 mg/kg, i.p.), the Mst-siRNA/ATCOL complex was injected into the masseter and biceps femoris muscles. The method of Kinouchi et al. may be applied to CRISPR Cas and injected into a human, for example, at a dosage of about 500 to 1000 ml of a 40 μ M solution into the muscle. Hagstrom et al. (Molecular Therapy Vol. 10, No. 2, August 2004) describe an intravascular, nonviral methodology that enables efficient and repeatable delivery of nucleic acids to muscle cells (myofibers) throughout the limb muscles of mammals. The procedure involves the injection of naked plasmid DNA or siRNA into a distal vein of a limb that is transiently isolated by a tourniquet or blood pressure cuff. Nucleic acid delivery to myofibers is facilitated by its rapid

injection in sufficient volume to enable extravasation of the nucleic acid solution into muscle tissue. High levels of transgene expression in skeletal muscle were achieved in both small and large animals with minimal toxicity. Evidence of siRNA delivery to limb muscle was also obtained. For plasmid DNA intravenous injection into a rhesus monkey, a threeway stopcock was connected to two syringe pumps (Model PHD 2000; Harvard Instruments), each loaded with a single syringe. Five minutes after a papaverine injection, pDNA (15.5 to 25.7 mg in 40 –100 ml saline) was injected at a rate of 1.7 or 2.0 ml/s. This could be scaled up for plasmid DNA expressing CRISPR Cas of the present invention with an injection of about 300 to 500 mg in 800 to 2000 ml saline for a human. For adenoviral vector injections into a rat, 2×10^9 infectious particles were injected in 3 ml of normal saline solution (NSS). This could be scaled up for an adenoviral vector expressing CRISPR Cas of the present invention with an injection of about 1×10^{13} infectious particles were injected in 10 liters of NSS for a human. For siRNA, a rat was injected into the great saphenous vein with 12.5 μg of a siRNA and a primate was injected into the great saphenous vein with 750 μg of a siRNA. This could be scaled up for a CRISPR Cas of the present invention, for example, with an injection of about 15 to about 50 mg into the great saphenous vein of a human.

[00628] See also, for example, WO2013163628 A2, Genetic Correction of Mutated Genes, published application of Duke University describes efforts to correct, for example, a frameshift mutation which causes a premature stop codon and a truncated gene product that can be corrected via nuclease mediated non-homologous end joining such as those responsible for Duchenne Muscular Dystrophy, ("DMD") a recessive, fatal, X-linked disorder that results in muscle degeneration due to mutations in the dystrophin gene. The majority of dystrophin mutations that cause DMD are deletions of exons that disrupt the reading frame and cause premature translation termination in the dystrophin gene. Dystrophin is a cytoplasmic protein that provides structural stability to the dystroglycan complex of the cell membrane that is responsible for regulating muscle cell integrity and function. The dystrophin gene or "DMD gene" as used interchangeably herein is 2.2 megabases at locus Xp21. The primary transcription measures about 2,400 kb with the mature mRNA being about 14 kb. 79 exons code for the protein which is over 3500 amino acids. Exon 51 is frequently adjacent to frame-disrupting deletions in DMD patients and has been targeted in clinical trials for oligonucleotide-based exon skipping. A clinical trial for the exon 51 skipping compound eteplirsen recently reported a

significant functional benefit across 48 weeks, with an average of 47% dystrophin positive fibers compared to baseline. Mutations in exon 51 are ideally suited for permanent correction by NHEJ-based genome editing.

[00629] The methods of US Patent Publication No. 20130145487 assigned to Collectis, which relates to meganuclease variants to cleave a target sequence from the human dystrophin gene (DMD), may also be modified to for the nucleic acid-targeting system of the present invention.

Treating Diseases of the Skin

[00630] The present invention also contemplates delivering the CRISPR-Cas system described herein, e.g. Cas9 effector protein systems, to the skin.

[00631] Hickerson et al. (Molecular Therapy—Nucleic Acids (2013) 2, e129) relates to a motorized microneedle array skin delivery device for delivering self-delivery (sd)-siRNA to human and murine skin. The primary challenge to translating siRNA-based skin therapeutics to the clinic is the development of effective delivery systems. Substantial effort has been invested in a variety of skin delivery technologies with limited success. In a clinical study in which skin was treated with siRNA, the exquisite pain associated with the hypodermic needle injection precluded enrollment of additional patients in the trial, highlighting the need for improved, more “patient-friendly” (i.e., little or no pain) delivery approaches. Microneedles represent an efficient way to deliver large charged cargos including siRNAs across the primary barrier, the stratum corneum, and are generally regarded as less painful than conventional hypodermic needles. Motorized “stamp type” microneedle devices, including the motorized microneedle array (MMNA) device used by Hickerson et al., have been shown to be safe in hairless mice studies and cause little or no pain as evidenced by (i) widespread use in the cosmetic industry and (ii) limited testing in which nearly all volunteers found use of the device to be much less painful than a flushot, suggesting siRNA delivery using this device will result in much less pain than was experienced in the previous clinical trial using hypodermic needle injections. The MMNA device (marketed as Triple-M or Tri-M by Bomtech Electronic Co, Seoul, South Korea) was adapted for delivery of siRNA to mouse and human skin. sd-siRNA solution (up to 300 μ l of 0.1 mg/ml RNA) was introduced into the chamber of the disposable Tri-M needle cartridge (Bomtech), which was set to a depth of 0.1 mm. For treating human skin, deidentified skin (obtained immediately following surgical procedures) was manually stretched and pinned to a cork platform before treatment. All intradermal injections were performed using an insulin syringe

with a 28-gauge 0.5-inch needle. The MMNA device and method of Hickerson et al. could be used and/or adapted to deliver the CRISPR Cas of the present invention, for example, at a dosage of up to 300 μ l of 0.1 mg/ml CRISPR Cas to the skin.

[00632] Leachman et al. (Molecular Therapy, vol. 18 no. 2, 442–446 Feb. 2010) relates to a phase Ib clinical trial for treatment of a rare skin disorder pachyonychia congenita (PC), an autosomal dominant syndrome that includes a disabling plantar keratoderma, utilizing the first short-interfering RNA (siRNA)-based therapeutic for skin. This siRNA, called TD101, specifically and potently targets the keratin 6a (K6a) N171K mutant mRNA without affecting wild-type K6a mRNA.

[00633] Zheng et al. (PNAS, July 24, 2012, vol. 109, no. 30, 11975–11980) show that spherical nucleic acid nanoparticle conjugates (SNA-NCs), gold cores surrounded by a dense shell of highly oriented, covalently immobilized siRNA, freely penetrate almost 100% of keratinocytes in vitro, mouse skin, and human epidermis within hours after application. Zheng et al. demonstrated that a single application of 25 nM epidermal growth factor receptor (EGFR) SNA-NCs for 60 h demonstrate effective gene knockdown in human skin. A similar dosage may be contemplated for CRISPR Cas immobilized in SNA-NCs for administration to the skin.

General Gene Therapy Considerations

[00634] Examples of disease-associated genes and polynucleotides and disease specific information is available from McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, Md.) and National Center for Biotechnology Information, National Library of Medicine (Bethesda, Md.), available on the World Wide Web.

[00635] Mutations in these genes and pathways can result in production of improper proteins or proteins in improper amounts which affect function. Further examples of genes, diseases and proteins are hereby incorporated by reference from US Provisional application 61/736,527 filed December 12, 2012. Such genes, proteins and pathways may be the target polynucleotide of a CRISPR complex of the present invention.

[00636] Embodiments of the invention also relate to methods and compositions related to knocking out genes, amplifying genes and repairing particular mutations associated with DNA repeat instability and neurological disorders (Robert D. Wells, Tetsuo Ashizawa, Genetic Instabilities and Neurological Diseases, Second Edition, Academic Press, Oct 13, 2011 – Medical). Specific aspects of tandem repeat sequences have been found to be responsible for

more than twenty human diseases (New insights into repeat instability: role of RNA•DNA hybrids. McIvor EI, Polak U, Napierala M. RNA Biol. 2010 Sep-Oct;7(5):551-8). The present effector protein systems may be harnessed to correct these defects of genomic instability.

[00637] Several further aspects of the invention relate to correcting defects associated with a wide range of genetic diseases which are further described on the website of the National Institutes of Health under the topic subsection Genetic Disorders (website at health.nih.gov/topic/GeneticDisorders). The genetic brain diseases may include but are not limited to Adrenoleukodystrophy, Agenesis of the Corpus Callosum, Aicardi Syndrome, Alpers' Disease, Alzheimer's Disease, Barth Syndrome, Batten Disease, CADASIL, Cerebellar Degeneration, Fabry's Disease, Gerstmann-Straussler-Scheinker Disease, Huntington's Disease and other Triplet Repeat Disorders, Leigh's Disease, Lesch-Nyhan Syndrome, Menkes Disease, Mitochondrial Myopathies and NINDS Colpocephaly. These diseases are further described on the website of the National Institutes of Health under the subsection Genetic Brain Disorders.

Selected Other Conditions

[00638] Cancer

[00639] Target genes suitable for the treatment or prophylaxis of cancer may include, in some embodiments, those described in WO2015048577 the disclosure of which is hereby incorporated by reference.

[00640] Usher Syndrome or retinitis pigmentosa-39

[00641] In some embodiments, the treatment, prophylaxis or diagnosis of Usher Syndrome or retinitis pigmentosa-39 is provided. The target is preferably the USH2A gene. In some embodiments, correction of a G deletion at position 2299 (2299delG) is provided. This is described in WO2015134812A1, the disclosure of which is hereby incorporated by reference.

[00642] Leber's Congenital Amaurosis 10

[00643] In some embodiments, the treatment, prophylaxis or diagnosis of Leber's Congenital Amaurosis 10 (LCA10). The target is preferably the CEP290 gene. This is described in WO2015138510A1, the disclosure of which is hereby incorporated by reference.

[00644] HIV and AIDS

[00645] In some embodiments, the treatment, prophylaxis or diagnosis of HIV and AIDS is provided. The target is preferably the CCR5 gene in HIV. This is described in WO2015148670A1, the disclosure of which is hereby incorporated by reference.

[00646] Beta Thalassaemia

[00647] In some embodiments, the treatment, prophylaxis or diagnosis of Beta Thalassaemia is provided. The target is preferably the BCL11A gene. This is described in WO2015148860, the disclosure of which is hereby incorporated by reference.

[00648] Sickle Cell Disease (SCD)

[00649] In some embodiments, the treatment, prophylaxis or diagnosis of Sickle Cell Disease (SCD) is provided. The target is preferably the HBB or BCL11A gene. This is described in WO2015148863, the disclosure of which is hereby incorporated by reference.

[00650] Herpes Simplex Virus 1 and 2

[00651] In some embodiments, the treatment, prophylaxis or diagnosis of HSV-1 (Herpes Simplex Virus 1) is provided. The target is preferably the UL19, UL30, UL48 or UL50 gene in HSV-1. This is described in WO2015153789, the disclosure of which is hereby incorporated by reference.

[00652] In other embodiments, the treatment, prophylaxis or diagnosis of HSV-2 (Herpes Simplex Virus 2) is provided. The target is preferably the UL19, UL30, UL48 or UL50 gene in HSV-2. This is described in WO2015153791, the disclosure of which is hereby incorporated by reference.

[00653] In some embodiments, the treatment, prophylaxis or diagnosis of Primary Open Angle Glaucoma (POAG) is provided. The target is preferably the MYOC gene. This is described in WO2015153780, the disclosure of which is hereby incorporated by reference.

[00654] The present invention may be further illustrated and extended based on aspect of CISPR-Cas9 development and use as set forth in the following articles hereby incorporated

herein by reference and particularly as relates to delivery of a CRISPR protein complex and uses of an RNA guided endonuclease in cells and organisms:

- *Multiplex genome engineering using CRISPR/Cas systems.* Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W., Marraffini, L.A., & Zhang, F. *Science* Feb 15;339(6121):819-23 (2013);
- *RNA-guided editing of bacterial genomes using CRISPR-Cas systems.* Jiang W., Bikard D., Cox D., Zhang F, Marraffini LA. *Nat Biotechnol* Mar;31(3):233-9 (2013);
- *One-Step Generation of Mice Carrying Mutations in Multiple Genes by CRISPR/Cas-Mediated Genome Engineering.* Wang H., Yang H., Shivalila CS., Dawlaty MM., Cheng AW., Zhang F., Jaenisch R. *Cell* May 9;153(4):910-8 (2013);
- *Optical control of mammalian endogenous transcription and epigenetic states.* Konermann S, Brigham MD, Trevino AE, Hsu PD, Heidenreich M, Cong L, Platt RJ, Scott DA, Church GM, Zhang F. *Nature*. 2013 Aug 22;500(7463):472-6. doi: 10.1038/Nature12466. Epub 2013 Aug 23;
- *Double Nicking by RNA-Guided CRISPR Cas9 for Enhanced Genome Editing Specificity.* Ran, FA., Hsu, PD., Lin, CY., Gootenberg, JS., Konermann, S., Trevino, AE., Scott, DA., Inoue, A., Matoba, S., Zhang, Y., & Zhang, F. *Cell* Aug 28. pii: S0092-8674(13)01015-5. (2013);
- *DNA targeting specificity of RNA-guided Cas9 nucleases.* Hsu, P., Scott, D., Weinstein, J., Ran, FA., Konermann, S., Agarwala, V., Li, Y., Fine, E., Wu, X., Shalem, O., Cradick, TJ., Marraffini, LA., Bao, G., & Zhang, F. *Nat Biotechnol* doi:10.1038/nbt.2647 (2013);
- *Genome engineering using the CRISPR-Cas9 system.* Ran, FA., Hsu, PD., Wright, J., Agarwala, V., Scott, DA., Zhang, F. *Nature Protocols* Nov;8(11):2281-308. (2013);
- *Genome-Scale CRISPR-Cas9 Knockout Screening in Human Cells.* Shalem, O., Sanjana, NE., Hartenian, E., Shi, X., Scott, DA., Mikkelsen, T., Heckl, D., Ebert, BL., Root, DE., Doench, JG., Zhang, F. *Science* Dec 12. (2013). [Epub ahead of print];
- *Crystal structure of cas9 in complex with guide RNA and target DNA.* Nishimasu, H., Ran, FA., Hsu, PD., Konermann, S., Shehata, SI, Dohmae, N., Ishitani, R., Zhang, F., Nureki, O. *Cell* Feb 27. (2014). 156(5):935-49;
- *Genome-wide binding of the CRISPR endonuclease Cas9 in mammalian cells.* Wu X., Scott DA., Kriz AJ., Chiu AC., Hsu PD., Dadon DB., Cheng AW., Trevino AE., Konermann S., Chen S., Jaenisch R., Zhang F., Sharp PA. *Nat Biotechnol*. (2014) Apr 20. doi: 10.1038/nbt.2889,

- *CRISPR-Cas9 Knockin Mice for Genome Editing and Cancer Modeling*, Platt et al., Cell 159(2): 440-455 (2014) DOI: 10.1016/j.cell.2014.09.014,
- *Development and Applications of CRISPR-Cas9 for Genome Engineering*, Hsu et al, Cell 157, 1262-1278 (June 5, 2014) (Hsu 2014),
- *Genetic screens in human cells using the CRISPR/Cas9 system*, Wang et al., Science. 2014 January 3; 343(6166): 80–84. doi:10.1126/science.1246981,
- *Rational design of highly active sgRNAs for CRISPR-Cas9-mediated gene inactivation*, Doench et al., Nature Biotechnology published online 3 September 2014; doi:10.1038/nbt.3026, and
- *In vivo interrogation of gene function in the mammalian brain using CRISPR-Cas9*, Swiech et al, Nature Biotechnology ; published online 19 October 2014; doi:10.1038/nbt.3055.
- *Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex*, Konermann S, Brigham MD, Trevino AE, Joung J, Abudayyeh OO, Barcena C, Hsu PD, Habib N, Gootenberg JS, Nishimasu H, Nureki O, Zhang F., Nature. Jan 29;517(7536):583-8 (2015).
- *A split-Cas9 architecture for inducible genome editing and transcription modulation*, Zetsche B, Volz SE, Zhang F., (published online 02 February 2015) Nat Biotechnol. Feb;33(2):139-42 (2015);
- *Genome-wide CRISPR Screen in a Mouse Model of Tumor Growth and Metastasis*, Chen S, Sanjana NE, Zheng K, Shalem O, Lee K, Shi X, Scott DA, Song J, Pan JQ, Weissleder R, Lee H, Zhang F, Sharp PA. Cell 160, 1246–1260, March 12, 2015 (multiplex screen in mouse), and
- *In vivo genome editing using Staphylococcus aureus Cas9*, Ran FA, Cong L, Yan WX, Scott DA, Gootenberg JS, Kriz AJ, Zetsche B, Shalem O, Wu X, Makarova KS, Koonin EV, Sharp PA, Zhang F., (published online 01 April 2015), Nature. Apr 9;520(7546):186-91 (2015).
- *High-throughput functional genomics using CRISPR-Cas9*, Shalem et al., Nature Reviews Genetics 16, 299-311 (May 2015).
- *Sequence determinants of improved CRISPR sgRNA design*, Xu et al., Genome Research 25, 1147-1157 (August 2015).

- *A Genome-wide CRISPR Screen in Primary Immune Cells to Dissect Regulatory Networks*, Parnas et al., Cell 162, 675-686 (July 30, 2015).
- *CRISPR/Cas9 cleavage of viral DNA efficiently suppresses hepatitis B virus*, Ramanan et al., Scientific Reports 5:10833. doi: 10.1038/srep10833 (June 2, 2015).
- *Crystal Structure of Staphylococcus aureus Cas9*, Nishimasu et al., Cell 162, 1113-1126 (Aug. 27, 2015).
- *BCL11A enhancer dissection by Cas9-mediated in situ saturating mutagenesis*, Canver et al., Nature 527(7577):192-7 (Nov. 12, 2015) doi: 10.1038/nature15521. Epub 2015 Sep 16.
- *Cpf1 Is a Single RNA-Guided Endonuclease of a Class 2 CRISPR-Cas System*, Zetsche et al., Cell 163, 759-71 (Sep 25, 2015).
- *Discovery and Functional Characterization of Diverse Class 2 CRISPR-Cas Systems*, Shmakov et al., Molecular Cell, 60(3), 385–397 doi: 10.1016/j.molcel.2015.10.008 Epub October 22, 2015.
- *Rationally engineered Cas9 nucleases with improved specificity*, Slaymaker et al., Science, DOI:10.1126/science.aad5227, Published online 1 December 2015.

each of which is incorporated herein by reference, and discussed briefly below:

- Cong *et al.* engineered type II CRISPR/Cas systems for use in eukaryotic cells based on both *Streptococcus thermophilus* Cas9 and also *Streptococcus pyogenes* Cas9 and demonstrated that Cas9 nucleases can be directed by short RNAs to induce precise cleavage of DNA in human and mouse cells. Their study further showed that Cas9 as converted into a nicking enzyme can be used to facilitate homology-directed repair in eukaryotic cells with minimal mutagenic activity. Additionally, their study demonstrated that multiple guide sequences can be encoded into a single CRISPR array to enable simultaneous editing of several at endogenous genomic loci sites within the mammalian genome, demonstrating easy programmability and wide applicability of the RNA-guided nuclease technology. This ability to use RNA to program sequence specific DNA cleavage in cells defined a new class of genome engineering tools. These studies further showed that other CRISPR loci are likely to be transplantable into mammalian cells and can also mediate mammalian genome cleavage. Importantly, it can be envisaged that

several aspects of the CRISPR/Cas system can be further improved to increase its efficiency and versatility.

- Jiang *et al.* used the clustered, regularly interspaced, short palindromic repeats (CRISPR)–associated Cas9 endonuclease complexed with dual-RNAs to introduce precise mutations in the genomes of *Streptococcus pneumoniae* and *Escherichia coli*. The approach relied on dual-RNA:Cas9-directed cleavage at the targeted genomic site to kill unmutated cells and circumvents the need for selectable markers or counter-selection systems. The study reported reprogramming dual-RNA:Cas9 specificity by changing the sequence of short CRISPR RNA (crRNA) to make single- and multinucleotide changes carried on editing templates. The study showed that simultaneous use of two crRNAs enabled multiplex mutagenesis. Furthermore, when the approach was used in combination with recombineering, in *S. pneumoniae*, nearly 100% of cells that were recovered using the described approach contained the desired mutation, and in *E. coli*, 65% that were recovered contained the mutation.
- Wang *et al.* (2013) used the CRISPR/Cas system for the one-step generation of mice carrying mutations in multiple genes which were traditionally generated in multiple steps by sequential recombination in embryonic stem cells and/or time-consuming intercrossing of mice with a single mutation. The CRISPR/Cas system will greatly accelerate the *in vivo* study of functionally redundant genes and of epistatic gene interactions.
- Konermann *et al.* addressed the need in the art for versatile and robust technologies that enable optical and chemical modulation of DNA-binding domains based CRISPR Cas9 enzyme and also Transcriptional Activator Like Effectors.
- Ran *et al.* (2013-A) described an approach that combined a Cas9 nickase mutant with paired guide RNAs to introduce targeted double-strand breaks. This addresses the issue of the Cas9 nuclease from the microbial CRISPR-Cas system being targeted to specific genomic loci by a guide sequence, which can tolerate certain mismatches to the DNA target and thereby promote undesired off-target mutagenesis. Because individual nicks in the genome are repaired with high fidelity, simultaneous nicking via appropriately offset guide RNAs is required for double-stranded breaks and extends the number of specifically recognized bases for target cleavage. The authors demonstrated that using

paired nicking can reduce off-target activity by 50- to 1,500-fold in cell lines and to facilitate gene knockout in mouse zygotes without sacrificing on-target cleavage efficiency. This versatile strategy enables a wide variety of genome editing applications that require high specificity.

- Hsu *et al.* (2013) characterized SpCas9 targeting specificity in human cells to inform the selection of target sites and avoid off-target effects. The study evaluated >700 guide RNA variants and SpCas9-induced indel mutation levels at >100 predicted genomic off-target loci in 293T and 293FT cells. The authors show that SpCas9 tolerates mismatches between guide RNA and target DNA at different positions in a sequence-dependent manner, sensitive to the number, position and distribution of mismatches. The authors further showed that SpCas9-mediated cleavage is unaffected by DNA methylation and that the dosage of SpCas9 and sgRNA can be titrated to minimize off-target modification. Additionally, to facilitate mammalian genome engineering applications, the authors reported providing a web-based software tool to guide the selection and validation of target sequences as well as off-target analyses.
- Ran *et al.* (2013-B) described a set of tools for Cas9-mediated genome editing *via* non-homologous end joining (NHEJ) or homology-directed repair (HDR) in mammalian cells, as well as generation of modified cell lines for downstream functional studies. To minimize off-target cleavage, the authors further described a double-nicking strategy using the Cas9 nickase mutant with paired guide RNAs. The protocol provided by the authors experimentally derived guidelines for the selection of target sites, evaluation of cleavage efficiency and analysis of off-target activity. The studies showed that beginning with target design, gene modifications can be achieved within as little as 1–2 weeks, and modified clonal cell lines can be derived within 2–3 weeks.
- Shalem *et al.* described a new way to interrogate gene function on a genome-wide scale. Their studies showed that delivery of a genome-scale CRISPR-Cas9 knockout (GeCKO) library targeted 18,080 genes with 64,751 unique guide sequences enabled both negative and positive selection screening in human cells. First, the authors showed use of the GeCKO library to identify genes essential for cell viability in cancer and pluripotent stem cells. Next, in a melanoma model, the authors screened for genes whose loss is involved in resistance to vemurafenib, a therapeutic that inhibits mutant protein kinase BRAF.

Their studies showed that the highest-ranking candidates included previously validated genes NF1 and MED12 as well as novel hits NF2, CUL3, TADA2B, and TADA1. The authors observed a high level of consistency between independent guide RNAs targeting the same gene and a high rate of hit confirmation, and thus demonstrated the promise of genome-scale screening with Cas9.

- Nishimasu *et al.* reported the crystal structure of *Streptococcus pyogenes* Cas9 in complex with sgRNA and its target DNA at 2.5 Å resolution. The structure revealed a bilobed architecture composed of target recognition and nuclease lobes, accommodating the sgRNA:DNA heteroduplex in a positively charged groove at their interface. Whereas the recognition lobe is essential for binding sgRNA and DNA, the nuclease lobe contains the HNH and RuvC nuclease domains, which are properly positioned for cleavage of the complementary and non-complementary strands of the target DNA, respectively. The nuclease lobe also contains a carboxyl-terminal domain responsible for the interaction with the protospacer adjacent motif (PAM). This high-resolution structure and accompanying functional analyses have revealed the molecular mechanism of RNA-guided DNA targeting by Cas9, thus paving the way for the rational design of new, versatile genome-editing technologies.
- Wu *et al.* mapped genome-wide binding sites of a catalytically inactive Cas9 (dCas9) from *Streptococcus pyogenes* loaded with single guide RNAs (sgRNAs) in mouse embryonic stem cells (mESCs). The authors showed that each of the four sgRNAs tested targets dCas9 to between tens and thousands of genomic sites, frequently characterized by a 5-nucleotide seed region in the sgRNA and an NGG protospacer adjacent motif (PAM). Chromatin inaccessibility decreases dCas9 binding to other sites with matching seed sequences; thus 70% of off-target sites are associated with genes. The authors showed that targeted sequencing of 295 dCas9 binding sites in mESCs transfected with catalytically active Cas9 identified only one site mutated above background levels. The authors proposed a two-state model for Cas9 binding and cleavage, in which a seed match triggers binding but extensive pairing with target DNA is required for cleavage.
- Platt *et al.* established a Cre-dependent Cas9 knockin mouse. The authors demonstrated *in vivo* as well as *ex vivo* genome editing using adeno-associated virus (AAV)-

lentivirus-, or particle-mediated delivery of guide RNA in neurons, immune cells, and endothelial cells.

- Hsu *et al.* (2014) is a review article that discusses generally CRISPR-Cas9 history from yogurt to genome editing, including genetic screening of cells.
- Wang *et al.* (2014) relates to a pooled, loss-of-function genetic screening approach suitable for both positive and negative selection that uses a genome-scale lentiviral single guide RNA (sgRNA) library.
- Doench *et al.* created a pool of sgRNAs, tiling across all possible target sites of a panel of six endogenous mouse and three endogenous human genes and quantitatively assessed their ability to produce null alleles of their target gene by antibody staining and flow cytometry. The authors showed that optimization of the PAM improved activity and also provided an on-line tool for designing sgRNAs.
- Swiech *et al.* demonstrate that AAV-mediated SpCas9 genome editing can enable reverse genetic studies of gene function in the brain.
- Konermann *et al.* (2015) discusses the ability to attach multiple effector domains, e.g., transcriptional activator, functional and epigenomic regulators at appropriate positions on the guide such as stem or tetraloop with and without linkers.
- Zetsche *et al.* demonstrates that the Cas9 enzyme can be split into two and hence the assembly of Cas9 for activation can be controlled.
- Chen *et al.* relates to multiplex screening by demonstrating that a genome-wide *in vivo* CRISPR-Cas9 screen in mice reveals genes regulating lung metastasis.
- Ran *et al.* (2015) relates to SaCas9 and its ability to edit genomes and demonstrates that one cannot extrapolate from biochemical assays. Shalem *et al.* (2015) described ways in which catalytically inactive Cas9 (dCas9) fusions are used to synthetically repress (CRISPRi) or activate (CRISPRa) expression, showing advances using Cas9 for genome-scale screens, including arrayed and pooled screens, knockout approaches that inactivate genomic loci and strategies that modulate transcriptional activity.
- Shalem *et al.* (2015) described ways in which catalytically inactive Cas9 (dCas9) fusions are used to synthetically repress (CRISPRi) or activate (CRISPRa) expression, showing advances using Cas9 for genome-scale screens, including arrayed and pooled screens,

knockout approaches that inactivate genomic loci and strategies that modulate transcriptional activity.

- Xu *et al.* (2015) assessed the DNA sequence features that contribute to single guide RNA (sgRNA) efficiency in CRISPR-based screens. The authors explored efficiency of CRISPR/Cas9 knockout and nucleotide preference at the cleavage site. The authors also found that the sequence preference for CRISPRi/a is substantially different from that for CRISPR/Cas9 knockout.
- Parnas *et al.* (2015) introduced genome-wide pooled CRISPR-Cas9 libraries into dendritic cells (DCs) to identify genes that control the induction of tumor necrosis factor (Tnf) by bacterial lipopolysaccharide (LPS). Known regulators of Tlr4 signaling and previously unknown candidates were identified and classified into three functional modules with distinct effects on the canonical responses to LPS.
- Ramanan *et al.* (2015) demonstrated cleavage of viral episomal DNA (cccDNA) in infected cells. The HBV genome exists in the nuclei of infected hepatocytes as a 3.2kb double-stranded episomal DNA species called covalently closed circular DNA (cccDNA), which is a key component in the HBV life cycle whose replication is not inhibited by current therapies. The authors showed that sgRNAs specifically targeting highly conserved regions of HBV robustly suppresses viral replication and depleted cccDNA.
- Nishimasu *et al.* (2015) reported the crystal structures of SaCas9 in complex with a single guide RNA (sgRNA) and its double-stranded DNA targets, containing the 5'-TTGAAT-3' PAM and the 5'-TTGGGT-3' PAM. A structural comparison of SaCas9 with SpCas9 highlighted both structural conservation and divergence, explaining their distinct PAM specificities and orthologous sgRNA recognition.
- Canver *et al.* (2015) demonstrated a CRISPR-Cas9-based functional investigation of non-coding genomic elements. The authors we developed pooled CRISPR-Cas9 guide RNA libraries to perform *in situ* saturating mutagenesis of the human and mouse BCL11A enhancers which revealed critical features of the enhancers.
- Zetsche *et al.* (2015) reported characterization of Cpf1, a class 2 CRISPR nuclease from *Francisella novicida* U112 having features distinct from Cas9. Cpf1 is a single RNA-

guided endonuclease lacking tracrRNA, utilizes a T-rich protospacer-adjacent motif, and cleaves DNA via a staggered DNA double-stranded break.

- Shmakov et al. (2015) reported three distinct Class 2 CRISPR-Cas systems. Two system CRISPR enzymes (C2c1 and C2c3) contain RuvC-like endonuclease domains distantly related to Cpf1. Unlike Cpf1, C2c1 depends on both crRNA and tracrRNA for DNA cleavage. The third enzyme (C2c2) contains two predicted HEPN RNase domains and is tracrRNA independent.
- Slaymaker et al (2015) reported the use of structure-guided protein engineering to improve the specificity of *Streptococcus pyogenes* Cas9 (SpCas9). The authors developed "enhanced specificity" SpCas9 (eSpCas9) variants which maintained robust on-target cleavage with reduced off-target effects.

[00655] Mention is also made of Tsai et al, "Dimeric CRISPR RNA-guided FokI nucleases for highly specific genome editing," *Nature Biotechnology* 32(6): 569-77 (2014) which is not believed to be prior art to the instant invention or application, but which may be considered in the practice of the instant invention. Mention is also made of Konermann et al., "Genome-scale transcription activation by an engineered CRISPR-Cas9 complex," doi:10.1038/nature14136, incorporated herein by reference.

[00656] Also, "Dimeric CRISPR RNA-guided FokI nucleases for highly specific genome editing", Shengdar Q. Tsai, Nicolas Wyvekens, Cyd Khayter, Jennifer A. Foden, Vishal Thapar, Deepak Reyon, Mathew J. Goodwin, Martin J. Aryee, J. Keith Joung *Nature Biotechnology* 32(6): 569-77 (2014), relates to dimeric RNA-guided FokI Nucleases that recognize extended sequences and can edit endogenous genes with high efficiencies in human cells.

[00657] With respect to general information on CRISPR-Cas Systems, components thereof, and delivery of such components, including methods, materials, delivery vehicles, vectors, particles, AAV, and making and using thereof, including as to amounts and formulations, all useful in the practice of the instant invention, reference is made to: US Patents Nos. 8,697,359, 8,771,945, 8,795,965, 8,865,406, 8,871,445, 8,889,356, 8,889,418, 8,895,308, 8,906,616, 8,932,814, 8,945,839, 8,993,233 and 8,999,641; US Patent Publications US 2014-0310830 (US App. Ser. No. 14/105,031), US 2014-0287938 A1 (U.S. App. Ser. No. 14/213,991), US 2014-0273234 A1 (U.S. App. Ser. No. 14/293,674), US2014-0273232 A1 (U.S. App. Ser. No. 14/290,575), US 2014-0273231 (U.S. App. Ser. No. 14/259,420), US 2014-0256046 A1 (U.S.

App. Ser. No. 14/226,274), US 2014-0248702 A1 (U.S. App. Ser. No. 14/258,458), US 2014-0242700 A1 (U.S. App. Ser. No. 14/222,930), US 2014-0242699 A1 (U.S. App. Ser. No. 14/183,512), US 2014-0242664 A1 (U.S. App. Ser. No. 14/104,990), US 2014-0234972 A1 (U.S. App. Ser. No. 14/183,471), US 2014-0227787 A1 (U.S. App. Ser. No. 14/256,912), US 2014-0189896 A1 (U.S. App. Ser. No. 14/105,035), US 2014-0186958 (U.S. App. Ser. No. 14/105,017), US 2014-0186919 A1 (U.S. App. Ser. No. 14/104,977), US 2014-0186843 A1 (U.S. App. Ser. No. 14/104,900), US 2014-0179770 A1 (U.S. App. Ser. No. 14/104,837) and US 2014-0179006 A1 (U.S. App. Ser. No. 14/183,486), US 2014-0170753 (US App Ser No 14/183,429); US 2015-0184139 (U.S. App. Ser. No. 14/324,960); 14/054,414 European Patent Applications EP 2 771 468 (EP13818570.7), EP 2 764 103 (EP13824232.6), and EP 2 784 162 (EP14170383.5); and PCT Patent Publications WO 2014/093661 (PCT/US2013/074743), WO 2014/093694 (PCT/US2013/074790), WO 2014/093595 (PCT/US2013/074611), WO 2014/093718 (PCT/US2013/074825), WO 2014/093709 (PCT/US2013/074812), WO 2014/093622 (PCT/US2013/074667), WO 2014/093635 (PCT/US2013/074691), WO 2014/093655 (PCT/US2013/074736), WO 2014/093712 (PCT/US2013/074819), WO 2014/093701 (PCT/US2013/074800), WO 2014/018423 (PCT/US2013/051418), WO 2014/204723 (PCT/US2014/041790), WO 2014/204724 (PCT/US2014/041800), WO 2014/204725 (PCT/US2014/041803), WO 2014/204726 (PCT/US2014/041804), WO 2014/204727 (PCT/US2014/041806), WO 2014/204728 (PCT/US2014/041808), WO 2014/204729 (PCT/US2014/041809), WO 2015/089351 (PCT/US2014/069897), WO 2015/089354 (PCT/US2014/069902), WO 2015/089364 (PCT/US2014/069925), WO 2015/089427 (PCT/US2014/070068), WO 2015/089462 (PCT/US2014/070127), WO 2015/089419 (PCT/US2014/070057), WO 2015/089465 (PCT/US2014/070135), WO 2015/089486 (PCT/US2014/070175), PCT/US2015/051691, PCT/US2015/051830. Reference is also made to US provisional patent applications 61/758,468; 61/802,174; 61/806,375; 61/814,263; 61/819,803 and 61/828,130, filed on January 30, 2013; March 15, 2013; March 28, 2013; April 20, 2013; May 6, 2013 and May 28, 2013 respectively. Reference is also made to US provisional patent application 61/836,123, filed on June 17, 2013. Reference is additionally made to US provisional patent applications 61/835,931, 61/835,936, 61/835,973, 61/836,080, 61/836,101, and 61/836,127, each filed June 17, 2013. Further reference is made to US provisional patent applications 61/862,468 and 61/862,355 filed on August 5, 2013; 61/871,301

filed on August 28, 2013; 61/960,777 filed on September 25, 2013 and 61/961,980 filed on October 28, 2013. Reference is yet further made to: PCT/US2014/62558 filed October 28, 2014, and US Provisional Patent Applications Serial Nos.: 61/915,148, 61/915,150, 61/915,153, 61/915,203, 61/915,251, 61/915,301, 61/915,267, 61/915,260, and 61/915,397, each filed December 12, 2013; 61/757,972 and 61/768,959, filed on January 29, 2013 and February 25, 2013; 62/010,888 and 62/010,879, both filed June 11, 2014; 62/010,329, 62/010,439 and 62/010,441, each filed June 10, 2014; 61/939,228 and 61/939,242, each filed February 12, 2014; 61/980,012, filed April 15, 2014; 62/038,358, filed August 17, 2014; 62/055,484, 62/055,460 and 62/055,487, each filed September 25, 2014; and 62/069,243, filed October 27, 2014. Reference is made to PCT application designating, inter alia, the United States, application No. PCT/US14/41806, filed June 10, 2014. Reference is made to US provisional patent application 61/930,214 filed on January 22, 2014. Reference is made to PCT application designating, inter alia, the United States, application No. PCT/US14/41806, filed June 10, 2014.

[00658] Mention is also made of US application 62/180,709, 17-Jun-15, PROTECTED GUIDE RNAS (PGRNAS); US application 62/091,455, filed, 12-Dec-14, PROTECTED GUIDE RNAS (PGRNAS); US application 62/096,708, 24-Dec-14, PROTECTED GUIDE RNAS (PGRNAS); US applications 62/091,462, 12-Dec-14, 62/096,324, 23-Dec-14, 62/180,681, 17-Jun-2015, and 62/237,496, 5-Oct-2015, DEAD GUIDES FOR CRISPR TRANSCRIPTION FACTORS; US application 62/091,456, 12-Dec-14 and 62/180,692, 17-Jun-2015, ESCORTED AND FUNCTIONALIZED GUIDES FOR CRISPR-CAS SYSTEMS; US application 62/091,461, 12-Dec-14, DELIVERY, USE AND THERAPEUTIC APPLICATIONS OF THE CRISPR-CAS SYSTEMS AND COMPOSITIONS FOR GENOME EDITING AS TO HEMATOPOETIC STEM CELLS (HSCs); US application 62/094,903, 19-Dec-14, UNBIASED IDENTIFICATION OF DOUBLE-STRAND BREAKS AND GENOMIC REARRANGEMENT BY GENOME-WISE INSERT CAPTURE SEQUENCING; US application 62/096,761, 24-Dec-14, ENGINEERING OF SYSTEMS, METHODS AND OPTIMIZED ENZYME AND GUIDE SCAFFOLDS FOR SEQUENCE MANIPULATION; US application 62/098,059, 30-Dec-14, 62/181,641, 18-Jun-2015, and 62/181,667, 18-Jun-2015, RNA-TARGETING SYSTEM; US application 62/096,656, 24-Dec-14 and 62/181,151, 17-Jun-2015, CRISPR HAVING OR ASSOCIATED WITH DESTABILIZATION DOMAINS; US application 62/096,697, 24-Dec-14, CRISPR HAVING OR ASSOCIATED WITH AAV; US application 62/098,158, 30-Dec-14,

ENGINEERED CRISPR COMPLEX INSERTIONAL TARGETING SYSTEMS; US application 62/151,052, 22-Apr-15, CELLULAR TARGETING FOR EXTRACELLULAR EXOSOMAL REPORTING; US application 62/054,490, 24-Sep-14, DELIVERY, USE AND THERAPEUTIC APPLICATIONS OF THE CRISPR-CAS SYSTEMS AND COMPOSITIONS FOR TARGETING DISORDERS AND DISEASES USING PARTICLE DELIVERY COMPONENTS; US application 61/939,154, 12-F EB-14, SYSTEMS, METHODS AND COMPOSITIONS FOR SEQUENCE MANIPULATION WITH OPTIMIZED FUNCTIONAL CRISPR-CAS SYSTEMS; US application 62/055,484, 25-Sep-14, SYSTEMS, METHODS AND COMPOSITIONS FOR SEQUENCE MANIPULATION WITH OPTIMIZED FUNCTIONAL CRISPR-CAS SYSTEMS; US application 62/087,537, 4-Dec-14, SYSTEMS, METHODS AND COMPOSITIONS FOR SEQUENCE MANIPULATION WITH OPTIMIZED FUNCTIONAL CRISPR-CAS SYSTEMS; US application 62/054,651, 24-Sep-14, DELIVERY, USE AND THERAPEUTIC APPLICATIONS OF THE CRISPR-CAS SYSTEMS AND COMPOSITIONS FOR MODELING COMPETITION OF MULTIPLE CANCER MUTATIONS IN VIVO; US application 62/067,886, 23-Oct-14, DELIVERY, USE AND THERAPEUTIC APPLICATIONS OF THE CRISPR-CAS SYSTEMS AND COMPOSITIONS FOR MODELING COMPETITION OF MULTIPLE CANCER MUTATIONS IN VIVO; US applications 62/054,675, 24-Sep-14 and 62/181,002, 17-Jun-2015, DELIVERY, USE AND THERAPEUTIC APPLICATIONS OF THE CRISPR-CAS SYSTEMS AND COMPOSITIONS IN NEURONAL CELLS/TISSUES; US application 62/054,528, 24-Sep-14, DELIVERY, USE AND THERAPEUTIC APPLICATIONS OF THE CRISPR-CAS SYSTEMS AND COMPOSITIONS IN IMMUNE DISEASES OR DISORDERS; US application 62/055,454, 25-Sep-14, DELIVERY, USE AND THERAPEUTIC APPLICATIONS OF THE CRISPR-CAS SYSTEMS AND COMPOSITIONS FOR TARGETING DISORDERS AND DISEASES USING CELL PENETRATION PEPTIDES (CPP); US application 62/055,460, 25-Sep-14, MULTIFUNCTIONAL-CRISPR COMPLEXES AND/OR OPTIMIZED ENZYME LINKED FUNCTIONAL-CRISPR COMPLEXES; US application 62/087,475, 4-Dec-14 and 62/181,690, 18-Jun-2015, FUNCTIONAL SCREENING WITH OPTIMIZED FUNCTIONAL CRISPR-CAS SYSTEMS; US application 62/055,487, 25-Sep-14, FUNCTIONAL SCREENING WITH OPTIMIZED FUNCTIONAL CRISPR-CAS SYSTEMS; US application 62/087,546, 4-Dec-14 and 62/181,687, 18-Jun-2015, MULTIFUNCTIONAL CRISPR COMPLEXES AND/OR

OPTIMIZED ENZYME LINKED FUNCTIONAL-CRISPR COMPLEXES; and US application 62/098,285, 30-Dec-14, CRISPR MEDIATED IN VIVO MODELING AND GENETIC SCREENING OF TUMOR GROWTH AND METASTASIS.

[00659] Mention is made of US applications 62/181,659, 18-Jun-2015 and 62/207,318, 19-Aug-2015, ENGINEERING AND OPTIMIZATION OF SYSTEMS, METHODS, ENZYME AND GUIDE SCAFFOLDS OF CAS9 ORTHOLOGS AND VARIANTS FOR SEQUENCE MANIPULATION. Mention is made of US applications 62/181,663, 18-Jun-2015 and 62/245,264, 22-Oct-2015, NOVEL CRISPR ENZYMES AND SYSTEMS, US application 62/232,067, 24-Sep-2015, US application 62/205,733, 16-Aug-2015, US application 62/201,542, 5-Aug-2015, US application 62/193,507, 16-Jul-2015, and US application 62/181,739, 18-Jun-2015, each entitled NOVEL CRISPR ENZYMES AND SYSTEMS. Mention is also made of US application 61/939,256, 12-Feb-2014, and WO 2015/089473 (PCT/US2014/070152), 12-Dec-2014, each entitled ENGINEERING OF SYSTEMS, METHODS AND OPTIMIZED GUIDE COMPOSITIONS WITH NEW ARCHITECTURES FOR SEQUENCE MANIPULATION. Mention is also made of PCT/US2015/045504, 15-Aug-2015, US application 62/180,699, 17-Jun-2015, and US application 62/038,358, 17-Aug-2014, each entitled GENOME EDITING USING CAS9 NICKASES.

[00660] Each of these patents, patent publications, and applications, and all documents cited therein or during their prosecution (“appln cited documents”) and all documents cited or referenced in the appln cited documents, together with any instructions, descriptions, product specifications, and product sheets for any products mentioned therein or in any document therein and incorporated by reference herein, are hereby incorporated herein by reference, and may be employed in the practice of the invention. All documents (e.g., these patents, patent publications and applications and the appln cited documents) are incorporated herein by reference to the same extent as if each individual document was specifically and individually indicated to be incorporated by reference.

[00661] In addition, mention is made of PCT application PCT/US14/70057, Attorney Reference 47627.99.2060 and BI-2013/107 entitled “DELIVERY, USE AND THERAPEUTIC APPLICATIONS OF THE CRISPR-CAS SYSTEMS AND COMPOSITIONS FOR TARGETING DISORDERS AND DISEASES USING PARTICLE DELIVERY COMPONENTS (claiming priority from one or more or all of US provisional patent

applications: 62/054,490, filed September 24, 2014; 62/010,441, filed June 10, 2014; and 61/915,118, 61/915,215 and 61/915,148, each filed on December 12, 2013) ("the Particle Delivery PCT"), incorporated herein by reference, with respect to a method of preparing an sgRNA-and-Cas9 protein containing particle comprising admixing a mixture comprising an sgRNA and Cas9 protein (and optionally HDR template) with a mixture comprising or consisting essentially of or consisting of surfactant, phospholipid, biodegradable polymer, lipoprotein and alcohol; and particles from such a process. For example, wherein Cas9 protein and sgRNA were mixed together at a suitable, e.g., 3:1 to 1:3 or 2:1 to 1:2 or 1:1 molar ratio, at a suitable temperature, e.g., 15-30C, e.g., 20-25C, e.g., room temperature, for a suitable time, e.g., 15-45, such as 30 minutes, advantageously in sterile, nuclease free buffer, e.g., 1X PBS. Separately, particle components such as or comprising: a surfactant, e.g., cationic lipid, e.g., 1,2-dioleoyl-3-trimethylammonium-propane (DOTAP); phospholipid, e.g., dimyristoylphosphatidylcholine (DMPC); biodegradable polymer, such as an ethylene-glycol polymer or PEG, and a lipoprotein, such as a low-density lipoprotein, e.g., cholesterol were dissolved in an alcohol, advantageously a C1-6 alkyl alcohol, such as methanol, ethanol, isopropanol, e.g., 100% ethanol. The two solutions were mixed together to form particles containing the Cas9-sgRNA complexes. Accordingly, sgRNA may be pre-complexed with the Cas9 protein, before formulating the entire complex in a particle. Formulations may be made with a different molar ratio of different components known to promote delivery of nucleic acids into cells (e.g. 1,2-dioleoyl-3-trimethylammonium-propane (DOTAP), 1,2-ditetradecanoyl-sn-glycero-3-phosphocholine (DMPC), polyethylene glycol (PEG), and cholesterol) For example DOTAP : DMPC : PEG : Cholesterol Molar Ratios may be DOTAP 100, DMPC 0, PEG 0, Cholesterol 0; or DOTAP 90, DMPC 0, PEG 10, Cholesterol 0; or DOTAP 90, DMPC 0, PEG 5, Cholesterol 5. DOTAP 100, DMPC 0, PEG 0, Cholesterol 0. That application accordingly comprehends admixing sgRNA, Cas9 protein and components that form a particle; as well as particles from such admixing. Aspects of the instant invention can involve particles; for example, particles using a process analogous to that of the Particle Delivery PCT, e.g., by admixing a mixture comprising sgRNA and/or Cas9 as in the instant invention and components that form a particle, e.g., as in the Particle Delivery PCT, to form a particle and particles from such admixing (or, of course, other particles involving sgRNA and/or Cas9 as in the instant invention).

[00662] The present invention will be further illustrated in the following Examples which are given for illustration purposes only and are not intended to limit the invention in any way.

Examples

Example 1: Targeted CRISPR gene activation without nuclease activity / without indel activity using dead guide sequence

[00663] sgRNA directed to Sp. Cas9 and comprising a dead guide sequence having a length of 13 nucleotides was designed to target IL1B. Using transcriptional analysis, IL1B/GAPDH activation was at least as strong as the positive control. The sgRNA included two MS2 loops paired with transcriptional activator. The positive control had dCas9, MS2-p65-HSF1, and an IL1B targeting sequence with 20 bp. The IL1B-13 group had Cas9, MS2-p65-HSF1, and an IL1B targeting sequence reduced to 13 bp.

Example 2: Dead guides direct CRISPR binding to target without nuclease activity / indel activity.

[00664] Shortened sgRNA sequences targeting different sequences within +/- 100 base pair of the transcriptional start site are designed (Konermann et al., "Genome-scale transcription activation by an engineered CRISPR-Cas9 complex," doi:10.1038/nature14136, incorporated herein by reference.). The sequence length is 20 base pair (control), or 16, 15, 14, 13, 12, 11, or 10 base pairs. 17 base pair constructs have been shown to produce insertions and deletions. Sequences are designed that have a G on their 5' end, in order to enhance their production in the cell.

[00665] One day after plating HEK293 cells in a 96 well plate, the cells are transfected with 100 ng (active) Sp Cas9 plasmid, 100 ng MS2 plasmid, and 100 ng deadGuides. Two days later, cellular DNA is isolated, and insertions and deletions are analyzed using surveyor analysis. Separately, cellular RNA is isolated, and quantify the transcriptional expression of the gene of interest (GeneX) and the control gene GAPDH. DeadRNAs have a high (GeneX/GAPDH) expression, and no insertions or deletions.

Example 3: Dead guide effect repeatable across different genes

[00666] Four controls: 1) untreated cells, 2) GFP plasmid to control for differences in cell behavior caused by Lipofectamine used for cell transfection, 3) positive control for activation (construct + dCas9 + MS2) (Konermann et al., "Genome-scale transcription activation by an

engineered CRISPR-Cas9 complex,” doi:10.1038/nature14136, incorporated herein by reference.), and 4) a positive control for indel formation (normal sgRNA + active Cas9).

[00667] Shortened sgRNA sequences targeting different sequences within +/- 100 base pair of the transcriptional start site of three genes (i.e. one target per gene, three genes) are generated. The sequence length is 20 base pair (control), or 16, 15, 14, 13, 12, 11, or 10 base pair.

[00668] One day after plating HEK293 cells in a 96 well plate, the cells are transfected with 100 ng (active) Cas9 plasmid, 100 ng MS2 plasmid, and 100 ng deadGuides. Two days later, the cellular DNA is isolated, and insertions and deletions are analyzed using either surveyor or next generation sequencing. Separately, cellular RNA is separated, and the transcriptional expression of gene X and the control gene GAPDH is quantified. Dead RNAs have a high (GeneX/GAPDH) expression, and no insertions or deletions using surveyor analysis.

Example 4: Dead guide multigene activation and deletion

[00669] Two constructs were selected. Two other normal sgRNAs that have been previously validated and which are directed to genes whose downregulation is easily measured *in vivo* are also used. Following the same HEK293 protocol, the cells are transfected with four different constructs, and transcriptional activation and insertions or deletions for all four genes is measured.

Example 5: Dead guide off target effects

[00670] Off-target effects are analyzed using BLESS, if necessary. Off-target activation is not particularly expected on account that off-target binding would have to take place very close to the transcriptional start site of the off-target gene.

Example 6: Dead guide in vivo multigene activation and deletion

[00671] The CRISPR-Cas9 knockin mouse (Platt et al., Cell 159, 440-455, Oct. 2014) is used to repeat in a mouse using virus apply to the liver. This experiment is repeated using local injection in the ear.

Example 7: Dead guide combinatorial biology.

[00672] Biology which is relatively quick, and biology where one can first delete a first gene of interest (gene X) and then compensate for it by increasing a second gene of interest (gene Y) is chosen. For instance, p53 is deleted, then this is compensated for with upregulating LKB1. Further, in the growth pathway Roman loves, a major gene at the top of that signaling pathway is knocked out and this is compensated for by upregulating immediately downstream factors.

Experiments are performed in cell lines as well in vivo.

Example 8: Orthogonal gene regulation

[00673] Orthogonal gene regulation utilizing expression of a single active Cas9 enzyme uses sgRNA scaffolds for activation and repression of target genes. Genes targeted for repression utilize 20 base pair guide RNAs and genes targeted for activation utilize shorter guide RNAs of 13 base pairs and 14 base pairs respectively. Not being bound by a theory the shorter guide RNAs allow recruitment of Cas9 to a target gene without cutting of the target. Genes targeted for activation additionally include stem loop structures, such as the MS2 aptamer sequence, for recruitment of adaptor proteins linked to an activator. The activator can be p65 or HSF1 as is shown in figure 5.

[00674] **Materials and Methods.** HEK.293 cells were plated in a 96 well plate. 24 hours later, cells were transfected with 100 ng component 1, 100ng component two, and in some cases, 100 ng of component three (see Table 1). 48 hours after transfection, the cells were lysed, and cellular DNA or cellular RNA was isolated. Cellular DNA was isolated using Quick Extract buffer, according to the manufacturer instructions. Cellular RNA was isolated using a Qiagen RNA isolation kit, per manufacturer instructions. The presence of indels was measured using Surveyor, as previously described (Cong et al, *Science* 2013). The transcription of the target gene IL1B, as well as the transcription of a control gene GapDH was measured using a Applied Biosystems qPCR kit, following the manufacturer instructions. The relative upregulation of IL1B was quantified as follows: the ratio of IL1B/GapDH was quantified for each well (N=4 wells/group for all treated groups, and N=16 for untreated cells). This ratio in untreated cells was defined as 1. The ratio for all other groups was normalized to this ratio. For example, if the average IL1B/GapDH ratio for the 16 untreated wells is 0.25, then a treated well with a IL1B/GapDH ratio equal to 250 is upregulated by 1000x.

Table 1.

	IL1B sgRNA Added (Component 1)	Cas9 Added (Component 2)	Activator Added (Component 3)
1	——AGCGAGGGAGAAAC + tracrRNA w/MS2 loop	Cas9+sgEMX1.3	MS2-p65-HSF1
2	———GCGAGGGAGAAAC + tracrRNA w/MS2 loop	Cas9+sgEMX1.3	MS2-p65-HSF1
3	AAAAACAGCGAGGGAGAAAC + regular tracrRNA	Cas9+sgEMX1.3	None
4	GAAAAACAGCGAGGGAGAAAC + tracrRNA w/ MS2 loop	dCas9	MS2-p65-HSF1
5	GFP Plasmid	None	None

[00675] *Example 9: Engineering Dead sgRNAs for Bimodal Gene Control*

[00676] Cells execute complex transcriptional programs with independent regulation at different genome loci. A variety of CRISPR/Cas9 systems have been developed for single gene perturbations, such as gene activation or inactivation. There remains a need to provide methods to recapitulate aspects of complex cell circuits, for example activating and inactivating alternative genes in a single system. The present example illustrates an approach to engineering sgRNAs so as to facilitate bimodal gene control, in some embodiments using only a single active Cas9. More specifically, Applicants use truncated sgRNA guides that can mediate binding of a Cas9 to a target DNA without cutting it. Applicants illustrate the modification of these truncated sgRNAs with MS2-loops on the scaffold, so as to recruit MS2-p65-HSF1 fusions, which promote targeted gene activation. When used with full-length 20bp sgRNAs with an unmodified scaffold, Applicants demonstrate bimodal gene control at multiple loci, and are able to illustrate effective combinations of tumor suppressors and oncogenes for synergistic resistance in melanoma.

[00677] As illustrated schematically in Figure 6A, dead guide RNAs can combine changes to the sgRNA that prevent cutting, and stem loop MS2 modifications that allow recruitment of transcriptional activators (HSF1/P65), to generate an active Cas9 complex that is capable of transcriptional activation. This example illustrates that gene *activation* can be achieved using an *active* Cas9, and in this way the dead guides described herein enable gene activation in a Cas9-expressing mouse which may be utilized in bimodal gene perturbation assays that require only a single Cas9 enzyme. In this example, dead-guide-mediated gene activation is achieved with four components: a 14-15bp guide sequence, MS2 loops on the tetraloop and stem loop 2, MS2-P65-HSF1 fusion protein, and an active Cas9 enzyme.

[00678] *Determining Optimal Truncation Length*

[00679] As an initial step, to determine an optimal truncation (or mismatch) length, Applicants designed a set of possible guides ranging in length from 20bp to 10bp targeting the upstream promoter region of HBG1 (as set out in the Table below). All sgRNA had MS2 loops and were delivered along with an active Cas9 in order to test for activation. As an alternative approach, Applicants additionally synthesized a similar set of constructs that had mismatched bases in place of truncations (mismatched guides function similarly to truncated guides).

HBG1-E20	<u>GTATCCAGT</u> GAGGCCAGGGGC
HBG1-E19	- <u>GATCCAGT</u> GAGGCCAGGGGC
HBG1-E20	-- <u>GTCCAGT</u> GAGGCCAGGGGC
HBG1-E17	--- <u>GCCAGT</u> GAGGCCAGGGGC
HBG1-E16	---- <u>GCAST</u> GAGGCCAGGGGC
HBG1-E15	----- <u>GAGT</u> GAGGCCAGGGGC
HBG1-E14	----- <u>GST</u> GAGGCCAGGGGC
HBG1-E13	----- <u>GT</u> GAGGCCAGGGGC
HBG1-E12	----- <u>GG</u> GAGGCCAGGGGC
HBG1-E11	----- <u>GAG</u> GCCAGGGGC
HBG1-E10	----- <u>GGG</u> CCAGGGGC
HBG1-E20 Mismatched	<u>GTATCCAGT</u> GAGGCCAGGGGC
HBG1-E19 Mismatched	<u>GCATCCAGT</u> GAGGCCAGGGGC
HBG1-E18 Mismatched	<u>GCGTCCAGT</u> GAGGCCAGGGGC
HBG1-E17 Mismatched	<u>GCGCCAGT</u> GAGGCCAGGGGC
HBG1-E16 Mismatched	<u>GCGCTCAGT</u> GAGGCCAGGGGC
HBG1-E15 Mismatched	<u>GCGCTTAGT</u> GAGGCCAGGGGC
HBG1-E14 Mismatched	<u>GCGCTTGST</u> GAGGCCAGGGGC
HBG1-E13 Mismatched	<u>GCGCTTGAT</u> GAGGCCAGGGGC
HBG1-E12 Mismatched	<u>GCGCTTGAC</u> GAGGCCAGGGGC
HBG1-E11 Mismatched	<u>GCGCTTGACA</u> AGGCCAGGGGC
HBG1-E10 Mismatched	<u>GCGCTTGACAG</u> GGGCCAGGGGC

[00680] As illustrated in Figure 6B, for three different prospective guides within 200bp upstream of HBG1, having the target sequences shown in Figure 6B above the respective results, Applicants achieved robust gene activation for guides less than 16bp in length. Applicants confirmed by next generation sequencing that activation at 16bp or longer was due to cutting at the locus. For further clarity, the results for truncated guides illustrated in the first column of Figure 6B are illustrated independently, apart from the results for mismatched guides, in the bar

graphs of Figure 6BB. As summarized above, these results were obtained by transfecting eighty sgRNA-MS2s targeting four DNA sequences within 200 bp of the transcriptional start site of HBG1 together with active Cas9 and the MS2-P65-HSF1 (MPH) activation complex. Applicants illustrate that guides from 20nt to 16nt resulted in indel formation, whereas shorter guides (11nt to 15nt) did not show detectable levels of indel formation in most cases (Fig. 6BB second graph). Notably, guides truncated to 11-15nt of complementarity to the target DNA were able to increase HBG1 mRNA expression by as much as 10,000 fold (Fig. 6BB).

[00681] As illustrated in the second, third and fourth row plots of Figure 6BB, Three different dRNAs targeting the HBG1 promoter region were designed. The length of the RNA targeting sequence was varied from 11nt to 20nt. HBG1 mRNA (normalized to GAPDH, and compared to cells transfected with GFP plasmid) was quantified, as well HBG1 indel frequency. In all cases, guides were designed with MS2 binding loops in the tetraloops and stem loop two, and were co-transfected with active Cas9 and the MPH transcriptional activation complex. Average +/- SEM is plotted, N=2-3 replicates group.

[00682] *Dead Guide Activation of Multiple Genes*

[00683] As illustrated in Figure 6C, Applicants designed 14 and 15bp sgRNAs with MS2 loops to target three different genes (IL1B, HBG1, and ZFP42) in order to demonstrate that the activation effect using a dead sgRNA was reproducible at different loci. The graphs of Figure 6C show that dead guides robustly work for these three genes, and in some cases the active Cas9 with a dead sgRNA mediates an activity that is similar to activation with a dead Cas9 - or in some cases better. Next generation sequencing shows that the effect of truncation is to eliminate cutting and that this effect is mostly due to truncation and not due to the addition of the MS2 loop.

[00684] The results illustrated in Figure 6C are illustrative of fourteen and fifteen nt dRNAs, when cotransfected into HEK293FT cells with active Cas9 and the MPH complex, showing increased target mRNA expression of all three human genes (HBG1, Interleukin 1 β (IL1 β), and Zinc Finger Protease 42 (ZFP42)) without inducing significant indel formation (Fig. 6C). Notably, dRNA activation was comparable to the recently reported system using dCas9 in combination with a 20nt sgRNA-MS211. At all three loci 20nt sgRNAs cut target DNA and did not activate gene expression when combined with active Cas9. This was true for sgRNAs with and without the MS2 binding loops (Fig. 6C). Taken together, these data demonstrate that

dRNAs can activate gene expression without forming indels at targeted DNA using an active Cas9 with comparable efficiency to the current dCas9 system.

[00685] *Whole genome specificity analysis of dead sgRNA*

[00686] Whole transcriptome RNA sequencing on 15bp deadRNA (with active Cas9 + MS2-P65-HSF1) and 20bp sgRNA (with dead Cas9 + MS2-P65-HSF1) was used to illustrate the degree of change in specificity caused by shorter sgRNAs. In this example, the sgRNAs targeted the promoter region of *HBG1*. As illustrated in Figure 7A, using the approach summarized below, specificity was not significantly changed for the truncated 15bp deadRNA, which evidenced specificity similar to the 20bp sgRNA with dCas9. More specifically, to illustrate the difference in specificity between 20nt sgRNA-MS2 and 15nt dRNAs Applicants compared whole transcriptome mRNA levels in HEK293FT cells. Cells were co-transfected with dCas9, the MPH complex, and a 20nt activator sgRNA-MS2, or active Cas9, the MPH complex, and 15nt dRNA targeting the same sequence in the human *HBG1/2* promoter. Applicants separately determined that *HBG1/2* upregulation induces limited downstream effects that could confound analysis in HEK293FT cells. RNA-seq results showed that both the sgRNA/dCas9 and dRNA systems significantly activated *HBG1/2* only, demonstrating that dRNAs can specifically upregulate target genes (Fig. 7a). Applicants next performed off-target analysis on a second 15nt dRNA and 20nt sgRNA targeting the same *HBG1/2* promoter. Surprisingly, Applicants found a significant number of perturbed transcripts for both the 15nt and 20nt guide RNAs (Fig. 7b).

[00687] Differential gene expression analysis yielded results shown in Figure 7c, showing that the off target genes have minimal gene expression fold differences when compared to the on target gene *HBG1/2*.

[00688] *Bimodal gene control to model tumor resistance - resistance to BRAF-mutant A375 cells.*

[00689] Bimodal gene control was illustrated by inducing resistance to BRAF-inhibition through combinations of tumor suppressor knockouts and oncogene activation. In summary, this involved the delivery of an active Cas9 and MS2-p65-HSF1 fusion protein, along with 15bp guides having MS2 loops, targeting oncogenes for activation; and, delivery of 20bp guides targeting tumor suppressors for cutting allows. Perturbations were made in pairwise combinations between tumor suppressors (*CUL3* and *MED12*) and oncogenes (*LPAR5*, *ITGA9*, and *EGFR*). Resistance was measured in the BRAF-mutant melanoma line A375 against the

BRAF inhibitor PLX4720. The gene targets were selected from GECKO knockout (CUL3 and MED12) and SAM (LPAR5, ITGA9, and EGFR) screens, and all pairwise combinations were tested. An A375 cell line expressing Cas9 and MS2-P65-HSF1 was generated via lentiviral transduction. This cell line was then transduced, as summarized above, with different combinations of active and dead sgRNAs using lentivirus. (either: a) single active sgRNA; or b) single dead sgRNA(MS2); or a combination of a) + b)).

[00690] Figure 8A illustrates successful bimodal control (cutting of one gene and activation of a separate gene in the same pool of cells using active Cas9), measured one week after lentiviral transduction and following antibiotic selection.

[00691] To illustrate that bimodal gene perturbations of this kind may be used to cause phenotypic effects, the increase in resistance conferred to A375 cells under PLX4720 BRAF inhibition was measured. The results, as shown in Figure 8B, indicate that each perturbation individually increased the resistance of these cells to PLX4720 and that the combinations further shifted resistance, with some combinations exhibiting synergistic behaviour (e.g. MED12 and LPAR4, which exhibited a perturbation index (P.I.) > 1, indicating synergistic behavior).

[00692] Definition of perturbation index:

$$P.I. = \frac{\frac{PC}{C}}{\frac{P1}{C} \frac{P2}{C}}$$

- P1 = IC50 for PLX under perturbation 1
- P2 = IC50 for PLX under perturbation 2
- C = IC50 for PLX on control line (A375)
- CP = IC50 for PLX under combination of perturbations
- Synergy if P.I. > 1
- Additive if P.I. = 1
- Antagonistic if P.I. < 1

[00693] Additional data related to this example is provided in Figure 9, illustrative of the fact that dRNAs in combination with sgRNAs can mediate orthogonal gene control (activation and knockout) using only active Cas9. Applicants separately used CRISPR-Cas9 loss-of-function (LOF)21 and gain-of-function (GOF)11 screens to identify genetic modifiers that promote

resistance of A375 melanoma cells to the BRAF inhibitor PLX-4720. Specifically, Applicants exemplify embodiments combining hits from the LOF and GOF screens to assess enhanced drug resistance. To do so, as summarized above, Applicants first transduced and selected A375 cells with two lentiviral constructs encoding active Cas9 and the MPH complex, respectively (Fig. 9a). Applicants then transduced these cells with dRNA targeting LPAR5 for activation and / or sgRNAs targeting MED12 or TADA2B for gene knockout. LPAR5 mRNA expression increased over 600-fold when cells were treated with dRNA targeting LPAR5, even when combined with sgRNAs targeting other genes. In all conditions, no significant LPAR5 indels were detected (Fig. 9b). By contrast, the loci targeted by MED12 and TADA2B showed robust indel formation even in the orthogonal conditions (Fig. 9c). Additionally, the activation and knockout perturbations – both individually and in combination - resulted in shifts of the A375 survival curves and increased resistance to PLX-4720 (Fig. 9d). Interestingly, the orthogonal conditions (LPAR5/MED12 and LPAR5/TADA2B) showed additional increases in resistance beyond the individual perturbations, as measured by the IC50 value (Fig. 9e).

[00694] *Example 10: Activation of HBG1 with different lengths of sgRNAs using Cas9 mutants.*

[00695] The example illustrates that mutations in Cas9 can change the length requirements for dead guide RNAs. To show this effect, HEK293 cells were transfected with alternative sgRNAs of length 15bp, 17bp, and 20bp, all targeting the upstream region of HBG1, each in combination with different Cas9 mutants. As illustrated in Figure 10, double Cas9 mutants (DM R780A/K810A and DMR780A/K855A) all acted as dead Cas9s, with all three lengths of guide RNA being capable of activating expression. In contrast, single Cas9 mutants SM K810A, SM848 and SM K855A acted in a manner analogous to a wildtype Cas9, with only the 15bp guide showing activation of expression. Of note, the SMR780A Cas9 had the length requirement for a dead guide RNA shifted, such that 17bp sgRNA could mediate activation of expression.

[00696] An example of a 15bp deadRNA with tetraloop and stem loop 2 MS2 insertions is as follows:

```

NNNNNNNNNNNNNNNGTTTTAGAGCTAGGCCAACATGAGGATCACCCATGTCTGCA
GGGCCTAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTTGGCCAACATGAGGAT
CACCCATGTCTGCAGGGCCAAGTGGCACCGAGTCGGTGCTTTTT

```

[00697] *Example 11: Using dead guides for efficient activation and bimodal control in Cas9 transgenic mice*

[00698] Dead guides can be delivered to transgenic mice to enable efficient activation without the need to deliver dCas9. For example, deadSgRNA(MS2) + MS2-P65-HSF1 can be packaged together into a single AAV vector. In alternative embodiments, both a dead sgRNA + an active sgRNA can be delivered at the same time (in a single vector) to enable bimodal control in vivo. This strategy can be utilized for a wide variety of Cas9 transgenic species. Different delivery methods (e.g. lentivirus etc.) may be used in alternative embodiments.

[00699] An example sequence of a pAAV-U6-sgRNA(MS2)-syn-MS2-P65-HSF1_2A_GFP (sequence is including Itrs and the sequence between Itrs) for transcriptional modulation in neurons is listed below:

```

cctgcaggcagctgcgcgctcgctcgctcactgaggccgcccgggctcgggacgaccttggtcgccggcctcagtga
gcgagcgagcgcgcagagagggagtgcccaactccatcactaggggtcctgcggccgcacgcgtgagggcctattccatgattcctt
catattgcatatacatacaaggctgtagagagataattggaattaattgactgtaaacacaaagatattagtaaaaatacgtgacgtaga
aagtaataattcttggtagtggcagttttaaattatgttttaaattggactatcatatgcttaccgtaactgaaagtatttcgatttcttggcttta
tatatcttGTGGAAAGGACGAAACACCggagaccactgtagtctctgttttagagctaggccAACATGAGGAT
CACCCATGTCTGCAGggcctagcaaggttaaataaggctagtcggttatcaacttgcccAACATGAGGATCACC
CATGTCTGCAGggccaagtggcaccgagtcggtgcTTTTTTTgtgtctagactgcagagggccctgcgtatgagtgcaag
tgggttttaggaccaggatgagggcggggtgggggtgacctgacgaccgaccccgaccactggacaagcacccaacccccattccc
caaattgcgcatcccctatcagagagggggaggggaaacaggatgcccgcaggcgcgtgcgcaactgccagcttcagcaccgcgga
gtgccttcgcccccgctggcggcgcgcgccaccgcccctcagcactgaaggcgcgctgacgtcactcgccgggtcccccgcaaactc
ccctcccggccaccttggctcgctcgcgccgcccggcccagccggaccgcaccacgcgagggcgcgagataggggggacggg
cgcgaccatctgcgctgcggcgcggcgactcagcgcgtgctcagctcgcggtgggcagcggaggagtcgtgtcgtgctgagagcgc
agtcgagaaggatcggccaccATGGCTTCAAACCTTTACTCAGTTCGTGCTCGTGGACAATGGTG
GGACAGGGGATGTGACAGTGGCTCCTTCTAATTTTCGCTAATGGGGTGGCAGAGTGG
ATCAGCTCCAACTCACGGAGCCAGGCCTACAAGGTGACATGCAGCGTCAGGCAGTC
TAGTGCCAGAAgAGAAAGTATAACCATCAAGGTGGAGGTCCCCAAAGTGGCTACCC
AGACAGTGGGCGGAGTCGAACTGCCTGTCGCCGCTTGGAGGTCTACCTGAACATG
GAGCTCACTATCCCAATTTTCGCTACCAATTCTGACTGTGAACTCATCGTGAAGGCA
ATGCAGGGGCTCCTCAAAGACGGTAATCCTATCCCTTCCGCCATCGCCGCTAACTCA
GGTATCTACagcgcgtGGAGGAGGTGGAAGCGGAGGAGGAGGAAGCGGAGGAGGAGGT

```

AGCggacctaagaaaaagaggaaggtggcggccgctggatccCCTTCAGGGCAGATCAGCAACCAGGCCC
TGGCTCTGGCCCCTAGCTCCGCTCCAGTGCTGGCCCAGACTATGGTGCCCTCTAGTG
CTATGGTGCCTCTGGCCCAGCCACCTGCTCCAGCCCCTGTGCTGACCCCAGGACCAC
CCCAGTCACTGAGCGCTCCAGTGCCCAAGTCTACACAGGCCGGCGAGGGGACTCTG
AGTGAAGCTCTGCTGCACCTGCAGTTCGACGCTGATGAGGACCTGGGAGCTCTGCTG
GGGAACAGCACCCGATCCCGGAGTGTTACAGATCTGGCCTCCGTGGACAACCTCTGA
GTTTCAGCAGCTGCTGAATCAGGGCGTGTCCATGTCTCATAAGTACAGCCGAACCAAT
GCTGATGGAGTACCCCGAAGCCATTACCCGGCTGGTGACCCGGCAGCCAGCGGCCCC
CCGACCCCGCTCCAACCTCCCCTGGGAACCAGCGGCCTGCCTAATGGGCTGTCCGGAG
ATGAAGACTTCTCAAGCATCGCTGATATGGACTTTAGTGCCCTGCTGTACAGATTT
CCTCTAGTGGGCAGGGAGGAGGTGGAAGCGGCTTCAGCGTGGACACCAGTGCCCTG
CTGGACCTGTTTCAGCCCCTCGGTGACCGTGCCCGACATGAGCCTGCCTGACCTTGAC
AGCAGCCTGGCCAGTATCCAAGAGCTCCTGTCTCCCCAGGAGCCCCCAGGCCTCCC
GAGGCAGAGAACAGCAGCCCGGATTCAGGGAAGCAGCTGGTGCCTACACAGCGC
AGCCGCTGTTCCCTGCTGGACCCCGGCTCCGTGGACACCCGGGAGCAACGACCTGCCG
GTGCTGTTTGAGCTGGGAGAGGGCTCCTACTTCTCCGAAGGGGACGGCTTCGCCGAG
GACCCACCATCTCCCTGCTGACAGGCTCGGAGCCTCCCAAAGCCAAGGACCCCACT
GTCTCCgctagcGGCAGTGGAGAGGGCAGAGGAAGTCTGCTAACATGCGGTGACGTCG
AGGAGAATCCTGGCCCAGTGAGCAAGGGCGAGGAGCTGTTACCCGGGGTGGTGCCC
ATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGA
GGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCG
GCAAGCTGCCCCGTGCCCTGGCCCACCCTCGTGACCACCCTGACCTACGGCGTGCAGT
GCTTCAGCCGCTACCCCGACCACATGAAGCAGCACGACTTCTTCAAGTCCGCCATGC
CCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAG
ACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAA
GGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACAACCT
ACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAACGGCATCAAGGTG
AACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGACCACTA
CCAGCAGAACACCCCATCGGGCAGCGGCCCCCGTGCTGCTGCCCGACAACCACTACC
TGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTC
CTGCTGGAGTTCGTGACCGCCGCGGGATCACTCTCGGCATGGACGAGCTGTACAA

GTAAGaattcgatatcaagcttatcgataatcaacctctggattacaaaattgtgaaagattgactggattcttaactatggtgctccttttac
gctatgtggatacgtgctttaatgcctttgtatcatgctattgcttcccgataggctttcattttctcctcctgtataaatcctgggtgctgtcttta
tgaggagttgtggcccggtgtaggcaacgtggcgtggtgtgcaactgtgtttgctgacgcaacccccactggttggggcattgccaccacct
gtcagctcctttccgggactttcgctttccccctcctattgccacggcggaactcatcgccgcctgcttggcccgctgctggacaggggctc
ggctgttgggcactgacaattccgtggtgtgtcggggaaatcatcgtcctttcctggctgctcgctAtgttgccacctggattctgcgcgg
gacgtccttctgctacgtcccttcggccctcaatccageggaccttccctcccgcggcctgctgceggctctgceggcctcttcgcgcttctgc
cttcgccctcagacgagtcggatctccctttggggccgctccccgcacgataccgagcgtgctcgagCTAGAGCTCGCTGA
TCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGCCCCCTCCCCCGTGC
CTTCCTTGACCCTGGAAGGTGCCACTCCCCTGTCCTTTTCTAATAAAAATGAGGAAA
TTGCATCGCATTGTCTGAGTAGGTGTCATTCTATTCTGGGGGGTGGGGTGGGGCAGG
ACAGCAAGGGGGAGGATTGGGAAGACAATAGCAGGCATGCTGGGGAggtaaccacgtgcgg
accgagcggccgcaggaacccttagtgatggagttggccactccctctctgcgcgctcgcctcactgagggccggcgaccaaaggt
cgccccagcggccggctttgcccggggcggcctcagtgagcgcgagcgcgcagctgcctgcagg.

[00700] *Example 15 – Enhanced Cas9 Mutants Have High Activity and Specificity*

[00701] Applicants generated SpCas9 mutants consisting of individual alanine substitutions at 29 positively-charged residues within the nt-groove and assessed changes to genome editing specificity. Point mutants were tested for specificity by targeting them to the *EMXI(1)* target site in human embryonic kidney (HEK) cells using a previously validated guide sequence; indel formation was assessed at the on-target site and a known genomic off-target (OT) site. Six of the 29 point mutants reduced off-target activity by at least 10-fold compared to wild-type (WT) SpCas9 while maintaining on-target cleavage efficiency, and 6 others improved specificity 2 to 5-fold. These mutants also exhibited improved specificity when tested on a second locus, *VEGFA(1)* (Fig. 12D). Although some point mutants were more specific than WT SpCas9 when targeting *EMXI(1)* and *VEGFA(1)*, off-target indels were still detectable (~0.1%) (Fig. 12D). To further improve specificity, Applicants performed combinatorial mutagenesis using the top point mutants identified in the initial screen. Eight out of 35 combination mutants retained wild-type on-target activity and displayed undetectable off-target indel levels at *EMXI(1)* OT1, *VEGFA(1)* OT1, and *VEGFA(2)* OT2 (Fig. 12E.) To ensure that the observed increased in specificity was not due to reduced on-target activity, Applicants measured on-target indel formation at 10 target loci using the top 16 mutants (Fig. 12F), as determined by a combination of on- and off-target activity. Applicants observed high efficiency and specificity for three mutants: SpCas9

(K855A), SpCas9 (K810A/K1003A/R1060A) (also referred to as eSpCas9(1.0)), and SpCas9 (K848A/K1003A/R1060A) (also referred to as eSpCas9(1.1)). These three variants were selected for further analysis.

[00702] To assess whether SpCas9 (K855A), eSpCas9(1.0), and eSpCas9(1.1) broadly retained efficient nuclease activity, Applicants measured on-target indel generation at 24 target sites spanning 10 different genomic loci (Fig. 13A). All three mutants generated similar indel levels as WT SpCas9 with the majority of target sites (Fig. 13B). To test whether improvements in specificity could be attributed to decreased Cas9 expression, Applicants performed a Western blot for SpCas9 and found that all three mutants were expressed equivalently or at higher levels than WT SpCas9 (Fig. 13C). This demonstrated that improvements in specificity were not due to decreased protein expression levels.

[00703] Applicants then compared the specificity of the three mutants to WT SpCas9 with truncated guide sequences (18nt for *EMXI*(1) and 17nt for *VEGFA*(1)), which have been shown to reduce off-target indel formation. All three mutants reduced cleavage at all off-target sites assessed. Moreover, eSpCas9(1.0) and eSpCas9(1.1) eliminated 20 of 24 of these sites. In contrast, WT SpCas9 with truncated guides eliminated 14 of 24 sites but also increased off-target activity at 5 sites compared to WT SpCas9 with full-length guides.

[00704] To assess tolerance of SpCas9 (K855A), eCas9(1.0), and eCas9(1.1) for mismatched target sites, Applicants systematically mutated the *VEGFA*(1) guide sequence to introduce single and double base mismatches at different positions (Fig. 14A-C). Compared to WT SpCas9, all three mutants induced lower levels of indels with mismatched guides. Of note, eSpCas9(1.0) and eSpCas9(1.1) induced lower indel levels even with single base mismatches located outside of the 7-12bp seed sequence. Given that Applicants did not observe any difference between eSpCas9(1.0) and eSpCas9(1.1) in terms of specificity, SpCas9 (K855A) and eSpCas9(1.1) were selected for further analysis based on on-target efficiency.

[00705] Genome-wide editing specificity of SpCas9 (K855A) and eSpCas9(1.1) was assessed using BLESS (direct *in situ* breaks labelling, enrichment on streptavidin and next-generation sequencing, which quantifies DNA double-stranded breaks (DSBs) across the genome (Fig. 14A). Applicants assayed the *EMXI*(1) and *VEGFA*(1) targets for both mutants and compared these results to WT SpCas9. (Fig. 14B). Both SpCas9(K855A) and eSpCas9(1.1) exhibited a

genome-wide reduction in off-target cleavage and did not generate any new off-target sites (Fig. 14C-D).

Algorithms

[00706] Algorithms have been developed to predict off-target indels and rationally improve sgRNA activity for Cas9 nuclease. To develop a similar algorithm for predicting Cas9 activator specificity, Applicants used guides with mismatches on the 5' end of the sgRNA analogous to the truncation experiments (Figure 15). In accordance with the results from truncated guides, Applicants observed that guides with only 15bp complementarity to the target DNA were still able to mediate efficient activation in all four cases. Given the results demonstrating differences between mismatch tolerance for Cas9 transcriptional control and Cas9 nuclease activity, Applicants provide a novel algorithm specific to Cas9-based activators. To create design rules for Cas9 activators, Applicants performed whole transcriptome analysis on ten additional sgRNAs targeting the proximal promoter of human HBG1/2 (Fig. 16a).

[00707] Based on the data from Figure 15 Applicants calculated a new activator off-target score that evaluates off-target matches of the first 15nt of the sgRNA only within a 2kb window of all refseq gene promoters. This activator off-target score was significantly correlated with the number of genome-wide off-targets for the set of guides as detected by RNAseq ($R = -0.6$, $p < 0.05$) (Fig. 16b). A second variable correlating with the detected specificity of an sgRNA was its GC content, which is known to affect Watson Crick binding energy to the DNA target. Specificity was greater for guides with lower GC content ($R = 0.6$, $p < 0.05$) (Fig. 3b). Overall, four out of 12 guides exhibited very high specificity (<3 significant genome-wide off-targets). The results illustrate that sgRNAs can be designed to minimize non-specific upregulation by minimizing GC content and avoiding off-target matches of the first 15nt in gene promoters. To optimize the selection of activator sgRNAs with high specificity, Applicants performed linear regression on the dataset. The combined model using both the new activator off-target score and GC content had a correlation of $R = 0.65$ with the number of off-target hits (Fig. 16c).

* * *

[00708] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to

those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

WHAT IS CLAIMED:

1. A non-naturally occurring or engineered composition comprising a CRISPR-Cas system, said system comprising a functional CRISPR Cas9 enzyme and single guide RNA (sgRNA);

wherein the sgRNA comprises a dead guide sequence;

whereby the sgRNA is capable of hybridizing to a target sequence;

whereby the CRISPR-Cas system is directed to the target sequence without detectable indel activity resultant from nuclease activity of a non-mutant Cas9 enzyme of the system as detected by a SURVEYOR assay.

2. The non-naturally occurring or engineered composition comprising the guide RNA (sgRNA) of claim 1, wherein the sgRNA is specific to Sp Cas9 and: the dead guide is 10 - 16 nucleotides in length, optionally 12-15 nucleotides in length; or, the dead guide comprises matching and mismatching sequences compared to the target sequence, and the contiguous matching sequences are 10 – 16 nucleotides in length, optionally 12 – 15 nucleotides in length.

3. The non-naturally occurring or engineered composition comprising a guide RNA (sgRNA) of claim 1, wherein the sgRNA is specific to Sp Cas9 and: the dead guide is 13 nucleotides in length; or, the dead guide comprises matching and mismatching sequences compared to the target sequence, and the contiguous matching sequences are 13 nucleotides in length.

4. The non-naturally occurring or engineered composition comprising a guide RNA (sgRNA) of claim 1, wherein the sgRNA is specific to Sa Cas9 and: the dead guide is 15 - 19 nucleotides in length, optionally 17-18 nucleotides in length; or, the dead guide comprises matching and mismatching sequences compared to the target sequence, and the contiguous matching sequences are 15 – 19 nucleotides in length, optionally 17 – 18 nucleotides in length.

5. The non-naturally occurring or engineered composition comprising a guide RNA (sgRNA) of claim 1, wherein the sgRNA is specific to Sa Cas9 and the dead guide is 17 nucleotides in length.

6. A non-naturally occurring or engineered CRISPR-Cas9 complex composition comprising the dead sgRNA of any one of claims 1-5 and a Cas9, wherein optionally the Cas9 comprises at least one mutation, and optionally one or more nuclear localization sequences.

7. The sgRNA of any one of claims 1-5 or the CRISPR-Cas9 complex of claim 6 including a non-naturally occurring or engineered composition comprising two or more adaptor proteins, wherein each protein is associated with one or more functional domains and wherein the adaptor protein binds to the distinct RNA sequence(s) inserted into the at least one loop of the sgRNA.

8. A non-naturally occurring or engineered composition comprising a guide RNA (sgRNA) comprising a dead guide sequence capable of hybridizing to a target sequence in a genomic locus of interest in a cell, wherein the dead guide sequence is according to the dead guide sequence of any one of claims 1-5,

a Cas9 comprising at least one or more nuclear localization sequences, wherein the Cas9 optionally comprises at least one mutation

wherein at least one loop of the sgRNA is modified by the insertion of distinct RNA sequence(s) that bind to one or more adaptor proteins, and wherein the adaptor protein is associated with one or more functional domains; or, wherein the sgRNA is modified to have at least one non-coding functional loop,

and wherein the composition comprises two or more adaptor proteins, wherein each protein is associated with one or more functional domains.

9. The composition of any one of claims 1-8, wherein the Cas9 comprises at least one mutation and has nuclease activity of at least 97%, or 100% as compared with the Cas9 not having the at least one mutation.

10. The composition of any one of claims 1-9, wherein the Cas9 comprises two or more mutations and has nuclease activity of at least 97%, or 100% as compared with the Cas9 not having the at least one mutation.

11. The composition of claim 10 wherein the Cas9 comprises three or more mutations and has nuclease activity of at least 97%, or 100% as compared with the Cas9 not having the at least one mutation.

12. The composition of any one of claims 1-9, wherein the Cas9 is an ortholog of SpCas9 protein.

13. The composition of any one of claims 1-12, wherein the Cas9 is associated with one or more functional domains.

14. The composition of claim 13, wherein the one or more functional domains associated with the adaptor protein is a heterologous functional domain.

15. The composition of claim 13, wherein the one or more functional domains associated with the Cas9 is a heterologous functional domain.

16. The composition of any one of claims 1-15, wherein the adaptor protein is a fusion protein comprising the functional domain, the fusion protein optionally comprising a linker between the adaptor protein and the functional domain, the linker optionally including a GlySer linker.

17. The composition of any one of claims 7-16, wherein the at least one loop of the sgRNA is not modified by the insertion of distinct RNA sequence(s) that bind to the two or more adaptor proteins.

18. The composition of any one of claims 7-17, wherein the one or more functional domains associated with the adaptor protein is a transcriptional activation domain.

19. The composition of any one of claims 13-18, wherein the one or more functional domains associated with the Cas9 is a transcriptional activation domain.

20. The composition of any one of claims 7-19, wherein the one or more functional domains associated with the adaptor protein is a transcriptional activation domain comprising VP64, p65, MyoD1, HSF1, RTA or SET7/9.

21. The composition of any one of claims 13-20, wherein the one or more functional domains associated with the Cas9 is a transcriptional activation domain comprises VP64, p65, MyoD1, HSF1, RTA or SET7/9.

22. The composition of any one of claims 7-17, wherein the one or more functional domains associated with the adaptor protein is a transcriptional repressor domain.

23. The composition of any one of claims 13-18, wherein the one or more functional domains associated with the Cas9 is a transcriptional repressor domain.

24. The composition of claim 22 or 23, wherein the transcriptional repressor domain is a KRAB domain.

25. The composition of claim 22 or 23, wherein the transcriptional repressor domain is a NuE domain, NcoR domain, SID domain or a SID4X domain.

26. The composition of any one of claims 7-17, wherein at least one of the one or more functional domains associated with the adaptor protein have one or more activities

comprising methylase activity, demethylase activity, transcription activation activity, transcription repression activity, transcription release factor activity, histone modification activity, DNA integration activity RNA cleavage activity, DNA cleavage activity or nucleic acid binding activity.

27. The composition of any one of claims 13-17, wherein the one or more functional domains associated with the Cas9 have one or more activities comprising methylase activity, demethylase activity, transcription activation activity, transcription repression activity, transcription release factor activity, histone modification activity, DNA integration activity RNA cleavage activity, DNA cleavage activity, nucleic acid binding activity, or molecular switch activity or chemical inducibility or light inducibility.

28. The composition of any one of claims 26-27, wherein the DNA cleavage activity comprises FokI nuclease activity.

29. The composition of any one of claims 7-28, wherein the one or more functional domains is attached to the Cas9 so that upon binding to the sgRNA and target the functional domain is in a spatial orientation allowing for the functional domain to function in its attributed function; or, optionally,

wherein the one or more functional domains is attached to the Cas9 via a linker, optionally a GlySer linker.

30. The composition of any one of claims 7-29, wherein the sgRNA is modified so that, after sgRNA binds the adaptor protein and further binds to the Cas9 and target, the functional domain is in a spatial orientation allowing for the functional domain to function in its attributed function.

31. The composition of any one of claims 13-29, wherein the one or more functional domains associated with the Cas9 is attached to the Rec1 domain, the Rec2 domain, the HNH domain, or the PI domain of the SpCas9 protein or any ortholog corresponding to these domains.

32. The composition of any one of claims 13-31, wherein the one or more functional domains associated with the Cas9 is attached to the Rec1 domain at position 553, Rec1 domain at 575, the Rec2 domain at any position of 175-306 or replacement thereof, the HNH domain at any position of 715-901 or replacement thereof, or the PI domain at position 1153 of the SpCas9 protein or any ortholog corresponding to these domains.

33. The composition of any one of claims 13-31, wherein the one or more functional domains associated with the Cas9 is attached to the Rec1 domain or the Rec2 domain, of the SpCas9 protein or any ortholog corresponding to these domains.

34. The composition of any one of claims 13-33, wherein the one or more functional domains associated with the Cas9 is attached to the Rec2 domain of the SpCas9 protein or any ortholog corresponding to this domain.

35. The composition of any one of claims 7-34, wherein the at least one loop of the sgRNA comprises a tetraloop and/or loop2.

36. The composition of any one of claims 7-35, wherein the tetraloop and loop 2 of the sgRNA are modified by the insertion of the distinct RNA sequence(s).

37. The composition of any one of claims 35 or 36, wherein the insertion of distinct RNA sequence(s) that bind to one or more adaptor proteins comprises an aptamer sequence.

38. The composition of claim 37, wherein the aptamer sequence comprises two or more aptamer sequences specific to the same adaptor protein.

39. The composition of claim 37, wherein the aptamer sequence comprises two or more aptamer sequences specific to different adaptor proteins.

40. The composition of any one of the claims above, wherein the adaptor protein comprises MS2, PP7, Q β , F2, GA, fr, JP501, M12, R17, BZ13, JP34, JP500, KU1, M11, MX1, TW18, VK, SP, FL, ID2, NL95, TW19, AP205, ϕ Cb5, ϕ Cb8r, ϕ Cb12r, ϕ Cb23r, 7s, or PRR1.

41. A cell comprising the non-naturally occurring or engineered composition of any one of the preceding claims.

42. The cell of claim 41, wherein the cell is a eukaryotic cell.

43. The cell of claim 42, wherein the eukaryotic cell is a mammalian cell, optionally a mouse cell.

44. The cell of claim 43, wherein the mammalian cell is a human cell.

45. The cell or composition of any one of the preceding claims, wherein a first adaptor protein is associated with a p65 domain and a second adaptor protein is associated with a HSF1 domain.

46. The cell or composition of any one of the preceding claims, wherein the composition comprises a Cas9 complex having at least three functional domains, at least one of which is associated with the Cas9 and at least two of which are associated with sgRNA.

47. The cell or composition of any one of claims 1-46, further comprising a second sgRNA, wherein the second sgRNA comprises a live sgRNA capable of hybridizing to a second target sequence such that a second Cas9 system is directed to a second genomic locus of interest in a cell with detectable indel activity at the second genomic locus resultant from nuclease activity of the Cas9 enzyme of the system.

48. The cell or composition of claim 47, further comprising a plurality of dead sgRNAs, and/or a plurality of live sgRNAs.

49. A method for introducing a genomic locus event comprising the administration to a host or expression in a host in vivo of one or more of the compositions from claims 1-48.

50. The method according to claim 49, wherein the genomic locus event comprises affecting gene activation, gene inhibition, or cleavage in the locus.

51. The method according to claims 49 or 50, wherein the host is a eukaryotic cell.

52. The method according to claim 51, wherein the host is a mammalian cell, optionally a mouse cell.

53. The method according to claims 49 or 50, wherein the host is a non-human eukaryote.

54. The method according to claim 53, wherein the non-human eukaryote is a non-human mammal.

55. The method according to claim 54, wherein the non-human mammal is a mouse.

56. A method of modifying a genomic locus of interest to change gene expression in a cell by introducing or expressing in a cell the composition of any of the preceding claims.

57. The method according to any one of claims 49-56 comprising the delivery of the composition or nucleic acid molecule(s) coding therefor, wherein said nucleic acid molecule(s) are operatively linked to regulatory sequence(s) and expressed in vivo.

58. The method according to claim 56 wherein the expression in vivo is via a lentivirus, an adenovirus, or an AAV.

59. A mammalian cell line derived from the cells as defined in claim 43, 51 or 52, wherein the cell line is, optionally, a human cell line or a mouse cell line.

60. A transgenic mammalian model, optionally a mouse, wherein the model has been transformed with the composition of any one of claims 1-40, or is a progeny of said transformant.

61. A nucleic acid molecule(s) encoding the sgRNA or the Cas9 complex or the composition of any of the preceding claims.

62. A vector system comprising: a nucleic acid molecule encoding the dead guide RNA (sgRNA) as defined in any one of claims 1 - 48.

63. The vector system of claim 62, further comprising a nucleic acid molecule(s) encoding the Cas9 as defined in any one of claims 1 - 48.

64. The vector system of claim 62 or 63, further comprising a nucleic acid molecule(s) encoding the live sgRNA of claim 47 or 48.

65. The nucleic acid molecule of claim 61 or the vector of claim 62 or 63, further comprising regulatory element(s) operable in a eukaryotic cell operably linked to the nucleic acid molecule encoding the guide sequence (sgRNA) and/or the nucleic acid molecule encoding the Cas9 and/or the optional nuclear localization sequence(s).

66. A method of screening for gain of function (GOF) or loss of function (LOF) comprising the cell line of claim 59 or cells of the model or progeny of claim 60 containing or expressing Cas9 and introducing the composition of any one of claim 1 – 48 into cells of the cell line or model, whereby the dead sgRNA includes either an activator or a repressor, and monitoring for GOF or LOF respectively as to those cells as to which the introduced dead sgRNA includes an activator or as to those cells as to which the introduced dead sgRNA includes a repressor.

67. The composition of any preceding claim wherein the Cas9 includes one or more functional domains.

68. The composition of any preceding claim wherein there is more than one dead sgRNA, and the dead sgRNAs target different sequences whereby when the composition is employed, there is multiplexing.

69. The composition of claim 68 wherein there is more than one dead sgRNA modified by the insertion of distinct RNA sequence(s) that bind to one or more adaptor proteins.

70. The composition of claim 66 or 67 wherein one or more adaptor proteins associated with one or more functional domains is present and bound to the distinct RNA sequence(s) inserted into the at least one loop of the sgRNA.

71. The composition of any preceding claim, wherein the target sequence(s) are non-coding or regulatory sequences.

72. The composition of claim 71, wherein the regulatory sequences are promoter, enhancer or silencer sequence(s).

73. The composition of any preceding claim wherein the sgRNA is modified to have at least one non-coding functional loop.

74. The composition of claim 73 wherein the at least one non-coding functional non-coding loop is repressive.

75. The composition of claim 74 wherein at least one non-coding functional non-coding loop comprises Alu.

76. A method of selecting a guide RNA targeting sequence for directing a functionalized CRISPR-Cas9 system to a gene locus in an organism, which comprises:

- a) locating one or more CRISPR motifs in the gene locus;
- b) analyzing the 20 nt sequence upstream of each CRISPR motif by:
 - i) determining the GC content of the sequence; and
 - ii) determining whether there are off-target matches of the first 15 nt of the sequence in the genome of the organism;
- c) selecting the sequence for use in a guide RNA if the GC content of the sequence is 70% or less and no off-target matches are identified.

77. The method of claim 76, wherein the sequence is selected if the GC content is 50% or less.

78. The method of claim 76, wherein the sequence is selected if the GC content is 40% or less.

79. The method of claim 76, wherein the sequence is selected if the GC content is 30% or less.

80. The method of claim 76, wherein two or more sequences are analyzed and the sequence having the lowest GC content is selected.

81. The method of claim 76, wherein off-target matches are determined in regulatory sequences of the organism.

82. The method of claim 76, wherein the gene locus is a regulatory region.

83. The method of claim 76, wherein the CRISPR motif is recognized by a SpCas9 enzyme.

84. The method of claim 76, wherein the organism is a eukaryotic organism.

85. The method of claim 84, wherein the eukaryotic organism is a human, a mouse, or a rat.

86. A guide RNA comprising the targeting sequence selected according to any one of claims 76 to 85.

87. A method of altering expression of at least one gene product comprising introducing into a cell an engineered CRISPR-Cas9 system comprising a guide RNA comprising a targeting sequence selected according to any one of claims 76 to 85.

88. A method of altering expression of at least two gene products comprising introducing into a cell an engineered CRISPR-Cas9 system comprising two or more guide RNAs comprising a targeting sequence selected according to any one of claims 76 to 85.

89. A cell comprising the CRISPR-Cas9 system of claim 88, wherein the expression of one or more gene products has been altered.

90. The cell of claim 89, wherein the expression of two or more gene products has been altered.

91. A cell line of the cell according to any one of claims 89 or 90.

92. A multicellular organism comprising one or more cells according to any one of claims 89 or 90.

93. A gene product from the cell of claim 89 or 90, from the cell line of claim 91, or from the multicellular organism of claim 92.

94. The gene product of claim 93, wherein the amount of gene product expressed is greater than or less than the amount of gene product expressed from a cell, cell line or a multicellular organism that does not have altered expression.

95. A guide RNA for directing a functionalized CRISPR-Cas9 system to a gene locus in an organism which comprises a targeting sequence, wherein the CG content of the target sequence is 70% or less, and the first 15 nt of the targeting sequence does not match an off-target sequence upstream from a CRISPR motif in the regulatory sequence of another gene locus in the organism.

96. A method of selecting a guide RNA targeting sequence for directing a functionalized CRISPR-Cas enzyme to a gene locus in an organism, which comprises:

- a) locating one or more CRISPR motifs in the gene locus;
- b) analyzing the sequence upstream of each CRISPR motif by:

- i) selecting 10 to 15 nt adjacent to the CRISPR motif
- ii) determining the GC content of the sequence; and
- c) selecting the 10 to 15 nt sequence as a targeting sequence for use in a guide RNA if the GC content of the sequence is 40% or more.

97. The method of claim 96, wherein the sequence is selected if the GC content is 50% or more.

98. The method of claim 96, wherein the sequence is selected if the GC content is 60% or more.

99. The method of claim 96, wherein the sequence is selected if the GC content is 70% or more.

100. The method of claim 96, wherein two or more sequences are analyzed and the sequence having the highest GC content is selected.

101. The method of any one of claims 96 to 100, which further comprises adding nucleotides to the 5' end of the selected sequence which do not match the sequence upstream of the CRISPR motif.

102. The method of claim 96, wherein the organism is a eukaryotic organism.

103. The method of claim 102, wherein the eukaryotic organism is a human, a mouse, or a rat.

104. A guide RNA comprising the targeting sequence selected according to any one of claims 96 to 101, which.

105. A method of altering expression of at least one gene product comprising introducing into a cell an engineered CRISPR-Cas9 system comprising a guide RNA comprising a targeting sequence selected according to any one of claims claims 96 to 101.

106. A method of altering expression of at least two gene products comprising introducing into a cell an engineered CRISPR-Cas9 system comprising a guide RNAs comprising a targeting sequence selected according to any one of claims claims 96 to 101.

107. The method of claim 106, wherein at each of the at least two gene loci are independently regulated by an activator or inhibitor associated with the CRISPR-Cas9 system.

108. The method of claim 106, wherein at least one gene locus is regulated by an activator or inhibitor associated with the CRISPR-Cas9 system, and the second gene locus is cleaved.

109. A cell comprising the CRISPR-Cas9 system of any one of claims 106 to 108, wherein the expression of one or more gene products has been altered.

110. The cell of claim 109, wherein the expression of two or more gene products has been altered.

111. A cell line of the cell according to any one of claims 109 or 110.

112. A multicellular organism comprising one or more cells according to any one of claims 109 or 110.

113. A gene product from the cell of claim 109 or 110, from the cell line of claim 111, or from the multicellular organism of claim 112.

114. The gene product of claim 113, wherein the amount of gene product expressed is greater than or less than the amount of gene product expressed from a cell, cell line or a multicellular organism that does not have altered expression.

115. A guide RNA for directing a functionalized CRISPR-Cas9 system to a gene locus in an organism wherein the targeting sequence of the guide RNA consists of 10 to 15 nucleotides adjacent to the CRISPR motif of the gene locus, wherein the CG content of the target sequence is 50% or more.

116. The guide RNA of claim 115, which further comprises nucleotides added to the 5' end of the targeting sequence which do not match the sequence upstream of the CRISPR motif of the gene locus.

IL1B-20	AAAAACAGCGAGGGAGAAAC
IL1B-15	-----CAGCGAGGGAGAAAC
IL1B-14	-----AGCGAGGGAGAAAC
IL1B-13	-----GCGAGGGAGAAAC
IL1B-12	-----CGAGGGAGAAAC
IL1B-11	-----GAGGGAGAAAC

IL1B-20	AAAAACAGCGAGGGAGAAAC
---------	----------------------

Fig. 1





Fig. 2




Fig. 3A




Fig. 3C




Fig. 3D




FIG. 4E




FIG. 4F

	IL1B sgRNA Added	Cas9 Added	Activator Added
1	-----AGCGAGGGGAGAAAC + tracrRNA w/MS2 loop	Cas9+sgEMX1.3	MS2-p65-HSF1
2	-----GCGAGGGGAGAAAC + tracrRNA w/MS2 loop	Cas9+sgEMX1.3	MS2-p65-HSF1
3	AAAAACAGCGAGGGGAGAAAC + regular tracrRNA	Cas9+sgEMX1.3	None
4	GAAAAACAGCGAGGGGAGAAAC + tracrRNA w/ MS2 loop	dcas9	MS2-p65-HSF1
5	GFP Plasmid	None	None

Fig. 5A




Fig. 5B




Fig. 5C




Fig. 5D




Fig. 6A




Fig. 6B




Fig. 6C




Fig. 6D

FIG. 7A





FIG. 7B






Fig. 7C







Fig. 8A




Drug	IC50 (μM)
A375	0.142
MED12	0.367
LPAR5	2.255
MED12+LPAR5	12.560
P.I.	1.80




Drug	IC50 (μM)
A375	0.142
MED12	0.367
ITGA9	0.342
MED12+ITGA9	0.960
P.I.	1.54




Drug	IC50 (μM)
A375	0.142
MED12	0.367
EGFR	0.356
MED12+EGFR	0.804
P.I.	0.87



Drug	IC50 (μM)
A375	0.142
CUL3	0.246
LPAR5	2.897
CUL3+LPAR5	3.914
P.I.	0.78



Drug	IC50 (μM)
A375	0.142
CUL3	0.246
ITGA9	0.342
CUL3+ITGA9	0.272
P.I.	0.64



Drug	IC50 (μM)
A375	0.142
CUL3	0.246
EGFR	0.344
CUL3+EGFR	0.350
P.I.	0.53

Fig. 8B

FIG. 9A




FIG. 9B




FIG. 9C




FIG. 9D





FIG. 9E








Fig. 10

Mutation of positively charged residues binding to non-targeted gDNA strand improves specificity.

Cas9 mutant	Indel %		
	ON Target (EMX1)	OFF Target 1(OT25)	OFF Target 2(OT46)
WT	24.8	10.5	8.8
R780	22.9	0.0	0.1
K810	23.3	0.1	0.1
K848	24.3	0.1	0.1
K855	25.1	0.2	0.3
R976	15.6	0.1	0.1
H982	20.9	0.5	0.4
K1003	24.6	4.1	2.8
R1060	20.4	1.3	1.8
GFP	0.1	0.0	0.1
untrans.	0.1	0.0	0.1



ALA mutations that improve specificity are distributed along the non-targeting strand groove



Mechanism proposal:

- A) Nuclease activity is inactive until non-targeted DNA strand sterically triggers **HNH conformational change**
- B) Non-targeted strand binding to the **groove** between HNH and RuvC depends on RNA:DNA pairing
- C) Mutating DNA binding residues in the **groove** place more energetic demand on proper RNA:DNA pairing




Fig. 11

FIG. 12A




FIG. 12B




FIG. 12C




FIG. 12D




FIG. 12E




FIG. 12F




FIG. 13A

gene(guide #)	5'	target	3'	PAM	WT SpCas9(k855A) esfCas9 (1.0)	esfCas9 (1.1)
EMX1(1)		GAGTCCGAGCAGAAAGAA	GGG			
EMX1(2)		GTCACCTCCAATGACTAGGG	TGG			
EMX1(3)		GGGCAACCAACAACCCACGA	GGG			
EMX1(4)		GCTTGTCCTCTGTCAATGG	CGG			
EMX1(5)		GCGCCACCGTTGATGTGAT	GGG			
EMX1(6)		GACATCGATGCTCTCCCAT	TGG			
EMX1(7)		GCTCCCAAAAGCCTGGCCA	GGG			
EMX1(8)		GCCCCGGGCTTCAAGCCCTG	TGG			
EMX1(9)		GGCAGAGTCTGCTTGTCTG	TGG			
VEGFA(1)		GGTGTGAGTGTGTGTGGTG	TGG			
VEGFA(2)		GCTAAAGAGGGGAATGGCTT	TGG			
VEGFA(3)		GTTTGGGAGGTCAGAAAATAG	GGG			
VEGFA(4)		GTTGGAGCGGGGAGAGGCC	AGG			
VEGFA(5)		GGGTGGGGGAGTTTGTCTCC	TGG			
Chr8.48907789(1)		GTAAGTGTGAGTGTGTGTG	TGG			
Chr8.9115779(1)		GAAGAAATGACAGAACTCTG	AGG			
DNMT1(1)		GATTCTGGTCCAGAAACA	GGG			
TPCN2(1)		GTTGGGTGAGTGTGCTGTC	GGG			
TPCN2(2)		GAGGCTGGGGTGGAGGTGTT	GGG			
Chr14.106029015(1)		GGTGTGAGTGTGTGTGTG	AGG			
Chr14.106029024(1)		GTTGTGTGTGTGAGGGTGTAA	GGG			
NRG2(1)		GGGGAGTTTGTCTCTGGCAC	AGG			
CACNA2D4(1)		GGAGAGAGGCTCCCATCACG	GGG			
ADAMTSL1(1)		GAGAAAGAGAAAGTGGGGTGGG	GGG			

FIG. 13B




FIG. 13C




FIG. 14B




FIG. 14A





FIG. 14C






Fig. 15

FIG. 16A




FIG. 16B

Guide variable	Correlation with # off-targets		Guide variable	Specificity model linear regression	
	R value	p-value		coeff.	p-value
% GC	0.600	<0.05	% GC	233.2	0.05
activator OT score	-0.661	<0.05	activator OT score	-329.8	0.01
nuclease OT score	0.120	0.71	model overall	R = 0.65	

FIG. 16C

Figure 17A





Figure 17B



INTERNATIONAL SEARCH REPORT

International application No
PCT/US2015/065393

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12N15/11
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal, WPI Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>YANFANG FU ET AL.: "Improving CRISPR-Cas nuclease specificity using truncated guide RNAs", NATURE BIOTECHNOLOGY, vol. 32, no. 3, 26 January 2014 (2014-01-26), pages 279-284, XP055194360, ISSN: 1087-0156, DOI: 10.1038/nbt.2808</p> <p>the whole document -& YANFANG FU ET AL.: "Supplementary Results", NATURE BIOTECHNOLOGY, vol. 32, no. 3, 26 January 2014 (2014-01-26), pages 279-284, XP055259718, US ISSN: 1087-0156, DOI: 10.1038/nbt.2808 the whole document</p> <p style="text-align: center;">-/--</p>	<p>1-3,6, 9-12, 41-44, 46, 49-63, 65,67, 70-72, 104, 113-115</p>

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Date of the actual completion of the international search 6 April 2016	Date of mailing of the international search report 15/04/2016
-------------------------------------------------------------------------------	----------------------------------------------------------------------

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Macchia, Giovanni
----------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2015/065393

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	-& YANFANG FU ET AL.: "Supplementary Table 7", 26 January 2014 (2014-01-26), XP055259722, Retrieved from the Internet: URL:http://www.nature.com/nbt/journal/v32/n3/extref/nbt.2808-S2.xlsx [retrieved on 2016-03-18] the whole document	
X	----- WO 2014/093718 A1 (THE BROAD INSTITUTE, INC. [US]; MASSACHUSETTS INSTITUT TECHNOLOGY [US]) 19 June 2014 (2014-06-19)	86-95
Y	page 4 - page 5 page 7 - page 8 page 23 - page 24; figures 15-22 page 26; figure 32	13,15, 19,21, 23, 27-29,31
A	page 42, paragraph [00112] page 78 - page 86; examples 5, 6 page 89; figures 27, 29 claims page 49, paragraph [00101] - page 50	76-85, 96-103, 105-112, 116
Y	----- FU Y ET AL.: "Targeted genome editing in human cells using CRISPR/Cas nucleases and truncated guide RNAsc", 1 January 2014 (2014-01-01), THE USE OF CRISPR/CAS9, ZFNs, AND TALENS IN GENERATING SITE-SPECIFIC GENOME ALTERATIONS; [METHODS IN ENZYMOLOGY; ISSN 1557-7988; VOL. 546], ELSEVIER, NL, PAGE(S) 21 - 45, XP008179585, ISBN: 978-0-12-801185-0 [retrieved on 2014-11-12]	7,8, 13-31, 35-40
A	the whole document	1-3,6, 9-11
Y	----- PRASHANT MALI ET AL.: "CAS9 transcriptional activators for target specificity screening and paired nickases for cooperative genome engineering", NATURE BIOTECHNOLOGY, vol. 31, no. 9, 1 August 2013 (2013-08-01), pages 833-838, XP55149275, ISSN: 1087-0156, DOI: 10.1038/nbt.2675 the whole document	7,8,14, 16-18, 20,22, 24-26, 28,30, 35-40
Y	-& PRASHANT MALI ET AL.: "CAS9 transcriptional activators for target specificity screening and paired nickases for cooperative genome engineering", NATURE BIOTECHNOLOGY, vol. 31, no. 9, 1 August 2013 (2013-08-01), pages 833-838, XP055153963, ISSN: 1087-0156, DOI: 10.1038/nbt.2675 the whole document	7,8,14, 16-18, 20,22, 24-26, 28,30, 35-40
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2015/065393

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	<p>WO 2015/113063 A1 (GEORGIA TECH RESEARCH CORPORATION [US]) 30 July 2015 (2015-07-30)</p> <p>page 66, line 26 page 83; table 5 page 99; figures 12B, 12C page 101 - page 102; example 5 page 108, line 27 - page 109, line 12</p> <p>-----</p>	76-85, 96-103, 105-112, 116
T	<p>NISHIMASU HIROSHI ET AL.: "Crystal structure of Cas9 in complex with guide RNA and target DNA", CELL, CELL PRESS, US, vol. 156, no. 5, 13 February 2014 (2014-02-13), pages 935-949, XP028667665, ISSN: 0092-8674, DOI: 10.1016/J.CELL.2014.02.001 the whole document</p> <p>-----</p>	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2015/065393

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2014093718	A1	19-06-2014	
		DK 2784162 T3	13-07-2015
		DK 2896697 T3	07-12-2015
		EP 2784162 A1	01-10-2014
		EP 2840140 A1	25-02-2015
		EP 2848690 A1	18-03-2015
		EP 2896697 A1	22-07-2015
		EP 2921557 A1	23-09-2015
		EP 2932421 A1	21-10-2015
		EP 2940140 A1	04-11-2015
		US 2014186843 A1	03-07-2014
		WO 2014093718 A1	19-06-2014

WO 2015113063	A1	30-07-2015	NONE
