



(86) **Date de dépôt PCT/PCT Filing Date:** 2006/05/18
 (87) **Date publication PCT/PCT Publication Date:** 2006/11/23
 (45) **Date de délivrance/Issue Date:** 2015/10/20
 (85) **Entrée phase nationale/National Entry:** 2007/11/13
 (86) **N° demande PCT/PCT Application No.:** US 2006/019351
 (87) **N° publication PCT/PCT Publication No.:** 2006/125132
 (30) **Priorités/Priorities:** 2005/05/18 (US60/682,530);
 2005/07/01 (US60/696,160); 2006/02/13 (US60/773,300)

(51) **Cl.Int./Int.Cl. A61K 31/5383** (2006.01),
A61K 31/538 (2006.01), **A61K 9/12** (2006.01),
A61P 11/00 (2006.01), **A61P 31/04** (2006.01)

(72) **Inventeurs/Inventors:**
 SURBER, MARK W., US;
 BOSTIAN, KEITH A., US;
 LOMOVSKAYA, OLGA, US;
 GRIFFITH, DAVID C., US;
 DUDLEY, MICHAEL N., US


(73) **Propriétaire/Owner:**
 MPEX PHARMACEUTICALS, INC., US

(74) **Agent:** SMART & BIGGAR

(54) **Titre : FLUOROQUINOLONES EN AEROSOLS ET LEURS UTILISATIONS**
 (54) **Title: AEROSOLIZED FLUOROQUINOLONES AND USES THEREOF**

(57) **Abrégé/Abstract:**

Disclosed herein are formulations of fluoroquinolones suitable for aerosolization and use of such formulations for aerosol administration of fluoroquinolone antimicrobials for the treatment of pulmonary bacterial infections. In particular, inhaled levofloxacin specifically formulated and delivered for bacterial infections of the lungs is described. Methods include inhalation protocols and manufacturing procedures for production and use of the compositions described.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
23 November 2006 (23.11.2006)

PCT

(10) International Publication Number
WO 2006/125132 A3

(51) International Patent Classification:

A61K 31/538 (2006.01) A61K 9/12 (2006.01)
A61K 31/4375 (2006.01) A61P 11/00 (2006.01)
A61K 31/47 (2006.01)

(21) International Application Number:

PCT/US2006/019351

(22) International Filing Date: 18 May 2006 (18.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/682,530	18 May 2005 (18.05.2005)	US
60/696,160	1 July 2005 (01.07.2005)	US
60/773,300	13 February 2006 (13.02.2006)	US

(71) Applicant (for all designated States except US): MPEX PHARMACEUTICALS, INC. [US/US]; 3030 Bunker Hill Street, Suite 200, San Diego, California 92109 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SURBER, Mark, W. [US/US]; 4746 Edison Street, San Diego, California 92117 (US). BOSTIAN, Keith, A. [US/US]; 6 Callado Way, Atherton, California 94027 (US). LOMOVSKAYA, Olga [US/US]; 46 Laverne Avenue, Mill Valley, California 94941 (US). GRIFFITH, David, C. [US/US]; 714 Bandak Court, San Marcos, California 92069 (US). DUDLEY, Michael N. [US/US]; 17257 Silver Gum Way, San Diego, California 92127 (US).

(74) Agent: MALLON, Joseph J.; KNOBBE MARTENS OLSON & BEAR LLP, 2040 Main Street, 14th Floor, Irvine, California 92614 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

28 December 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: AEROSOLIZED FLUOROQUINOLONES AND USES THEREOF

(57) Abstract: Disclosed herein are formulations of fluoroquinolones suitable for aerosolization and use of such formulations for aerosol administration of fluoroquinolone antimicrobials for the treatment of pulmonary bacterial infections. In particular, inhaled levofloxacin specifically formulated and delivered for bacterial infections of the lungs is described. Methods include inhalation protocols and manufacturing procedures for production and use of the compositions described.



WO 2006/125132 A3

AEROSOLIZED FLUOROQUINOLONES AND USES THEREOF

BACKGROUND OF THE INVENTION

Description of the Related Art

[0002] Antibiotics have been effective tools in the treatment of infectious diseases during the last half-century. From the development of antimicrobial therapy to the late 1980s, most bacterial infections occurring in patients in developed countries could be controlled unless the infection occurred in an organ or environment where antibiotics were difficult to deliver or were ineffective, such as bacterial infections of the circulatory system in sepsis patients, or bacterial infections of the lungs in cystic fibrosis. However, even in ordinary infections, in response to the pressure of antimicrobial usage, multiple resistance mechanisms have become widespread and are threatening the clinical utility of even the most aggressive antibacterial therapy. The increase in antimicrobial resistant strains has been particularly common in major hospitals and care centers. The consequences of the increase in resistant strains include higher morbidity and mortality, longer patient hospitalization, and an increase in treatment costs.

[0003] Bacteria have developed several different mechanisms to overcome the action of antimicrobials. These mechanisms of resistance can be specific for a molecule or a family of antimicrobials, or can be non-specific and be involved in resistance to unrelated antimicrobials. Several mechanisms of resistance can exist in a single bacterial strain, and those mechanisms may act independently or they may act synergistically to overcome the action of an antimicrobial or a combination of antimicrobials. Specific mechanisms include degradation of the drug, inactivation of the drug by enzymatic

[0004] Modification, and alteration of the drug target. There are, however, more general mechanisms of drug resistance, in which access of the antimicrobial to the target is prevented or reduced by decreasing the transport of the antimicrobial into the cell or by increasing the efflux of the drug from the cell to the outside medium. Both mechanisms can lower the concentration of drug at the target site and allow bacterial survival in the presence of one or more antimicrobials that would otherwise inhibit or kill the bacterial cells. Some bacteria utilize both mechanisms, combining a low permeability of the cell wall (including membranes) with an active efflux of antimicrobials.

SUMMARY

[0005] Various embodiments provide compositions and methods for optimal antimicrobial activity for the treatment of respiratory tract and pulmonary infections in human and/or veterinary subjects using short-term, rapid aerosol administration, and through the delivery of high- concentration drug exposure directly to the affected tissue. Specifically, in some embodiments, concentrated doses of agents from the fluoroquinolone class of antibiotic are delivered to produce maximum concentrations of active drug to the respiratory, pulmonary, and other non-oral topical compartments including, but not limited to the skin, rectum, vagina, urethra, urinary bladder, eye, and ear. Because different drug products are known to produce different antimicrobial effects depending on the dose, form, concentration and delivery profile, some embodiments provide specific formulation and delivery parameters that produce antimicrobial results that are therapeutically significant. This disclosure includes, but is not limited to, specific fluoroquinolone antibiotics, such as levofloxacin, formulated to enable aerosol administration meeting specific concentrations and antimicrobial criteria necessary to treat patients with distinct bacterial infections. These formulations and methods are useful with commercially available inhalation devices for one or more aerosol therapeutic product opportunities.

[0006] Aerosol administration directly to the nasal, sinus, respiratory tract and pulmonary compartments through intra-nasal or oral inhalation enables high concentration drug delivery to a site of respiratory infection with decreased risk of extra-respiratory toxicity associated with non-respiratory routes of drug delivery. Furthermore,

direct administration to the site of infection permits very high local drug levels, a property that enables “rapid administration, high concentration, local exposure” killing effect special to this class of antibiotic. Accordingly, because the microbial killing effect of a particular antibiotic compound and therapeutic composition varies depending on the formulation and delivery parameters, newer compositions and delivery methods can be developed for existing drug compounds that are re-formulated and administered through novel delivery techniques. Other topical infections may also benefit from this discovery through high concentration, direct exposure of fluoroquinolone to infected skin, rectum, vagina, urethra, urinary bladder, eye, and ear.

[0007] Members of the fluoroquinolone drug class exhibit unique pharmacologic properties, including bioavailability (F), mean absorption time (MAT) from the lung, maximal drug concentrations in the epithelial lining fluid, bronchial lavage fluid, sputum and/or lung tissue (C_{max}) following aerosol administration, pulmonary retention time, area under the curve (AUC), minimal inhibitory concentrations (MIC) of the antibiotic required for antibacterial activity, AUC/MIC ratio, and local and systemic safety. Specific to the invention is the use short-term, rapid aerosol administration, delivering high concentration drug exposure directly to the affected tissue (ELF, sputum, BAL, tissue) via aerosol delivery for treatment of bacterial infection in animals and humans.

[0008] In addition to the clinical and pharmacological requirements present in any composition intended for therapeutic administration, many physicochemical factors unique to a drug compound must also be considered. These include, but are not limited to aqueous solubility, viscosity, partitioning coefficient (LogP), predicted stability in various formulations, osmolality, surface tension, pH, pK_a, pK_b, dissolution rate, sputum permeability, sputum binding/inactivation, taste, throat irritability and acute tolerability.

[0009] Other factors to consider when designing the product form include fluoroquinolone physical chemistry and antibacterial activity, disease indication, clinical acceptance, and patient compliance. By non-limiting example, if desired the aerosol fluoroquinolone product may be in the form of a simple liquid (e.g. soluble fluoroquinolone with non-encapsulating soluble excipients/salts), complex liquid (e.g.

fluoroquinolone encapsulated or complexed with soluble excipients such as lipids, liposomes, cyclodextrins, microencapsulations, and emulsions), complex suspension (e.g. fluoroquinolone as a low-solubility, stable nanosuspension alone, co-crystal/co-precipitate complexes, and mixtures with low solubility lipids such as solid-lipid nanoparticles), or dry powder (dry powder fluoroquinolone alone or in co-crystal/co-precipitate/spray-dried complex or mixture with low solubility excipients/salts or readily soluble blends such as lactose).

[0010] Combined with product form is a packaging consideration. By non-limiting example, considerations for packaging include intrinsic product stability, the need for stability-providing lyophilization, device selection (e.g. liquid nebulizer, dry-powder inhaler, meter-dose inhaler), and packaging form (e.g. simple liquid or complex liquid formulation in a vial as liquid or lyophilisate to be dissolved prior to or upon insertion into the device; complex suspension formulations in a vial as a liquid or lyophilisate with or without a soluble salt/excipient component to be dissolved prior to or upon insertion into the device, or separate packaging of liquid and solid components; dry powder formulations in a vial, capsule or blister pack; and other formulations packaged as readily soluble or low-solubility solid agents in separate containers alone or together with readily soluble or low-solubility solid agents. Any separately packaged agent will be manufactured to be mixed prior to or upon insertion into the delivery device).

[0011] In some aspects, **this disclosure** relates to the aerosol and topical delivery of fluoroquinolone antimicrobials, such as levofloxacin. Levofloxacin has favorable solubility characteristics enabling dosing of clinically-desirable fluoroquinolone levels by aerosol (e.g. through liquid nebulization, dry powder dispersion or meter-dose administration) or topically (e.g. aqueous suspension, oily preparation or the like or as a drip, spray, suppository, salve, or an ointment or the like) and can be used in methods for acute or prophylactic treatment of an infected vertebrate, e.g. a bacterial infection, or a subject at risk of an infection.

[0012] Others include: ofloxacin, lomefloxacin, pefloxacin, ciprofloxacin, gatifloxacin, gemifloxacin, moxifloxacin, tosufloxacin, pazufloxacin, rufloxacin, fleroxacin, balofloxacin, sparfloxacin, trovafloxacin, enoxacin, norfloxacin,

clinafloxacin, grepafloxacin, sitafloxacin, marbofloxacin, orbifloxacin, sarafloxacin, danofloxacin, difloxacin, enrofloxacin, garenoxacin, prulifloxacin, olamufloxacin, DX-619, TG-873870 and DW-276.

[0013] In a preferred embodiment, the method treats a bacterial infection in a subject using concentrated aerosol levofloxacin administered to a subject infected with a pathogenic bacteria in the lungs.

[0014] The therapeutic method may also include a diagnostic step, such as identifying a patient infected with a particular pathogenic bacteria, or a resistant bacteria. In some embodiments, the method further includes identifying a patient as colonized with a bacteria that is capable of developing resistance to fluoroquinolone antimicrobials. In some embodiments, the delivered amount of aerosol levofloxacin is sufficient to overcome resistance or prevent resistance development to levofloxacin. In one embodiment, the MIC of the fluoroquinolone antibacterial compound for the microbe is greater than about 2 ug/ml.

[0015] In another embodiment, the delivered amount of aerosol levofloxacin is sufficient to overcome resistance or prevent further resistance of an organism exhibiting an MIC of the fluoroquinolone antibacterial compound that is greater than about 4 ug/ml.

[0016] In another embodiment, the delivered amount of aerosol fluoroquinolone is sufficient to overcome resistance or prevent further resistance of an organism exhibiting an MIC of the fluoroquinolone antibacterial compound that is greater than about 8 ug/ml.

[0017] In another embodiment, the delivered amount of aerosol fluoroquinolone is sufficient to overcome resistance or prevent further resistance of an organism exhibiting an MIC of the fluoroquinolone antibacterial compound that is greater than about 16 ug/ml.

[0018] In another embodiment, the delivered amount of aerosol fluoroquinolone is sufficient to overcome resistance or prevent further resistance of an organism exhibiting an MIC of the fluoroquinolone antibacterial compound that is greater than about 32 ug/ml.

[0019] In another embodiment, a method is provided for prophylactic treatment of a subject, including administering to a subject, susceptible to microbial infection or a chronic carrier of an asymptomatic or low symptomatic microbial infection, a fluoroquinolone antimicrobial to achieve a minimal inhibitory concentration of antimicrobial at a site of potential or current infection. In one embodiment, the method further comprising identifying a subject as a subject at risk of a bacterial infection or at risk for an exacerbation of an infection.

[0020] In another embodiment, a method is provided for acute or prophylactic treatment of a patient through aerosol administration of fluoroquinolone to produce and maintain threshold drug concentrations in the lung, which may be measured as drug levels in epithelial lining fluid (ELF), sputum, lung tissue or bronchial lavage fluid (BAL). One embodiment includes the use of short-term, rapid aerosol administration, delivering high concentration drug exposure directly to the affected tissue for treatment of bacterial infections in animals and humans.

[0021] In another embodiment, a method is provided for treating a microbial infection in a subject, including administering to a subject infected with a microbe a fluoroquinolone antimicrobial to achieve a minimal inhibitory concentration of antimicrobial at a site of infection. In one embodiment, the method further comprising identifying the subject as infected with a microbe that is resistant to an antimicrobial agent.

[0022] In another embodiment, a method is provided for acute or prophylactic treatment of a patient through non-oral or non-nasal topical administration of fluoroquinolone to produce and maintain threshold drug concentrations at the site of infection or at risk of infection. One embodiment includes the use of short-term, rapid aerosol administration, delivering high concentration drug exposure directly to the affected tissue for treatment or prevention of bacterial infections in skin, rectal, vaginal, urethral, ocular, and auricular tissues.

[0023] In another embodiment, a method is provided for administering a fluoroquinolone antimicrobial by inhalation, wherein the inhaled liquid or dry powder aerosol has a mean particle size from about 1 micron to 10 microns mass median

aerodynamic diameter and a particle size geometric standard deviation of less than or equal to about 3 microns. In another embodiment, the particle size is 2 microns to about 5 microns mass median aerodynamic diameter and a particle size geometric standard deviation of less than or equal to about 2 microns. In one embodiment, the particle size geometric standard deviation is less than or equal to about 1.8 microns.

[0024] In some embodiments of the methods described above, fluoroquinolone antimicrobial minimal inhibitory concentration remains at the site of infection for at least about a 5 minute period, at least about a 10 min period, at least about a 20 min period, at least about a 30 min period, at least about a 1 hour period, 2 hour period, at least about a 4 hour period or other time values on the quarter hour interval. The effective fluoroquinolone antimicrobial minimal inhibitory concentration (MIC) is sufficient to cause a therapeutic effect and the effect may be localized to the site of infection. In some embodiments, one or more levofloxacin administrations achieve an ELF, BAL, and/or sputum fluoroquinolone concentration of at least 1-fold to 5000-fold the infecting or potentially infecting organisms MIC, including all integral values therein such as 2-fold, 4-fold, 8-fold, 16-fold, 32-fold, 64-fold, 128-fold, 256-fold, 512-fold, 1028-fold, 2056-fold, and 4112-fold the microbials MIC.

[0025] In some embodiments, such as a pulmonary site, the fluoroquinolone antimicrobial is administered in one or more administrations so as to achieve a respirable delivered dose daily of at least about 5 mg to about 50 mg, including all integral values therein such as 10, 15, 20, 25, 30, 35, 40 and 45 milligrams. Similarly, the fluoroquinolone antimicrobial is administered in one or more administrations so as to achieve a respirable delivered dose daily of at least about 50 to about 100 mg including all integral values therein, such as 55, 60, 65, 70, 75, 80, 85, 90, and 95 mg. In some embodiments of the methods described above, the fluoroquinolone antimicrobial is administered in one or more administrations so as to achieve a respirable delivered daily dose of up to 150 mg including all integral values therein such as 105, 110, 115, 120, 125, 130, 135, 140 and 145 mg. The fluoroquinolone antimicrobial is administered in the described respirable delivered dose in less than 20 minutes, less than 10 minutes, less than 7 minutes, less than 5 minutes, in less than 3 minutes and in less than 2 minutes. In some embodiments of the methods described above, the antimicrobial agent is selected

from the group consisting of ofloxacin, lomefloxacin, pefloxacin, ciprofloxacin, gatifloxacin, gemifloxacin, moxifloxacin, tosufloxacin, pazufloxacin, rufloxacin, fleroxacin, balofloxacin, sparfloxacin, trovafloxacin, enoxacin, norfloxacin, clinafloxacin, grepafloxacin, sitafloxacin, marbofloxacin, orbifloxacin, sarafloxacin, danofloxacin, difloxacin, enrofloxacin, garenoxacin, prulifloxacin, olamufloxacin, DX-619, TG-873870 and DW-276, although levofloxacin is preferred.

[0026] In some embodiments of the methods described above, the bacteria is a gram-negative bacteria such as *Pseudomonas aeruginosa*, *Pseudomonas fluorescens*, *Pseudomonas acidovorans*, *Pseudomonas alcaligenes*, *Pseudomonas putida*, *Stenotrophomonas maltophilia*, *Burkholderia cepacia*, *Aeromonas hydrophilia*, *Escherichia coli*, *Citrobacter freundii*, *Salmonella typhimurium*, *Salmonella typhi*, *Salmonella paratyphi*, *Salmonella enteritidis*, *Shigella dysenteriae*, *Shigella flexneri*, *Shigella sonnei*, *Enterobacter cloacae*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Klebsiella oxytoca*, *Serratia marcescens*, *Francisella tularensis*, *Morganella morganii*, *Proteus mirabilis*, *Proteus vulgaris*, *Providencia alcalifaciens*, *Providencia rettgeri*, *Providencia stuartii*, *Acinetobacter calcoaceticus*, *Acinetobacter haemolyticus*, *Yersinia enterocolitica*, *Yersinia pestis*, *Yersinia pseudotuberculosis*, *Yersinia intermedia*, *Bordetella pertussis*, *Bordetella parapertussis*, *Bordetella bronchiseptica*, *Haemophilus influenzae*, *Haemophilus parainfluenzae*, *Haemophilus haemolyticus*, *Haemophilus parahaemolyticus*, *Haemophilus ducreyi*, *Pasteurella multocida*, *Pasteurella haemolytica*, *Branhamella catarrhalis*, *Helicobacter pylori*, *Campylobacter fetus*, *Campylobacter jejuni*, *Campylobacter coli*, *Borrelia burgdorferi*, *Vibrio cholerae*, *Vibrio parahaemolyticus*, *Legionella pneumophila*, *Listeria monocytogenes*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Kingella*, *Moraxella*, *Gardnerella vaginalis*, *Bacteroides fragilis*, *Bacteroides distasonis*, *Bacteroides 3452A* homology group, *Bacteroides vulgatus*, *Bacteroides ovalus*, *Bacteroides thetaiotaomicron*, *Bacteroides uniformis*, *Bacteroides eggerthii*, and *Bacteroides splanchnicus*. In some embodiments of the methods described above, the bacteria is a gram-negative anaerobic bacteria, by non-limiting example these include *Bacteroides fragilis*, *Bacteroides distasonis*, *Bacteroides 3452A* homology group, *Bacteroides vulgatus*, *Bacteroides ovalus*, *Bacteroides thetaiotaomicron*, *Bacteroides uniformis*, *Bacteroides eggerthii*, and *Bacteroides splanchnicus*. In some embodiments

of the methods described above, the bacteria is a gram-positive bacteria, by non-limiting example these include: *Corynebacterium diphtheriae*, *Corynebacterium ulcerans*, *Streptococcus pneumoniae*, *Streptococcus agalactiae*, *Streptococcus pyogenes*, *Streptococcus milleri* ; *Streptococcus* (Group G); *Streptococcus* (Group C/F); *Enterococcus faecalis*, *Enterococcus faecium*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, *Staphylococcus intermedius*, *Staphylococcus hyicus* subsp. *hyicus*, *Staphylococcus haemolyticus*, *Staphylococcus hominis*, and *Staphylococcus saccharolyticus*. In some embodiments of the methods described above, the bacteria is a gram-positive anaerobic bacteria, by non-limiting example these include *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, and *Clostridium botulinum*. In some embodiments of the methods described above, the bacteria is a acid-fast bacteria, by non-limiting example these include *Mycobacterium tuberculosis*, *Mycobacterium avium*, *Mycobacterium intracellulare*, and *Mycobacterium leprae*. In some embodiments of the methods described above, the bacteria is a atypical bacteria, by non-limiting example these include *Chlamydia pneumoniae* and *Mycoplasma pneumoniae*.

[0027] In some embodiments of the methods described above, the subject is a human. In some embodiments of the methods described above, the subject is a human with cystic fibrosis. In some embodiments of the methods described above, the subject is a human with pneumonia, a chronic obstructive pulmonary disease, or sinusitis, or a human being mechanically ventilated.

[0028] In another embodiment, a pharmaceutical composition is provided that includes a simple liquid fluoroquinolone antimicrobial formulation (e.g. soluble fluoroquinolone with non-encapsulating water soluble excipients) as described above having an osmolality from about 200 mOsmol/kg to about 1250 mOsmol/kg. In one such embodiment, the solution has a permeant ion concentration from about 30 mM to about 300 mM. In one embodiment, the osmolality is from about 250 mOsmol/kg to about 1050 mOsmol/kg. In one embodiment, the osmolality is from preferably from about 350 mOsmol/kg and about 750 mOsmol/kg and most preferably approximately 300 mOsmol/kg.

[0029] In another embodiment, a pharmaceutical composition is provided that includes a simple liquid fluoroquinolone antimicrobial formulation having a permeant ion concentration between from about 30 mM to about 300 mM and preferably between from about 50mM to 200 mM. In one such embodiment, one or more permeant ions in the composition are selected from the group consisting of chloride and bromide.

[0030] In another embodiment, a pharmaceutical composition is provided that includes a complex liquid fluoroquinolone antimicrobial formulation (e.g. fluoroquinolone encapsulated or complexed with water soluble excipients such as lipids, liposomes, cyclodextrins, microencapsulations, and emulsions) as described above having a solution osmolality from about 200 mOsmol/kg to about 1250 mOsmol/kg. In one such embodiment, the solution has a permeant ion concentration from about 30 mM to about 300 mM. In one embodiment, the osmolality is from about 250 mOsmol/kg to about 1050 mOsmol/kg. In one embodiment, the osmolality is from preferably from about 350 mOsmol/kg and about 750 mOsmol/kg and most preferably approximately 300 mOsmol/kg.

[0031] In another embodiment, a pharmaceutical composition is provided that includes a complex liquid fluoroquinolone antimicrobial formulation having a permeant ion concentration from about 30 mM to about 300 mM. In one such embodiment, one or more permeant ions in the composition are selected from the group consisting of chloride and bromide.

[0032] In another embodiment, a pharmaceutical composition is provided that includes a complex liquid fluoroquinolone antimicrobial formulation having a permeant ion concentration from about 50 mM to about 200 mM. In one such embodiment, one or more permeant ions in the composition are selected from the group consisting of chloride and bromide.

[0033] In another embodiment, a pharmaceutical composition is provided that includes a complex liquid fluoroquinolone antimicrobial formulation (e.g. fluoroquinolone as a low water soluble stable nanosuspension alone or in co-crystal/co-precipitate complexes, or mixtures with low solubility lipids, such as lipid nanosuspensions) as described above having a solution osmolality from about 200

mOsmol/kg to about 1250 mOsmol/kg. In one such embodiment, the solution has a permeant ion concentration from about 30 mM to about 300 mM. In one embodiment, the osmolality is from about 250 mOsmol/kg to about 1050 mOsmol/kg. In one embodiment, the osmolality is from preferably from about 350 mOsmol/kg and about 750 mOsmol/kg and most preferably approximately 300 mOsmol/kg.

[0034] In another embodiment, a pharmaceutical composition is provided that includes a complex suspension fluoroquinolone antimicrobial formulation having a permeant ion concentration from about 30 mM to about 300 mM. In one such embodiment, one or more permeant ions in the composition are selected from the group consisting of chloride and bromide.

[0035] In another embodiment, a pharmaceutical composition is provided that includes a complex suspension fluoroquinolone antimicrobial formulation having a permeant ion concentration from about 50 mM to about 200 mM. In one such embodiment, one or more permeant ions in the composition are selected from the group consisting of chloride and bromide.

[0036] In another embodiment, a pharmaceutical composition is provided that includes a taste-masking agent. By non-limiting example a taste-masking agent may include a sugar, a divalent or trivalent cation that complexes with a fluoroquinolone, optimized osmolality, and/or an optimized permeant ion concentration.

[0037] In another embodiment, a pharmaceutical composition is provided that includes a simple dry powder fluoroquinolone antimicrobial compound (e.g. fluoroquinolone alone in dry powder form with or without a blending agent such as lactose).

[0038] In another embodiment, a pharmaceutical composition is provided that includes a complex dry powder fluoroquinolone antimicrobial formulation (e.g. fluoroquinolone in co-crystal/co-precipitate/spray dried complex or mixture with low water soluble excipients/salts in dry powder form with or without a blending agent such as lactose).

[0039] In another embodiment, a system is provided for administering a

fluoroquinolone antimicrobial that includes a container comprising a solution of a fluoroquinolone antimicrobial and a nebulizer physically coupled or co-packaged with the container and adapted to produce an aerosol of the solution having a particle size from about 2 microns to about 5 microns mean mass aerodynamic diameter and a particle size geometric standard deviation of less than or equal to about 2.5 microns mean mass aerodynamic diameter. In one embodiment, the particle size geometric standard deviation is less than or equal to about 2.0 microns. In one embodiment, the particle size geometric standard deviation is less than or equal to about 1.8 microns.

[0040] In another embodiment, a system is provided for administering a fluoroquinolone antimicrobial that includes a container comprising a dry powder of a fluoroquinolone antimicrobial and a dry powder inhaler coupled to the container and adapted to produce a dispersed dry powder aerosol having a particle size from about 2 microns to about 5 microns mean mass aerodynamic and a particle size standard deviation of less than or equal to about 3.0 microns. In one embodiment, the particle size standard deviation is less than or equal to about 2.5 microns. In one embodiment, the particle size standard deviation is less than or equal to about 2.0 microns.

[0041] In another embodiment, a kit is provided that includes a container comprising a pharmaceutical formulation comprising a quinolone antimicrobial agent and an aerosolizer adapted to aerosolize the pharmaceutical formulation and deliver it to the lower respiratory tract and pulmonary compartment following intraoral administration. The formulation may also be delivered as a dry powder or through a metered-dose inhaler.

[0042] In another embodiment, a kit is provided that includes a container comprising a pharmaceutical formulation comprising a quinolone antimicrobial agent and an aerosolizer adapted to aerosolize the pharmaceutical formulation and deliver it to nasal cavity following intranasal administration. The formulation may also be delivered as a dry powder or through a metered-dose inhaler.

[0043] It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

[043a] Various embodiments of the claimed invention relate to a pharmaceutical composition, comprising a solution of levofloxacin or ofloxacin and a divalent or trivalent cation. The composition may comprise another antimicrobial and the solution may be one with that is suitable for inhalation into a lung. Also claimed is a sterile, single use container comprising such a solution and kits comprising such a container. Such a kit may further comprise a nebulizer adapted to aerosolize the solution for delivery to the lower respiratory tract through oral inhalation.

[043b] Various embodiments of the claimed invention relate to use of levofloxacin or ofloxacin in combination with a divalent or trivalent cation for treating a lung infection, a chronic obstructive pulmonary disease, cystic fibrosis, or sinusitis. The use may be in preparation for an aerosolize for such treating.

[043c] Various embodiments of the claimed invention relate to an aerosol of a solution comprising levofloxacin or ofloxacin and a divalent or trivalent cation.

[043d] Various embodiments of the claimed invention relate to an aerosol dose of a levofloxacin and magnesium solution comprised of: a concentration of levofloxacin greater than 50 mg/ml and a taste-masking concentration of magnesium wherein the aerosol is comprised of a mist having a mean particle size of between 2 and 5 microns or a particle size geometric standard deviation of less than or equal to about 2 microns.

[043e] Various embodiments of the claimed invention relate to a system, comprising: a reservoir comprising the pharmaceutical composition as claimed herein; and a nebulizer configured to aerosolize the solution for delivery to the lung through oral inhalation.

DESCRIPTION OF FIGURES

[0044] Figure 1 is a graph showing the dose:MIC relationship of fluoroquinolones and other antibiotics to bacterial killing.

[0045] Figure 2 is a graph showing ciprofloxacin serum concentrations following oral dosing in both CF patients vs healthy controls.

[0046] Figure 3 is a graph showing ciprofloxacin sputum and serum concentrations following oral dosing.

[0047] Figure 4A is a graph showing Levofloxacin time-kill affects on logarithmic PAM1020 cells.

[0048] Figure 4B is a graph showing Levofloxacin time-kill affects on logarithmic PAM1032 cells.

[0049] Figure 5A is a graph showing Levofloxacin time-kill affects on stationary phase PAM1020 cells.

[0050] Figure 5B is a graph showing Levofloxacin time-kill affects on stationary phase PAM1032 cells.

[0051] Figure 6A is a graph showing PAM 1020 re-growth following a 10 minute Levofloxacin exposure.

[0052] Figure 6B is a graph showing PAM 1020 re-growth following a 160 minute Levofloxacin exposure.

[0053] Figure 6C is a graph showing PAM 1032 re-growth following a 10 minute Levofloxacin exposure.

[0054] Figure 6D is a graph showing PAM 1032 re-growth following a 160 minute Levofloxacin exposure.

[0055] Figure 7A is a graph showing Levofloxacin time-kill affects on late-logarithmic PAM 1020 cells under oxygen limiting conditions.

[0056] Figure 7B is a graph showing Levofloxacin time-kill affects on late-logarithmic PAM 1032 cells under oxygen limiting conditions.

[0057] Figure 8A is a graph showing Levofloxacin killing kinetics of PAM1032 in Mueller-Hinton broth (MHB).

[0058] Figure 8B is a graph showing Levofloxacin killing kinetics of PAM1032 in cystic fibrosis sputum.

[0059] Figure 9 is a graph showing levofloxacin killing affects on *Pseudomonas* biofilms.

[0060] Figure 10 is a graph showing the bactericidal effects of levofloxacin with a Cmax of 1000 µg/ml and a 10 minute half-life in a hollow fiber model.

[0061] Figure 11 is a graph showing the bactericidal effects of levofloxacin with a Cmax of 600 µg/ml and a 10 minute half-life in a hollow fiber model.

[0062] Figure 12 is a graph relating the micronization pressure used to micronized dry powder levofloxacin vs. mean Levofloxacin dry powder particle size.

[0063] Figure 13 is a graph showing the DSC profile of pre-micronized and micronized dry powder Levofloxacin.

[0064] Figure 14A is a graph showing SEM photomicrographs of pre-micronized dry powder Levofloxacin.

[0065] Figure 14B is a graph showing SEM photomicrographs of micronized dry powder Levofloxacin.

[0066] Figure 15 is a graph showing X-ray diffraction of pre-micronized and micronized dry powder Levofloxacin.

[0067] Figure 16 is a graph showing the pH solubility profile of Levofloxacin by acid titration.

[0068] Figure 17 is a graph measuring pH while titrating Levofloxacin with HCl.

[0069] Figure 18 is a graph showing the $V_t[\text{OH}]$ vs. V_t of Levofloxacin.

[0070] Figure 19 is a graph measuring pH while titrating Levofloxacin with NaOH.

[0071] Figure 20 is a graph measuring dpH/dV vs volume of NaOH titrant (Vt) for titration of Levofloxacin.

[0072] Figure 21 is a graph measuring the absorbance of a Levofloxacin solution at 257 nm vs pH.

[0073] Figure 22 depicts graphs showing DSC scans of pamoic acid, Levofloxacin, Levofloxacin pamoic acid co-crystallized precipitate, and Levofloxacin-pamoic acid physical mixture.

[0074] Figure 23 depicts graphs showing FTIR spectra of pamoic acid, Levofloxacin, Levofloxacin Pamoic Acid Co-Crystallized Precipitate, and Levofloxacin-Pamoic Acid Physical Mixture.

[0075] Figure 24 depicts graphs showing DSC scans of xinafoic acid, and Levofloxacin xinafoic acid co-crystallized precipitate.

[0076] Figure 25 depicts graphs showing FTIR spectra of xinafoic acid and Levofloxacin xinafoate co-crystals.

[0077] Figure 26 depicts graphs showing DSC scans of stearic acid, Levofloxacin stearic acid co-crystallized precipitate, and a physical mixture of Levofloxacin and stearic acid.

[0078] Figure 27 depicts graphs showing FTIR spectra of stearic acid, Levofloxacin stearic acid co-crystallized precipitate, and a physical mixture of Levofloxacin and stearic acid.

[0079] Figure 28 depicts graphs showing DSC scans of oleic acid, Levofloxacin oleic acid co-crystallized precipitate, physical mixture of Levofloxacin and oleic acid (50:50), physical mixture of Levofloxacin and oleic acid (10:90), and physical mixture of Levofloxacin and oleic acid (90:10).

[0080] Figure 29 depicts graphs showing FTIR spectra of oleic Acid, Levofloxacin oleic acid co-crystallized precipitate, Levofloxacin oleic acid co-crystallized precipitate as compared to an equimolar physical mixture of levofloxacin and oleic acid.

[0081] Figure 30 is a graph showing the kinetic solubility of the co-crystallized precipitate of Levofloxacin with oleic acid at room temperature, 40°C, and equimolar physical mixture at 40°C.

[0082] Figure 31 is a graph showing the dissolution profile of Levofloxacin xinafoate.

[0083] Figure 32 is a graph showing the dissolution profile of Levofloxacin xinafoate focused on the period between two and ten minutes.

[0084] Figure 33 is a graph showing the dissolution profile of Levofloxacin xinafoate focused on the period between ten and thirty minutes.

[0085] Figure 34 is a graph showing the dissolution profile of Levofloxacin base.

[0086] Figure 35 is a graph showing the dissolution profile of Levofloxacin pamoate.

[0087] Figure 36 is a graph showing the dissolution profile of Levofloxacin pamoate focused on the period between two and ten minutes.

[0088] Figure 37 is a graph showing the dissolution profile of Levofloxacin pamoate focused on the period between ten and sixty minutes.

[0089] Figure 38 is a graph showing the dissolution profile of Levofloxacin stearate.

[0090] Figure 39 is a graph showing the dissolution profile of Levofloxacin stearate focused on the period between two and ten minutes.

[0091] Figure 40 is a graph showing the dissolution profile of Levofloxacin stearate focused on the period between ten and thirty minutes.

[0092] Figure 41 is a graph showing the complexation of Levofloxacin with divalent and trivalent cations.

[0093] Figure 42 is a graph showing the dual titration complexation of Levofloxacin with Mg²⁺.

[0094] Figure 43 is a graph showing the dual titration complexation of Levofloxacin with Fe^{2+} .

[0095] Figure 44 is a graph showing the dual titration complexation of Levofloxacin with Ca^{2+} .

[0096] Figure 45 is a graph showing the dual titration complexation of Levofloxacin with Zn^{2+} .

[0097] Figure 46 is a graph showing Levofloxacin complexed with Ca^{2+} vs. free Levofloxacin.

[0098] Figure 47 is a graph showing Levofloxacin complexed with Mg^{2+} vs. free Levofloxacin.

[0099] Figure 48 is a graph showing Levofloxacin complexed with Fe^{2+} vs. free Levofloxacin.

[00100] Figure 49 is a graph showing Levofloxacin complexed with Zn^{2+} vs. free Levofloxacin.

[00101] Figure 50 is a graph showing the solubility of Levofloxacin in the presence of Mg^{2+} .

[00102] Figure 51 is a graph showing the solubility of Levofloxacin in the presence of Mg^{2+} at constant ionic strength.

[00103] Figure 52 is a graph showing complexation of Levofloxacin with Fe^{2+} as measured by spectrofluorometry.

[00104] Figure 53 is a graph showing complexation of Levofloxacin with Zn^{2+} as measured by spectrofluorometry.

DETAILED DESCRIPTION

[00105] Many of the problems associated with antimicrobial-resistant pathogens could be alleviated if the concentration of the antimicrobial could be safely increased at the site of infection. For example, pulmonary infections may be treated by administration of the antimicrobial agent directly, at high concentrations directly to the site of infection without incurring large systemic concentrations of the antimicrobial. Accordingly, some

embodiments disclosed herein are improved methods for delivering drug compositions to treat pulmonary bacterial infections. More specifically, as described herein, it has been discovered that aerosol levofloxacin and other fluoroquinolones can be safely delivered by inhalation at levels sufficient to kill susceptible bacterial infections, to decrease the frequency of antimicrobial resistance and to increase efficacy against resistant pulmonary infections.

Definitions

[00106] The term “administration” or “administering” refers to a method of giving a dosage of an antimicrobial pharmaceutical composition to a vertebrate. The preferred method of administration can vary depending on various factors, e.g., the components of the pharmaceutical composition, the site of the potential or actual bacterial infection, the microbe involved, and the severity of an actual microbial infection.

[00107] A “carrier” or “excipient” is a compound or material used to facilitate administration of the compound, for example, to increase the solubility of the compound. Solid carriers include, e.g., starch, lactose, dicalcium phosphate, sucrose, and kaolin. Liquid carriers include, e.g., sterile water, saline, buffers, non-ionic surfactants, and edible oils such as oil, peanut and sesame oils. In addition, various adjuvants such as are commonly used in the art may be included. These and other such compounds are described in the literature, e.g., in the Merck Index, Merck & Company, Rahway, NJ. Considerations for the inclusion of various components in pharmaceutical compositions are described, e.g., in Gilman et al. (Eds.) (1990); Goodman and Gilman’s: The Pharmacological Basis of Therapeutics, 8th Ed., Pergamon Press.

[00108] A “diagnostic” as used herein is a compound, method, system, or device that assists in the identification and characterization of a health or disease state. The diagnostic can be used in standard assays as is known in the art.

[00109] The term “mammal” is used in its usual biological sense. Thus, it specifically includes humans, cattle, horses, dogs, and cats, but also includes many other species.

[00110] The term “microbial infection” refers to the undesired proliferation or presence of invasion of pathogenic microbes in a host organism. This includes the excessive growth of microbes that are normally present in or on the body of a mammal or other organism. More generally, a microbial infection can be any situation in which the presence of a microbial population(s) is damaging to a host mammal. Thus, a microbial infection exists when excessive numbers of a microbial population are present in or on a mammal’s body, or when the effects of the presence of a microbial population(s) is damaging the cells or other tissue of a mammal.

[00111] The term “pharmaceutically acceptable carrier” or “pharmaceutically acceptable excipient” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

[00112] The term “pharmaceutically acceptable salt” refers to salts that retain the biological effectiveness and properties of the compounds of this invention and, which are not biologically or otherwise undesirable. In many cases, the compounds of this invention are capable of forming acid and/or base salts by virtue of the presence of amino and/or carboxyl groups or groups similar thereto. Pharmaceutically acceptable acid addition salts can be formed with inorganic acids and organic acids. Inorganic acids from which salts can be derived include, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like. Organic acids from which salts can be derived include, for example, acetic acid, propionic acid, naphtoic acid, oleic acid, palmitic acid, pamoic (emboic) acid, stearic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, ascorbic acid, glucoheptonic acid, glucuronic acid, lactic acid, lactobioic acid, tartaric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like. Pharmaceutically acceptable base addition salts can be formed with inorganic and organic bases. Inorganic bases from which salts can be derived include, for example, sodium, potassium, lithium, ammonium,

calcium, magnesium, iron, zinc, copper, manganese, aluminum, and the like; particularly preferred are the ammonium, potassium, sodium, calcium and magnesium salts. Organic bases from which salts can be derived include, for example, primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, basic ion exchange resins, and the like, specifically such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, histidine, arginine, lysine, benethamine, N-methyl-glucamine, and ethanolamine. Other acids include dodecylsulfuric acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, and saccharin.

[00113] "Solvate" refers to the compound formed by the interaction of a solvent and fluoroquinolone antimicrobial, a metabolite, or salt thereof. Suitable solvates are pharmaceutically acceptable solvates including hydrates.

[00114] In the context of the response of a microbe, such as a bacterium, to an antimicrobial agent, the term "susceptibility" refers to the sensitivity of the microbe for the presence of the antimicrobial agent. So, to increase the susceptibility means that the microbe will be inhibited by a lower concentration of the antimicrobial agent in the medium surrounding the microbial cells. This is equivalent to saying that the microbe is more sensitive to the antimicrobial agent. In most cases the minimum inhibitory concentration (MIC) of that antimicrobial agent will have been reduced.

[00115] By "therapeutically effective amount" or "pharmaceutically effective amount" is meant a fluoroquinolone antimicrobial agent, as disclosed for this invention, which has a therapeutic effect. The doses of fluoroquinolone antimicrobial agent which are useful in treatment are therapeutically effective amounts. Thus, as used herein, a therapeutically effective amount means those amounts of fluoroquinolone antimicrobial agent which produce the desired therapeutic effect as judged by clinical trial results and/or model animal infection studies. In particular embodiments, the fluoroquinolone antimicrobial agent are administered in a pre-determined dose, and thus a therapeutically effective amount would be an amount of the dose administered. This amount and the amount of the fluoroquinolone antimicrobial agent can be routinely determined by one of skill in the art, and will vary, depending on several factors, such as the particular

microbial strain involved. This amount can further depend upon the patient's height, weight, sex, age and medical history. For prophylactic treatments, a therapeutically effective amount is that amount which would be effective to prevent a microbial infection.

[00116] A "therapeutic effect" relieves, to some extent, one or more of the symptoms of the infection, and includes curing an infection. "Curing" means that the symptoms of active infection are eliminated, including the total or substantial elimination of excessive members of viable microbe of those involved in the infection to a point at or below the threshold of detection by traditional measurements. However, certain long-term or permanent effects of the infection may exist even after a cure is obtained (such as extensive tissue damage). As used herein, a "therapeutic effect" is defined as a statistically significant reduction in bacterial load in a host, emergence of resistance, or improvement in infection symptoms as measured by human clinical results or animal studies.

[00117] "Treat," "treatment," or "treating," as used herein refers to administering a pharmaceutical composition for prophylactic and/or therapeutic purposes. The term "prophylactic treatment" refers to treating a patient who is not yet infected, but who is susceptible to, or otherwise at risk of, a particular infection. The term "therapeutic treatment" refers to administering treatment to a patient already suffering from an infection. Thus, in preferred embodiments, treating is the administration to a mammal (either for therapeutic or prophylactic purposes) of therapeutically effective amounts of a fluoroquinolone antimicrobial agent.

[00118] Pharmacokinetics (PK) is concerned with the time course of antimicrobial concentration in the body. Pharmacodynamics (PD) is concerned with the relationship between pharmacokinetics and the antimicrobial efficacy *in vivo*. PK/PD parameters correlate antimicrobial exposure with antimicrobial activity. The rate of killing by antimicrobial is dependent on antimicrobial mode of action and is determined by either the length of time necessary to kill (time-dependent) or the effect of increasing concentrations (concentration-dependent). Accordingly, to predict the therapeutic efficacy of antimicrobials with diverse mechanisms of action different PK/PD parameters

may be used.

[00119] “AUC/MIC ratio” is one example of a PK/PD parameter. AUC is defined as the area under the plasma or site-of-infection concentration-time curve of an antimicrobial agent *in vivo* (in animal or human). AUC/MIC ratio is determined by dividing the 24-hour-AUC for an individual antimicrobial by the MIC for the same antimicrobial determined *in vitro*. Activity of antimicrobials with the dose-dependent killing (such as fluoroquinolones) is well predicted by the magnitude of the AUC/MIC ratio.

[00120] “Cmax:MIC” ratio is another PK:PD parameter. It describes the maximum drug concentration in plasma or tissue relative to the MIC. Fluoroquinolones and aminoglycosides are examples where Cmax:MIC may predict *in vivo* bacterial killing where resistance can be suppressed.

[00121] “Time above MIC” (T>MIC) is another PK/PD parameter. It is expressed as a percentage of a dosage interval in which the plasma or site-of-infection level exceeds the MIC. Activity of antimicrobials with the time-dependent killing (such as beta-lactams or oxazolidinones) is well predicted by the magnitude of the T>MIC ratio.

[00122] The term “dosing interval” refers to the time between administrations of the two sequential doses of a pharmaceutical’s during multiple dosing regimens. For example, in the case of ciprofloxacin, which is administered twice daily (traditional regimen of 400 mg b.i.d) and levofloxacin, which is administered once a day (500mg or 750mg q.d.), the dosing intervals are 12 hours and 24 hours, respectively.

[00123] As used herein, the “peak period” of a pharmaceutical’s *in vivo* concentration is defined as that time of the pharmaceutical dosing interval when the pharmaceutical concentration is not less than 50% of its maximum plasma or site-of-infection concentration. In some embodiments, “peak period” is used to describe an interval of antimicrobial dosing.

[00124] The “respirable delivered dose” is the amount of drug inhaled during the inspiratory phase of the breath simulator that is equal to or less than 5 microns using a simulator programmed to the European Standard pattern of 15 breaths per minute, with

an inspiration to expiration ratio of 1:1.

Advantages of Inhaled Aerosol and Topical (Non-Oral) Fluoroquinolone Delivery

[00125] The antibiotic rate of killing is dependent upon antibiotic mode of action and is determined by either the length of time necessary for the antibiotic to kill (time-dependent) or the effect of increasing the antibiotic concentration (concentration-dependent). Fluoroquinolones are characterized by concentration-dependent, time-kill activity where a therapeutic effect requires a high local peak concentration above the MICs of the infecting pathogen.

[00126] Fluoroquinolone efficacy in humans, animals and *in vitro* models of infection is linked to AUC:MIC ratio and C_{max}:MIC ratio. Given the prior uncertainty of the pharmacokinetics of fluoroquinolones in pulmonary tissue, a number of *in vitro* studies have been conducted to determine if high doses of levofloxacin with extremely short half-lives (as predicted from a rat and human PK model) result in bacterial killing superior to that seen under conditions with more prolonged residence times. In these studies, levofloxacin concentrations that were 0.018-fold – 1024-fold the MIC were evaluated in a standard kill-curve and *in vitro* hollow fiber assay. In both of these assays, high concentrations of levofloxacin were rapidly bactericidal and reached their maximum level of killing in 10-20 minutes. This level of killing was sustained whether levofloxacin was maintained at that level or given a half-life of 10 minutes. Accordingly, high doses and rapid delivery of specially formulated levofloxacin, such as a rapidly delivered 20-50 mg respirable deposited aerosol levofloxacin dose (which will produce initial ELF concentrations of 800 – 1600 ug/mL) is rapidly bactericidal for susceptible organisms and resistant organisms with MICs up to 32 ug/ml. It is expected that these unique antimicrobial properties of fluoroquinolones will also translate to topical administrations, including, but not limited to infections or prophylaxis of the skin, eye, ear, rectum, vagina, or urinary tract.

[00127] To measure the efficacy of different delivery models, AUC shape-enhancing levofloxacin formulations were prepared and measured *in vivo* in comparison to non-AUC shape-enhancing levofloxacin formulations and other antibiotics using both rat PK and mouse efficacy following intratracheal administration. As was previously

shown in a rat system, there were differences among drugs in pulmonary pharmacokinetics, with some agents showing lower AUCs (e.g., levofloxacin), while others such as gemifloxacin or tobramycin show higher concentrations resulting from slower pulmonary clearance. Studies in a single dose mouse model of infection with aerosol doses have shown variable efficacy among the compounds. Referring to Figure 1, analysis of the data by dividing the aerosolized dose by the MIC indicated a strong correlation between Dose:MIC ratio and bactericidal activity ($R^2 = 0.89$). These data suggest that the initial bactericidal activity in this model is not affected by the pulmonary clearance of the drug. Although pulmonary clearances have not been estimated in mice, transformation of dose to AUC using scaled rat values would be expected to degrade the relationship. Therefore, this data suggests that optimizing the AUC shape for levofloxacin may not be necessary for aerosol levofloxacin to be effective in treating respiratory tract and pulmonary infections.

[00128] Recent investigations with fluoroquinolones resulted in development of the concept of a “mutant selection window” (MSW) for bacterial resistance arising during therapy. This concept assists in identifying a concentration range where mutants are selected more frequently *in vitro* and *in vivo*. The lower boundary of the window is the lowest concentration that kills the majority of infecting cells (approximated by the MIC), while the upper boundary of the window is the drug concentration that blocks the growth of the least susceptible first-step mutant. Above the upper boundary concentration the growth of the infecting bacteria requires the presence of at least two resistance mutations. This upper boundary is designated the mutant prevention concentration (MPC). The values of MPC vary depending on bacteria and fluoroquinolone, and may be 10- to 20-fold higher than the MIC. Several modeling studies have demonstrated that the longer the drug concentration exceeds the MPC at the site of infection, the more effectively the treatment will prevent resistance development. Conversely, the longer the antibiotic concentration stays within the MSW, the higher the probability to select resistant mutants. Importantly, the currently approved dosing regimen for oral or intravenous levofloxacin has placed this antibiotic within the MSW for more than 20% of the dosing interval for such pathogens as *P. aeruginosa* (*Pa*) and *S. pneumonia*. Accordingly, a high level of levofloxacin resistance is reported for both of these pathogens.

[00129] Therefore, in one embodiment, the concentration of levofloxacin at the site of infection is increased by delivering it directly to the lung using inhalation therapy, thereby decreasing the amount of time levofloxacin is in the MSW. Such a therapeutic approach achieves broader coverage of pathogens (including levofloxacin resistant strains), prevents further resistance development, and results in shorter courses of levofloxacin therapy.

Pharmacokinetics of Orally Administered Fluoroquinolones in Non-CF and CF

Populations

Sputum Concentrations in CF Patients

[00130] The pharmacokinetics of ciprofloxacin has been extensively studied in CF patients after oral administration. In fact, it has been demonstrated that the serum PK profile of ciprofloxacin is very similar in CF patients to healthy volunteers (Figure 2).

[00131] Moreover, the sputum vs time profile of ciprofloxacin is very similar to its serum profile after oral administration (Figure 3). After a 750 mg oral dose, peak concentrations of ~ 4.2 $\mu\text{g/ml}$ and ~ 3.5 $\mu\text{g/ml}$ were achieved for serum and sputum, respectively. Serum and sputum drug concentrations peaked at 1.5 and 4 hours, respectively. While the the total amount of ciprofloxacin into sputum is high relative to serum concentrations, the absolute concentrations are low relative to MICs of target organisms such as Pa. This data is consistent with poor clinical outcome due to resistance development to these low drug concentrations.

[00132] While data for levofloxacin intrapulmonary pharmacokinetics in cystic fibrosis are not available, data on the closely-related ofloxacin were published in the 1980's and 1990s. Ofloxacin is comprised of a racemic mixture of the dextro- (microbiologically inactive) and levo-rotatory (levofloxacin-microbiologically active). Studies have shown that the pharmacokinetic properties of the 2 components are similar. In comparative studies with ciprofloxacin, ofloxacin had a longer half-life and higher distribution into sputum (79% vs 21%) than ciprofloxacin.

Lung Epithelial Lining Fluid

[00133] More recent emphasis on the use and development of fluoroquinolones

in community acquired gram-positive infections has focused on intrapulmonary pharmacokinetic studies in lung epithelial lining fluid (ELF). Although the relevance of drug distribution into this fluid is not clear in the setting of cystic fibrosis, insights in the drug pharmacology can be obtained from these studies. Levofloxacin penetrates well into lung tissues. Lung tissue concentrations are generally 2- to 5-fold higher than plasma concentrations. Several recent studies (summarized in Table 1) demonstrated that ELF concentrations of levofloxacin in healthy subjects following an oral dose of 750 mg reach a maximum concentration around 20 µg/mL. Similar peak concentrations are expected in the sputum of CF patients after oral or IV administration of 750 mg of levofloxacin. In contrast, ciprofloxacin penetrates lung tissues much less efficiently than levofloxacin. Based upon studies of the mutant selection window (MSW), these ELF fluoroquinolone drug levels are insufficient to achieve the required mutant prevention concentration of 10- to 20-fold the MIC for the infecting organism.

Table 1. Concentration of Levofloxacin in Epithelial Lining Fluid in Man.

Drug	Dose	Route	ELF Drug Concentration (µg/ml)						
			0.5 hr	1 hr	2 hr	4 hr	6 hr	12 hr	24 hr
levofloxacin	500 mg	IV				11		2.5	1.24
levofloxacin	500 mg	oral				9.9		6.5	0.7
levofloxacin	500 mg	oral				9.94		6.46	0.7
levofloxacin	500 mg	oral	4.74	10.8	9	10.9	9.6		
levofloxacin	750 mg	IV				12.94		6.04	1.73
levofloxacin	750 mg	oral				22.1		9.2	1.5
levofloxacin	750 mg	oral				22.13		9.19	1.55
ciprofloxacin	500 mg	oral				1.9		0.4	

Quinolones

[00134] Non-limiting examples of quinolones for use as described herein include amifloxacin, cinoxacin, ciprofloxacin, enoxacin, fleroxacin, flumequine, lomefloxacin, nalidixic acid, norfloxacin, ofloxacin, levofloxacin, lomefloxacin, oxolinic acid, pefloxacin, rosoxacin, temafloxacin, tosufloxacin, sparfloxacin, clinafloxacin, gatifloxacin, moxifloxacin; gemifloxacin; garenoxacin; olamufloxacin, clinofloxacin,

trovafloxacin, balofloxacin, prulifloxacin, moxifloxacin, gemifloxacin, rufloxacin, sitafloxacin (Sato, K, et al., 1992, Antimicrob Agents Chemother. 37:1491-98).

marbofloxacin, orbifloxacin, sarafloxacin, danofloxacin, difloxacin, enrofloxacin, TG-873870, DX-619, DW-276, ABT-492, DV-7751a (Tanaka, M, et al., 1992, Antimicrob. Agents Chemother. 37:2212-18), and F-1061 (Kurosaka et al., Interscience Conference on Antimicrobial Agents and Chemotherapy, 2003, 43rd:Chicago).

Methods of Treatment or Prophylaxis

[00135] In some embodiments, a method is provided for treating a microbial infection in an animal, specifically including in a mammal, by treating an animal suffering from such an infection with a fluoroquinolone antimicrobial. In some embodiments, fluoroquinolone antimicrobials may be administered following aerosol formation and inhalation. Thus, this method of treatment is especially appropriate for the treatment of pulmonary infections involving microbial strains that are difficult to treat using an antimicrobial agent delivered parenterally due to the need for high parenteral dose levels (which can cause undesirable side effects), or due to lack of any clinically effective antimicrobial agents. In one such embodiment, this method may be used to administer a fluoroquinolone antimicrobial directly to the site of infection. Such a method may reduce systemic exposure and maximizes the amount of antimicrobial agent to the site of microbial infection. This method is also appropriate for treating infections involving microbes that are susceptible to fluoroquinolone antimicrobials as a way of reducing the frequency of selection of resistant microbes. This method is also appropriate for treating infections involving microbes that are otherwise resistant to fluoroquinolone antimicrobials as a way of increasing the amount of antimicrobial at the site of microbial infection. A subject may be identified as infected with bacteria that are capable of developing resistance by diagnosing the subject as having symptoms that are characteristic of a bacterial infection with a bacteria species known to have resistant strains or a with a bacteria that is a member of group that are known to have resistant strains. Alternatively, the bacteria may be cultured and identified as a species known to

have resistant strains or a bacteria that is a member of group that are known to have resistant strains.

[00136] In some embodiments, the aerosol fluoroquinolone antimicrobial agent is administered at a level sufficient to overcome the emergence resistance in bacteria or increase killing efficiency such that resistance does not have the opportunity to develop.

[00137] In some embodiments, the aerosol fluoroquinolone therapy may be administered as a treatment or prophylaxis in combination or alternating therapeutic sequence with other aerosol, oral or parenteral antibiotics. By non-limiting example this may include aerosol tobramycin and/or other aminoglycoside, aerosol aztreonam and/or other beta or mono-bactam, aerosol ciprofloxacin and/or other fluoroquinolones, aerosol azithromycin and/or other macrolides or ketolides, tetracycline and/or other tetracyclines, quinupristin and/or other streptogramins, linezolid and/or other oxazolidinones, vancomycin and/or other glycopeptides, and chloramphenicol and/or other phenicols, and colistin and/or other polymyxins.

Pharmaceutical Compositions

[00138] For purposes of the method described herein, a fluoroquinolone antimicrobial agent may be administered using an inhaler. In some embodiments, a fluoroquinolone antimicrobial disclosed herein is produced as a pharmaceutical composition suitable for aerosol formation, good taste, storage stability, and patient safety and tolerability.

[00139] In some embodiments, the isoform content of the manufactured fluoroquinolone may be optimized for tolerability, antimicrobial activity and stability.

Administration

[00140] The fluoroquinolone antimicrobials disclosed herein can be administered at a therapeutically effective dosage, e.g., a dosage sufficient to provide treatment for the disease states previously described. While optimum human dosage levels have yet to be determined for aerosol delivery, generally a daily aerosol dose of levofloxacin (and for most fluoroquinolone antimicrobial agents described herein) is from about 0.1 to 10 mg/kg of body weight, preferably about 0.20 to 5.0 mg/kg of body weight, and most

preferably about 0.4 to 4.0 mg/kg of body weight. Thus, for administration to a 70 kg person, the dosage range would be about 7.0 to 700.0 mg per day, preferably about 14.0 to 350.0 mg per day, and most preferably about 28.0 to 280.0 mg per day. The amount of active compound administered will, of course, be dependent on the subject and disease state being treated, the severity of the affliction, the manner and schedule of administration, and the judgment of the prescribing physician; for example, a likely dose range for aerosol administration of levofloxacin would be about 20 to 400 mg per day.

[00141] Administration of the fluoroquinolone antimicrobial agents disclosed herein or the pharmaceutically acceptable salts thereof can be via any of the accepted modes of administration for agents that serve similar utilities including, but not limited to, aerosol inhalation.

[00142] Pharmaceutically acceptable compositions include solid, semi-solid, liquid and aerosol dosage forms, such as, e.g., powders, liquids, suspensions, complexations, liposomes, particulates, or the like. Preferably, the compositions are provided in unit dosage forms suitable for single administration of a precise dose. The unit dosage form can also be assembled and packaged together to provide a patient with a weekly or monthly supply and can also incorporate other compounds such as saline, taste masking agents, pharmaceutical excipients, and other active ingredients or carriers.

[00143] The fluoroquinolone antimicrobial agent can be administered either alone or more typically in combination with a conventional pharmaceutical carrier, excipient or the like (e.g., mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, sodium crosscarmellose, glucose, gelatin, sucrose, magnesium carbonate, magnesium chloride, magnesium sulfate, calcium chloride, lactose, sucrose, glucose and the like). If desired, the pharmaceutical composition can also contain minor amounts of nontoxic auxiliary substances such as wetting agents, emulsifying agents, solubilizing agents, pH buffering agents and the like (e.g., sodium acetate, sodium citrate, cyclodextrin derivatives, sorbitan monolaurate, triethanolamine acetate, triethanolamine oleate, and the like). Generally, depending on the intended mode of administration, the pharmaceutical formulation will contain about 0.005% to 95%, preferably about 0.5% to 50% by weight of a compound of the invention. Actual

methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pennsylvania.

[00144] In one preferred embodiment, the compositions will take the form of a unit dosage form such as vial containing a liquid, solid to be suspended, dry powder, lyophilisate, or other composition and thus the composition may contain, along with the active ingredient, a diluent such as lactose, sucrose, dicalcium phosphate, or the like; a lubricant such as magnesium stearate or the like; and a binder such as starch, gum acacia, polyvinylpyrrolidone, gelatin, cellulose, cellulose derivatives or the like.

[00145] Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, etc. an active compound as defined above and optional pharmaceutical adjuvants in a carrier (e.g., water, saline, aqueous dextrose, glycerol, glycols, ethanol or the like) to form a solution or suspension. Solutions to be aerosolized can be prepared in conventional forms, either as liquid solutions or suspensions, as emulsions, or in solid forms suitable for dissolution or suspension in liquid prior to aerosol production and inhalation. The percentage of active compound contained in such aerosol compositions is highly dependent on the specific nature thereof, as well as the activity of the compound and the needs of the subject. However, percentages of active ingredient of 0.01% to 90% in solution are employable, and will be higher if the composition is a solid, which will be subsequently diluted to the above percentages. In some embodiments, the composition will comprise 1.0%-50.0% of the active agent in solution.

[00146] Fluoroquinolone formulations can be separated into two groups; those of simple formulation and complex formulations providing taste-masking properties, improved tolerability and/or an AUC shape-enhancing formulation. Simple formulations can be further separated into three groups. 1. Simple formulations may include water-based liquid formulations for nebulization. By non-limiting example water-based liquid formulations may consist of fluoroquinolone alone or with non-encapsulating water soluble excipients. 2. Simple formulations may also include organic-based liquid formulations for nebulization or meter-dose inhaler. By non-limiting example organic-

based liquid formulations may consist of fluoroquinolone or with non-encapsulating organic soluble excipients. 3. Simple formulations may also include dry powder formulations for administration with a dry powder inhaler. By non-limiting example dry powder formulations may consist of fluoroquinolone alone or with either water soluble or organic soluble non-encapsulating excipients with or without a blending agent such as lactose. Complex formulations can be further separated into five groups. 1. Complex formulations may include water-based liquid formulations for nebulization. By non-limiting example water-based liquid complex formulations may consist of fluoroquinolone encapsulated or complexed with water-soluble excipients such as lipids, liposomes, cyclodextrins, microencapsulations, and emulsions. 2. Complex formulations may also include organic-based liquid formulations for nebulization or meter-dose inhaler. By non-limiting example organic-based liquid complex formulations may consist of fluoroquinolone encapsulated or complexed with organic-soluble excipients such as lipids, microencapsulations, and reverse-phase water-based emulsions. 3. Complex formulations may also include low-solubility, water-based liquid formulations for nebulization. By non-limiting example low-solubility, water-based liquid complex formulations may consist of fluoroquinolone as a low-water soluble, stable nanosuspension alone or in co-crystal/co-precipitate excipient complexes, or mixtures with low solubility lipids, such as lipid nanosuspensions. 4. Complex formulations may also include low-solubility, organic-based liquid formulations for nebulization or meter-dose inhaler. By non-limiting example low-solubility, organic-based liquid complex formulations may consist of fluoroquinolone as a low-organic soluble, stable nanosuspension alone or in co-crystal/co-precipitate excipient complexes, or mixtures with low solubility lipids, such as lipid nanosuspensions. 5. Complex formulations may also include dry powder formulations for administration using a dry powder inhaler. By non-limiting example, complex dry powder formulations may consist of fluoroquinolone in co-crystal/co-precipitate/spray dried complex or mixture with low-water soluble excipients/salts in dry powder form with or without a blending agent such as lactose. Specific methods for simple and complex formulation preparation are described herein.

Aerosol Delivery

[00147] Fluoroquinolone antimicrobial agents as described herein, are preferably directly administered as an aerosol to a site of infection in the respiratory tract. In some embodiments, aerosol delivery is used to treat an infection in the lungs, such as a *Pseudomonas* lung infection.

[00148] Several device technologies exist to deliver either dry powder or liquid aerosolized products. Dry powder formulations generally require less time for drug administration, yet longer and more expensive development efforts. Conversely, liquid formulations have historically suffered from longer administration times, yet have the advantage of shorter and less expensive development efforts. The fluoroquinolone antimicrobial agents disclosed herein range in solubility, are generally stable and have a range of tastes. In one such embodiment, the fluoroquinolone antimicrobial levofloxacin is water soluble at neutral pH, is stable in aqueous solution and has limited to no taste.

[00149] Accordingly, in one embodiment, a particular formulation of fluoroquinolone antimicrobial agent disclosed herein is combined with a particular aerosolizing device to provide an aerosol for inhalation that is optimized for maximum drug deposition at a site of infection and maximal tolerability. Factors that can be optimized include solution or solid particle formulation, rate of delivery, and particle size and distribution produced by the aerosolizing device.

Particle Size and Distribution

[00150] Generally, inhaled particles are subject to deposition by one of two mechanisms: impaction, which usually predominates for larger particles, and sedimentation, which is prevalent for smaller particles. Impaction occurs when the momentum of an inhaled particle is large enough that the particle does not follow the air stream and encounters a physiological surface. In contrast, sedimentation occurs primarily in the deep lung when very small particles which have traveled with the inhaled air stream encounter physiological surfaces as a result of random diffusion within the air stream.

[00151] For pulmonary administration, the upper airways are avoided in favor of the middle and lower airways. Pulmonary drug delivery may be accomplished by inhalation of an aerosol through the mouth and throat. Particles having a mass median aerodynamic diameter (MMAD) of greater than about 5 microns generally do not reach the lung; instead, they tend to impact the back of the throat and are swallowed and possibly orally absorbed. Particles having diameters of about 2 to about 5 microns are small enough to reach the upper- to mid-pulmonary region (conducting airways), but are too large to reach the alveoli. Smaller particles, i.e., about 0.5 to about 2 microns, are capable of reaching the alveolar region. Particles having diameters smaller than about 0.5 microns can also be deposited in the alveolar region by sedimentation, although very small particles may be exhaled. Measures of particle size can be referred to as volumetric mean diameter (VMD), mass median diameter (MMD), or MMAD. These measurements may be made by impaction (MMD and MMAD) or by laser (VMD). For liquid particles, VMD, MMD and MMAD may be the same if environmental conditions are maintained, e.g. standard humidity. However, if humidity is not maintained, MMD and MMAD determinations will be smaller than VMD due to dehydration during impaction measurements. For the purposes of this description, VMD, MMD and MMAD measurements are considered to be under standard conditions such that descriptions of VMD, MMD and MMAD will be comparable. Similarly, dry powder particle size determinations in MMD, and MMAD are also considered comparable.

[00152] In some embodiments, the particle size of the aerosol is optimized to maximize fluoroquinolone antimicrobial agent deposition at the site of infection and to maximize tolerability. Aerosol particle size may be expressed in terms of the mass median aerodynamic diameter (MMAD). Large particles (e.g., MMAD >5 μm) may deposit in the upper airway because they are too large to navigate the curvature of the upper airway. Small particles (e.g., MMAD < 2 μm) may be poorly deposited in the lower airways and thus become exhaled, providing additional opportunity for upper airway deposition. Hence, intolerability (e.g., cough and bronchospasm) may occur from upper airway deposition from both inhalation impaction of large particles and settling of small particles during repeated inhalation and expiration. Thus, in one embodiment, an optimum particle size is used (e.g., MMAD = 2-5 μm) in order to maximize deposition at

a mid-lung site of infection and to minimize intolerability associated with upper airway deposition. Moreover, generation of a defined particle size with limited geometric standard deviation (GSD) may optimize deposition and tolerability. Narrow GSD limits the number of particles outside the desired MMAD size range. In one embodiment, an aerosol containing one or more compounds disclosed herein is provided having a MMAD from about 2 microns to about 5 microns with a GSD of less than or equal to about 2.5 microns. In another embodiment, an aerosol having an MMAD from about 2.8 microns to about 4.3 microns with a GSD less than or equal to 2 microns is provided. In another embodiment, an aerosol having an MMAD from about 2.5 microns to about 4.5 microns with a GSD less than or equal to 1.8 microns is provided.

[00153] Fluoroquinolone antimicrobial agents disclosed herein intended for respiratory delivery (for either systemic or local distribution) can be administered as aqueous formulations, as suspensions or solutions in halogenated hydrocarbon propellants, or as dry powders. Aqueous formulations may be aerosolized by liquid nebulizers employing either hydraulic or ultrasonic atomization. Propellant-based systems may use suitable pressurized metered-dose inhalers (pMDIs). Dry powders may use dry powder inhaler devices (DPIs), which are capable of dispersing the drug substance effectively. A desired particle size and distribution may be obtained by choosing an appropriate device.

Liquid Nebulizer

[00154] In one embodiment, a nebulizer is selected on the basis of allowing the formation of an aerosol of a fluoroquinolone antimicrobial agent disclosed herein having an MMAD predominantly between about 2 to about 5 microns. In one embodiment, the delivered amount of fluoroquinolone antimicrobial agent provides a therapeutic effect for respiratory infections.

[00155] Previously, two types of nebulizers, jet and ultrasonic, have been shown to be able to produce and deliver aerosol particles having sizes between 2 and 4 μm . These particle sizes have been shown as being optimal for treatment of pulmonary bacterial infection cause by gram-negative bacteria such as *Pseudomonas aeruginosa*, *Escherichia coli*, *Enterobacter* species, *Klebsiella pneumoniae*, *K. oxytoca*, *Proteus*

mirabilis, *Pseudomonas aeruginosa*, *Serratia marcescens*, *Haemophilus influenzae*, *Burkholderia cepacia*, *Stenotrophomonas maltophilia*, *Alcaligenes xylosoxidans*, and multidrug resistant *Pseudomonas aeruginosa*. However, unless a specially formulated solution is used, these nebulizers typically need larger volumes to administer sufficient amount of drug to obtain a therapeutic effect. A jet nebulizer utilizes air pressure breakage of an aqueous solution into aerosol droplets. An ultrasonic nebulizer utilizes shearing of the aqueous solution by a piezoelectric crystal. Typically, however, the jet nebulizers are only about 10% efficient under clinical conditions, while the ultrasonic nebulizer is only about 5% efficient. The amount of pharmaceutical deposited and absorbed in the lungs is thus a fraction of the 10% in spite of the large amounts of the drug placed in the nebulizer.

[00156] Accordingly, in one embodiment, a vibrating mesh nebulizer is used to deliver an aerosol of the fluoroquinolone antimicrobial agent disclosed herein. A vibrating mesh nebulizer consists of a liquid storage container in fluid contact with a diaphragm and inhalation and exhalation valves. In one embodiment, about 1 to about 5 ml of the fluoroquinolone antimicrobial agent is placed in the storage container and the aerosol generator is engaged producing atomized aerosol of particle sizes selectively between about 1 and about 5 μm .

[00157] By non-limiting example, a fluoroquinolone antimicrobial agent disclosed herein is placed in a liquid nebulization inhaler and prepared in dosages to deliver from about 7 to about 700 mg from a dosing solution of about 1 to about 5 ml, preferably from about 14 to about 350 mg in about 1 to about 5 ml, and most preferably from about 28 to about 280 mg in about 1 to about 5 ml with MMAD particles sizes between about 2 to about 5 μm being produced.

[00158] By non-limiting example, a nebulized fluoroquinolone antimicrobial may be administered in the described respirable delivered dose in less than about 20 min, preferably less than about 10 min, more preferably less than about 7 min, more preferably less than about 5 min, more preferably less than about 3 min, and in some cases most preferable if less than about 2 min.

[00159] By non-limiting example, in other circumstances, a nebulized fluoroquinolone antimicrobial may achieve improved tolerability and/or exhibit an AUC shape-enhancing characteristic when administered over longer periods of time. Under these conditions, the described respirable delivered dose in more than about 2 min, preferably more than about 3 min, more preferably more than about 5 min, more preferably more than about 7 min, more preferably more than about 10 min, and in some cases most preferable from about 10 to about 20 min.

[00160] For aqueous and other non-pressurized liquid systems, a variety of nebulizers (including small volume nebulizers) are available to aerosolize the formulations. Compressor-driven nebulizers incorporate jet technology and use compressed air to generate the liquid aerosol. Such devices are commercially available from, for example, Healthdyne Technologies, Inc.; Invacare, Inc.; Mountain Medical Equipment, Inc.; Pari Respiratory, Inc.; Mada Medical, Inc.; Puritan-Bennet; Schuco, Inc., DeVilbiss Health Care, Inc.; and Hospitak, Inc. Ultrasonic nebulizers rely on mechanical energy in the form of vibration of a piezoelectric crystal to generate respirable liquid droplets and are commercially available from, for example, Omron Healthcare, Inc. and DeVilbiss Health Care, Inc. Vibrating mesh nebulizers rely upon either piezoelectric or mechanical pulses to respirable liquid droplets generate. Other examples of nebulizers for use with fluoroquinolone antimicrobial agents described herein are described in U.S. Patent Nos. 4,268,460; 4,253,468; 4,046,146; 3,826,255; 4,649,911; 4,510,929; 4,624,251; 5,164,740; 5,586,550; 5,758,637; 6,644,304; 6,338,443; 5,906,202; 5,934,272; 5,960,792; 5,971,951; 6,070,575; 6,192,876; 6,230,706; 6,349,719; 6,367,470; 6,543,442; 6,584,971; 6,601,581; 4,263,907; 5,709,202; 5,823,179; 6,192,876; 6,644,304; 5,549,102; 6,083,922; 6,161,536; 6,264,922; 6,557,549; and 6,612,303.

Commercial examples of nebulizers that can be used with the fluoroquinolone antimicrobial agents described herein include Respigard II®, Aeroneb®, Aeroneb® Pro, and Aeroneb® Go produced by Aerogen; AERx® and AERx Essence™ produced by Aradigm; Porta-Neb®, Freeway Freedom™, Sidestream,, Ventstream and I-neb produced by Respironics, Inc.; and PARI LC-Plus®, PARI LC-Star®, and e-Flow™ produced by PARI, GmbH. By further non-

limiting example, U.S. Patent No. 6,196,219 is hereby referenced.

[00161] In some embodiments, the drug solution is formed prior to use of the nebulizer by a patient. In other embodiments, the drug is stored in the nebulizer in solid form. In this case, the solution is mixed upon activation of the nebulizer, such as described in U.S. Patent No. 6,427,682 and PCT Publication No. WO 03/035030.

In these nebulizers, the solid drug, optionally combined with excipients to form a solid composition, is stored in a separate compartment from a liquid solvent.

[00162] The liquid solvent is capable of dissolving the solid composition to form a liquid composition, which can be aerosolized and inhaled. Such capability is, among other factors, a function of the selected amount and, potentially, the composition of the liquid. To allow easy handling and reproducible dosing, the sterile aqueous liquid may be able to dissolve the solid composition within a short period of time, possibly under gentle shaking. In some embodiments, the final liquid is ready to use after no longer than about 30 seconds. In some cases, the solid composition is dissolved within about 20 seconds, and advantageously, within about 10 seconds. As used herein, the terms "dissolve(d)", "dissolving", and "dissolution" refer to the disintegration of the solid composition and the release, i.e. the dissolution, of the active compound. As a result of dissolving the solid composition with the liquid solvent a liquid composition is formed in which the active compound is contained in the dissolved state. As used herein, the active compound is in the dissolved state when at least about 90 wt.-% are dissolved, and more preferably when at least about 95 wt.-% are dissolved.

[00163] With regard to basic separated-compartment nebulizer design, it primarily depends on the specific application whether it is more useful to accommodate the aqueous liquid and the solid composition within separate chambers of the same container or primary package, or whether they should be provided in separate containers. If separate containers are used, these are provided as a set within the same secondary package. The use of separate containers is especially preferred for nebulizers containing two or more doses of the active compound. There is no limit to the total number of

containers provided in a multi-dose kit. In one embodiment, the solid composition is provided as unit doses within multiple containers or within multiple chambers of a container, whereas the liquid solvent is provided within one chamber or container. In this case, a favorable design provides the liquid in a metered-dose dispenser, which may consist of a glass or plastic bottle closed with a dispensing device, such as a mechanical pump for metering the liquid. For instance, one actuation of the pumping mechanism may dispense the exact amount of liquid for dissolving one dose unit of the solid composition.

[00164] In another embodiment for multiple-dose separated-compartment nebulizers, both the solid composition and the liquid solvent are provided as matched unit doses within multiple containers or within multiple chambers of a container. For instance, two-chambered containers can be used to hold one unit of the solid composition in one of the chambers and one unit of liquid in the other. As used herein, one unit is defined by the amount of drug present in the solid composition, which is one unit dose. Such two-chambered containers may, however, also be used advantageously for nebulizers containing only one single drug dose.

[00165] In one embodiment of a separated-compartment nebulizer, a blister pack having two blisters is used, the blisters representing the chambers for containing the solid composition and the liquid solvent in matched quantities for preparing a dose unit of the final liquid composition. As used herein, a blister pack represents a thermoformed or pressure-formed primary packaging unit, most likely comprising a polymeric packaging material that optionally includes a metal foil, such as aluminum. The blister pack may be shaped to allow easy dispensing of the contents. For instance, one side of the pack may be tapered or have a tapered portion or region through which the content is dispensable into another vessel upon opening the blister pack at the tapered end. The tapered end may represent a tip.

[00166] In some embodiments, the two chambers of the blister pack are connected by a channel, the channel being adapted to direct fluid from the blister containing the liquid solvent to the blister containing the solid composition. During storage, the channel is closed with a seal. In this sense, a seal is any structure that

prevents the liquid solvent from contacting the solid composition. The seal is preferably breakable or removable; breaking or removing the seal when the nebulizer is to be used will allow the liquid solvent to enter the other chamber and dissolve the solid composition. The dissolution process may be improved by shaking the blister pack. Thus, the final liquid composition for inhalation is obtained, the liquid being present in one or both of the chambers of the pack connected by the channel, depending on how the pack is held.

[00167] According to another embodiment, one of the chambers, preferably the one that is closer to the tapered portion of the blister pack, communicates with a second channel, the channel extending from the chamber to a distal position of the tapered portion. During storage, this second channel does not communicate with the outside of the pack but is closed in an air-tight fashion. Optionally, the distal end of the second channel is closed by a breakable or removable cap or closure, which may e.g. be a twist-off cap, a break-off cap, or a cut-off cap.

[00168] In one embodiment, a vial or container having two compartments is used, the compartment representing the chambers for containing the solid composition and the liquid solvent in matched quantities for preparing a dose unit of the final liquid composition. The liquid composition and a second liquid solvent may be contained in matched quantities for preparing a dose unit of the final liquid composition (by non-limiting example in cases where two soluble excipients or the fluoroquinolone and an excipient are unstable for storage, yet desired in the same mixture for administration).

[00169] In some embodiments, the two compartments are physically separated but in fluid communication such as when so the vial or container are connected by a channel or breakable barrier, the channel or breakable barrier being adapted to direct fluid between the two compartments to enable mixing prior to administration. During storage, the channel is closed with a seal or the breakable barrier intact. In this sense, a seal is any structure that prevents mixing of contents in the two compartments. The seal is preferably breakable or removable; breaking or removing the seal when the nebulizer is to be used will allow the liquid solvent to enter the other chamber and dissolve the solid composition or in the case of two liquids permit mixing. The dissolution or mixing

process may be improved by shaking the container. Thus, the final liquid composition for inhalation is obtained, the liquid being present in one or both of the chambers of the pack connected by the channel or breakable barrier, depending on how the pack is held.

[00170] The solid composition itself can be provided in various different types of dosage forms, depending on the physicochemical properties of the drug, the desired dissolution rate, cost considerations, and other criteria. In one of the embodiments, the solid composition is a single unit. This implies that one unit dose of the drug is comprised in a single, physically shaped solid form or article. In other words, the solid composition is coherent, which is in contrast to a multiple unit dosage form, in which the units are incoherent.

[00171] Examples of single units which may be used as dosage forms for the solid composition include tablets, such as compressed tablets, film-like units, foil-like units, wafers, lyophilized matrix units, and the like. In a preferred embodiment, the solid composition is a highly porous lyophilized form. Such lyophilizates, sometimes also called wafers or lyophilized tablets, are particularly useful for their rapid disintegration, which also enables the rapid dissolution of the active compound.

[00172] On the other hand, for some applications the solid composition may also be formed as a multiple unit dosage form as defined above. Examples of multiple units are powders, granules, microparticles, pellets, beads, lyophilized powders, and the like. In one embodiment, the solid composition is a lyophilized powder. Such a dispersed lyophilized system comprises a multitude of powder particles, and due to the lyophilization process used in the formation of the powder, each particle has an irregular, porous microstructure through which the powder is capable of absorbing water very rapidly, resulting in quick dissolution.

[00173] Another type of multiparticulate system which is also capable of achieving rapid drug dissolution is that of powders, granules, or pellets from water-soluble excipients which are coated with the drug, so that the drug is located at the outer surface of the individual particles. In this type of system, the water-soluble low molecular weight excipient is useful for preparing the cores of such coated particles, which can be subsequently coated with a coating composition comprising the drug and,

preferably, one or more additional excipients, such as a binder, a pore former, a saccharide, a sugar alcohol, a film-forming polymer, a plasticizer, or other excipients used in pharmaceutical coating compositions.

[00174] In another embodiment, the solid composition resembles a coating layer that is coated on multiple units made of insoluble material. Examples of insoluble units include beads made of glass, polymers, metals, and mineral salts. Again, the desired effect is primarily rapid disintegration of the coating layer and quick drug dissolution, which is achieved by providing the solid composition in a physical form that has a particularly high surface-to-volume ratio. Typically, the coating composition will, in addition to the drug and the water-soluble low molecular weight excipient, comprise one or more excipients, such as those mentioned above for coating soluble particles, or any other excipient known to be useful in pharmaceutical coating compositions.

[00175] To achieve the desired effects, it may be useful to incorporate more than one water-soluble low molecular weight excipient into the solid composition. For instance, one excipient may be selected for its drug carrier and diluent capability, while another excipient may be selected to adjust the pH. If the final liquid composition needs to be buffered, two excipients that together form a buffer system may be selected.

[00176] In one embodiment, the liquid to be used in a separated-compartment nebulizer is an aqueous liquid, which is herein defined as a liquid whose major component is water. The liquid does not necessarily consist of water only; however, in one embodiment it is purified water. In another embodiment, the liquid contains other components or substances, preferably other liquid components, but possibly also dissolved solids. Liquid components other than water which may be useful include propylene glycol, glycerol, and polyethylene glycol. One of the reasons to incorporate a solid compound as a solute is that such a compound is desirable in the final liquid composition, but is incompatible with the solid composition or with a component thereof, such as the active ingredient.

[00177] Another desirable characteristic for the liquid solvent is that it is sterile. An aqueous liquid would be subject to the risk of considerable microbiological contamination and growth if no measures were taken to ensure sterility. In order to

provide a substantially sterile liquid, an effective amount of an acceptable antimicrobial agent or preservative can be incorporated or the liquid can be sterilized prior to providing it and to seal it with an air-tight seal. In one embodiment, the liquid is a sterilized liquid free of preservatives and provided in an appropriate air-tight container. However, according to another embodiment in which the nebulizer contains multiple doses of the active compound, the liquid may be supplied in a multiple-dose container, such as a metered-dose dispenser, and may require a preservative to prevent microbial contamination after the first use.

Meter Dose Inhaler (MDI)

[00178] A propellant driven inhaler (pMDI) releases a metered dose of medicine upon each actuation. The medicine is formulated as a suspension or solution of a drug substance in a suitable propellant such as a halogenated hydrocarbon. pMDIs are described in, for example, Newman, S. P., *Aerosols and the Lung*, Clarke et al., eds., pp. 197-224 (Butterworths, London, England, 1984).

[00179] In some embodiments, the particle size of the drug substance in an MDI may be optimally chosen. In some embodiments, the particles of active ingredient have diameters of less than about 50 microns. In some embodiments, the particles have diameters of less than about 10 microns. In some embodiments, the particles have diameters of from about 1 micron to about 5 microns. In some embodiments, the particles have diameters of less than about 1 micron. In one advantageous embodiment, the particles have diameters of from about 2 microns to about 5 microns.

[00180] The propellants for use with the MDIs may be any propellants known in the art. Examples of propellants include chlorofluorocarbons (CFCs) such as dichlorodifluoromethane, trichlorofluoromethane, and dichlorotetrafluoroethane; hydrofluoroalkanes (HFAs); and carbon dioxide. It may be advantageous to use HFAs instead of CFCs due to the environmental concerns associated with the use of CFCs. Examples of medicinal aerosol preparations containing HFAs are presented in U.S. Patent Nos. 6,585,958; 2,868,691 and 3,014,844.

In some embodiments, a co-solvent is mixed with the propellant to facilitate dissolution or suspension of the drug substance.

[00181] In some embodiments, the propellant and active ingredient are contained in separate containers, such as described in U.S. Patent No. 4,534,345.

[00182] In some embodiments, the MDI used herein is activated by a patient pushing a lever, button, or other actuator. In other embodiments, the release of the aerosol is breath activated such that, after initially arming the unit, the active compound aerosol is released once the patient begins to inhale, such as described in U.S. Patent Nos. 6,672,304; 5,404,871; 5,347,998; 5,284,133; 5,217,004; 5,119,806; 5,060,643; 4,664,107; 4,648,393; 3,789,843; 3,732,864; 3,636,949; 3,598,294; 3,565,070; 3,456,646; 3,456,645; and 3,456,644.

Such a system enables more of the active compound to get into the lungs of the patient. Another mechanism to help a patient get adequate dosage with the active ingredient may include a valve mechanism that allows a patient to use more than one breath to inhale the drug, such as described in U.S. Patent Nos. 4,470,412 and 5,385,140.

[00183] Additional examples of MDIs known in the art and suitable for use herein include U.S. Patent Nos. 6,435,177; 6,585,958; 5,642,730; 6,223,746; 4,955,371; 5,404,871; 5,364,838; and 6,523,536.

Dry Powder Inhaler (DPI)

[00184] There are two major designs of dry powder inhalers. One design is the metering device in which a reservoir for the drug is placed within the device and the patient adds a dose of the drug into the inhalation chamber. The second is a factory-metered device in which each individual dose has been manufactured in a separate container. Both systems depend upon the formulation of drug into small particles of mass median diameters from about 1 to about 5 μm , and usually involve co-formulation with larger excipient particles (typically 100 μm diameter lactose particles). Drug powder is placed into the inhalation chamber (either by device metering or by breakage of a factory-metered dosage) and the inspiratory flow of the patient accelerates the powder out of the device and into the oral cavity. Non-laminar flow characteristics of the

powder path cause the excipient-drug aggregates to decompose, and the mass of the large excipient particles causes their impaction at the back of the throat, while the smaller drug particles are deposited deep in the lungs.

[00185] As with liquid nebulization and MDIs, particle size of the fluoroquinolone antimicrobial agent aerosol formulation may be optimized. If the particle size is larger than about 5 μm MMAD then the particles are deposited in upper airways. If the particle size of the aerosol is smaller than about 1 μm then it is delivered into the alveoli and may get transferred into the systemic blood circulation.

[00186] By non-limiting example, in dry powder inhalers, the fluoroquinolone antimicrobial agents disclosed herein are prepared in dosages to deliver from about 7 to about 700 mg from a dosing solution of about 1 to about 5 ml, preferably from about 14 to about 350 mg in about 1 to about 5 ml, and most preferably from about 28 to about 280 mg in about 1 to about 5 ml with MMAD particles sizes between about 2 to about 5 μm being produced.

[00187] In some embodiments, a dry powder inhaler (DPI) is used to dispense the fluoroquinolone antimicrobial agents described herein. DPIs contain the drug substance in fine dry particle form. Typically, inhalation by a patient causes the dry particles to form an aerosol cloud that is drawn into the patient's lungs. The fine dry drug particles may be produced by any technique known in the art. Some well-known techniques include use of a jet mill or other comminution equipment, precipitation from saturated or super saturated solutions, spray drying, in situ micronization (Hovione), or supercritical fluid methods. Typical powder formulations include production of spherical pellets or adhesive mixtures. In adhesive mixtures, the drug particles are attached to larger carrier particles, such as lactose monohydrate of size about 50 to about 100 microns in diameter. The larger carrier particles increase the aerodynamic forces on the carrier/drug agglomerates to improve aerosol formation. Turbulence and/or mechanical devices break the agglomerates into their constituent parts. The smaller drug particles are then drawn into the lungs while the larger carrier particles deposit in the mouth or throat. Some examples of adhesive mixtures are described in U.S. Patent No. 5,478,578 and PCT Publication Nos. WO 95/11666, WO 87/05213, WO 96/23485, and WO 97/03649.

Additional excipients may also be included with the drug substance.

[00188] There are three common types of DPIs, all of which may be used with the fluoroquinolone antimicrobial agents described herein. In a single-dose DPI, a capsule containing one dose of dry drug substance/excipients is loaded into the inhaler. Upon activation, the capsule is breached, allowing the dry powder to be dispersed and inhaled using a dry powder inhaler. To dispense additional doses, the old capsule must be removed and an additional capsule loaded. Examples of single-dose DPIs are described in U.S. Patent Nos. 3,807,400; 3,906,950; 3,991,761; and 4,013,075.

In a multiple unit dose DPI, a package containing multiple single dose compartments is provided. For example, the package may comprise a blister pack, where each blister compartment contains one dose. Each dose can be dispensed upon breach of a blister compartment. Any of several arrangements of compartments in the package can be used. For example, rotary or strip arrangements are common. Examples of multiple unit dose DPIs are described in EPO Patent Application Publication Nos. 0211595A2, 0455463A1, and 0467172A1.

In a multi-dose DPI, a single reservoir of dry powder is used. Mechanisms are provided that measure out single dose amounts from the reservoir to be aerosolized and inhaled, such as described in U.S. Patent Nos. 5,829,434; 5,437,270; 2,587,215; 5,113,855; 5,840,279; 4,688,218; 4,667,668; 5,033,463; and 4,805,811 and PCT Publication No. WO 92/09322.

[00189] In some embodiments, auxiliary energy in addition to or other than a patient's inhalation may be provided to facilitate operation of a DPI. For example, pressurized air may be provided to aid in powder de-agglomeration, such as described in U.S. Patent Nos. 3,906,950; 5,113,855; 5,388,572; 6,029,662 and PCT Publication Nos. WO 93/12831, WO 90/07351, and WO 99/62495.

Electrically driven impellers may also be provided, such as described in U.S. Patent Nos. 3,948,264; 3,971,377; 4,147,166; 6,006,747 and PCT Publication No. WO 98/03217.

Another mechanism is an electrically powered tapping piston, such as described

in PCT Publication No. WO 90/13327.

Other DPIs use a vibrator, such as described in U.S. Patent Nos. 5,694,920 and 6,026,809.

Finally, a scraper system may be employed, such as described in PCT Publication No. WO 93/24165.

[00190] Additional examples of DPIs for use herein are described in U.S. Patent Nos. 4,811,731; 5,113,855; 5,840,279; 3,507,277; 3,669,113; 3,635,219; 3,991,761; 4,353,365; 4,889,144, 4,907,538; 5,829,434; 6,681,768; 6,561,186; 5,918,594; 6,003,512; 5,775,320; 5,740,794; and 6,626,173.

[00191] In some embodiments, a spacer or chamber may be used with any of the inhalers described herein to increase the amount of drug substance that gets absorbed by the patient, such as is described in U.S. Patent Nos. 4,470,412; 4,790,305; 4,926,852; 5,012,803; 5,040,527; 5,024,467; 5,816,240; 5,027,806; and 6,026,807.

For example, a spacer may delay the time from aerosol production to the time when the aerosol enters a patient's mouth. Such a delay may improve synchronization between the patient's inhalation and the aerosol production. A mask may also be incorporated for infants or other patients that have difficulty using the traditional mouthpiece, such as is described in U.S. Patent Nos. 4,809,692; 4,832,015; 5,012,804; 5,427,089; 5,645,049; and 5,988,160.

[00192] Dry powder inhalers (DPIs), which involve deaggregation and aerosolization of dry powders, normally rely upon a burst of inspired air that is drawn through the unit to deliver a drug dosage. Such devices are described in, for example, U.S. Pat. No. 4,807,814, which is directed to a pneumatic powder ejector having a suction stage and an injection stage; SU 628930 (Abstract), describing a hand-held powder disperser having an axial air flow tube; Fox et al., Powder and Bulk Engineering, pages 33-36 (March 1988), describing a venturi eductor having an axial air inlet tube upstream of a venturi restriction; EP 347 779, describing a hand-held powder disperser

having a collapsible expansion chamber, and U.S. Pat. No. 5,785,049, directed to dry powder delivery devices for drugs.

Solution/Dispersion Formulations

[00193] In one embodiment, aqueous formulations containing soluble or nanoparticulate drug particles are provided. For aqueous aerosol formulations, the drug may be present at a concentration of about 1 mg/mL up to about 700 mg/mL. Such formulations provide effective delivery to appropriate areas of the lung, with the more concentrated aerosol formulations having the additional advantage of enabling large quantities of drug substance to be delivered to the lung in a very short period of time. In one embodiment, a formulation is optimized to provide a well tolerated formulation. Accordingly, in one embodiment, fluoroquinolone antimicrobial agents disclosed herein are formulated to have good taste, pH from about 5.5 to about 7, osmolarity from about 200 to about 1250 mOsmol/kg, permeant ion concentration from about 30 to about 300 mM.

[00194] In one embodiment, the solution or diluent used for preparation of aerosol formulations has a pH range from about 4.5 to about 7.5, preferably from about 5.5 to about 7.0. This pH range improves tolerability. When the aerosol is either acidic or basic, it can cause bronchospasm and cough. Although the safe range of pH is relative and some patients may tolerate a mildly acidic aerosol, while others will experience bronchospasm. Any aerosol with a pH of less than about 4.5 typically induces bronchospasm. Aerosols with a pH from about 4.5 to about 5.5 will cause bronchospasm occasionally. Any aerosol having pH greater than about 7.5 may have low tolerability because body tissues are generally unable to buffer alkaline aerosols. Aerosols with controlled pH below about 4.5 and over about 7.5 typically result in lung irritation accompanied by severe bronchospasm cough and inflammatory reactions. For these reasons as well as for the avoidance of bronchospasm, cough or inflammation in patients, the optimum pH for the aerosol formulation was determined to be between about pH5.5 to about pH 7.0. Consequently, in one embodiment, aerosol formulations for use as described herein are adjusted to pH between about 4.5 and about 7.5 with preferred pH

range from about about 5.5 to about 7.5. Most preferred pH range is from about 5.5 to about 7.5.

[00195] By non-limiting example, compositions may also include a buffer or a pH adjusting agent, typically a salt prepared from an organic acid or base. Representative buffers include organic acid salts of citric acid, ascorbic acid, gluconic acid, carbonic acid, tartaric acid, succinic acid, acetic acid, or phthalic acid, Tris, tromethamine, hydrochloride, or phosphate buffers.

[00196] Many patients have increased sensitivity to various chemical tastes, including bitter, salt, sweet, metallic sensations. To create well-tolerated drug products, by non-limiting example taste masking may be accomplished through the addition of taste-masking excipients, adjusted osmolality, and sweeteners.

[00197] Many patients have increased sensitivity to various chemical agents and have high incidence of bronchospastic, asthmatic or other coughing incidents. Their airways are particularly sensitive to hypotonic or hypertonic and acidic or alkaline conditions and to the presence of any permanent ion, such as chloride. Any imbalance in these conditions or a presence of chloride above certain value leads to bronchospastic or inflammatory events and/or cough which greatly impair treatment with inhalable formulations. Both these conditions prevent efficient delivery of aerosolized drugs into the endobronchial space.

[00198] In some embodiments, the osmolality of aqueous solutions of the fluoroquinolone antimicrobial agent disclosed herein are adjusted by providing excipients. In some cases, a certain amount of chloride or another anion is needed for successful and efficacious delivery of aerosolized antibiotic. However, it has been discovered that such amounts are lower than amounts provided and typically used for aerosols of other compounds.

[00199] Bronchospasm or cough reflexes do not respond to the same osmolality of the diluent for aerosolization. However, they can be sufficiently controlled and/or suppressed when the osmolality of the diluent is in a certain range. A preferred solution for aerosolization of therapeutic compounds which is safe and tolerated has a total osmolality from about 200 to about 1250 mOsmol/kg with a range of chloride

concentration of from about 30 mM to about 300 mM and preferably from about 50 mM to about 150 mM. This osmolality controls bronchospasm, the chloride concentration, as a permeant anion, controls cough. Because they are both permeant ions, both bromine or iodine anions may be substituted for chloride. In addition, bicarbonate may substituted for chloride ion.

[00200] By non-limiting example, the formulation for an aerosol fluoroquinolone antimicrobial agent may comprise from about 7 to about 700 mg, preferably from about 14 to about 300 mg, or most preferably from about 28 to about 280 mg fluoroquinolone antimicrobial agent per about 1 to about 5ml of dilute saline (between 1/10 to 1/1 normal saline). Accordingly, the concentration of a levofloxacin solution may be greater than about 25mg/ml, greater than about 35 mg/ml and is preferably greater than about 40 mg/ml, and is as high or greater than 50/ml.

[00201] In one embodiment, solution osmolality is from about 100 mOsmol/kg to about 600 mOsmol/kg. In various other embodiments, the solution osmolality is from about 2000 mOsmol/kg to about 1250 mOsmol/kg; from about 250 mOsmol/kg to about 1050 mOsmol/kg; and from about 350 mOsmol/kg to about 750 mOsmol/kg.

[00202] In one embodiments, permeant ion concentration is from about 25 mM to about 400 mM. In various other embodiments, permeant ion concentration is from about 30 mM to about 300 mM; from about 40 mM to about 200 mM; and from about 50 mM to about 150 mM.

Solid Particle Formulations

[00203] In some embodiments, solid drug nanoparticles are provided for use in generating dry aerosols or for generating nanoparticles in liquid suspension. Powders comprising nanoparticulate drug can be made by spray-drying aqueous dispersions of a nanoparticulate drug and a surface modifier to form a dry powder which consists of aggregated drug nanoparticles. In one embodiment, the aggregates can have a size of about 1 to about 2 microns which is suitable for deep lung delivery. The aggregate particle size can be increased to target alternative delivery sites, such as the upper

bronchial region or nasal mucosa by increasing the concentration of drug in the spray-dried dispersion or by increasing the droplet size generated by the spray dryer.

[00204] Alternatively, an aqueous dispersion of drug and surface modifier can contain a dissolved diluent such as lactose or mannitol which, when spray dried, forms respirable diluent particles, each of which contains at least one embedded drug nanoparticle and surface modifier. The diluent particles with embedded drug can have a particle size of about 1 to about 2 microns, suitable for deep lung delivery. In addition, the diluent particle size can be increased to target alternate delivery sites, such as the upper bronchial region or nasal mucosa by increasing the concentration of dissolved diluent in the aqueous dispersion prior to spray drying, or by increasing the droplet size generated by the spray dryer.

[00205] Spray-dried powders can be used in DPIs or pMDIs, either alone or combined with freeze-dried nanoparticulate powder. In addition, spray-dried powders containing drug nanoparticles can be reconstituted and used in either jet or ultrasonic nebulizers to generate aqueous dispersions having respirable droplet sizes, where each droplet contains at least one drug nanoparticle. Concentrated nanoparticulate dispersions may also be used in these aspects of the invention.

[00206] Nanoparticulate drug dispersions can also be freeze-dried to obtain powders suitable for nasal or pulmonary delivery. Such powders may contain aggregated nanoparticulate drug particles having a surface modifier. Such aggregates may have sizes within a respirable range, e.g., about 2 to about 5 microns MMAD.

[00207] Freeze dried powders of the appropriate particle size can also be obtained by freeze drying aqueous dispersions of drug and surface modifier, which additionally contain a dissolved diluent such as lactose or mannitol. In these instances the freeze dried powders consist of respirable particles of diluent, each of which contains at least one embedded drug nanoparticle.

[00208] Freeze-dried powders can be used in DPIs or pMDIs, either alone or combined with spray-dried nanoparticulate powder. In addition, freeze-dried powders containing drug nanoparticles can be reconstituted and used in either jet or ultrasonic

nebulizers to generate aqueous dispersions that have respirable droplet sizes, where each droplet contains at least one drug nanoparticle.

[00209] One embodiment of the invention is directed to a process and composition for propellant-based systems comprising nanoparticulate drug particles and a surface modifier. Such formulations may be prepared by wet milling the coarse drug substance and surface modifier in liquid propellant, either at ambient pressure or under high pressure conditions. Alternatively, dry powders containing drug nanoparticles may be prepared by spray-drying or freeze-drying aqueous dispersions of drug nanoparticles and the resultant powders dispersed into suitable propellants for use in conventional pMDIs. Such nanoparticulate pMDI formulations can be used for either nasal or pulmonary delivery. For pulmonary administration, such formulations afford increased delivery to the deep lung regions because of the small (e.g., about 1 to about 2 microns MMAD) particle sizes available from these methods. Concentrated aerosol formulations can also be employed in pMDIs.

[00210] Another embodiment is directed to dry powders which contain nanoparticulate compositions for pulmonary or nasal delivery. The powders may consist of respirable aggregates of nanoparticulate drug particles, or of respirable particles of a diluent which contains at least one embedded drug nanoparticle. Powders containing nanoparticulate drug particles can be prepared from aqueous dispersions of nanoparticles by removing the water via spray-drying or lyophilization (freeze drying). Spray-drying is less time consuming and less expensive than freeze-drying, and therefore more cost-effective. However, certain drugs, such as biologicals benefit from lyophilization rather than spray-drying in making dry powder formulations.

[00211] Conventional micronized drug particles used in dry powder aerosol delivery having particle diameters of from about 2 to about 5 microns MMAD are often difficult to meter and disperse in small quantities because of the electrostatic cohesive forces inherent in such powders. These difficulties can lead to loss of drug substance to the delivery device as well as incomplete powder dispersion and sub-optimal delivery to the lung. Many drug compounds, particularly proteins and peptides, are intended for deep lung delivery and systemic absorption. Since the average particle sizes of

conventionally prepared dry powders are usually in the range of from about 2 to about 5 microns MMAD, the fraction of material which actually reaches the alveolar region may be quite small. Thus, delivery of micronized dry powders to the lung, especially the alveolar region, is generally very inefficient because of the properties of the powders themselves.

[00212] The dry powder aerosols which contain nanoparticulate drugs can be made smaller than comparable micronized drug substance and, therefore, are appropriate for efficient delivery to the deep lung. Moreover, aggregates of nanoparticulate drugs are spherical in geometry and have good flow properties, thereby aiding in dose metering and deposition of the administered composition in the lung or nasal cavities.

[00213] Dry nanoparticulate compositions can be used in both DPIs and pMDIs. As used herein, "dry" refers to a composition having less than about 5% water.

[00214] In one embodiment, compositions are provided containing nanoparticles which have an effective average particle size of less than about 1000 nm, more preferably less than about 400 nm, less than about 300 nm, less than about 250 nm, or less than about 200 nm, as measured by light-scattering methods. By "an effective average particle size of less than about 1000 nm" it is meant that at least 50% of the drug particles have a weight average particle size of less than about 1000 nm when measured by light scattering techniques. Preferably, at least 70% of the drug particles have an average particle size of less than about 1000 nm, more preferably at least 90% of the drug particles have an average particle size of less than about 1000 nm, and even more preferably at least about 95% of the particles have a weight average particle size of less than about 1000 nm.

[00215] For aqueous aerosol formulations, the nanoparticulate agent may be present at a concentration of about 5.0 mg/mL up to about 700 mg/mL. For dry powder aerosol formulations, the nanoparticulate agent may be present at a concentration of about 5.0 mg/g up to about 1000 mg/g, depending on the desired drug dosage. Concentrated nanoparticulate aerosols, defined as containing a nanoparticulate drug at a concentration of about 5.0 mg/mL up to about 700 mg/mL for aqueous aerosol formulations, and about 5.0 mg/g up to about 1000 mg/g for dry powder aerosol

formulations, are specifically provided. Such formulations provide effective delivery to appropriate areas of the lung or nasal cavities in short administration times, ie., less than about 3-15 seconds per dose as compared to administration times of up to 4 to 20 minutes as found in conventional pulmonary nebulizer therapies.

[00216] Nanoparticulate drug compositions for aerosol administration can be made by, for example, (1) nebulizing a dispersion of a nanoparticulate drug, obtained by either grinding or precipitation; (2) aerosolizing a dry powder of aggregates of nanoparticulate drug and surface modifier (the aerosolized composition may additionally contain a diluent); or (3) aerosolizing a suspension of nanoparticulate drug or drug aggregates in a non-aqueous propellant. The aggregates of nanoparticulate drug and surface modifier, which may additionally contain a diluent, can be made in a non-pressurized or a pressurized non-aqueous system. Concentrated aerosol formulations may also be made via such methods.

[00217] Milling of aqueous drug to obtain nanoparticulate drug may be performed by dispersing drug particles in a liquid dispersion medium and applying mechanical means in the presence of grinding media to reduce the particle size of the drug to the desired effective average particle size. The particles can be reduced in size in the presence of one or more surface modifiers. Alternatively, the particles can be contacted with one or more surface modifiers after attrition. Other compounds, such as a diluent, can be added to the drug/surface modifier composition during the size reduction process. Dispersions can be manufactured continuously or in a batch mode.

[00218] Another method of forming nanoparticle dispersion is by microprecipitation. This is a method of preparing stable dispersions of drugs in the presence of one or more surface modifiers and one or more colloid stability enhancing surface active agents free of any trace toxic solvents or solubilized heavy metal impurities. Such a method comprises, for example, (1) dissolving the drug in a suitable solvent with mixing; (2) adding the formulation from step (1) with mixing to a solution comprising at least one surface modifier to form a clear solution; and (3) precipitating the formulation from step (2) with mixing using an appropriate nonsolvent. The method can be followed by removal of any formed salt, if present, by dialysis or diafiltration and

concentration of the dispersion by conventional means. The resultant nanoparticulate drug dispersion can be utilized in liquid nebulizers or processed to form a dry powder for use in a DPI or pMDI.

[00219] In a non-aqueous, non-pressurized milling system, a non-aqueous liquid having a vapor pressure of about 1 atm or less at room temperature and in which the drug substance is essentially insoluble may be used as a wet milling medium to make a nanoparticulate drug composition. In such a process, a slurry of drug and surface modifier may be milled in the non-aqueous medium to generate nanoparticulate drug particles. Examples of suitable non-aqueous media include ethanol, trichloromonofluoromethane, (CFC-11), and dichlorotetrafluoroethane (CFC-114). An advantage of using CFC-11 is that it can be handled at only marginally cool room temperatures, whereas CFC-114 requires more controlled conditions to avoid evaporation. Upon completion of milling the liquid medium may be removed and recovered under vacuum or heating, resulting in a dry nanoparticulate composition. The dry composition may then be filled into a suitable container and charged with a final propellant. Exemplary final product propellants, which ideally do not contain chlorinated hydrocarbons, include HFA-134a (tetrafluoroethane) and HFA-227 (heptafluoropropane). While non-chlorinated propellants may be preferred for environmental reasons, chlorinated propellants may also be used in this aspect of the invention.

[00220] In a non-aqueous, pressurized milling system, a non-aqueous liquid medium having a vapor pressure significantly greater than 1 atm at room temperature may be used in the milling process to make nanoparticulate drug compositions. If the milling medium is a suitable halogenated hydrocarbon propellant, the resultant dispersion may be filled directly into a suitable pMDI container. Alternately, the milling medium can be removed and recovered under vacuum or heating to yield a dry nanoparticulate composition. This composition can then be filled into an appropriate container and charged with a suitable propellant for use in a pMDI.

[00221] Spray drying is a process used to obtain a powder containing nanoparticulate drug particles following particle size reduction of the drug in a liquid medium. In general, spray-drying may be used when the liquid medium has a vapor

pressure of less than about 1 atm at room temperature. A spray-dryer is a device which allows for liquid evaporation and drug powder collection. A liquid sample, either a solution or suspension, is fed into a spray nozzle. The nozzle generates droplets of the sample within a range of about 20 to about 100 μm in diameter which are then transported by a carrier gas into a drying chamber. The carrier gas temperature is typically from about 80 to about 200° C. The droplets are subjected to rapid liquid evaporation, leaving behind dry particles which are collected in a special reservoir beneath a cyclone apparatus.

[00222] If the liquid sample consists of an aqueous dispersion of nanoparticles and surface modifier, the collected product will consist of spherical aggregates of the nanoparticulate drug particles. If the liquid sample consists of an aqueous dispersion of nanoparticles in which an inert diluent material was dissolved (such as lactose or mannitol), the collected product will consist of diluent (e.g., lactose or mannitol) particles which contain embedded nanoparticulate drug particles. The final size of the collected product can be controlled and depends on the concentration of nanoparticulate drug and/or diluent in the liquid sample, as well as the droplet size produced by the spray-dryer nozzle. Collected products may be used in conventional DPIs for pulmonary or nasal delivery, dispersed in propellants for use in pMDIs, or the particles may be reconstituted in water for use in nebulizers.

[00223] In some instances it may be desirable to add an inert carrier to the spray-dried material to improve the metering properties of the final product. This may especially be the case when the spray dried powder is very small (less than about 5 μm) or when the intended dose is extremely small, whereby dose metering becomes difficult. In general, such carrier particles (also known as bulking agents) are too large to be delivered to the lung and simply impact the mouth and throat and are swallowed. Such carriers typically consist of sugars such as lactose, mannitol, or trehalose. Other inert materials, including polysaccharides and cellulose, may also be useful as carriers.

[00224] Spray-dried powders containing nanoparticulate drug particles may be used in conventional DPIs, dispersed in propellants for use in pMDIs, or reconstituted in a liquid medium for use with nebulizers.

[00225] For compounds that are denatured or destabilized by heat, such as compounds having a low melting point (i.e., about 70 to about 150° C.), or for example, biologics, sublimation is preferred over evaporation to obtain a dry powder nanoparticulate drug composition. This is because sublimation avoids the high process temperatures associated with spray-drying. In addition, sublimation, also known as freeze-drying or lyophilization, can increase the shelf stability of drug compounds, particularly for biological products. Freeze-dried particles can also be reconstituted and used in nebulizers. Aggregates of freeze-dried nanoparticulate drug particles can be blended with either dry powder intermediates or used alone in DPIs and pMDIs for either nasal or pulmonary delivery.

[00226] Sublimation involves freezing the product and subjecting the sample to strong vacuum conditions. This allows for the formed ice to be transformed directly from a solid state to a vapor state. Such a process is highly efficient and, therefore, provides greater yields than spray-drying. The resultant freeze-dried product contains drug and modifier(s). The drug is typically present in an aggregated state and can be used for inhalation alone (either pulmonary or nasal), in conjunction with diluent materials (lactose, mannitol, etc.), in DPIs or pMDIs, or reconstituted for use in a nebulizer.

Liposomal Compositions

[00227] In some embodiments, fluoroquinolone antimicrobial agents disclosed herein may be formulated into liposome particles, which can then be aerosolized for inhaled delivery. Lipids which are useful in the present invention can be any of a variety of lipids including both neutral lipids and charged lipids. Carrier systems having desirable properties can be prepared using appropriate combinations of lipids, targeting groups and circulation enhancers. Additionally, the compositions provided herein can be in the form of liposomes or lipid particles, preferably lipid particles. As used herein, the term "lipid particle" refers to a lipid bilayer carrier which "coats" a nucleic acid and has little or no aqueous interior. More particularly, the term is used to describe a self-assembling lipid bilayer carrier in which a portion of the interior layer comprises cationic lipids which form ionic bonds or ion-pairs with negative charges on the nucleic acid (e.g., a plasmid phosphodiester backbone). The interior layer can also comprise neutral or

fusogenic lipids and, in some embodiments, negatively charged lipids. The outer layer of the particle will typically comprise mixtures of lipids oriented in a tail-to-tail fashion (as in liposomes) with the hydrophobic tails of the interior layer. The polar head groups present on the lipids of the outer layer will form the external surface of the particle.

[00228] Liposomal bioactive agents can be designed to have a sustained therapeutic effect or lower toxicity allowing less frequent administration and an enhanced therapeutic index. Liposomes are composed of bilayers that entrap the desired pharmaceutical. These can be configured as multilamellar vesicles of concentric bilayers with the pharmaceutical trapped within either the lipid of the different layers or the aqueous space between the layers.

[00229] By non-limiting example, lipids used in the compositions may be synthetic, semi-synthetic or naturally-occurring lipids, including phospholipids, tocopherols, steroids, fatty acids, glycoproteins such as albumin, negatively-charged lipids and cationic lipids. Phospholipids include egg phosphatidylcholine (EPC), egg phosphatidylglycerol (EPG), egg phosphatidylinositol (EPI), egg phosphatidylserine (EPS), phosphatidylethanolamine (EPE), and egg phosphatidic acid (EPA); the soya counterparts, soy phosphatidylcholine (SPC); SPG, SPS, SPI, SPE, and SPA; the hydrogenated egg and soya counterparts (e.g., HEPC, HSPC), other phospholipids made up of ester linkages of fatty acids in the 2 and 3 of glycerol positions containing chains of 12 to 26 carbon atoms and different head groups in the 1 position of glycerol that include choline, glycerol, inositol, serine, ethanolamine, as well as the corresponding phosphatidic acids. The chains on these fatty acids can be saturated or unsaturated, and the phospholipid can be made up of fatty acids of different chain lengths and different degrees of unsaturation. In particular, the compositions of the formulations can include dipalmitoylphosphatidylcholine (DPPC), a major constituent of naturally-occurring lung surfactant as well as dioleoylphosphatidylcholine (DOPC) and dioleoylphosphatidylglycerol (DOPG). Other examples include dimyristoylphosphatidylcholine (DMPC) and dimyristoylphosphatidylglycerol (DMPG) dipalmitoylphosphatidylcholine (DPPC) and dipalmitoylphosphatidylglycerol (DPPG) distearoylphosphatidylcholine (DSPC) and distearoylphosphatidylglycerol (DSPG), dioleoylphosphatidylethanolamine (DOPE) and mixed phospholipids like

palmitoylstearylphosphatidylcholine (PSPC) and palmitoylstearylphosphatidylglycerol (PSPG), and single acylated phospholipids like mono-oleoyl-phosphatidylethanolamine (MOPE).

[00230] In a preferred embodiment, PEG-modified lipids are incorporated into the compositions of the present invention as the aggregation-preventing agent. The use of a PEG-modified lipid positions bulky PEG groups on the surface of the liposome or lipid carrier and prevents binding of DNA to the outside of the carrier (thereby inhibiting cross-linking and aggregation of the lipid carrier). The use of a PEG-ceramide is often preferred and has the additional advantages of stabilizing membrane bilayers and lengthening circulation lifetimes. Additionally, PEG-ceramides can be prepared with different lipid tail lengths to control the lifetime of the PEG-ceramide in the lipid bilayer. In this manner, "programmable" release can be accomplished which results in the control of lipid carrier fusion. For example, PEG-ceramides having C₂₀ -acyl groups attached to the ceramide moiety will diffuse out of a lipid bilayer carrier with a half-life of 22 hours. PEG-ceramides having C₁₄ - and C₈ -acyl groups will diffuse out of the same carrier with half-lives of 10 minutes and less than 1 minute, respectively. As a result, selection of lipid tail length provides a composition in which the bilayer becomes destabilized (and thus fusogenic) at a known rate. Though less preferred, other PEG-lipids or lipid-polyoxyethylene conjugates are useful in the present compositions. Examples of suitable PEG-modified lipids include PEG-modified phosphatidylethanolamine and phosphatidic acid, PEG-modified diacylglycerols and dialkylglycerols, PEG-modified dialkylamines and PEG-modified 1,2-diacyloxypropan-3-amines. Particularly preferred are PEG-ceramide conjugates (e.g., PEG-Cer-C₈, PEG-Cer-C₁₄ or PEG-Cer-C₂₀) which are described in U.S. Pat. No. 5,820,873 .

[00231] The compositions of the present invention can be prepared to provide liposome compositions which are about 50 nm to about 400 nm in diameter. One with skill in the art will understand that the size of the compositions can be larger or smaller depending upon the volume which is encapsulated. Thus, for larger volumes, the size distribution will typically be from about 80 nm to about 300 nm.

Surface Modifiers

[00232] Fluoroquinolone antimicrobial agents disclosed herein may be prepared in a pharmaceutical composition with suitable surface modifiers which may be selected from known organic and inorganic pharmaceutical excipients. Such excipients include low molecular weight oligomers, polymers, surfactants and natural products. Preferred surface modifiers include nonionic and ionic surfactants. Two or more surface modifiers can be used in combination.

[00233] Representative examples of surface modifiers include cetyl pyridinium chloride, gelatin, casein, lecithin (phosphatides), dextran, glycerol, gum acacia, cholesterol, tragacanth, stearic acid, benzalkonium chloride, calcium stearate, glycerol monostearate, cetostearyl alcohol, cetomacrogol emulsifying wax, sorbitan esters, polyoxyethylene alkyl ethers (e.g., macrogol ethers such as cetomacrogol 1000), polyoxyethylene castor oil derivatives, polyoxyethylene sorbitan fatty acid esters (e.g., the commercially available Tweens.RTM, such as e.g., Tween 20.RTM, and Tween 80.RTM, (ICI Specialty Chemicals)); polyethylene glycols (e.g., Carbowaxs 3350.RTM, and 1450.RTM., and Carbopol 934.RTM, (Union Carbide)), dodecyl trimethyl ammonium bromide, polyoxyethylenestearates, colloidal silicon dioxide, phosphates, sodium dodecylsulfate, carboxymethylcellulose calcium, hydroxypropyl cellulose (HPC, HPC-SL, and HPC-L), hydroxypropyl methylcellulose (HPMC), carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethyl-cellulose phthalate, noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol (PVA), polyvinylpyrrolidone (PVP), 4-(1,1,3,3-tetraamethylbutyl)-phenol polymer with ethylene oxide and formaldehyde (also known as tyloxapol, superione, and triton), poloxamers (e.g., Pluronics F68.RTM, and F108.RTM., which are block copolymers of ethylene oxide and propylene oxide); poloxamnines (e.g., Tetronic 908.RTM., also known as Poloxamine 908.RTM., which is a tetrafunctional block copolymer derived from sequential addition of propylene oxide and ethylene oxide to ethylenediamine (BASF Wyandotte Corporation, Parsippany, N.J.)); a charged phospholipid such as dimyristoyl phosphatidyl glycerol, dioctylsulfosuccinate (DOSS); Tetronic 1508.RTM; (T-1508) (BASF Wyandotte Corporation), dialkylesters of sodium sulfosuccinic acid (e.g., Aerosol OT.RTM., which

is a dioctyl ester of sodium sulfosuccinic acid (American Cyanamid)); Duponol P.RTM., which is a sodium lauryl sulfate (DuPont); Tritons X-200.RTM., which is an alkyl aryl polyether sulfonate (Rohm and Haas); Crodestas F-110.RTM., which is a mixture of sucrose stearate and sucrose distearate (Croda Inc.); p-isononylphenoxypoly-(glycidol), also known as Olin-log.RTM, or Surfactant 10-G.RTM, (Olin Chemicals, Stamford, Conn.); Crodestas SL-40.RTM, (Croda, Inc.); and SA9OHCO, which is C.sub.18 H.sub.37 CH.sub.2 (CON(CH.sub.3)-CH.sub.2 (CHOH).sub.4 (CH.sub.2 OH).sub.2 (Eastman Kodak Co.); decanoyl-N-methylglucamide; n-decyl .beta.-D-glucopyranoside; n-decyl .beta.-D-maltopyranoside; n-dodecyl .beta.-D-glucopyranoside; n-dodecyl .beta.-D-maltoside; heptanoyl-N-methylglucamide; n-heptyl-.beta.-D-glucopyranoside; n-heptyl .beta.-D-thioglucoside; n-hexyl .beta.-D-glucopyranoside; nonanoyl-N-methylglucamide; n-nonyl .beta.-D-glucopyranoside; octanoyl-N-methylglucamide; n-octyl-.beta.-D-glucopyranoside; octyl .beta.-D-thioglucopyranoside; and the like. Tyloxapol is a particularly preferred surface modifier for the pulmonary or intranasal delivery of steroids, even more so for nebulization therapies.

[00234] Examples of surfactants for use in the solutions disclosed herein include, but are not limited to, ammonium laureth sulfate, cetamine oxide, cetrimonium chloride, cetyl alcohol, cetyl myristate, cetyl palmitate, cocamide DEA, cocamidopropyl betaine, cocamidopropylamine oxide, cocamide MEA, DEA lauryl sulfate, di-stearyl phthalic acid amide, dicetyl dimethyl ammonium chloride, dipalmitoylethyl hydroxethylmonium, disodium laureth sulfosuccinate, di(hydrogenated) tallow phthalic acid, glyceryl dilaurate, glyceryl distearate, glyceryl oleate, glyceryl stearate, isopropyl myristate nf, isopropyl palmitate nf, lauramide DEA, lauramide MEA, lauramide oxide, myristamine oxide, octyl isononanoate, octyl palmitate, octyldodecyl neopentanoate, olealkonium chloride, PEG-2 stearate, PEG-32 glyceryl caprylate/caprinate, PEG-32 glyceryl stearate, PEG-4 and PEG-150 stearate & distearate, PEG-4 to PEG-150 laurate & dilaurate, PEG-4 to PEG-150 oleate & dioleate, PEG-7 glyceryl cocoate, PEG-8 beeswax, propylene glycol stearate, sodium C14-16 olefin sulfonate, sodium lauryl sulfoacetate, sodium lauryl sulphate, sodium trideceth sulfate, stearammonium chloride, stearamide oxide, TEA-dodecylbenzene sulfonate, TEA lauryl sulfate

[00235] Most of these surface modifiers are known pharmaceutical excipients and are described in detail in the Handbook of Pharmaceutical Excipients, published jointly by the American Pharmaceutical Association and The Pharmaceutical Society of Great Britain (The Pharmaceutical Press, 1986).

The surface modifiers are commercially available and/or can be prepared by techniques known in the art. The relative amount of drug and surface modifier can vary widely and the optimal amount of the surface modifier can depend upon, for example, the particular drug and surface modifier selected, the critical micelle concentration of the surface modifier if it forms micelles, the hydrophilic-lipophilic-balance (HLB) of the surface modifier, the melting point of the surface modifier, the water solubility of the surface modifier and/or drug, the surface tension of water solutions of the surface modifier, etc.

[00236] In the present invention, the optimal ratio of drug to surface modifier is ~0.1% to ~99.9% fluoroquinolone antimicrobial agent, more preferably about 10% to about 90%.

Microspheres

[00237] Microspheres can be used for pulmonary delivery of fluoroquinolones by first adding an appropriate amount of drug compound to be solubilized in water. For example, an aqueous levofloxacin solution may be dispersed in methylene chloride containing a predetermined amount (0.1-1% w/v) of poly(DL-lactide-co-glycolide) (PLGA) by probe sonication for 1-3 min on an ice bath. Separately, levofloxacin will be solubilized in methylene chloride containing PLGA (0.1-1% w/v). The resulting water-in-oil primary emulsion or the polymer/drug solution will be dispersed in an aqueous continuous phase consisting of 1-2% polyvinyl alcohol (previously cooled to 4°C) by probe sonication for 3-5 min on an ice bath. The resulting emulsion will be stirred continuously for 2-4 hours at room temperature to evaporate methylene chloride. Microparticles thus formed will be separated from the continuous phase by centrifuging at 8000-10000 rpm for 5-10 min. Sedimented particles will be washed thrice with distilled water and freeze dried. Freeze-dried levofloxacin microparticles will be stored at -20°C.

[00238] By non-limiting example, a spray drying approach will be employed to prepare levofloxacin microspheres. An appropriate amount of levofloxacin will be solubilized in methylene chloride containing PLGA (0.1-1%). This solution will be spray dried to obtain the microspheres.

[00239] By non-limiting example, levofloxacin microparticles will be characterized for size distribution (requirement: 90% <5 μm , 95% <10 μm), shape, drug loading efficiency and drug release using appropriate techniques and methods.

[00240] By non-limiting example, this approach may also be used to sequester and improve the water solubility of solid, AUC shape-enhancing formulations, such as low-solubility levofloxacin salt forms for nanoparticle-based formulations.

[00241] A certain amount of fluoroquinolone can be first dissolved in the minimal quantity of ethanol 96% necessary to maintain the fluoroquinolone in solution when diluted with water from 96 to 75%. This solution can then be diluted with water to obtain a 75% ethanol solution and then a certain amount of paracetamol can be added to obtain the following w/w drug/polymer ratios: 1:2, 1:1, 2:1, 3:1, 4:1, 6:1, 9:1, and 19:1. These final solutions are spray-dried under the following conditions: feed rate, 15 mL/min; inlet temperature, 110°C; outlet temperature, 85°C; pressure 4 bar and throughput of drying air, 35m³/hr. Powder is then collected and stored under vacuum in a desiccator.

Solid Lipid Particles

[00242] Preparation of fluoroquinolone solid lipid particles may involve dissolving the drug in a lipid melt (phospholipids such as phosphatidyl choline and phosphatidyl serine) maintained at least at the melting temperature of the lipid, followed by dispersion of the drug-containing melt in a hot aqueous surfactant solution (typically 1-5% w/v) maintained at least at the melting temperature of the lipid. The coarse dispersion will be homogenized for 1-10 min using a Microfluidizer® to obtain a nanoemulsion. Cooling the nanoemulsion to a temperature between 4-25°C will re-solidify the lipid, leading to formation of solid lipid nanoparticles. Optimization of formulation parameters (type of lipid matrix, surfactant concentration and production

parameters) will be performed so as to achieve a prolonged drug delivery. By non-limiting example, this approach may also be used to sequester and improve the water solubility of solid, AUC shape-enhancing formulations, such as low-solubility levofloxacin salt forms for nanoparticle-based formulations.

Melt-Extrusion AUC Shape-Enhancing Formulation

[00243] Melt-Extrusion AUC shape-enhancing fluoroquinolone formulations may be prepared by dissolving the drugs in micelles by adding surfactants or preparing micro-emulsion, forming inclusion complexes with other molecules such as cyclodextrins, forming nanoparticles of the drugs, or embedding the amorphous drugs in a polymer matrix. Embedding the drug homogeneously in a polymer matrix produces a solid dispersion. Solid dispersions can be prepared in two ways: the solvent method and the hot melt method. The solvent method uses an organic solvent wherein the drug and appropriate polymer are dissolved and then (spray) dried. The major drawbacks of this method are the use of organic solvents and the batch mode production process. The hot melt method uses heat in order to disperse or dissolve the drug in an appropriate polymer. The melt-extrusion process is an optimized version of the hot melt method. The advantage of the melt-extrusion approach is lack of organic solvent and continuous production process. As the melt-extrusion is a novel pharmaceutical technique, the literature dealing with it is limited. The technical set-up involves a mixture and extrusion of fluoroquinolone, hydroxypropyl- β -cyclodextrin (HP- β -CD), and hydroxypropylmethylcellulose (HPMC), in order to, by non-limiting example create a AUC shape-enhancing formulation of levofloxacin or other fluoroquinolone. Cyclodextrin is a toroidal-shaped molecule with hydroxyl groups on the outer surface and a cavity in the center. Cyclodextrin sequesters the drug by forming an inclusion complex. The complex formation between cyclodextrins and drugs has been investigated extensively. It is known that water-soluble polymer interacts with cyclodextrin and drug in the course of complex formation to form a stabilized complex of drug and cyclodextrin co-complexed with the polymer. This complex is more stable than the classic cyclodextrin-drug complex. As one example, HPMC is water soluble; hence using this polymer with HP- β -CD in the melt is expected to create an aqueous soluble AUC shape-enhancing

formulation. By non-limiting example, this approach may also be used to sequester and improve the water solubility of solid, AUC shape-enhancing formulations, such as low-solubility levofloxacin salt forms for nanoparticle-based formulations.

Co-Precipitates

[00244] Co-precipitate fluoroquinolone formulations may be prepared by formation of co-precipitates with pharmacologically inert, polymeric materials. It has been demonstrated that the formation of molecular solid dispersions or co-precipitates to create an AUC shape-enhancing formulations with various water-soluble polymers can significantly slow their *in vitro* dissolution rates and/or *in vivo* absorption. In preparing powdered products, grinding is generally used for reducing particle size, since the dissolution rate is strongly affected by particle size. Moreover, a strong force (such as grinding) may increase the surface energy and cause distortion of the crystal lattice as well as reducing particle size. Co-grinding drug with hydroxypropylmethylcellulose, β -cyclodextrin, chitin and chitosan, crystalline cellulose, and gelatine, may enhance the dissolution properties such that AUC shape-enhancement is obtained for otherwise readily bioavailable fluoroquinolones. By non-limiting example, this approach may also be used to sequester and improve the water solubility of solid, AUC shape-enhancing formulations, such as low-solubility levofloxacin salt forms for nanoparticle-based formulations.

Dispersion-Enhancing Peptides

[00245] Compositions may include one or more di- or tripeptides containing two or more leucine residues. By further non-limiting example, U.S. Patent No. 6,835,372 disclosing dispersion-enhancing peptides, is hereby referenced.

This patent describes the discovery that di-leucyl-containing dipeptides (e.g., dileucine) and tripeptides are superior in their ability to increase the dispersibility of powdered composition.

[00246] In another embodiment, highly dispersible particles including an amino acid are administered. Hydrophobic amino acids are preferred. Suitable amino acids include naturally occurring and non-naturally occurring hydrophobic amino acids. Some

naturally occurring hydrophobic amino acids, including but not limited to, non-naturally occurring amino acids include, for example, beta-amino acids. Both D, L and racemic configurations of hydrophobic amino acids can be employed. Suitable hydrophobic amino acids can also include amino acid analogs. As used herein, an amino acid analog includes the D or L configuration of an amino acid having the following formula: --NH--CHR--CO--, wherein R is an aliphatic group, a substituted aliphatic group, a benzyl group, a substituted benzyl group, an aromatic group or a substituted aromatic group and wherein R does not correspond to the side chain of a naturally-occurring amino acid. As used herein, aliphatic groups include straight chained, branched or cyclic C1-C8 hydrocarbons which are completely saturated, which contain one or two heteroatoms such as nitrogen, oxygen or sulfur and/or which contain one or more units of desaturation. Aromatic groups include carbocyclic aromatic groups such as phenyl and naphthyl and heterocyclic aromatic groups such as imidazolyl, indolyl, thienyl, furanyl, pyridyl, pyranyl, oxazolyl, benzothienyl, benzofuranyl, quinolinyl, isoquinolinyl and acridintyl.

[00247] Suitable substituents on an aliphatic, aromatic or benzyl group include --OH, halogen (--Br,--Cl,--I and --F)--O(aliphatic, substituted aliphatic, benzyl, substituted benzyl, aryl or substituted aryl group),--CN, --NO₂, --COOH, --NH₂, --NH(aliphatic group, substituted aliphatic, benzyl, substituted benzyl, aryl or substituted aryl group), --N(aliphatic group, substituted aliphatic, benzyl, substituted benzyl, aryl or substituted aryl group)₂, --COO(aliphatic group, substituted aliphatic, benzyl, substituted benzyl, aryl or substituted aryl group), --CONH₂, --CONH(aliphatic, substituted aliphatic group, benzyl, substituted benzyl, aryl or substituted aryl group)), --SH,--S(aliphatic, substituted aliphatic, benzyl, substituted benzyl, aromatic or substituted aromatic group) and --NH--C(=NH)--NH₂. A substituted benzylic or aromatic group can also have an aliphatic or substituted aliphatic group as a substituent. A substituted aliphatic group can also have a benzyl, substituted benzyl, aryl or substituted aryl group as a substituent. A substituted aliphatic, substituted aromatic or substituted benzyl group can have one or more substituents. Modifying an amino acid substituent can increase, for example, the lipophilicity or hydrophobicity of natural amino acids which are hydrophilic.

[00248] A number of the suitable amino acids, amino acids analogs and salts thereof can be obtained commercially. Others can be synthesized by methods known in the art.

[00249] Hydrophobicity is generally defined with respect to the partition of an amino acid between a nonpolar solvent and water. Hydrophobic amino acids are those acids which show a preference for the nonpolar solvent. Relative hydrophobicity of amino acids can be expressed on a hydrophobicity scale on which glycine has the value 0.5. On such a scale, amino acids which have a preference for water have values below 0.5 and those that have a preference for nonpolar solvents have a value above 0.5. As used herein, the term hydrophobic amino acid refers to an amino acid that, on the hydrophobicity scale, has a value greater or equal to 0.5, in other words, has a tendency to partition in the nonpolar acid which is at least equal to that of glycine.

[00250] Examples of amino acids which can be employed include, but are not limited to: glycine, proline, alanine, cysteine, methionine, valine, leucine, tyrosine, isoleucine, phenylalanine, tryptophan. Preferred hydrophobic amino acids include leucine, isoleucine, alanine, valine, phenylalanine and glycine. Combinations of hydrophobic amino acids can also be employed. Furthermore, combinations of hydrophobic and hydrophilic (preferentially partitioning in water) amino acids, where the overall combination is hydrophobic, can also be employed.

[00251] The amino acid can be present in the particles of the invention in an amount of at least 10 weight %. Preferably, the amino acid can be present in the particles in an amount ranging from about 20 to about 80 weight %. The salt of a hydrophobic amino acid can be present in the particles of the invention in an amount of at least 10 weight percent. Preferably, the amino acid salt is present in the particles in an amount ranging from about 20 to about 80 weight %. In preferred embodiments the particles have a tap density of less than about 0.4 g/cm.³.

[00252] Methods of forming and delivering particles which include an amino acid are described in U.S. Patent No. 6,586,008, entitled Use of Simple Amino Acids to Form Porous Particles During Spray Drying.

Proteins/Amino Acids

[00253] Protein excipients may include albumins such as human serum albumin (HSA), recombinant human albumin (rHA), gelatin, casein, hemoglobin, and the like. Suitable amino acids (outside of the dileucyl-peptides of the invention), which may also function in a buffering capacity, include alanine, glycine, arginine, betaine, histidine, glutamic acid, aspartic acid, cysteine, lysine, leucine, isoleucine, valine, methionine, phenylalanine, aspartame, tyrosine, tryptophan, and the like. Preferred are amino acids and polypeptides that function as dispersing agents. Amino acids falling into this category include hydrophobic amino acids such as leucine, valine, isoleucine, tryptophan, alanine, methionine, phenylalanine, tyrosine, histidine, and proline. Dispersibility-enhancing peptide excipients include dimers, trimers, tetramers, and pentamers comprising one or more hydrophobic amino acid components such as those described above.

Carbohydrates

[00254] By non-limiting example, carbohydrate excipients may include monosaccharides such as fructose, maltose, galactose, glucose, D-mannose, sorbose, and the like; disaccharides, such as lactose, sucrose, trehalose, cellobiose, and the like; polysaccharides, such as raffinose, melezitose, maltodextrins, dextrans, starches, and the like; and alditols, such as mannitol, xylitol, maltitol, lactitol, xylitol sorbitol (glucitol), pyranosyl sorbitol, myoinositol, isomalt, trehalose and the like.

Polymers

[00255] By non-limiting example, compositions may also include polymeric excipients/additives, e.g., polyvinylpyrrolidones, derivatized celluloses such as hydroxymethylcellulose, hydroxyethylcellulose, and hydroxypropylmethylcellulose, Ficolls (a polymeric sugar), hydroxyethylstarch, dextrans (by non-limiting example cyclodextrins may include, 2-hydroxypropyl-beta-cyclodextrin, 2-hydroxypropyl-gamma-cyclodextrin, randomly methylated beta-cyclodextrin, dimethyl-alpha-cyclodextrin, dimethyl-beta-cyclodextrin, maltosyl-alpha-cyclodextrin, glucosyl-1-alpha-cyclodextrin, glucosyl-2-alpha-cyclodextrin, alpha-cyclodextrin, beta-cyclodextrin,

gamma-cyclodextrin, and sulfobutylether-beta-cyclodextrin), polyethylene glycols, and pectin may also be used.

[00256] Highly dispersible particles administered comprise a bioactive agent and a biocompatible, and preferably biodegradable polymer, copolymer, or blend. The polymers may be tailored to optimize different characteristics of the particle including: i) interactions between the agent to be delivered and the polymer to provide stabilization of the agent and retention of activity upon delivery; ii) rate of polymer degradation and, thereby, rate of drug release profiles; iii) surface characteristics and targeting capabilities via chemical modification; and iv) particle porosity.

[00257] Surface eroding polymers such as polyanhydrides may be used to form the particles. For example, polyanhydrides such as poly[(p-carboxyphenoxy)hexane anhydride] (PCPH) may be used. Biodegradable polyanhydrides are described in U.S. Pat. No. 4,857,311. Bulk eroding polymers such as those based on polyesters including poly(hydroxy acids) also can be used. For example, polyglycolic acid (PGA), polylactic acid (PLA), or copolymers thereof may be used to form the particles. The polyester may also have a charged or functionalizable group, such as an amino acid. In a preferred embodiment, particles with controlled release properties can be formed of poly(D,L-lactic acid) and/or poly(DL-lactic-co-glycolic acid) ("PLGA") which incorporate a surfactant such as dipalmitoyl phosphatidylcholine (DPPC).

[00258] Other polymers include polyamides, polycarbonates, polyalkylenes such as polyethylene, polypropylene, poly(ethylene glycol), poly(ethylene oxide), poly(ethylene terephthalate), poly vinyl compounds such as polyvinyl alcohols, polyvinyl ethers, and polyvinyl esters, polymers of acrylic and methacrylic acids, celluloses and other polysaccharides, and peptides or proteins, or copolymers or blends thereof. Polymers may be selected with or modified to have the appropriate stability and degradation rates *in vivo* for different controlled drug delivery applications.

[00259] Highly dispersible particles can be formed from functionalized polyester graft copolymers, as described in Hrkach et al., *Macromolecules*, 28: 4736-4739 (1995); and Hrkach et al., "Poly(L-Lactic acid-co-amino acid) Graft Copolymers: A Class of Functional Biodegradable Biomaterials" in *Hydrogels and Biodegradable Polymers for*

Bioapplications, ACS Symposium Series No. 627, Raphael M, Ottenbrite et al., Eds., American Chemical Society, Chapter 8, pp. 93-101, 1996.

[00260] In a preferred embodiment of the invention, highly dispersible particles including a bioactive agent and a phospholipid are administered. Examples of suitable phospholipids include, among others, phosphatidylcholines, phosphatidylethanolamines, phosphatidylglycerols, phosphatidylserines, phosphatidylinositols and combinations thereof. Specific examples of phospholipids include but are not limited to phosphatidylcholines dipalmitoyl phosphatidylcholine (DPPC), dipalmitoyl phosphatidylethanolamine (DPPE), distearoyl phosphatidylcholine (DSPC), dipalmitoyl phosphatidyl glycerol (DPPG) or any combination thereof. Other phospholipids are known to those skilled in the art. In a preferred embodiment, the phospholipids are endogenous to the lung.

[00261] The phospholipid, can be present in the particles in an amount ranging from about 0 to about 90 weight %. More commonly it can be present in the particles in an amount ranging from about 10 to about 60 weight %.

[00262] In another embodiment of the invention, the phospholipids or combinations thereof are selected to impart controlled release properties to the highly dispersible particles. The phase transition temperature of a specific phospholipid can be below, around or above the physiological body temperature of a patient. Preferred phase transition temperatures range from 30 degrees C to 50 degrees C (e.g., within +/-10 degrees of the normal body temperature of patient). By selecting phospholipids or combinations of phospholipids according to their phase transition temperature, the particles can be tailored to have controlled release properties. For example, by administering particles which include a phospholipid or combination of phospholipids which have a phase transition temperature higher than the patient's body temperature, the release of dopamine precursor, agonist or any combination of precursors and/or agonists can be slowed down. On the other hand, rapid release can be obtained by including in the particles phospholipids having lower transition temperatures.

Taste Masking, Flavor, Other

[00263] By non-limiting example, compositions may further include flavoring agents, taste-masking agents, inorganic salts (e.g., sodium chloride), antimicrobial agents (e.g., benzalkonium chloride), sweeteners, antioxidants, antistatic agents, surfactants (e.g., polysorbates such as "TWEEN 20" and "TWEEN 80"), sorbitan esters, saccharin, cyclodextrins, lipids (e.g., phospholipids such as lecithin and other phosphatidylcholines, phosphatidylethanolamines), fatty acids and fatty esters, steroids (e.g., cholesterol), and chelating agents (e.g., EDTA, zinc and other such suitable cations). Other pharmaceutical excipients and/or additives suitable for use in the compositions according to the invention are listed in "Remington: The Science & Practice of Pharmacy", 19.sup.th ed., Williams & Williams, (1995), and in the "Physician's Desk Reference", 52.sup.nd ed., Medical Economics, Montvale, N.J. (1998).

[00264] By non-limiting example, classes of taste-masking agents for fluoroquinolone formulation include the addition of flavorings, sweeteners, and other various coating strategies. By non-limiting examples these may be chosen from sugars such as sucrose, dextrose, and lactose), carboxylic acids, salts such as magnesium and calcium (non-specific or chelation-based fluoroquinolone taste masking), menthol, amino acids or amino acid derivatives such as arginine, lysine, and monosodium glutamate, and synthetic flavor oils and flavoring aromatics and/or natural oils, extracts from plants, leaves, flowers, fruits, etc. and combinations thereof. These may include cinnamon oils, oil of wintergreen, peppermint oils, clover oil, bay oil, anise oil, eucalyptus, vanilla, citrus oil such as lemon oil, orange oil, grape and grapefruit oil, fruit essences including apple, peach, pear, strawberry, raspberry, cherry, plum, pineapple, apricot, etc. Additional sweeteners include sucrose, dextrose, aspartame (NutraSweet®), acesulfame-K, sucralose and saccharin, organic acids (by non-limiting example citric acid and aspartic acid). Such flavors may be present at about 0.05 to about 4 percent. Another approach to improve or mask the taste of unpleasant inhaled drugs is to decrease the drugs solubility, e.g. drugs must dissolve to interact with taste receptors. Hence, to deliver solid forms of the drug may avoid the taste response and acquire the desired improved taste affect. Non-limiting methods to decrease fluoroquinolone solubility are described in this document, e.g. salt forms of levofloxacin or gemifloxacin with xinafoic

acid, oleic acid, stearic acid and pamoic acid. Additional co-precipitating agents include dihydropyridines and a polymer such as polyvinyl pyrrolidone. Moreover, taste-masking may be accomplished by creation of lipophilic vesicles. Additional coating or capping agents include dextrans (by non-limiting example cyclodextrins may include, 2-hydroxypropyl-beta-cyclodextrin, 2-hydroxypropyl-gamma-cyclodextrin, randomly methylated beta-cyclodextrin, dimethyl-alpha-cyclodextrin, dimethyl-beta-cyclodextrin, maltosyl-alpha-cyclodextrin, glucosyl-1-alpha-cyclodextrin, glucosyl-2-alpha-cyclodextrin, alpha-cyclodextrin, beta-cyclodextrin, gamma-cyclodextrin, and sulfobutylether-beta-cyclodextrin), modified celluloses such as ethyl cellulose, methyl cellulose, hydroxypropyl cellulose, hydroxyl propyl methyl cellulose, polyalkylene glycols, polyalkylene oxides, sugars and sugar alcohols, waxes, shellacs, acrylics and mixtures thereof. By non-limiting example, other methods to deliver non-dissolved forms of fluoroquinolones are to administer the drug alone or in simple, non-solubility affecting formulation as a crystalline micronized, dry powder, spray-dried, and nanosuspension formulation. However, an alternative method is to include taste-modifying agents. These include taste-masking substance that is mixed with, coated onto or otherwise combined with the fluoroquinolone active medicament. However, this addition may also serve to improve the taste of another chosen drug product addition, e.g. a mucolytic agent. Non-limiting examples of such substances include acid phospholipids, lysophospholipid, tocopherol polyethyleneglycol succinate, and embonic acid (pamoate). Many of these agents can be used alone or in combination with fluoroquinolones for aerosol administration.

Mucolytic Agents

[00265] Methods to produce formulations that combine agents to reduce sputum viscosity during aerosol treatment with a fluoroquinolone include the following. These agents can be prepared in fixed combination or be administered in succession with aerosol fluoroquinolone therapy.

[00266] The most commonly prescribed agent is N-acetylcysteine (NAC), which depolymerizes mucus in vitro by breaking disulphide bridges between macromolecules. It is assumed that such reduction of sputum tenacity facilitates its removal from the

respiratory tract. In addition, NAC may act as an oxygen radical scavenger. NAC can be taken either orally or by inhalation. Differences between these two methods of administration have not been formally studied. After oral administration, NAC is reduced to cysteine, a precursor of the antioxidant glutathione, in the liver and intestine. The antioxidant properties could be useful in preventing decline of lung function in cystic fibrosis (CF). Nebulized NAC is commonly prescribed to patients with CF, in particular in continental Europe, in order to improve expectoration of sputum by reducing its tenacity. The ultimate goal of this is to slow down the decline of lung function in CF.

[00267] L-lysine-N-acetylcysteinate (ACC) or Nacystelyn (NAL) is a novel mucoactive agent possessing mucolytic, antioxidant, and anti-inflammatory properties. Chemically, it is a salt of ACC. This drug appears to present an activity superior to its parent molecule ACC because of a synergistic mucolytic activity of L-lysine and ACC. Furthermore, its almost neutral pH (6.2) allows its administration in the lungs with a very low incidence of bronchospasm, which is not the case for the acidic ACC (pH 2.2). NAL is difficult to formulate in an inhaled form because the required lung dose is very high (approximately 2 mg) and the micronized drug is sticky and cohesive and it is thus problematic to produce a redispersible formulation. NAL was first developed as a chlorofluorocarbon (CFC) containing metered-dose inhaler (MDI) because this form was the easiest and the fastest to develop to begin the preclinical and the first clinical studies. NAL MDI delivered 2 mg per puff, from which approximately 10% was able to reach the lungs in healthy volunteers. One major inconvenience of this formulation was patient compliance because as many as 12 puffs were necessary to obtain the required dose. Furthermore, the progressive removal of CFC gases from medicinal products combined with the problems of coordination met in a large proportion of the patient population (12) have led to the development of a new galenic form of NAL. A dry powder inhaler (DPI) formulation was chosen to resolve the problems of compliance with MDIs and to combine it with an optimal, reproducible, and comfortable way to administer the drug to the widest possible patient population, including young children.

[00268] The DPI formulation of NAL involved the use of a nonconventional lactose (usually reserved for direct compression of tablets), namely, a roller-dried (RD) anhydrous β -lactose. When tested in vitro with a monodose DPI device, this powder

formulation produces a fine particle fraction (FPF) of at least 30% of the nominal dose, namely three times higher than that with MDIs. This approach may be used in combination with a fluoroquinolone antibiotic for either co-administration or fixed combination administration antibiotic therapy.

[00269] In addition to mucolytic activity, excessive neutrophil elastase activity within airways of cystic fibrosis (CF) patients results in progressive lung damage. Disruption of disulfide bonds on elastase by reducing agents may modify its enzymatic activity. Three naturally occurring dithiol reducing systems were examined for their effects on elastase activity: 1) *Escherichia coli* thioredoxin (Trx) system, 2) recombinant human thioredoxin (rhTrx) system, and 3) dihydrolipoic acid (DHLA). The Trx systems consisted of Trx, Trx reductase, and NADPH. As shown by spectrophotometric assay of elastase activity, the two Trx systems and DHLA inhibited purified human neutrophil elastase as well as the elastolytic activity present in the soluble phase (sol) of CF sputum. Removal of any of the three Trx system constituents prevented inhibition. Compared with the monothiols N-acetylcysteine and reduced glutathione, the dithiols displayed greater elastase inhibition. To streamline Trx as an investigational tool, a stable reduced form of rhTrx was synthesized and used as a single component. Reduced rhTrx inhibited purified elastase and CF sputum sol elastase without NADPH or Trx reductase. Because Trx and DHLA have mucolytic effects, we investigated changes in elastase activity after mucolytic treatment. Unprocessed CF sputum was directly treated with reduced rhTrx, the Trx system, DHLA, or DNase. The Trx system and DHLA did not increase elastase activity, whereas reduced rhTrx treatment increased sol elastase activity by 60%. By contrast, the elastase activity after DNase treatment increased by 190%. The ability of Trx and DHLA to limit elastase activity combined with their mucolytic effects makes these compounds potential therapies for CF.

[00270] In addition, bundles of F-actin and DNA present in the sputum of cystic fibrosis (CF) patients but absent from normal airway fluid contribute to the altered viscoelastic properties of sputum that inhibit clearance of infected airway fluid and exacerbate the pathology of CF. One approach to alter these adverse properties is to remove these filamentous aggregates using DNase to enzymatically depolymerize DNA to constituent monomers and gelsolin to sever F-actin to small fragments. The high

densities of negative surface charge on DNA and F-actin suggest that the bundles of these filaments, which alone exhibit a strong electrostatic repulsion, may be stabilized by multivalent cations such as histones, antimicrobial peptides, and other positively charged molecules prevalent in airway fluid. Furthermore, As a matter-a-fact, it has been observed that bundles of DNA or F-actin formed after addition of histone H1 or lysozyme are efficiently dissolved by soluble multivalent anions such as polymeric aspartate or glutamate. Addition of poly-aspartate or poly-glutamate also disperses DNA and actin-containing bundles in CF sputum and lowers the elastic moduli of these samples to levels comparable to those obtained after treatment with DNase I or gelsolin. Addition of poly-aspartic acid also increased DNase activity when added to samples containing DNA bundles formed with histone H1. When added to CF sputum, poly-aspartic acid significantly reduced the growth of bacteria, suggesting activation of endogenous antibacterial factors. These findings suggest that soluble multivalent anions have potential alone or in combination with other mucolytic agents to selectively dissociate the large bundles of charged biopolymers that form in CF sputum.

[00271] Hence, NAC, unfractionated heparin, reduced glutathione, dithiols, Trx, DHLA, other monothiols, DNase, dornase alfa, hypertonic formulations (e.g. osmolalities greater than about 350 mOsmol/kg), multivalent anions such as polymeric aspartate or glutamate, glycosidases and other examples listed above can be combined with fluoroquinolone antibiotics and other mucolytic agents for aerosol administration to improve antibacterial activity through better distribution from reduced sputum viscosity, and improved clinical outcome through improved pulmonary function (from improved sputum mobility and mucociliary clearance) and decreased lung tissue damage from the immune inflammatory response.

EXAMPLES

[00272] The following examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these examples in no way serve to limit the true scope of this invention, but rather are presented for

illustrative purposes.

Example 1 – High Local Concentration with Short Duration Aerosol Fluoroquinolone Exposure.

[00273] Aerosol administration of fluoroquinolones such as levofloxacin produces high concentrations in the epithelial lining fluid (ELF) of rats and humans. However, this dose has been observed to rapidly decline following administration.

[00274] In order to determine if short duration, high concentrations of levofloxacin could be effective in treatment of *P. aeruginosa* (PA), studies were conducted to measure their bactericidal activity on various strains of this organism which were grown at different conditions. Those were chosen based on what is known about conditions and growth of PA in a cystic fibrosis (CF) lung. Four isogenic strains of *P. aeruginosa* were used for these experiments (Table 2).

Table 2. Strains of PA Used in Time-Kill Experiments.

Strain	Genotype	Levofloxacin MIC (ug/ml)
PAM1020	wt	0.25
PAM1032	<i>nalB</i>	1

[00275] PAM1020 is the parent wild-type strain, PAM1032 contains *nalB* mutation which results in increased levofloxacin resistance due to overexpression of the MexAB-OprM efflux pump which can extrude levofloxacin out of cells.

Experiment 1. Activity of Levofloxacin Against Exponentially Grown Cells.

Methods

Inoculum Preparation

[00276] Strains were grown aerobically overnight in Mueller-Hinton Broth (MHB) at 35° C. Next, cultures were diluted 1:1000 into 100 ml of fresh MHB and grown to OD₆₀₀~0.3 to reach CFU/ml ~ 10⁸. 10 ml of this culture was moved to 50 ml flasks, each containing 10 ml of pre-warmed MHB broth with appropriate concentrations

of levofloxacin (2X as compared to the exposure concentrations).

Exposure

[00277] All strains were treated for 10 min., 20 min., 40 min., 80 min. and 160 minutes. The following concentrations of levofloxacin (ug/ml) were used for the exposure of PAM1020 and PAM1032: 16, 32, 64, 128 and 256. All strains were treated at each concentration for 10 min., 20 min., 40 min., 80 min. and 160 minutes.

Determination of viable cell numbers

[00278] At appropriate times, 1 ml of each exposure culture was centrifuged for 2 minutes, the pellet was washed twice with 1 ml of drug-free MHB, and re-suspended in 1 ml of MHB. The viable cell numbers were enumerated by plating serially diluted samples (in duplicates) on MHB plates by the drop (10 ul) plating method. The limit of detection was 100 CFU/ml. Killing is reported as the log reduction calculated relative to cell count at the time of initiation of antibiotic exposure. Relative antibiotic concentrations (relative to MIC of the corresponding strains) are used. Cell numbers at initiation of antibiotic exposure are shown in Table 3.

Table 3. Bacterial Numbers at Time of Initial Bacterial Exposure.

Strain	CFU/ml
PAM1020	4.03E+07
PAM1032	5.60E+07

Results

[00279] For the most susceptible strain, PAM1020, maximum killing (5.5 log decrease in viable cell counts) was achieved after incubation for 10 minutes with the concentration of levofloxacin corresponding to 256-fold MIC (64 ug/ml tested). 5-logs of killing were achieved already with the lowest concentration tested (16 ug/ml or 64-fold MIC) (Figure 4A). For the strain PAM1032, as long as the concentration above 128-fold the MIC (128 ug/ml) was reached, 10 minute of exposure was sufficient to result in maximum killing (more than 5 logs). At short duration exposures (10 or 20 minutes), less killing was observed at concentrations below 128-fold of MICs. At longer exposure times, concentration corresponding to 16-fold MICs and above resulted in similar maximum killing (Figure 4B). These results indicate that logarithmic cells of *P.*

aeruginosa are efficiently killed after short duration exposures to high concentrations of levofloxacin.

Experiment 2. Activity of Levofloxacin Against Stationary Phase Cells.

Methods

Inoculum Preparation

[00280] Strains were grown aerobically overnight in Mueller-Hinton Broth (MHB) at 35°C (350 ml total). The spent medium was obtained after centrifugation of overnight cultures and filtering the supernatant. Cultures were diluted to OD=0.3 into spent medium. The same medium was also used to prepare antibiotic concentrations (the same as in Experiment 1).

Exposure

[00281] Antibiotic concentrations, time of exposure as determination of viable cell counts were the same as in Experiment 1.

Results

[00282] Cell numbers at initiation of antibiotic exposure are shown in Table 4.

Table 4. Bacterial Numbers at Time of Initial Bacterial Exposure.

Strain	CFU/ml
PAM1020	8.0E+08
PAM1032	8.50E+08

[00283] For stationary phase cells of PAM1020, maximum killing was observed at the lowest concentration corresponding to 64-fold above MIC (16 ug/ml) and the shortest duration of exposure, 10 minutes (Figure 5A). However, PAM1032 demonstrated clear dose-dependent killing with the maximum killing (4 logs) at concentrations 64 the MIC at a short exposure time. Extending the exposure times did not result in larger extent of killing. However, lower concentrations of drug were required to achieve the same killing at longer exposure times (Figure 5B).

[00284] Next, we have compared the re-growth of PAM1020 and PAM1032 after either 10 minutes or 160 minutes of treatment with various concentrations of levofloxacin. After the corresponding treatments, cells were washed twice with antibiotic-free medium. 150 μ l of cells was placed into 96-well plate and the growth was continuously monitored at A_{660} using SpectraMax (Molecular Devices). The results are shown in Figure 6A-6D.

[00285] The results demonstrate that the re-growth of both strains was observed at approximately the same time whether cells were treated with high concentrations of levofloxacin for 10 minutes or 160 minutes. These results further support the efficiency of short duration treatment with high concentrations of levofloxacin.

Experiment 3. Activity of Levofloxacin Against Cells Grown Under Oxygen-Limiting Conditions

Methods

Inoculum Preparation

[00286] Overnight cultures were grown aerobically overnight in Mueller-Hinton Broth and next diluted 1:10000 in MHB which filled growth flasks to the very top. Cultures were grown without shaking to $OD \sim 0.3$ at $37^{\circ}C$. Under these conditions an average of ~ 20 hours was required to reach an $OD=0.3$ as compared to ~ 5 hours under aerated conditions (50 ml of medium in 250 ml flasks, vigorous shaking). Upon analysis, it appeared that an $OD=0.3$ corresponded to a late-logarithmic phase of growth. Other than decreased aeration, antibiotic concentration, time of exposure, and determination of viable cell counts were the same as in Experiments 1 and 2.

Results

[00287] Cell numbers at initiation of antibiotic exposure are shown in Table 5.

Table 5. Bacterial Numbers at Time of Initial Bacterial Exposure.

Strain	CFU/ml
PAM1020	7.5E+07
PAM1032	8.5E+07

[00288] In the case of PAM1020 near maximum killing (4 logs vs 4.5 logs observed under normal aeration) was achieved after exposure with the lowest concentration of levofloxacin for the shortest duration of time (10 minutes) (Figure 7A). In the case of PAM1032 dose-dependent killing was observed for 10 minutes or 20 minutes of exposure with the highest killing observed at concentrations corresponding to 128 to 256-fold the MIC. Slightly stronger (less than 1 log difference) killing was observed for longer exposure intervals (Figure 7B). These data indicate that under conditions of oxygen limitation cells which are at the late logarithmic phase of growth are efficiently killed after short duration of exposure with high concentrations of levofloxacin.

Experiment 4. Activity of Levofloxacin Against PAM1032 in CF Sputum.

Methods

[00289] Cells of strain PAM1032 (MIC = 1 ug/ml) were grown to OD=1 (late-exponential/early stationary phase of growth) in MHB and next concentrated 10-fold in 10-fold concentrated MHB. 10 ul of cells were then added to 90 ul of sputum or water in 96-well round bottom plates, restoring MHB to its original concentration. Quantitation plates were pre-warmed for 5 minutes at 37°C and different concentrations of levofloxacin (512 ug/ml, 128 ug/ml, 32 ug/ml, 8 ug/ml, 2 ug/ml, and 0.5 ug/ml) were added. At appropriate times, 10 ul of each treatment culture was diluted 100-fold in MHB to minimize the carryover of levofloxacin. Viable cell numbers were enumerated by plating serially diluted samples on MHB plates by the drop (10 ul) plating method. The limit of detection was 10⁴ CFU/ml. Killing is reported as the percentage of the starting inoculum survived after the levofloxacin treatment. Results are shown in Figures 8A and 8B.

Results

[00290] The results indicate that while sputum slightly affected the degree of killing by levofloxacin, rapid and extensive (up to five orders of magnitude) killing by levofloxacin in sputum was still observed after short duration of treatment at high concentrations of antibiotic.

Experiment 4. Activity of Levofloxacin Against Colony Biofilms of PAM1032.**Methods*****Biofilm Preparation***

[00291] Colony biofilms were grown on polycarbonate membrane filters (diameter, 25 mm; Poretics, Livermore, CA) resting on MHB plates. Overnight culture of PAM1032 was diluted to OD=0.3, and then diluted 1:100 in fresh MHB. 5 ul of this culture was spotted on the membrane filter. Bacteria were incubated at 37°C for 48 hours (mature biofilms).

Exposure

[00292] After growth filters were placed into tubes containing 3 ml saline or saline and levofloxacin at 128 ug/ml and 1024 ug/ml. Each tube was treated for 10 minutes and 80 minutes. At about 5 min. before incubation time elapsed, tubes were vigorously vortexed (A) or sonicated and vortexed (B) to detach cells. 1 ml of each exposure culture was centrifuged for 2 minutes, the pellet was washed twice with 1 ml of drug-free MHB, and re-suspended in 1 ml of MHB. The viable cell numbers were enumerated by plating serially diluted samples (in duplicates) on MHB plates by the drop (10 ul) plating method. Results are shown in Figure 9.

Results

[00293] The data demonstrate that maximum killing (~2 logs) is obtained after 10 min with the lowest concentration of levofloxacin tested (128-fold MIC). No additional killing was observed at the higher levofloxacin concentration. These data indicate that colony biofilms are more resistant to killing as compared to logarithmic or stationary phase cells. However, the maximum observed bactericidal activity against biofilms (99% under these conditions) was achieved after 10 minutes of levofloxacin exposure.

Experiment 5. Simulated Short-Term, Rapid Aerosol Administration, Delivering High Concentration Drug Exposure in in vitro Pharmacodynamic Model.

[00294] In vitro pharmacodynamic models of infection allow for exposure of a

growing bacterial inoculum to changing concentrations of drug as would occur in vivo. The strength of this approach is that the serum concentration vs. time profile of a drug in man can be simulated in the laboratory in vitro to determine the optimal exposure profile (i.e., dose and dosing interval) for a drug and target pathogen.

[00295] The following report describes experiments designed to determine the C_{max} and AUC that will provide maximum bactericidal effects after an aerosol dose of a fluoroquinolone.

Material and Methods

In Vitro Pharmacodynamic Model of Infection

[00296] The in vitro pharmacodynamic model consists of a central (analogous “serum” compartment) and peripheral (“extravascular”) compartment. The peripheral compartments consist of artificial capillary units (Unisyn, Hopkinton, MA) arranged in series with the central compartment. Each capillary unit has a bundle of small semi-permeable fibers with a molecular size retention of ca. 10,000 MW to allow passage of nutrients but not bacteria. The entire system is set up in a dry heat incubator adjusted to 37°C.

[00297] Both the central and peripheral compartments were filled with Mueller-Hinton broth. Each peripheral compartment (capillary unit and tubing) contained ca. 23 ml of growth medium.

[00298] Bacteria were introduced into the peripheral chamber of the model and allowed to grow for 2 hours prior to the first “dose” of drug. Drug doses were administered into the central compartment and pumped to peripheral chambers by a peristaltic pump. Concentrations in the model declined according to first order elimination (half-life) by dilution of the central compartment with drug free medium introduced by an additional peristaltic pump adjusted to the desired clearance.

[00299] Samples (0.3 ml) were collected from peripheral compartments at various intervals for determination of drug and bacterial concentrations. Samples were collected from the peripheral compartments and assayed for drug concentrations by HPLC.

Bacterial Test Strains

[00300] *Pseudomonas aeruginosa* PAM1032 and PAM1582. The MICs of these strains to levofloxacin were 1.0 and 32 ug/ml, respectively.

Inoculum Preparation

[00301] Strains were grown aerobically overnight in Mueller-Hinton Broth (MHB) at 35°C and subcultured to fresh MHB and reincubated at 35°C for 2 hours. After 2 hours, the inoculum was further diluted 1:1000 to a final concentration of approx. 1.0×10^6 CFU/ml. Of the resulting dilution, 2.3 ml was injected into each peripheral chamber of the hollow-fiber bioreactors (Unisyn, Hopkinton, MA).

Pharmacokinetics

[00302] The half-life of levofloxacin was adjusted to be 10 minutes to be equivalent to that observed following aerosol delivery of levofloxacin to the pulmonary compartment of man. The targeted C_{max} was 1000 and 600 ug/ml over two experiments.

Results

[00303] As targeted, the model exhibited a levofloxacin half-life of 10 minutes and the C_{max} of 1000 ug/ml for Experiment 5. By comparison, Experiment 6 was modified to achieve the same half-life as Experiment 5, but with a targeted C_{max} of 600 µg/ml.

[00304] The bactericidal effects of these two regimens correlated with the C_{max}. In Experiment 5 with a C_{max} of 1000 ug/ml, the maximum bactericidal effect was observed as a 5 log reduction in bacterial counts within 10 minutes with PAM1032 and a 4 log reduction in bacterial counts within 20 minutes with PAM1582 and no re-growth observed over the remaining 2 hours of the experiment (Figure 10). In contrast, while the C_{max} of 600 ug/ml used in Experiment 6 maintained the 5-log reduction in bacterial counts for PAM1032, albeit taking 30 min instead of 10 min observed in Experiment 1, only a 3-log reduction in bacterial counts was observed for PAM1582 after 45 min (Figure 11). Moreover, PAM1582 exhibited initial re-growth before the end of the 2 hour experimental window.

Conclusions

[00305] Levofloxacin can produce up to a 99.9999% bacterial reduction with a Cmax of both 600 and 1000 ug/ml against a strain with an MIC of 1 ug/ml. However, maximal bactericidal activity requires 3X more time at a Cmax of 600 ug/ml. Levofloxacin can also produce up to 99.99% bacterial reduction with a Cmax of both 600 ug/ml against a strain with an MIC of 32ug/ml. However, the time to reach the maximum effect is 45 minutes. In contrast, levofloxacin can produce up to 99.999% bacterial reduction of this resistant strain with a Cmax of 1000 ug/ml and the time to maximum effect is reduced to 20 minutes. From these results, extremely high, but short duration exposures of levofloxacin produce rapid and sustained bacterial killing in both flask and hollow fiber models. Taken together, the above results indicate that achieving an initial 800 ug/ml levofloxacin or other fluoroquinolone human ELF or sputum concentration is sufficient to achieve the above antibiotic affects for the MIC99 population as represented by PAM1582 (MIC of 32 ug/ml).

Example 2 – Determination of the Aerosol Properties of Antibacterial Fluoroquinolones.

Introduction

[00306] Objective. The purpose of these studies was to evaluate the ability to formulate and deliver by nebulization a variety of fluoroquinolones for treatment of pulmonary bacterial infections by aerosol administration. The fluoroquinolones evaluated are shown in Table 6.

Table 6. Fluoroquinolones Tested.

Fluoroquinolone	Sp MIC₉₀ (ug/mL)	MSSA MIC₉₀ (ug/mL)	MRSA MIC₉₀ (ug/mL)	Pa MIC₉₀ (ug/mL)	Approval Status
Ciprofloxacin	2	1	64	8	Approved
Gemifloxacin	0.06	0.06	2	8	Approved
Levofloxacin	2	0.5	16	8	Approved
Marbofloxacin	2	2	ND	8	Veterinary
Gatifloxacin	0.5	0.125	4	16	Approved

Ofloxacin	2	1	>32	16	Approved
Tosufloxacin	0.5	0.125	>16	16	Japan
Lomefloxacin	16	2	>32	32	Approved
Moxifloxacin	0.25	0.125	2	32	Approved
Sparfloxacin	0.5	0.125	16	32	Withdrawn
Orbifloxacin	2	2	ND	>32	Veterinary
Pefloxacin	32	2	>32	>32	Europe
Trovafloxacin	0.25	0.06	8	>32	Withdrawn

[00307] These fluoroquinolones were chosen based on their availability, approval status and antimicrobial properties. All tested fluoroquinolones are either currently approved in the United States or have been approved but later withdrawn due to various adverse reactions. In addition, several fluoroquinolones, which are in use for veterinary applications, have also been evaluated. Among bacterial pathogens responsible for respiratory tract infections, *Pseudomonas aeruginosa* (Pa) and methicillin resistant *Staphylococcus aureus* (MRSA) are the most refractory to treatment with fluoroquinolones. *Streptococcus pneumoniae* (Sp) is probably the most important pathogen responsible for respiratory tract infections and numerous reports demonstrate high rates of fluoroquinolone resistance in these bacteria. MIC₉₀ for Pa ranges from 4 ug/ml to 32 ug/ml and from 2 ug/ml to >32 ug/ml for Pa and MRSA, respectively. Ciprofloxacin, levofloxacin, gemifloxacin and gatifloxacin vs gemifloxacin and moxifloxacin are the most potent against Pa and MRSA, respectively.

[00308] Table 7 contains a list of additional fluoroquinolones for potential evaluation. The most microbiologically interesting compounds in the list are clinafloxacin and olamufloxacin, which were discontinued due to adverse reactions, and sitofloxacin, which is in Phase III clinical trials.

Table 7. Fluoroquinolones For Potential Evaluation.

Fluoroquinolone	Sp MIC₉₀ (ug/mL)	Sa MIC₉₀ (ug/mL)	MRSA MIC₉₀ (ug/mL)	Pa MIC₉₀ (ug/mL)	Market Status
Clinafloxacin	0.06	0.06	2	4	Discontinued
Sitafloxacin	0.06	0.125	4	8	Phase III

Olamufloxacin	0.06	1	2	16	Discontinued
Norfloxacin	16	1	>4	16	Approved
Prulifloxacin	1	0.25	32	16	Phase III
Danofloxacin	NA	0.125	NA	>16	Veterinary
Enrofloxacin	1	0.125	8	>16	Veterinary
Sarafloxacin	NA	0.25	>16	>16	Veterinary
Balofloxacin	0.5	0.25	8	32	Korea
Fleroxacin	8	1	>4	32	Europe
Difloxacin	2	0.5	NA	32	Veterinary
Rufloxacin	32	2	64	32	Europe, China
Enoxacin	16	1	>4	>32	Withdrawn
Garenoxacin	0.06	0.06	8	>32	Phase III
Grepafloxacin	0.5	0.125	32	>32	Withdrawn
Pazufloxacin	4	0.5	>16	>32	Japan

[00309] The fluoroquinolones in these two tables represent one field of options for an aerosol fluoroquinolone candidate. Several potent fluoroquinolones such as DX-619 and DW-286, which are at the early stage of clinical development, might also be of interest for future studies.

[00310] Specific physico-chemical considerations for nebulization include aqueous solubility, viscosity and surface tension. The aqueous solubility of the drug should advantageously be sufficient to meet or exceed the minimal dosing requirement. The loading drug concentration also affects delivery time. Longer delivery times may be commercially unacceptable or lead to poor patient compliance. Although longer delivery times may in effect modify the AUC shape, by non-limiting example, the PARI eFlow device has been discovered to administer 4 ml of aqueous levofloxacin in less than 5 min. Moreover, using such an efficient device, high concentration levofloxacin may be able to deliver the effective doses described herein in a timeframe further enabling the rapid administration, high concentration drug requirements needed for optimal fluoroquinolone therapy.

[00311] In the case of fluoroquinolones, pH directly affects solubility. In general, solubility decreases significantly with increase in pH in the range 1.5 to 6.5.

Because pH also affects patient tolerability (see below), the optimal choice of fluoroquinolone for pulmonary delivery via aerosol has certain solubility and pH levels.

[00312] For the purpose of this feasibility study, the target solubility was set at 10 mg/mL or higher at a pH of 4.5 or greater, based on calculations of therapeutic dose and the delivery metrics for available nebulizers. In order to be above the mutant prevention concentration (MPC), peak concentration of fluoroquinolone after aerosol administration advantageously reaches about 100 ug/ml to about 1000 ug/ml at the site of infection, pending the MIC of the infecting organism. Based on these considerations, the minimal dose to be in this therapeutically relevant range was projected to be at least about 30-40 mg Respirable Delivered Dose (RDD). Given the relative half-life of levofloxacin in the human lung, the practical achievement of this dose by nebulization may be obtained with a loading dose of at least about 100 mg in a volume of about 2 mL (about 50 mg/mL) in a high-efficiency vibrating-mesh device operating at its maximum performance efficiency delivering this dose in less than 4 min. A standard ultrasonic or jet nebulizer may require a loading dose of at least about 400 mg in a volume of about 5 mL (about 80 mg/mL). However, the rate of administration by these less efficient devices may not be sufficient to achieve high local concentration with short duration exposure. Similar efficacious doses may also be achieved by administration of levofloxacin as a dry powder, where the rapid solubility properties of levofloxacin may permit a quick dissolution resulting in these desired soluble drug concentrations. However, alternative concentrations or alteration of the fluoroquinolone AUC shape profile may be desirable.

[00313] Alternatively, although aqueous solubility is important, it is reasonable to predict a formulation utilizing particle or complexation technology to enable nebulization of less soluble fluoroquinolones. Unfortunately, more intricate formulations increase both the complexity and cost of drug development, and in the case of jet and ultrasonic nebulizers, a significant reduction in efficiency of delivery, and limit the ability to introduce other design elements into a final drug product.

[00314] In addition to drug solubility, for vibrating mesh devices nebulization is also sensitive to the surface tension of the drug formulation. Therefore, in one

embodiment, the surface tension is adjusted during formulation by modifying drug concentration, excipient concentration and/or the addition of surfactant.

[00315] In addition to factors that affect efficient nebulization, other factors may be considered for patient tolerability and compliance. By non-limiting example, these factors may include osmolality, pH and taste. Osmolality affects acute tolerability in the respiratory tract and can be optimized for most drugs during formulation. Similarly, the pH of an aerosol also contributes to tolerability, yet only negatively when the formulation pH is less than 4.5. Therefore, because pH directly contributes to fluoroquinolone solubility, fluoroquinolones that require a pH less than 4.5 for solubility are likely to be poorly tolerated. Finally, fluoroquinolone taste can affect good patient compliance. Fluoroquinolones are known generally to be associated with an unpleasant, sometimes very intense taste. While there are technologies available that may mask poor drug taste, these technologies increase development complexity and cost, and may not be totally effective in the case of fluoroquinolones. Thus, similar to pH, taste may be considered in identifying a fluoroquinolone suitable for nebulization.

Preparation and Characterization of the Test Solutions

[00316] Antibiotics were purchased from one of several sources as shown in Table 8.

Table 8. Preparation of Fluoroquinolone Test Solutions.

No.	Fluoro-quinolone	Source ^a	Purity ^b	Amount	Volume H ₂ O	Final Conc.
1	Gatifloxacin	LKT	99.6	8.7 mg	0.87 mL	10 mg/mL
2	Gemifloxacin	LG	99.6	9.5 mg	0.95 mL	10 mg/mL
3	Levofloxacin	LKT	99.2	10.3 mg	1.03 mL	10 mg/mL
4	Moxifloxacin	LKT	99.5	12.5 mg	1.25 mL	10 mg/mL
5	Ciprofloxacin	LKT	99.3	19.5 mg	1.95 mL	10 mg/mL
6	Ofloxacin	LKT	99.1	11.7 mg	1.17 mL	10 mg/mL
7	Lomefloxacin	MPI	NA	17.0 mg	1.70 mL	10 mg/mL
8	Marbofloxacin	Vetoquino	NA	4.8 mg	0.48 mL	10 mg/mL
9	Orbifloxacin	MPI	NA	4.2 mg	0.42 mL	10 mg/mL
10	Pefloxacin	MPI	NA	15.0 mg	1.50 mL	10 mg/mL

11	Sparfloxacin	MPI	NA	14.5 mg	1.45 mL	10 mg/mL
12	Tosufloxacin	MPI	NA	15.2 mg	1.52 mL	10 mg/mL
13	Trovafloxacin	MPI	NA	2.0 mg	0.20 mL	10 mg/mL

- a. LKT: LKT Laboratories. LG: LG Chem. NA. Source unavailable.
- b. Purity of material tested. Described as GMP or in percent API.
- c. 25 mg/ml solution.

[00317] A 2 to 20 mg sample of each antibiotic was weighed into sterile plastic tubes and brought up with a volume of sterile water to make a 10 mg/mL solution or suspension of the antibiotic. Samples were incubated for approximately 10 minutes at room temperature with occasional mixing, prior to further handling.

[00318] Following the incubation period, the antibiotic solutions were observed for their visible appearance, with results as shown in Table 9.

[00319] Five of the fluoroquinolones tested were visibly soluble, and either colorless, or a shade of yellow. Eight were visibly insoluble, appearing either cloudy (fine particulate), opaque (dense fine to medium particulate), or turbid (thick, large particulate slurry), in all cases with a visible sediment. The pH of these initial solutions were determined, and ranged from 3.5 to 7.0. The insoluble solutions were titrated with 1N HCl to the point of visible solubility, and the pH of the solubilized solution determined. In three cases, marbofloxacin, sparfloxacin and tosufloxacin, solubility was not reached by pH 1.5, and further addition of acid was stopped. With the exception of ofloxacin, the pH of these titrated solutions was in the range of 1.5 to 3.0.

Table 9. Fluoroquinolone Solution Characteristics.

No.	Fluoro-quinolone	Initial Solution		After pH Adjustment		
		Appearance	pH	1N HCl (uL)	Appearance ^a	pH
1	Gatifloxacin	white, cloudy, visible sediment	7.0	5	slight yellow color, transparent, no sediment	3.0
2	Gemifloxacin	colorless, transparent, no sediment	4.7		NR	-
3	Levofloxacin	slight yellow color, transparent, no sediment	4.7		NR	-

4	Moxifloxacin	bright yellow color, transparent, no sediment	4.7		NR	-
5	Ciprofloxacin	white, opaque (very dense), visible sediment	5.5	60	colorless, transparent, no sediment	2.0
6	Ofloxacin	cloudy, visible sediment	6.5	10	slightly yellow color, transparent, no sediment	5.2
7	Lomefloxacin	cloudy, visible sediment	4.2	-	transparent, no sediment, after 10 min. at rm. temp.	-
8	Marbofloxacin	white, very turbid, visible sediment	6.5	40	white, turbid, visible sediment	1.5
9	Orbifloxacin	white, cloudy, visible sediment		20	colorless, transparent, no sediment	1.7
10	Pefloxacin	colorless, transparent, no precipitate	4.5		NR	-
11	Sparfloxacin	bright yellow, turbid, visible sediment	5.0	20	bright yellow, densely turbid, visible sediment	1.5
12	Tosufloxacin	white, turbid, visible sediment	3.5	20	white, cloudy, less turbid, visible sediment	1.5
13	Trovafloxacin	colorless, slightly cloudy, no sediment	4.2		NR	-

- a. NR: pH adjustment not required. Fluoroquinolone was soluble at a pH ≥ 4 in the initial solution.

[00320] After the pH adjustment, and following a further 10 minute incubation period with occasional mixing, the final appearance of the solutions was determined, just prior to the aerosol tolerability and taste test. Results are shown in Table 10.

Table 10. Appearance of Fluoroquinolone Final Solution.

No.	Fluoroquinolone	pH	Solubility	Color	Sediment	Opaqueness
1	Gatifloxacin	3.0	+	C	none	none to very slight
2	Gemifloxacin		+	C	none	none to very slight
3	Levofloxacin	4.7	+	VLY	none	none
4	Moxifloxacin	4.7	+	Y	+/-	none
5	Ciprofloxacin	2.0	+	C	none	none
6	Ofloxacin	5.2	+	LY	+/-	none
7	Lomefloxacin	4.2	+	C	+/-	none to very slight
8	Marbofloxacin	1.5	--	W	++	++
9	Orbifloxacin	1.7	+	C	none	slight

10	Pefloxacin	4.5	+	C	none	slight
11	Sparfloxacin	1.5	---	DY	+++	++++
12	Tosufloxacin	1.5	--	W	++	+++
13	Trovafloxacin	4.2	+	C	+	slight

Y=yellow; LY= light yellow; VLY=very light yellow; DY=dark yellow; C= colorless; W = white.

[00321] The compounds exhibiting preferred solubility for solutions suitable for delivery by inhalation (10 mg/mL at a pH of 4.5 or above), were levofloxacin, gemifloxacin, moxifloxacin, ofloxacin and pefloxacin. Levofloxacin, ofloxacin and moxifloxacin exhibited the best solubility/pH characteristics.

Taste and Tolerability Evaluation

[00322] Two evaluations were done to determine the suitability of the fluoroquinolone solutions with respect to taste and tolerability.

[00323] First, in an oral taste test the taste of a 20 uL portion of test sample was determined in a single, healthy human volunteer by placing the material directly onto the center front part of the tongue. Taste was then monitored over a 1 minute period. This test was performed on the initial solutions prepared as well as the final solutions following pH adjustment. Data are shown in Table 11.

Table 11. Oral Fluoroquinolone Taste Test.

No.	Fluoro-quinolone	Initial Solution	Final Solution
1	Gatifloxacin	moderate bitter unpleasant taste, slightly aromatic	strong bitter unpleasant almond-like taste, strong aftertaste
2	Gemifloxacin	very bitter unpleasant taste with strong aftertaste, all the way into throat	not performed
3	Levofloxacin	slight chemical taste, slightly bitter, slight almond-like taste	not performed
4	Moxifloxacin	moderate bitter-sweet unpleasant taste, slightly aromatic	not performed
5	Ciprofloxacin	sweet almond-like taste	very strong bitter taste all the way into the throat
6	Ofloxacin	bitter unpleasant, almond-like taste	moderate bitter unpleasant,

			almond-like taste
7	Lomefloxacin	moderate to strong almond-like taste, not very unpleasant	not performed
8	Marbofloxacin	bitter unpleasant almond-like taste	moderate to strong bitter unpleasant almond-like taste
9	Orbifloxacin	strong unpleasant taste	very strong, very unpleasant bitter taste
10	Pefloxacin	strong bitter unpleasant almond-like taste	not performed
11	Sparfloxacin	slight taste	strong almond-like taste
12	Tosufloxacin	mild to moderate almond-like taste	strong almond-like taste
13	Trovafoxacin	very strong bitter unpleasant almond-like taste, strong aftertaste	not done

[00324] Lowering of the pH generally had the effect of enhancing the taste properties of the solution. Gatifloxacin, gemifloxacin, ciprofloxacin, orbifloxacin and trovafloxacin were the least desirable in taste testing. Of the fluoroquinolones tested, Levofloxacin was the only fluoroquinolone that was tolerable with respect to taste, at the concentration tested. Lomefloxacin had a moderately strong almond-like taste, and the taste was slightly unpleasant.

[00325] In the second test, the tolerability and taste of a small aerosol sample from a 0.5 ml aliquot of the test formulation was determined in a single healthy human volunteer, following nebulization in a PARI eFlow nebulizer (Table 12).

Table 12. Aerosol Fluoroquinolone Tolerability and Taste Test.

No.	Fluoro-quinolone	Aerosol Tolerability and Taste
1	Gatifloxacin	moderate bitter unpleasant taste, mild cough sensation
2	Gemifloxacin	strong unpleasant bitter taste, strong aftertaste, mild cough sensation
3	Levofloxacin	chemical taste, somewhat bitter, mild cough sensation
4	Moxifloxacin	moderate bitter unpleasant taste, some cough, strong bitter aftertaste
5	Ciprofloxacin	very strong, bitter unpleasant taste, immediate coughing
6	Ofloxacin	bitter chemical taste, mild cough sensation
7	Lomefloxacin	chemical taste, somewhat bitter, mild cough sensation
8	Marbofloxacin	too insoluble to test
9	Orbifloxacin	very acidic, strong bitter unpleasant taste, strong cough
10	Pefloxacin	chemical taste, some cough

11	Sparfloxacin	too insoluble to test
12	Tosufloxacin	too insoluble to test
13	Trovafloxacin	bitter unpleasant taste, no cough or cough sensation, no aftertaste

[00326] In the case of orbifloxacin, marbofloxacin and trovafloxacin, smaller portions were tested, due to solubility limitations. In a calibration experiment, the inhaler produced an aerosol output of 4.1 microns VMD, with a geometric standard deviation (GSD) of 1.64 micron VMD. In addition to these measurements, the inhaler produced a fine particle dose (FPD) of 54.9% (percent of emitted dose in particles less than 5 microns). The tolerability and taste of the drug during a very brief administration period and for a period of 10 minutes post administration were monitored. Tolerability parameters were of the following types: (i) cough, cough sensation, or sneezing (ii) irritation, burning or tightness of throat, (ii) irritation or runniness in nasal passages or eyes, (iii) irritation, burning or tightness of the lungs or shortness of breath, and (iv) dizziness, headache, nausea or other systemic effects.

[00327] Marbofloxacin, sparfloxacin and tosufloxacin were too insoluble to evaluate in this test. For the remaining fluoroquinolones tested, no tolerability effects were observed during or after aerosol exposure in categories ii, iii or iv (above). Gatifloxacin, moxifloxacin ciprofloxacin, orbifloxacin and pefloxacin were all associated with cough. In the case of ciprofloxacin and orbifloxacin this may have been associated with the low-pH of the solution. Of the fluoroquinolones tested, Levofloxacin at 10 mg/ml had the best taste characteristics. Ofloxacin, lomefloxacin and pefloxacin had a more discernible taste than levofloxacin, which were also acceptable during the short course of administration.

Summary and Conclusions From the Fluoroquinolone Taste Test

[00328] Of the thirteen fluoroquinolones tested in this study, levofloxacin had preferred physical-chemical properties for aerosol administration and a demonstration of best acute tolerability of the fluoroquinolones tested (Table 13). Levofloxacin is also recognized as having one of the best antimicrobial profiles for respiratory pathogens and has the highest *in vivo* efficacy, comparable to ciprofloxacin, for treatment of *Pseudomonas aeruginosa* infections.

Table 13. Overall Suitability for Nebulization.

No.	Fluoro-quinolone	Assessment	Overall Score	Limitation
1	Gatifloxacin	poor solubility and pH, moderately strong bitter aerosol taste	--	Solubility, Taste
2	Gemifloxacin	sufficient solubility and pH, strong bitter aerosol taste, strong aftertaste	---	Taste
3	Levofloxacin	excellent solubility and pH, chemical aerosol taste, somewhat bitter	+	Taste
4	Moxifloxacin	sufficient solubility and pH, moderately strong bitter aerosol taste, strong aftertaste	--	Taste, Pa Activity
5	Ciprofloxacin	poor solubility and pH, very strong bitter aerosol taste, coughing	---	Solubility, Taste
6	Ofloxacin	minimally acceptable solubility and pH, bitter chemical aerosol taste	-/+	Taste
7	Lomefloxacin	minimally acceptable solubility and pH, chemical aerosol taste, strong liquid taste	-/+	Pa Activity
8	Marbofloxacin	very poor solubility even at low pH, unable to test	---	Solubility
9	Orbifloxacin	very poor solubility even at low pH, strong bitter unpleasant aerosol taste, strong cough	---	Solubility, Taste, Pa Activity
10	Pefloxacin	sufficient solubility and pH, aerosol chemical taste, strong unpleasant liquid taste	-/+	Pa Activity
11	Sparfloxacin	very poor solubility even at low pH, unable to test	---	Solubility, Pa Activity
12	Tosufloxacin	very poor solubility even at low pH, unable to test	---	Solubility
13	Trovafloxacin	moderate solubility and pH, bitter aerosol taste	--	Taste, Pa Activity

[00329] Ofloxacin, lomefloxacin and pefloxacin exhibited lower solubility and stronger taste at 10 mg/mL than levofloxacin. Ofloxacin is 2-fold less potent than levofloxacin, and lomefloxacin and pefloxacin are 4-fold less potent. Higher concentrations of these antibiotics have the preferred potency and administration times under 15 minutes.

[00330] In a separate study, conducted in a similar manner, norfloxacin was tested and found to have a solubility, taste and potency profile very similar to gatifloxacin, with the exception of significantly less activity against the gram-positive pathogens.

Taste Testing of Aerosol Salt Formulations of Levofloxacin and Gemifloxacin

[00331] Based on the results of the above studies, levofloxacin, and its racemate ofloxacin, as well as gemifloxacin, and to a lesser extent gatifloxacin and norfloxacin are amenable to aerosol administration for pulmonary antibacterial treatment. To further test the taste and acute tolerability (cough sensation and cough) properties of levofloxacin and gemifloxacin, several formulations were prepared with different organic and inorganic acids and tested in the manner described above. Solutions were prepared by first adding 500 mg levofloxacin to 10 ml of water or adding 500 mg of gemifloxacin to 20 ml of saline (due to solubility limitations), titrating the pH to ~6.5 with HCl or organic acid, then adjusting the osmolality of levofloxacin containing solutions to ~300 mOsm/kg with sodium chloride. The formulations tested are shown in Table 14.

[00332] These formulations were tested by a total of three healthy human volunteers in the same manner as described above, at a levofloxacin concentration of 50 mg/mL, and a gemifloxacin concentration of 25 mg/mL, in a carefully controlled, head-to-head, fully blinded test. Results are shown in Table 15 and 16.

[00333] These results demonstrate that hydrochloric acid, citric acid and ascorbic acid formulations of levofloxacin have superior taste and tolerability compared to the acetic acid, lactic acid and tartaric acid formulations of levofloxacin. Furthermore, these levofloxacin formulations have superior taste and tolerability over the equivalent gemifloxacin formulations. With respect to gemifloxacin, the citric acid formulation had superior taste and tolerability compared to the HCl and ascorbic acid formulations of gemifloxacin, and with further formulation refinement, would be amenable to aerosol administration.

Table 14. Levofloxacin and Gemifloxacin Formulations.

Fluoroquinolone	Acid	Conc (mg/mL)	pH	Osmolality (mOsm/kg)
Levofloxacin	HCl	50	6.5	181
Levofloxacin	Acetic	50	6.41	273
Levofloxacin	Citric	50	6.45	286
Levofloxacin	Lactic	50	6.42	286
Levofloxacin	Ascorbic	50	6.50	278

Levofloxacin	Tartaric	50	6.35	286
Gemifloxacin	HCl	25	5.6	330
Gemifloxacin	Citric	25	5.7	363
Gemifloxacin	Ascorbic	25	5.9	347

Table 15. Aerosol Taste and Tolerability of Levofloxacin Formulations at 50 mg/mL.

Acid	Taster		
	1	2	3
HCl	Moderate bitter taste	Bitter taste, cough sensation	Bitter taste
Acetic Acid	Very acidic taste	Strong acidic taste, cough sensation	Acidic taste, after taste
Citric Acid	Mild after taste, slightly sweet	Mild taste, sweet	Mild after taste
Lactic Acid	Strong bitter taste, aftertaste	Mild taste, after taste, slight cough	Bitter, mild after taste
Ascorbic Acid	Mild taste, slight acidity	Mild taste	Little taste or after-taste
Tartaric Acid	Very bitter, strong after taste	Strong bitter taste, bitter after taste	Bitter taste

Table 16. Taste and Tolerability of Gemifloxacin Formulations at 25 mg/mL.

Acid	Taster		
	1	2	3
Hydrochloric Acid	Metallic taste, strong after taste	Slight bitter taste	Bitter taste, slightly metallic
Citric Acid	Slightly sweet	Mild cough, slightly bitter	Very mild taste, no after taste
Ascorbic Acid	Mild taste	Cough, mild bitter after taste	Slightly bitter, mild after taste

Taste Testing of Additional Aerosol Levofloxacin Formulations

[00334] To further test the taste and tolerability properties of additional levofloxacin excipient combinations in a systematic manner, a series of formulations were prepared and tested. The formulations are listed in Table 17. They included sugars, salts, sweeteners and other excipients prepared by mixing levofloxacin with water, adding the excipients listed in Table 17, and titrating if necessary to the desired pH with dilute HCl. Osmolality was not optimized for these studies. However, osmolality was determined using an Advanced Instruments Model 3250 Osmometer. This measurement, made on 250 μ L of sample, relies on freezing point depression to determine osmolality.

[00335] These formulations were tested in a total of three healthy human volunteers in a series of tests (A-G) in the same manner as described above, in a carefully controlled, head-to-head, fully blinded manner. All tests were performed in a fully blinded fashion. Results of the tests (Tables 19-25) are described below. The following scoring system was used (Table 18).

[00336] Test A: Taste Testing of Sweeteners, Divalent Metal Salts, and Surface Active Agents. This test included sweeteners, calcium and magnesium salts, and surface active agents (i.e., glycerin and PS-80). As shown in Table 17, formulations containing the sweeteners shown are mildly bitter and have metallic taste. The artificial sweeteners appeared to produce a bitter taste that is distinct from the bitterness otherwise observed. Most significantly, the formulation containing CaCl_2 had the most improved taste relative to control (MgCl_2 was not tested in this experiment) (Table 19).

[00337] Test B: Taste Testing of Mono- and Disaccharides in the Presence of Calcium Chloride. All of the formulations screened in this experiment were well tolerated and tasted better than the control sample. Formulations containing both the calcium salt and sugar performed better than either one alone, suggesting that these compounds improve taste through different mechanisms. Of these formulations, 5% CaCl_2 + 7.5 % glucose performed best. Note that the lactose is present at a lower concentration than the other sugars (Table 20).

Table 17. Levofloxacin Formulations Containing Various Excipients.

Fluoroquinolone	Conc (mg/mL)	Excipients	pH	Osmolality (mOsm/kg)
Levofloxacin	50	Control A (0.225% NaCl)	6.50	180
Levofloxacin	50	Aspartame (0.1%)	6.49	175
Levofloxacin	50	Sucrose (0.1%)	6.49	178
Levofloxacin	50	Glucose (5%)	6.5	380
Levofloxacin	50	Sucrose (7.5%); NaCl (0.225%)	6.51	329
Levofloxacin	50	Glycerin (5%)	6.48	880
Levofloxacin	50	PS-80 (0.1%)	6.51	189
Levofloxacin	50	CaCl_2 (5%)	6.10	784
Levofloxacin	50	MgSO_4 (5%)	6.41	73
Levofloxacin	50	Control – B-E (0.225% NaCl)	6.51	182

Levofloxacin	50	CaCl ₂ (5%)	6.1	735
Levofloxacin	50	CaCl ₂ (5%), Sucrose (7.5%)	6.10	958
Levofloxacin	50	CaCl ₂ (5%), Glucose (7.5%)	6.10	1174
Levofloxacin	50	CaCl ₂ (5%), Glucose (7.5%)	5.25	1246
Levofloxacin	50	CaCl ₂ (5%), Lactose (5%)	6.07	864
Levofloxacin	50	MgCl ₂ (5%)	5.90	600
Levofloxacin	50	MgCl ₂ (5%), Sucrose (7.5%)	5.98	815
Levofloxacin	50	MgCl ₂ (5%), Glucose (7.5%)	5.98	999
Levofloxacin	50	MgCl ₂ (5%), Glucose (7.5%)	5.04	1035
Levofloxacin	50	MgCl ₂ (5%), Lactose (5%)	5.96	697
Levofloxacin	50	MgSO ₄ (5%), Sucrose (7.5%)	6.20	433
Levofloxacin	50	MgSO ₄ (5%), Glucose (7.5%)	6.21	625
Levofloxacin	50	MgSO ₄ (5%), Glucose (7.5%)	5.40	660
Levofloxacin	50	MgSO ₄ (5%), Lactose (5%)	6.18	387
Levofloxacin	50	Control F- G (0.45% NaCl)	6.5	221
Levofloxacin	50	Glucose (5%)	6.5	376
Levofloxacin	50	Sucrose (5%)	6.5	240
Levofloxacin	50	Lactose (5%)	6.62	241
Levofloxacin	50	Lactose (2.5%)	6.55	170
Levofloxacin	50	CaCl ₂ (5%)	6.10	735
Levofloxacin	50	CaCl ₂ (5%), Lactose (5%)	6.21	1037
Levofloxacin	50	CaCl ₂ (2.5%), Lactose (5%)	6.36	565
Levofloxacin	50	CaCl ₂ (2.5%), Lactose (2.5%)	6.41	370
Levofloxacin	50	CaCl ₂ (1.25%), Lactose (2.5%)	6.64	227
Levofloxacin	50	CaCl ₂ (0.625%), Lactose (2.5%)	6.06	163

Table 18. Taste Test Scoring System.

Score	Taste	Tolerability
1	Comparable to saline	No cough sensation, no cough
1.25	Slightly more taste than saline	Slight cough sensation, no cough
1.5	Mild bitter/metallic taste	Cough sensation, slight cough
1.75	Between 1.5 and 2	-
2	Moderate bitter/metallic taste	Cough sensation, moderate cough
2.25	Between 2 and 2.5	-
2.5	Strong bitter/metallic taste	-
2.75	Between 2.5 and 3	-
3	Very strong bitter/metallic taste	Cough sensation and strong cough
4	Very strong bitter/metallic taste and other unacceptable taste	Cough sensation, strong cough and other irritation

Table 19. Taste and Tolerability of Levofloxacin Formulations Containing Sweeteners, Divalent Metal Salts, and Surface Active Agent.

Excipients	Taster							
	1		2		3		Median	
	Taste	Tol.	Taste	Tol.	Taste	Tol.	Taste	Tol.
Aspartame (0.1%)	2	1.25	2	1	2	1	2	1
Sucrose (0.1%)	2	1	1.75	1	2	1	2	1
Sucrose (7.5%); NaCl (0.225%)	2	1	2.25	1	2	1	2	1
Glucose (5%)	1.5	2	2.5	1	2	1	2	1
Glycerin (5%)	2.25	1	2.25	1	2.5	1	2.3	1
PS- 80 (0.1%)	1.75	1	2.25	1	2.5	1	2.3	1
CaCl ₂ (5%)	1.25	1	1.5	1.5	2	1	1.5	1
MgSO ₄ (5%)	1.5	1.5	2.5	2.5	2.5	1	2.5	1.5
Control -A (0.225% NaCl)	3	1	3	1	2.5	1	3	1

Table 20. Taste and Tolerability of Levofloxacin CaCl₂ Formulations.

Excipients	Taster							
	1		2		3		Median	
	Taste	Tol.	Taste	Tol.	Taste	Tol.	Taste	Tol.
CaCl ₂ (5%)	1.75	1	2	1	2.75	1	2	1
Sucrose (5%)	2	1	2	1	2	1	2	1
CaCl ₂ (5%), Sucrose (7.5%)	1.75	1	1.75	1	1.5	1	1.8	1
CaCl ₂ (5%), Glucose (7.5%)	1.5	1	1.5	1	2	1	1.5	1
CaCl ₂ (5%), Lactose (5%)	1	1	1.75	1	2	1	1.8	1
Control B-E (0.225% NaCl)	3	1	2.5	1	3	1	3	1

[00338] Test C: Taste Testing of Mono- and Disaccharides in the Presence of Magnesium Chloride. As above, all of the formulations screened in this experiment were well tolerated and tasted better than the control sample. Formulations containing both the magnesium salt and lactose appeared to perform slightly better than either one alone. This experiment confirms that combining divalent metal salts and simple sugars are effective at improving taste (Table 21).

Table 21. Taste and Tolerability of Levofloxacin MgCl₂ Formulations.

Excipients	Taster							
	1		2		3		Median	
	Taste	Tol.	Taste	Tol.	Taste	Tol.	Taste	Tol.
MgCl ₂ (5%)	1.5	1	--	-	1.75	1	1.6	1
MgCl ₂ (5%), Sucrose (7.5%)	1.5	1	1.75	1	2	1	1.8	1
MgCl ₂ (5%), Glucose (7.5%)	1.25	1	2.25	1	2	1	2	1
MgCl ₂ (5%), Lactose (5%)	1	1	1.5	1	1.5	1	1.5	1
Control B-E (0.225% NaCl)	2.25	1	-	-	2.75	1	2.5	1

[00339] Test D: Taste Testing of Mono- and Disaccharides in the Presence of Magnesium Sulfate. As with calcium and magnesium chloride, formulations containing magnesium sulfate and glucose, sucrose or lactose tasted better than the control sample. This experiment reconfirms that combining divalent metal salts and simple sugars improve taste (Table 22).

Table 22. Taste and Tolerability of Levofloxacin MgSO₄ Formulations.

Excipients	Taster							
	1		2		3		Median	
	Taste	Tol.	Taste	Tol.	Taste	Tol.	Taste	Tol.
MgSO ₄ , Sucrose	1.5	2	1.5	1.25	1.5	1	1.5	1.3
MgSO ₄ , Glucose	1.5	2.75	2	2.5	1.5	1.5	1.5	2.5
MgSO ₄ , Lactose	1.25	2.25	1.75	1.25	1.75	1	1.8	1.3
Control B-E (0.225% NaCl)	2.25	1	-	-	3	1	2.6	1

[00340] Test E: Taste Testing of Divalent Metal Salts in the Presence of Glucose at Low and High pH. In this experiment, the effect of glucose in combination with each of the three divalent cation salts on taste and tolerability was tested at low (≤ 5.5) and high (≥ 6.0) pH. Small but consistent improvements in taste were observed at the higher pH (Table 23).

Table 23. Taste and Tolerability of Levofloxacin CaCl₂ Formulations at Low Versus High pH.

Excipients	Taster							
	1		2		3		Median	
	Taste	Tol.	Taste	Tol.	Taste	Tol.	Taste	Tol.
CaCl ₂ (5%), Glucose (7.5%), pH6.1	1	1	1.5	1	2	1	1.5	1
CaCl ₂ (5%), Glucose (7.5%), pH 5.5	1.25	1	1.75	1	2.5	1	1.8	1
MgCl ₂ (5%), Glucose (7.5%), pH6.0	1.25	1	2	1	2	1	2	1
MgCl ₂ (5%), Glucose (7.5%), pH 5.0	1.75	1	1.75	1	1.5	1	1.8	1
MgSO ₄ (5%), Glucose (7.5%), pH 6.2	1.25	2.25	2.25	1.75	1.5	1	1.5	1.8
MgSO ₄ (5%), Glucose (7.5%), pH 5.4	1.5	1.75	1.75	1.5	2	1	1.8	1.5
Control B-E (0.225% NaCl)	2	1	-	-	-	-	-	-

[00341] Test F. Taste Testing of Mono- and Disaccharides. All of the formulations screened in this experiment were well tolerated and tasted better than the control sample. All three sugars at 5% were better than the control, lactose at 2.5% tasted better than the control, but not as good as at 5%. This experiment reconfirms that simple sugars improve taste (Table 24).

Table 24. Taste and Tolerability of Levofloxacin Sugar Formulations.

Excipients	Taster					
	1		3		Median	
	Taste	Tol.	Taste	Tol.	Taste	Tol.
Glucose (5%)	1.5	1.5	2	1	1.8	1.3
Sucrose (5%)	1.5	1.5	1.5	1	1.5	1.3
Lactose (5%)	1.75	1.25	2	1	1.9	1.1
Lactose (2.5%)	2.25	1.5	2	1	2.1	1.3
Control F-G (0.45%NaCl)	2.5	1	2.5	1	2.5	1

[00342] Test G. Taste and Tolerability of Levofloxacin CaCl₂ Formulations In the Presence of Lactose. In this experiment, levofloxacin was formulated with varying concentrations of calcium chloride and lactose (Table 25). As noted through this series of experiments, all formulations containing divalent metal salts and sugar were improved with respect to taste and tolerability relative to the control formulation. Most importantly, 5% calcium chloride or 2.5% calcium chloride in the presence of 5% lactose

were most effective at decreasing levofloxacin bitterness. Further decreases in the concentration of these excipients were less effective.

Table 25. Taste and Tolerability of Levofloxacin CaCl₂ Formulations in the Presence of Lactose.

Excipients	Taster					
	1		3		Median	
	Taste	Tol.	Taste	Tol.	Taste	Tol.
CaCl ₂ (5%)	1.25	1	1.5	1	1.4	1
CaCl ₂ (5%), Lactose (5%)	1.25	1	2	1	1.6	1
CaCl ₂ (2.5%), Lactose (5%)	1.25	1	2	1	1.6	1
CaCl ₂ (2.5%), Lactose (2.5%)	1.5	1	2.5	1	2	1
CaCl ₂ (1.25%), Lactose (2.5%)	1.75	1	2	1	1.9	1
CaCl ₂ (0.625%), Lactose (2.5%)	1.75	1.25	2	1	1.9	1.1
Control F-G (0.45%NaCl)	3	1	2.5	1	2.8	1

Example 3 – Aerosol Levofloxacin Characterization in PARI LC Plus Jet Nebulizer.

[00343] The following studies describe the potential for aerosolized delivery of levofloxacin to be administered to a patient via a jet nebulizer. To accomplish this task, a simple levofloxacin formulation was prepared and the aerosol was characterized in a jet nebulizer. The results of these studies are shown in the summary below.

[00344] Levofloxacin inhalation solution (55 mg/ml) was evaluated using a PARI LC Plus Air Jet Nebulizer with ProNeb Compressor. The emitted dose, particle size distribution and fine particle fraction were measured by cascade impaction using a Marple Miller Impactor. The above-mentioned parameters were used for evaluating the *in vitro* performance of aerosolized medications.

Marple Miller Study

[00345] Objective. To determine the particle size distribution and estimate the amount of drug that a patient is likely to inhale (respirable fraction). A secondary objective was to estimate emitted dose, which is the amount of levofloxacin that exited the nebulizer.

[00346] Methods. Formulation: 55 mg/ml levofloxacin, 120 mM chloride, 70 mM sodium, pH 6.7. Formulation established from maximum solubility permitting a 300 mg dosage in 6 ml and neutral pH. 5.5 ml of levofloxacin formulation was added to a PARI LC Plus Air-Jet Nebulizer with ProNeb Compressor. The nebulizer cup contained a total of 302 mg of levofloxacin. The nebulizer was connected inline with a Marple Miller Impactor (MMI), which was run with an airflow rate of 60 l/min. Each nebulizer (n=2) was run to dryness (no aerosol produced as judged by visual inspection for 15 minutes. Following aerosolization, the MMI was disassembled and levofloxacin was quantitatively extracted with mobile phase (90/10 ACN:water) from the USP entry port, each of the impactor collection cups (stages) and the glass fiber filter. Any remaining formulation in the nebulizer after aerosolization (cup and mouthpiece) was also quantified.

Results

[00347] As shown in Table 26, the total average amount recovered from the MMI experiments was 170.2 mg. The expected recovery was 302 mg. This represents a total recovery of ~57%, which does not meet the generally accepted specifications for impaction-based studies (85%-115% total recovery). This difference was found to be due to non-specific adherence of levofloxacin to the LC Plus nebulizer device. The average percent of drug exiting the nebulizer in fine particles was ~72%. Thus, the respirable emitted dose was 89.7 mg. Assuming that ~50% is not inhaled during normal tidal breathing, a total of ~40 mg may be deposited in the lung with this 300 mg dose. However, given the slow administration time with this device, competition with pulmonary clearance would likely prevent the accumulation of sufficient levofloxacin to meet the required minimal concentration for "rapid administration, high concentration" dosing needed for maximum fluoroquinolone antimicrobial activity and resistance prevention.

Table 26. Marple Miller Impactor Data Set.

Sample ID	A Emitted Dose (mg)	B Amount of Drug Remaining in Nebulizer Cup (mg)	A+B Amount of Drug Recovered (mg)	Percent of Total Drug Exiting Nebulizer in Fine Particle Fraction (% < 5 μ m)
Levo Run 1	134.70	45.00	179.70	73.5
Levo Run 2	114.40	46.30	160.70	70.4
Average	124.6	45.7	170.2	72.0

Example 4 – Animal Models and Evaluation of Fluoroquinolones and Fuoroquinolone Formulations.

Pharmacokinetic Model

[00348] Six rats per study are given a single slow bolus intravenous dose of 10 mg/kg via the lateral tail vein or are given a single microspray aerosol dose of 10 mg/kg using a microspray aerosol generation device (PennCentury, Philadelphia, PA). Blood samples are taken at various times over 3 hours to determine the plasma pharmacokinetic parameters. Two rats are sacrificed at 0.5, 1.5 and 3 hours after dosing to determine lung, bronchoalveolar lavage (BAL), and epithelial lining fluid (ELF) levels. The plasma and tissue concentrations are determined by an HPLC method and the data are then fit using WinNonlin. Data are shown in Table 27.

Efficacy Model

[00349] *P. aeruginosa* strain PAM 1723 is grown in Mueller-Hinton Broth (MHB) at 35°C under constant aeration, after 16 hours, the inoculum is sub-cultured into fresh MHB and allowed to regrow at 35°C, under constant aeration, for 4 hours. The inoculum is adjusted to ca. 5×10^6 CFU/ml by correlation of absorbance at 600 nm with predetermined plate counts. Male CFW mice (4 – 6 weeks old, N= 4/group) are made neutropenic by the intraperitoneal injection of 150 mg/kg cyclophosphamide (Cytosan, Mead Johnson, Princeton, NJ) on days 1 and 4. On day 5, mice are infected by an intratracheal instillation of 0.05 ml of inoculum while under isoflurane anesthesia (5% isoflurane in oxygen running at 4 L/min). Two hours after infection, mice are given either intraperitoneal or intratracheal doses of each fluoroquinolone at a dose of 25 mg/kg.

Mice are sacrificed 1 and 4 hours after treatment, their lungs removed, homogenized and plated to determine colony counts. Data are shown in Table 28.

Table 27. Pharmacokinetic Modeling.

Drug	Route	Dose (mg/kg)	Serum AUC (0-inf)	Serum t _{1/2}	ELF AUC (0.5-3h)	F, % from Lung vs. IV
Levofloxacin	IV	10	3.8	0.5	10.5	NA
Levofloxacin	IT	10	3.28	0.4	12.07	86%
Ciprofloxacin	IV	10	2.56	0.53	ND	NA
Ciprofloxacin	IT	3.3	0.8	0.93	194	82%
Clinafloxacin	IT	10	3.2	0.74	30.8	
Gatifloxacin	IV	10	5.31	1.06	5.32	
Gatifloxacin	IT	10	5.83	1.13	54.7	100%
Norfloxacin	IV	10	4.65	1.21	3.27	
Norfloxacin	IT	10	4.46	1.13	41.7	100%
Gemifloxacin	IV	8	4.54	1.04	3.72	
Gemifloxacin	IT	10	5.86	1.68	536.5	86%
Tobramycin	IV	10	15.7	0.5	27.6	NA
Tobramycin	IT	10	13.82	1.0	5152.0	81%

[00350] In rat pharmacokinetic studies, aerosol administration of fluoroquinolones results in increased ELF AUCs from 0.5 – 3 hours for all fluoroquinolones tested, as well as tobramycin, suggesting that the aerosol route of administration will produce increased efficacy against lung infections.

[00351] In a mouse lung infection model, the increased efficacy, suggested by the pharmacokinetic studies rats, was confirmed. For all fluoroquinolones tested, the aerosol route of administration (intratracheal, or IT), produced larger reductions in bacterial counts than the intraperitoneal (IP) route of administration, suggesting that observed increased efficacy was due to high local concentrations produced by direct aerosol administration.

Table 28. Efficacy Modeling.

Drug	Route ^a	Dose (mg/kg)	DeltaLOG CFU 1 hr ^b	DeltaLOG CFU 4 hr ^b
Levofloxacin	IP	25	-1.00	-0.52
Levofloxacin	IT	25	-1.97	-1.28
Gemifloxacin	IP	25	-0.28	-0.32
Gemifloxacin	IT	25	-2.45	-1.81
Levofloxacin	IP	25	-1.40	-1.14
Levofloxacin	IT	25	-2.48	-1.45
Gemifloxacin	IP	25	-0.74	-0.71
Gemifloxacin	IT	25	-3.20	-2.28
Clinafloxacin	IP	25	-1.32	-1.33
Clinafloxacin	IT	25	-2.86	-2.47
Tobramycin	IP	5	-0.70	0.29
Tobramycin	IT	5	-1.59	-0.94
Ciprofloxacin	IP	25	-1.59	-0.41
Ciprofloxacin	IT	25	-2.32	-1.45
Gatifloxacin	IP	25	-0.34	-0.02
Gatifloxacin	IT	25	-1.48	-2.11
Clinafloxacin	IP	10	-0.96	-1.39
Clinafloxacin	IT	10	-2.71	-2.40
Sparfloxacin	IP	25	-0.85	0.09
Sparfloxacin	IT	25	-1.56	-0.81
Tosufloxacin	IP	25	0.00	1.33
Tosufloxacin	IT	25	-0.48	-0.24

a. Route of drug administration.

b. Time post-drug administration.

Example 5 – Aerosol Levofloxacin Characterization in the PARI eFlow Nebulizer.

Laser Particle Sizing

[00352] Device performance was characterized by measuring the size of the particles emitted. By non-limiting example, particle sizing of emitted aerosol of Levofloxacin solution may be conducted with a Malvern Spraytec particle sizer under the following conditions. Ambient conditions are controlled to maintain a room temperature

of between 23.0°C and 24.0°C and relative humidity of 42% to 45%. Levofloxacin at 25 mg/ml was loaded into 2 PARI eFlow Nebulizers fitted with the "40" nebulizing heads. Software for the Malvern Spraytec particle sizer is programmed to calculate the following information. A) Volume Mean Diameter (VMD), the volume mean of the particles passing across the beam of the laser. B) Geometric Standard Deviation (GSD), diameter 84th percentile/diameter 50th percentile. C) % of particles ≤ 5 microns, the percent of the number of particles less than 5 microns or % of particle >1 micron and < 7 microns, the percent of the number of particles between 1 and 7 microns.

[00353] The device was loaded with 2 ml Levofloxacin at 25 mg/ml. The mouthpiece of the device was positioned with the tip of the mouthpiece 2 cm from the center of the beam on the x axis and as close to the optical lens of the laser as possible on the y axis. Ambient conditioned, bias flow was provided through the nebulizer in an amount to obtain a total nebulizer flow of 20 LPM. Ambient conditioned, bias flow was provided through the nebulizer in an amount to obtain a total nebulizer flow of 20 LPM. The nebulizer was turned on and allowed to run continuously for 1 minute prior to measuring. The measurement sequence was begun after 1 minute and measurements are made continuously for 1 minute in 1 second intervals. At the end of the measurement phase, these 60 records are averaged for VMD, GSD and % ≤ 5 micron and % >1 and <7 micron. Finally, the nebulizer was weighed for determination of output rate.

Breath Simulation Studies

[00354] Device performance was measured under conditions similar to natural inhalation by using a breath simulator PARI Compas breath Simulator programmed to use the European Standard pattern of 15 breaths per minute with an inspiration to expiration ratio of 1:1. Such measurements was performed under ambient conditions that can be controlled to maintain a room temperature of between 23.0°C and 24.0°C and relative humidity of 42% to 45%. For this experiment, the PARI eFlow device was loaded with 4 ml Levofloxacin solution at 25 mg/ml.

[00355] Breathing simulation was commenced, and the nebulizers begun. The devices were allowed to run continuously until nebulization ceases. The duration is timed from the beginning of nebulization. Following nebulization, the inspiratory and

expiratory filters were individually washed in a known amount of solvent (dH₂O). The nebulizer cup is also washed individually. For quantitation, the individual washings were assayed via spectrophotometry at a wavelength of 290 nanometers and the resultant concentration converted to content. Using this quantitative data, the following analysis was made. A) Inspired dose (ID), the total amount of drug assayed from the inspiratory filter. B) Residual dose (RD), the amount of drug assayed from the nebulizer at the end of nebulization. C) Fine Particle Dose (FPD), the ID multiplied by the respirable fraction (for example, % particles \leq 5 microns VMD depending on the method used to determine the size of the particles emitted from the selected device). D) Duration, time from the beginning to the end of nebulization. E) Respirable Delivered Dose (RDD), % ID that is, for example, \leq 5 microns VMD.

[00356] The results in Table 29 indicate that a 100 mg dose of levofloxacin likely deposits ~34 mg fluoroquinolone in the pulmonary compartment in ~4 min using the PARI eFlow device (Table 29) compared to the 300 mg dose from the PAR LC Plus device delivering an equivalent dose in >15 min. From the “rapid administration, high concentration” dosing and delivery model described herein, while the 15 min delivery time from the LC Plus will likely fail, a 4 min administration time for 35-40 mg levofloxacin may meet the criteria for maximum fluoroquinolone activity. However, increasing the drug concentration to enable more rapid administration (e.g. 50 mg/ml in a 2 ml dosing delivering 35-40 mg levofloxacin in ~2 min) will more likely meet these minimal requirements. Moreover, shorter administration times will improve patient dosing compliance. In addition, it should be noted that hypotonic solutions of levofloxacin at concentrations greater than 10 mg/ml are poorly tolerated for inhalation.

Table 29. Levofloxacin Aerosol Properties (100 mg Loading Dose).

Duration (minutes)	Residual Dose	Inspired Dose	FPD (%)		RDD (mg)		VMD um	GSD um	Osmo mOs/kg
			\leq 5u	1-7u	\leq 5u	1-7u			
3.9 \pm 0.1	24.8 \pm 3.4	61.1 \pm 1.6	54.9	73.8	33.5	45.1	4.7	1.6	67 \pm 1.0

Example 6 – Tolerability of Aerosol Levofloxacin in a Healthy Human Subject.**Methods**

[00357] In a single subject, healthy volunteer the feasibility of delivering levofloxacin as an aerosol was established using either an Aerogen Clinical vibrating mesh device creating a 3.4 micron volumetric mean diameter (VMD) particle, or ~2 micron MMAD (hereinafter “Aerogen Small”), or using a PARI eFlow nebulizer producing a ~4.7 micron VMD particle (hereinafter “PARI Large”). Levofloxacin was tested at a concentration of 4.25 mg/mL or 18.75 mg/mL at doses of 10 mg, 35 mg and 55 mg, in isotonic solution.

Results

[00358] In the first test, 6 mL of the 4.25 mg/mL solution was inhaled using the Aerogen Small nebulizer. The estimated RDD based on separate in vitro device characterization studies using breath simulation was estimated to be 10 mg. Delivery time was 22 minutes. No discernable adverse effects were observed in the throat, airway or lungs, during or after administration, including cough sensation or cough, and there was only a slight chemical taste during and after administration. No adverse effects or taste were observed over a 30 minute monitoring period following drug administration. At this low concentration and dose, and slow rate of administration, levofloxacin was well tolerated.

[00359] In the second test, 4 ml of the 18.75 mg/mL solution was inhaled using the Aerogen Small nebulizer. The estimated RDD based on separate in vitro device characterization studies using a breath simulator was 35 mg. Delivery time for administration of the drug was 14 minutes. Despite the increased dose, the acute tolerability was very comparable to the first test both during and after administration. The taste, which was stronger, was the solution had a more the bitter/metallic chemical taste characteristic of levofloxacin. The taste was most discernible for a period of a few minutes after the end of administration, again a characteristic of levofloxacin.

[00360] In the third test, 4 mL of the 18.75 mg/mL solution was inhaled using the PARI Large device. The estimated RDD based on separate in vitro device

characterization studies was ~55 mg (using the <5 microns FPD definition). Delivery time for administration of the drug was ~5 minutes. Despite the significantly increased particle size and delivery rate for drug compared to test 2, no adverse effects in the throat, airway or lungs, other than the acute effects of taste noted above, were experienced, including cough sensation or coughing, throughout the dosing period and for a 30 minute observation period following delivery of the dose. Urinary recovery of the drug, which is an accurate measure of exposure, confirms that the projected respirable dose of approximately 55 mg was successfully delivered.

[00361] These results demonstrate the feasibility of aerosol delivery of levofloxacin in a human subject at the intermediate concentrations tested, and suggest that higher concentrations and doses, properly formulated for tolerability and taste are achievable.

Example 7 – Levofloxacin Micronization.

Levofloxacin Micronization

[00362] Dry powder levofloxacin base may be micronized for high local concentration exposure therapy, taste-masking or AUC shape-enhanced delivery of levofloxacin using dry powder pulmonary administration. Other approaches currently being investigated include spray-dry and in situ micronization techniques. This approach may also be used with other fluoroquinolone antibiotics including, without limitation ofloxacin, lomefloxacin, pefloxacin, ciprofloxacin, gatifloxacin, gemifloxacin, moxifloxacin, tosufloxacin, pazufloxacin, rufloxacin, fleroxacin, balofloxacin, sparfloxacin, trovafloxacin, enoxacin, norfloxacin, clinafloxacin, grepafloxacin, sitafloxacin, marbofloxacin, orbifloxacin, sarafloxacin, danofloxacin, difloxacin, enrofloxacin, garenoxacin, prulifloxacin, olamufloxacin, DX-619, TG-873870 and DW-276.

Description

[00363] To characterize the feasibility of micronizing levofloxacin base, the following studies were performed.

Micronization

[00364] Levofloxacin drug powder was micronized using a jet mill. Following micronization the drug powder was collected in two fractions, one between 5-6 micron and a finer fraction.

Powder Characterization

[00365] The drug was characterized for particle size and particle size distribution, before and after milling using laser diffraction technology. Any changes in the physical form of the drug were evaluated by Differential Scanning Calorimetry (DSC) and X-Ray diffraction (XRD). Particle morphology was studied using Scanning Electron Microscopy (SEM). The equilibrium moisture content of the drug powder before and after micronization was determined by Thermogravimetric Analysis (TGA) or Karl Fischer. Any degradation of the drug substance during micronization was evaluated by HPLC. The separation conditions were used to determine if any new peaks are formed after micronization.

Micronization

Experimental Methodology

[00366] Two batches of Levofloxacin were micronized using a jet mill (Glen Mills). Method development was performed to determine the micronization pressure required to achieve the required size fractions between a) 5-6 micron and b) 2-3 micron. The particle size of levofloxacin was determined by Sympatec HELOS laser diffraction particle size analyzer.

Results

[00367] Figure 12 shows the plot of mean particle diameter (X50) Vs micronization pressure. The first batch of Levofloxacin showed a mean particle diameter of 10.6 micron before micronization. From the plot it is seen that particle size decreased as the micronization pressure increased. A pressure of about 120 psi was required to achieve a size of 2.5 micron. With the second batch of levofloxacin, having a mean

particle size of 12.99 micron before micronization, a pressure of 30 psi was required to achieve a particle size of 5.2 micron.

Powder Characterization

Differential Scanning calorimetry

Experimental Methodology

[00368] Differential Scanning Calorimetry of pre-micronized as well as micronized levofloxacin (mean particle size 2.5 micron) was performed using TA Instrument DSC Q1000. 1-2 mg of each sample was weighed into pan, sealed and heated at 10°C/min from 25°C to 300°C under nitrogen.

Results

[00369] The DSC profiles of the pre-micronized and micronized levofloxacin are shown as Figure 13. There was no difference in the DSC profiles of micronized compared to pre- micronized levofloxacin.

Experimental Methodology

[00370] The powders (micronized and pre-micronized) were adhered to double-sided carbon tabs on aluminum stubs, which were then coated with gold-palladium. Photomicrographs were taken of several different areas of the powder on the stub using a Scanning electron microscope.

Results

[00371] Representative Scanning electron micrographs of the pre-micronized and micronized levofloxacin are shown in Figure 14A and 14B. Crystals of levofloxacin are plate like before micronization. This shape is retained after micronization.

Experimental Methodology

[00372] A thin layer of powder sample was mounted on a zero background plate in an XRD sample holder. Each sample was analyzed using a Scintag XDS 2000 Diffractometer under the following conditions:

Excitation source: Copper K α X-rays; Scan rate: 1° per minute
Voltage: 40 KV; Curent: 35 mA.

Results

[00373] The X-ray diffraction plots of pre-micronized and micronized Levofloxacin are shown as Figure 15. Intensity of diffracted peak at 9° is reduced after micronization. These results are in accordance to those reported in the literature for micronization of olanzapine (Stephenson G,A. The Rigaku Journal, 22 (2005): 2-15). The reduction in the relative intensities of the diffracted peaks might be due to formation of new faces to a crystal. The most developed face after micronization would be the one for which the intensity is maximally reduced.

Experimental Methodology

[00374] 15-25 mg of micronized and pre-micronized levofloxacin samples were dissolved in methanol (having predetermined moisture content) and the moisture content in the samples were determined by Aquastar 3000 Coulometric Karl Fisher Titrator.

Results

[00375] The results of Karl Fisher analysis are shown in Table 30.

Table 30. Moisture Content of Pre and Post Micronized Levofloxacin.

	Moisture content (%)
Pre- micronized	6.16
Post- micronized	5.42

Example 8 - Preformulation of Levofloxacin Base.

[00376] The goal of this study was to characterize levofloxacin base to understand the physico-chemical capabilities and restrictions of levofloxacin base for various formulation approaches. The purpose of this study was to characterize the physicochemical properties of levofloxacin base.

Preformulation**pH-Solubility Studies**

[00377] The solubility of levofloxacin was determined as a function of pH. Buffers were first prepared in the pH range 2-10. Small aliquots of each buffer (~200-

250 μ L) were saturated with drug and agitated to achieve equilibrium solubility. The samples were then becentrifuged and the supernatant analyzed for dissolved drug by UV or HPLC. The buffers used in this study were shown to affect the solubility result (because different buffer counter-ions can form different levo salt forms in solution). Hence, pH-solubility will also be assessed in the absence of buffers (via titration).

pKa Determination

[00378] The pKa of levofloxacin was determined by titrimetry. Obtained pKa values were confirmed by UV spectrophotometry. This information was used to aid in salt selection for levofloxacin and to determine the charge on levofloxacin under the pH conditions in the lung.

Preformulation for Liquid System

[00379] The feasibility of a liquid formulation was investigated using (a) solubility and (b) surface tension as baseline parameters for formulation in saline alone.

Preformulation Studies on Levofloxacin

HPLC Method Transfer

Experimental Methodology

[00380] A HPLC method was used to evaluate the linearity, accuracy and precision of the levofloxacin assay. The column used was a 50 X 4.6 mm, Onyx Monolithic C18 (Phenomenex) at 30o C. The mobile phase consisted of 85% of 0.1% TFA in water and 15 % 0.1% TFA acetonitrile. The flow rate was adjusted to 3 ml/min. Samples were injected into the chromatographic system and the effluent monitored at 277 nm.

Results

[00381] The retention time for levofloxacin was approximately 0.82 min. The assay was found to be linear over a range of 5-15 μ g/ml, with a correlation coefficient of 1.000. RSD (relative standard deviation) was less than 0.5 % and accuracy was within 98-102%.

pH- Solubility Studies

By Titration

Experimental Methodology

[00382] A saturated solution of levofloxacin in 0.1 N HCl was titrated with NaOH. After each addition of the base, the solution was shaken by vortexing. An aliquot of the sample solution was removed, centrifuged and the supernatant analyzed by UV spectroscopy at 288 nm. The same solution was back titrated with HCl.

Results

[00383] The pH-solubility profile of levofloxacin is shown in Figure 16. By titrimetry levofloxacin exhibited a solubility of 25.4 mg/ml at pH 7.3. However contrary to the results of the shaking experiments, the solubility by titrimetry decreased below pH 6.5 which can be attributed to the common ion effect. Since a solution of levofloxacin was prepared in HCl, a hydrochloride salt of levofloxacin would have formed in solution. Further addition of chloride ions in the form of hydrochloric acid would suppress the solubility of the hydrochloride salt.

pKa determination

By Titrimetry

Experimental Methodology

[00384] A solution of levofloxacin (18 mg/g) was prepared in water (18.45 mg/g). The initial pH of the solution was 7.36. This solution was titrated with 1 N HCl. Measured aliquots of HCl were added and the pH recorded after each addition. Titration was continued till a pH of 1.

[00385] In order to determine the acidic pKa, a solution of levofloxacin (18.38 mg/g) was prepared in 0.1 N HCl. The initial pH of the solution was 1.32. The solution was titrated with 1 N NaOH. Titration was continued till a pH of 6.55.

Results

[00386] Figure 17 shows a plot of pH Vs volume of titrant added for the titration of levofloxacin with HCL. This data was fit into the following equation:

$$V_t [\text{OH}^-] = K_b \cdot V_{ep} - K_b \cdot V_t$$

where,

$$V_t = \text{Volume of titrant added}$$

V_{ep} = volume of titrant added till the equivalence point

$$[\text{OH}^-] = \text{hydroxide ion concentration} = K_w / [\text{H}^+]$$

$$[\text{H}^+] = \text{hydronium ion concentration} = 10^{-\text{pH}}$$

[00387] A plot of $V_t [\text{OH}^-]$ Vs V_t gave a straight line (Figure 18). Data shown is from the pre-equivalence point region. From the slope we get

$$\text{Slope: } K_b = 2.09 \times 10^{-8}$$

$$\text{p}K_b = -\log K_b = 7.7$$

$$\text{p}K_a = 14 - \text{p}K_b = 6.3$$

[00388] Figure 19 shows a plot of pH Vs volume of titrant added for the titration of levofloxacin with NaOH. Acidic pKa was difficult to calculate because it was quite low (< 2.0). However, a rough approximation of pKa can be made as the pH at half the equivalence point. From the plot dpH/dV vs volume of titrant (V_t) (Figure 20), the equivalence point is at 250 μl . The pH at half the equivalence point (i.e. when $V_t = 125 \mu\text{l}$) is 1.6. So the acidic pKa \sim 1.6.

By UV spectroscopy

Experimental Methodology

[00389] Dilute solutions of levofloxacin (0.013 mg/ml) were prepared in several buffers. The buffers used were HCL (pH 1,2), acetate (pH 4,5), phosphate (pH 6,7,8) and borate (9,10). The levofloxacin solutions were analyzed by UV spectroscopy at 257 nm.

Results

[00390] A plot of pH Vs Absorbance of the levofloxacin solution at 257 nm is shown as Figure 21. This data was fitted into a modified Henderson Hasselbach equation:

$$\text{Abs}_{\text{observed}} = \text{Abs}_{\text{HA}} \frac{[\text{H}^+]}{K_a + [\text{H}^+]} + \text{Abs}_{\text{A}^-} \frac{[\text{H}^+]}{K_a + [\text{H}^+]}$$

where,

$$\text{Abs}_{\text{observed}} = \text{Absorbance of levofloxacin solution}$$

Abs_{HA} = Absorbance of levofloxacin solution pH 1.2 ;

Abs_{A^-} = Absorbance of levofloxacin solution at pH 7.8;

$[H^+]$ = hydronium ion concentration = 10^{-pH}

The fitted equation provided an estimate of $pK_a = 5.91$.

Example 9 – Levofloxacin Salt Formation.

[00391] The goal of this study was to prepare various salt forms of levofloxacin that may gain AUC shape-enhancement properties through decreased solubility and/or dissolution. These benefits may alter the pharmacodynamic properties of levofloxacin following pulmonary administration using nanoparticle suspension or dry powder inhalation. These formulations may be optimized to prolong the release of levofloxacin from decreased solubility salt forms. These properties may also be incorporated into other fluoroquinolone antibiotics including, without limitation gemifloxacin, gatifloxacin, norfloxacin, tosufloxacin, sitafloxacin sarafloxacin, prulifloxacin, and pazufloxacin. Studies are underway to characterize various salt forms and co-precipitates of gemifloxacin for taste masking, AUC shape-enhancement, nanoparticle suspension and dry powder inhalation administration. Other approaches currently being investigated include spray-dry and in situ micronization techniques.

[00392] For suspension and powder formulations, a specific salt form can provide important physical and chemical characteristics that may have impacts on the product performance. For a AUC shape-enhancing formulation, the objective of salt selection was to decrease the solubility and/or reduce the dissolution rate of levofloxacin. The acid counterions can be selected by:

[00393] Manipulation of melting point: An increase in melting point is usually accompanied by a reduction in salt solubility. Salts formed from planar, high melting aromatic acids generally yield crystalline salts of high melting point.

[00394] Manipulation of hydrophobicity: Salts formed with hydrophobic conjugate acids are hydrophobic and difficult to wet, and may ultimately lead to prolonged dissolution. Examples of acids have been selected for salt preparation are listed as follows:

- a) Pamoic acid (embonic acid)
- b) 2-naphthalene sulfonic acid (napsylic acid)
- c) Oleic acid
- d) Xinafoic acid
- e) Stearic acid
- f) Lauryl Sulfonate (estolate)

[00395] Other factors considered include salt surface properties, polymorph and chemical stability.

Description

[00396] The objective of the study was to prepare salt forms of levofloxacin in order to reduce its solubility and/or dissolution rate. The goal was to:

(a) make levofloxacin less soluble through salt formation with suitable excipient(s).

(b) prepare salt forms of levofloxacin that will have a lower solubility and/or dissolution rate than the free base.

[00397] To accomplish these tasks, efforts were concentrated on preparing salts at the basic site of the molecule (pKa ~ 6.8).

[00398] Pamoic acid (mp = 280°C) and napsylic acid (mp = 125°C) possess planar, hydrophobic structures that were expected to provide hydrophobic character to the salt. The high melting point of pamoic acid may yield a high melting crystalline salt form. Oleic acid was chosen primarily because it is approved for lung delivery. It has a low melting point (4°C) which may not satisfy the condition under (1), however it was hoped that the long aliphatic chain may impart sufficient hydrophobicity to decrease solubility. Xinafoic acid (mp= 195°C) was also selected for salt formation as it also possess planar, hydrophobic structures that is expected to provide hydrophobic character to the salt. The rationale for choosing stearic acid and lauryl sulfonate (estolate) was similar to oleic acid, only their lung toxicities are unknown. Estolate is approved for oral delivery (erythromycin estolate has approximately 1/12th the solubility of the free base, and is formulated as an oral suspension).

Salt Formation

[00399] In general, levofloxacin base and the acid were dissolved in a suitable volatile, organic solvent (1:1 molar ratio), and stirred at room temperature. Any crystallized product formed was filtered, dried, and characterized. Characterization would consist of DSC, FTIR, and elemental analysis.

Formation and Characterization of Co-Crystals of Levofloxacin with Pamoic Acid

Experimental Methodology

Formation of Co-Crystals of Levofloxacin with Pamoic Acid

[00400] 0.31 g (0.8 mM) of pamoic acid was dissolved by stirring in 100 ml of tetrahydrofuran (THF). To this, 0.30 g (0.8 mM) of levofloxacin was added, dissolved by stirring and the resulting solution refluxed for 2.5 hrs. The suspension formed was cooled to room temperature, filtered and the precipitate obtained was dried in vacuum at about 70°C for 3 hours.

Characterization

[00401] Thermal analysis. Thermal analysis of (a) pamoic acid (b) levofloxacin (c) levofloxacin pamoate co-crystallized precipitate (d) physical mixture of pamoic acid and levofloxacin was performed using a Differential Scanning Calorimeter (TA Instrument DSC Q1000). 2-5 mg of each sample was weighed into pan, sealed and heated at 10°C/min from 25°C to 300°C under nitrogen.

[00402] Fourier transform Infrared (FT-IR) spectroscopy. FT-IR spectroscopy of (a) pamoic acid (b) levofloxacin (c) levofloxacin pamoate co-crystallized precipitate (d) physical mixture of pamoic acid and levofloxacin was carried out using a FTIR spectrometer (Model IRPrestige-21, Shimadzu).

[00403] Saturation solubility. Saturation solubility of levofloxacin and levofloxacin pamoic acid co-crystallized precipitate was determined by equilibrating excess amount of solid with water. The suspensions were adjusted to pH's 4, 5, 6 and 7 with HCL, shaken, centrifuged and the supernatant analyzed by UV spectroscopy at 288 nm.

Results

[00404] Thermal analysis. DSC scans of (a) pamoic acid (b) levofloxacin (c) levofloxacin pamoate co-crystal precipitate (d) physical mixture of pamoic acid and levofloxacin are shown as Figure 22. Pamoic acid and levofloxacin show sharp endotherms at 330°C and 239 °C, respectively, which most likely would be due to the melting of pamoic acid and levofloxacin, respectively. The DSC profile of levofloxacin pamoate co-crystals showed one major endotherm at 210° C, while a 1:1 molar admixture of levofloxacin and pamoic acid displayed broad endotherms at 129°C and 220° C.

[00405] FTIR. FTIR spectra obtained from (a) pamoic acid (b) levofloxacin (c) levofloxacin pamoate co-crystallized precipitate (d) physical mixture of pamoic acid and levofloxacin are displayed as Figure 23. The high intensity absorption bands at 1650 cm⁻¹ in the FTIR spectra of pamoic acid, which is due to the stretching of C=O group is greatly reduced in the co crystal.

[00406] Saturation solubility. Table 31 displays the saturation solubility data of levofloxacin and levofloxacin pamoate at different pH's. Solubility was determined in water, since buffer acids have an effect on the solubility of levofloxacin. However after shaking levofloxacin or salt solutions in water, the pH shifted, especially the levofloxacin solution at pH 5 shifted to pH 1.6. As the solution at pH 5 is between the two pKa's of levofloxacin (~1.6 and ~6), such a solution will have a lower buffer capacity and hence the pH shift. Solutions at pH's near the pKa's of the drug have a high buffer capacity and resist pH changes. Solubility of levofloxacin pamoate was considerably less than that of levofloxacin at all pH's.

Table 31. Saturation Solubility Data of Levofloxacin and Levofloxacin Pamoate.

Levofloxacin	pH before shaking	3.97	5.06	5.96	
	pH after shaking	4.21	1.63	5.70	
	Solubility (mg/ml)	158.28	225.04	297.20	
Levofloxacin pamoate	pH before shaking	4.06	4.98	5.98	7.00
	pH after shaking	5.63	6.30	6.66	7.32
	Solubility (mg/ml)	0.34	0.29	0.24	0.44

Interpretation

[00407] Since the co-crystallized precipitate of levofloxacin pamoate has a different melting point and FTIR spectra from that of levofloxacin, pamoic acid or their physical mixture, it is possible that the equimolar complex of the levofloxacin with pamoic acid might be the salt levofloxacin pamoate, having a considerably less solubility than levofloxacin.

Formation and Characterization of Co-Crystals of Levofloxacin with Xinafoic Acid.**Experimental Methodology*****Formation of Co-Crystals of Levofloxacin with Xinafoic Acid***

[00408] 1.004 g (2.7 mM) of levofloxacin was dissolved by refluxing in 80 ml ethyl acetate. To this, 0.51 g (2.7 mM) of xinafoic acid dissolved in 35 ml of ethyl acetate was added and the solution cooled overnight under stirring conditions to ambient temperature. The suspension obtained was filtered, and the precipitate dried under vacuum at 75°C for about 3.5 hours

Characterization

[00409] Thermal analysis. Thermal analysis of (a) xinafoic acid (b) levofloxacin xinafoate co-crystallized precipitate was performed using a Differential Scanning

Calorimeter (TA Instrument DSC Q1000). 2-5 mg of each sample was weighed into pan, sealed and heated at 10°C/min from 25°C to 300°C under nitrogen.

[00410] Fourier transform Infrared (FT-IR) spectroscopy. FT-IR spectroscopy of (a) xinafoic acid (b) levofloxacin xinafoate co-crystallized precipitate was carried out using a FTIR spectrometer (Model IRPrestige-21, Shimadzu).

[00411] Saturation solubility. Saturation solubility of levofloxacin xinafoic acid co-crystallized precipitate was determined by equilibrating excess amount of solid with water. The suspensions were adjusted to pH's 4, 5, 6 and 7 with HCL, shaken, centrifuged and the supernatant analyzed by UV spectroscopy at 288 nm.

Results

[00412] Thermal analysis. DSC profiles of (a) xinafoic acid (b) levofloxacin xinafoate co-crystallized precipitate are shown as Figure 24. Levofloxacin xinafoate co-crystallized precipitate exhibits a melting endotherm at 196°C, which is different than that of xinafoic acid (216°C) and levofloxacin (239°C).

[00413] FTIR. FTIR spectra obtained from (a) xinafoic acid (b) levofloxacin xinafoate co-crystallized precipitate are displayed as Figure 25. The FTIR spectrum of the co crystal exhibit transmittance minima's at wave numbers different from that of xinafoic acid and levofloxacin.

[00414] Saturation solubility. Table 32 shows the saturation solubility data of levofloxacin xinafoate at different pH's. Solubility of the xinafoate salt was intermediate between that of the levofloxacin base and levofloxacin pamoate co crystal.

Table 32. Saturation Solubility Data of Levofloxacin Xinafoate Co-Crystal.

Levofloxacin xinafoate	pH before shaking	4.05	4.94	6.00	6.95
	pH after shaking	4.78	5.37	6.30	7.17
	Solubility (mg/ml)	21.69	12.95	4.59	5.95

Interpretation

[00415] The co-crystallized precipitate of levofloxacin pamoate has a different melting point and FTIR spectra than that of levofloxacin and xinafoic acid, suggesting a possible formation of a levofloxacin xinafoate salt. This salt has intermediate solubility between levofloxacin and levofloxacin pamoate.

Formation and Characterization of Co-Crystals of Levofloxacin with Stearic Acid

Experimental Methodology

Formation of Co-Crystals of Levofloxacin with Stearic Acid

[00416] 0.77 g (2.07 mM) of stearic acid was dissolved by heating and sonication in 40 ml of methanol. To this, 1.00 g (2.07 mM) of levofloxacin dissolved in 60 ml of methanol was added. The resulting solution was heated at 55°C for about 15 minutes, followed by cooling to room temperature and then at -20°C. The suspension obtained was filtered.

Characterization

[00417] Thermal analysis. Thermal analysis of (a) stearic acid (b) levofloxacin stearate co-crystallized precipitate (c) physical mixture of stearic acid and levofloxacin was performed using a Differential Scanning Calorimeter (TA Instrument DSC Q1000). 2-5 mg of each sample was weighed into pan, sealed and heated at 10°C/min from 25°C to 250°C under nitrogen.

[00418] Fourier transform Infrared (FT-IR) spectroscopy. FT-IR spectroscopy of (a) stearic acid (b) levofloxacin stearic acid co-crystallized precipitate (d) physical mixture of stearic acid and levofloxacin was carried out using a FTIR spectrometer (Model IRPrestige-21, Shimadzu).

[00419] Saturation solubility. Saturation solubility of levofloxacin and levofloxacin stearic acid co-crystallized precipitate was determined by equilibrating excess amount of solid with water. The suspensions were adjusted to pH's 4, 5, 6 and 7 with HCL, shaken, centrifuged and the supernatant analyzed by UV spectroscopy at 288 nm.

Results

[00420] Thermal analysis. DSC scans of (a) stearic acid (b) levofloxacin stearate co-crystal precipitate (d) physical mixture of stearic acid and levofloxacin are shown as Figure 26. Stearic acid and levofloxacin show sharp endotherms at 76.4 °C and 239 °C, respectively, which most likely would be due to the melting of stearic acid and levofloxacin, respectively. The DSC profile of levofloxacin stearic acid co-crystals showed two sharp endotherms at 88.03°C and 138.54°C and minor endotherms at 231°C and 242.72°C. The minor endotherms might be due to melting of trace quantities of residual levofloxacin in the original sample. A 1:1 molar admixture of levofloxacin and stearic acid displayed endotherms at 68.87° C, 134.43°C and 240.74°C and minor endotherms at 79.73°C and 86.74°C.

[00421] FTIR. FTIR spectra obtained from (a) stearic acid (b) levofloxacin stearic acid co-crystallized precipitate (c) physical mixture of stearic acid and levofloxacin are displayed as Figure 27. The C=O stretch band is seen at 1700, 1705 and 1721 cm⁻¹ in stearic acid, co-crystallized precipitate and physical mixture, respectively.

[00422] Saturation solubility. Table 33 shows the saturation solubility data of levofloxacin stearic acid co-crystals at different pH's.

Table 33. Saturation Solubility Data of Levofloxacin Stearic Acid Co-Crystals.

Levofloxacin stearate	pH before shaking	4.05	5.02	6.01	6.96
	pH after shaking	3.36	5.05	6.02	6.98
	Solubility (mg/ml)	0.86	1.30	2.20	1.64

Interpretation

[00423] The DSC profile of levofloxacin stearic acid co-crystal precipitate shows two endotherms. One of these endotherms might be due to melting of the cocrystals. The nature of the second endotherm has to be investigated. Since the levofloxacin stearic

acid co-crystal precipitate has a solubility values lower than of levofloxacin, it is possible that the precipitate might be the salt, levofloxacin stearate salt.

Formation and Characterization of Co-Crystals of Levofloxacin with Oleic Acid

Experimental Methodology

Formation of Co-crystals of Levofloxacin with Oleic Acid

[00424] 0.78 g (2.76 mM) of oleic acid was dissolved in 10 ml chloroform. To this, 1.025 (2.76 mM) of levofloxacin dissolved in 10 ml of chloroform was added. The resulting solution was mixed thoroughly and evaporated at 40°C.

Characterization

[00425] Thermal analysis. Thermal analysis of (a) oleic acid (b) levofloxacin oleate co-crystal precipitate (c) physical mixture of oleic acid and levofloxacin (50:50) (d) physical mixture of oleic acid and levofloxacin (10:90) and (e) physical mixture of oleic acid and levofloxacin (90:10) was performed using a Differential Scanning Calorimeter (TA Instrument DSC Q1000). 2-5 mg of each sample was weighed into pan, sealed and heated at 1°C/min or 10°C/min from 25°C to 250°C under nitrogen.

[00426] Fourier transform Infrared (FT-IR) spectroscopy. FT-IR spectroscopy of (a) oleic acid (b) levofloxacin oleic acid co-crystallized precipitate (d) physical mixture of oleic acid and levofloxacin was carried out using a FTIR spectrometer (Model IRPrestige-21, Shimadzu).

[00427] Kinetic solubility determination. Levofloxacin oleate co-crystallized precipitate (50 mg) was suspended in 2 ml of water. The suspension was adjusted to pH 7 with HCL and shaken. The solubility of these co-crystals was determined at various time intervals. This study was performed at room temperature and at 40°C. The kinetic solubility of an equimolar physical mixture of levofloxacin and oleic acid was also performed and compared to that of the co-crystals at 40°C.

Results

[00428] Thermal analysis. DSC scans of (a) oleic acid (b) levofloxacin oleate co-crystal precipitate (c) physical mixture of oleic acid and levofloxacin (50:50) (d) physical mixture of oleic acid and levofloxacin (10:90) and (e) physical mixture of oleic acid and levofloxacin (90:10) are shown as Figure 28. Oleic acid thermogram shows endotherms at -6.15°C and 13.05°C . The endotherm at -6.15°C corresponds to gamma-alpha oleic acid phase transition (Crowley K.J, 1999). Levofloxacin oleate co-crystallized precipitate shows endotherm at 127.69°C , while an equimolar physical mixture of levofloxacin with oleic acid shows endotherms at 123.69°C , 179.35°C and 224°C . The equimolar physical mixture is showing an endotherm which is close to melting point of the co-crystals, suggesting a possible reaction between oleic acid and levofloxacin in the solid state. To investigate this phenomenon, DSC on physical mixtures of levofloxacin and oleic acid (90:10) and (10: 90) were performed. Physical mixture of levofloxacin and oleic acid (10:90) shows major endotherm at 10.33°C (possible melting of oleic acid) and at 281°C . It does not show endotherm near the melting point of co-crystals. A levofloxacin oleic acid (90: 10) physical mixture shows no melting endotherm at 10°C for oleic acid. It exhibits endotherms at 79.77°C and at 128°C (close to the melting point of the co-crystals), suggesting a possible reaction of levofloxacin and oleic in presence of high amounts of levofloxacin.

[00429] FTIR. The FTIR spectra of oleic acid shows a C=O stretch intense peak at 1710 cm^{-1} and O-H in-plane and out-of-plane bands at 1462 and 937 cm^{-1} , respectively.

[00430] The high intensity absorption bands at 1710 cm^{-1} in the FTIR spectra of oleic acid, which is due to the stretching of C=O group is slightly reduced in the co crystal. The O-H in plane and out of plane bands at 1462 and 937 cm^{-1} in the oleic acid are absent in the co-crystal. Also the FTIR spectrum of the physical mixture is different from that of the salt (Figure 29).

Kinetic Solubility Determinations

[00431] Figure 30 displays the data for the kinetic solubility experiments performed with the co-crystallized precipitate at room temperature and at 40°C . The

solubility of the co-crystals at room temperature is about 0.9 mg/ml, which remains constant throughout the period of the study. At 40° C, the solubility increased from 1.17 mg/ml at 15 minutes to 1.86 mg/ml at 4 hrs which remained almost constant till 24 hrs. The solubility profile of the equimolar physical mixture at 40°C looks different from that of the co-crystals. The physical mixture has a higher solubility (9.16 mg/ml at 24 hrs) as compared to the co-crystals (1.89 mg/ml at 24 hrs).

Interpretation

[00432] The DSC data of the equimolar physical mixture exhibits an endotherm near the melting endotherm of the co-crystallized precipitate. However the FTIR and solubility data of the co-crystals is different than that of physical mixture, with the co-crystals having a low saturation solubility. The saturation solubility of the co-crystals is 0.9 mg/ml temperature as opposed to 25 mg/ml for levofloxacin base.

[00433] However the levofloxacin oleate salt is waxy in nature which might be difficult to grind/micronize and thereby formulate. It is reported that the tacky and deformable properties of a wax- like drug fatty acid salt, propranolol oleate made particle size reduction difficult (Crowley. J., et al, International journal of Pharmaceutics, 2000, 211 (1-2): 9-17.

Dissolution Rate Studies

Levofloxacin Xinafoate

Experimental Methodology

[00434] 50 mg of the levofloxacin xinafoate salt was suspended in a dissolution bath containing 500 ml of pH 7.4 Tris buffer at 37 °C and rotated by means of paddles at 100 rpm. 5 ml samples were removed at periodic time intervals and replaced with same volume of plain buffer.

Results

[00435] The dissolution profile of levofloxacin xinafoate is shown as Figure 31. It is seen that the dissolution rate of levofloxacin xinafoate in the earlier stages of 2-10 minutes is faster than that seen from 10- 30 minutes. When levofloxacin xinafoate is added to the dissolution media, it gets dispersed as a fine powder, and the dissolution

from these fine particles is faster, approximately 1.24 mg/min (Figure 32). With time, the powder gets agglomerated and moves in a vortex created by the paddle, thereby decreasing the dissolution rate to 0.28 mg/min (Figure 33).

Levofloxacin

Experimental methodology

[00436] 200 mg of levofloxacin was suspended in a dissolution bath containing 500 ml of pH 7.4 Tris buffer at 37°C and rotated by means of paddles at 100 rpm. 5 ml samples were removed at periodic time intervals and replaced with same volume of plain buffer.

Results

[00437] The dissolution profile of levofloxacin is shown as Figure 34. As levofloxacin has a higher solubility than the salts its dissolution rate is very fast. With levofloxacin too, the earlier dissolution was from finely dispersed particles and hence a faster dissolution rate. In the later stages the particles agglomerated and reduced its dissolution rate.

Levofloxacin Pamoate

Experimental Methodology

[00438] 10 mg of the levofloxacin pamoate salt was suspended in a dissolution bath containing 500 ml of pH 7.4 Tris buffer at 37°C and rotated by means of paddles at 100 rpm. 5 ml samples were removed at periodic time intervals (2, 5, 10, 15, 20, 25, 30, 45, 60, 120, 240, 1320 and 1440 minutes) and replaced with same volume of plain buffer. The study was performed in duplicate.

Results

[00439] The dissolution profile of levofloxacin pamoate is shown as Figure 35. It is seen that the dissolution rate of levofloxacin pamoate in the earlier stages of 2-10 minutes is faster than that seen from 10- 60 minutes. When levofloxacin pamoate, is added to the dissolution media, it gets dispersed as a fine powder, and the dissolution from these fine particles is faster, approximately 0.146 mg/min (Figure 36). With time,

the powder gets agglomerated and moves in a vortex created by the paddle, thereby decreasing the dissolution rate to 0.0331 mg/min (Figure 37).

Levofloxacin Stearate

Experimental Methodology

[00440] 25 mg of the levofloxacin stearate salt was suspended in a dissolution bath containing 500 ml of pH 7.4 Tris buffer at 37°C and rotated by means of paddles at 100 rpm. 5 ml samples were removed at periodic time intervals (2,5,10,15,20,25,30,45,60, 120,240,1320 and 1440 minutes) and replaced with same volume of plain buffer.

Results

[00441] The dissolution profile of levofloxacin stearate is shown as Figure 38. It is seen that the dissolution rate of levofloxacin stearate in the earlier stages of 2-10 minutes is 0.499 mg/min (Figure 39) which is faster than that seen from 10- 30 minutes (0.161 mg/min) (Figure 40).

[00442] Dissolution of levofloxacin and salts was carried out with quantities such that the concentration of the dissolved solute in the dissolution bath never reached more than 10% of its saturation solubility. This was done in an attempt to maintain sink conditions.

[00443] Pending the dissolution rate of these and other salt forms and co-precipitates, these AUC shape-enhancement forms of levofloxacin, gemifloxacin and other fluoroquinolone antibiotics these forms may be best suited for nanoparticle suspension (solubilities <100 ug/ml, slow dissolution rates) or micron-size dry powders (solubilities >100 ug/ml, quicker dissolution rate than that best for nanosuspension. The nanoparticle suspensions may be administered by nebulization using jet, ultrasonic, or vibrating mesh technologies, while dry powder formulations may be administered using either active or passive dry powder inhalers.

Example 11 – Levofloxacin Solid Lipid Nanoparticles.

[00444] The goal of this study was to prepare solid lipid nanoparticles of

levofloxacin to obtain AUC shape-enhancing properties through decreased solubility and dissolution. These benefits may enhance the pharmacodynamic properties of levofloxacin following pulmonary administration using nanoparticle suspension or dry powder inhalation formulations. These formulations are being optimized to prolong the release of levofloxacin from decreased solubility salt forms. These properties may also be incorporated into other fluoroquinolone antibiotics including, without limitation gemifloxacin, gatifloxacin, norfloxacin, tosufloxacin, sitafloxacin sarafloxacin, prulifloxacin, and pazufloxacin. Studies are also underway to characterize various lipid nanoparticle forms of gemifloxacin for taste masking, AUC shape-enhancement, nanoparticle suspension and dry powder inhalation administration. Other approaches for solid lipid nanoparticle currently being investigated include spray-dry and in situ micronization techniques.

Preformulation Studies

[00445] Partitioning of each compound (including levofloxacin salts and metal cation complexes) into 1-octanol were determined at different relevant pH values. Partitioning as a function of time may also be evaluated to determine whether dissociation of levofloxacin occurs (from both salts and complexes), and in the case of salts, also to determine if selective partitioning of the fatty acid component occurs over time. Compound(s) with significant partitioning ($\log P > 2.0$) was evaluated for their solubility in various lipid melts. Additionally, partitioning of a lipophilic fluoroquinolone (if available) is also being studied, and its solubility in different lipid melts will be evaluated. A lipid in which drug is sufficiently soluble will be selected for formulation of solid lipid nanoparticles. A prerequisite to obtain a sufficient loading capacity of the drug in solid lipid nanoparticles was a high solubility of drug in the lipid melt.

Formulation of Solid Lipid Nanoparticles

[00446] Formulation of solid lipid nanoparticles typically involves dissolving the drug in a lipid melt, followed by dispersion of the drug-containing melt in a hot aqueous surfactant solution. The coarse dispersion is homogenized using a Microfluidizer[®] to obtain a nanoemulsion. Cooling the nanoemulsion to room temperature will re-solidify the lipid, leading to formation of solid lipid nanoparticles. Optimization of formulation

parameters (type of lipid matrix, surfactant concentration and production parameters) will be performed so as to achieve a prolonged drug delivery.

Characterization of Solid Lipid Nanoparticles

[00447] The nanoparticles are being characterized for size and zeta potential using a Dynamic Light Scattering instrument while Laser diffraction will be used for the detection of large microparticles.

[00448] Upon completion of the synthesis, differential scanning calorimetric studies will be performed to investigate any possible modifications induced in the physical form of the lipid.

[00449] *In vitro* drug release testing will be done using appropriate methodology.”

Example 10 – Levofloxacin Metal Ion Complexes.

[00450] The goal of this study was to prepare levofloxacin of various chelate salt forms to obtain gain taste-masking properties, AUC shape-enhancing properties through changes in solubility, dissolution and/or bioavailability. These benefits may enhance the pharmacodynamic properties of levofloxacin following pulmonary administration using nanoparticle suspension, dry powder inhalation or simple liquid formulations. These formulations may be optimized to create AUC shape-enhancing formulations of levofloxacin from altered solubility, or slow-release or low bioavailability chelates. These properties may also be incorporated into other fluoroquinolone antibiotics including, without limitation gemifloxacin, gatifloxacin, norfloxacin, tosufloxacin, sitafloxacin sarafloxacin, prulifloxacin, and pazufloxacin. Studies are also underway to characterize various and chelate forms of gemifloxacin for taste masking, AUC shape-enhancement, nanoparticle suspension and dry powder inhalation administration.

Preparation of Levofloxacin-Metal Ion Complexes

Preliminary Studies

[00451] A mixture of levofloxacin and a salt of a given cation was solubilized in deionized water and titrated with sodium hydroxide. The titration curve was compared

against one obtained for levofloxacin alone to assess formation of levofloxacin-metal complex as described in Physical Pharmacy (4th Edition) by Alfred Martin (pp 261-263). Salts of various metal cations (e.g. Ca²⁺, Mg²⁺, etc) were then evaluated to identify suitable candidate(s) for subsequent evaluations. Different molar ratios of cations and levofloxacin were also evaluated.

Preparation of Complexes

[00452] Levofloxacin solutions were titrated against aqueous solutions of selected metal salts. Titrations were carried out at a constant pH. Formation of complexes were monitored by different methods including titrimetry, spectrofluorometry, solubility, etc. as applicable. The end point of the complexation reaction depended on the method adopted.

Characterization of Levofloxacin Complexes

[00453] Levofloxacin-metal cation complexes were characterized for stoichiometry, formation constants and dissociation kinetics using appropriate methodology.

Goals

[00454] To formulate and characterize levofloxacin complexes with metal cations (di- and tri-valent).

Assessment of Complexation

[00455] Preliminary investigations suggested that levofloxacin forms soluble complexes with metal cations. As a result, evaluation of the complexation process by precipitation was not possible. Other approaches that were attempted are described below.

Titrimetry

[00456] This approach was based on the assumption that the carboxylic acid moiety of levofloxacin is involved in complex formation with a given metal cation and that complexation results in the release of a proton from levofloxacin. The concentration of released protons would thus be proportional to the extent of complexation (depending

on the binding constant) and the stoichiometry of the complex (Physical Pharmacy: 4th Edition by Alfred Martin; pp-261-263).

Experimental Methodology

[00457] About 0.35 mmoles of levofloxacin (in 16 mL of deionized water) were titrated with 6N NaOH in the presence and absence of salt of a metal cation (equimolar). Levofloxacin solutions were acidified to pH values less than 2.0 with 6N HCl prior to titration with NaOH. Salts of metal cations used include calcium chloride, magnesium chloride, ferrous chloride, zinc chloride, aluminum sulfate and aluminum chloride.

Results

[00458] As shown in Figure 41, titrations performed in the presence of metal cations resulted in a positive shift of the titration curves as compared to the one obtained with levofloxacin alone suggesting that additional NaOH (titrant) is required to obtain a specific pH of the solution in the presence of metal cation. The magnitude of the shift in titration curve at any point would represent moles of proton released due to complexation and hence moles of complexed levofloxacin.

[00459] Extent of complexation (binding and/or stoichiometry) appears to increase in the order $\text{Ca}^+ < \text{Mg}^{2+} < \text{Zn}^{2+} = \text{Fe}^{2+} < \text{Al}^{3+}$, which is in reasonable agreement with existing literature.

[00460] Note: It was noted from the literature that aluminum chloride and aluminum sulfate have acid-like properties and would lower the pH of aqueous solutions. Consequently, the titration curves obtained with AlCl_3 and $\text{Al}_2(\text{SO}_4)_3$ may not provide conclusive information on complexation with levofloxacin.

Dual Titration

[00461] In this approach levofloxacin solution was titrated with a solution of a given metal cation to observe a drop in pH presumably due to release of protons through complexation. This was followed by addition of NaOH to revert back to the initial pH of the levofloxacin solution (prior to addition of solution of cation). This enables determination of the fraction of levofloxacin in the complexed form at a given pH.


Experimental Methodology

[00462] About 1.55-1.72 mmoles of levofloxacin were solubilized in deionized water and the resulting solution was acidified with 6N HCl to the desired initial pH. This acidified levofloxacin solution was titrated with a known volume of concentrated solution of a given metal cation (Ca^{2+} , Mg^{2+} , Fe^{2+} and Zn^{2+}). The change in pH was neutralized (to the initial pH) by the addition of 6N NaOH and volume of NaOH solution added was recorded. Addition of solution of metal cation followed by neutralization with NaOH was continued until further addition of solution of metal cation failed to result in pH change of the levofloxacin solution, which would indicate endpoint of complexation. The cumulative amounts of metal cation added were plotted against cumulative amounts of NaOH required to neutralize the change in pH (Figures 42-45).

Results

[00463] From Figures 42-45, the plateau regions were extrapolated to obtain total amount of NaOH required to neutralize the change in pH due to complexation. These values also represent the amounts of levofloxacin in the complexed form (assuming that complexation of levofloxacin results in an equimolar release of protons). Amounts of levofloxacin in the complexed form with Ca^{2+} , Mg^{2+} , Fe^{2+} and Zn^{2+} are 0.8, 1.0, 1.3 and 1.1 mmoles, respectively. These represent 46.5, 64.5, 77.8 and 64.5% complexation for Ca^{2+} , Mg^{2+} , Fe^{2+} and Zn^{2+} , respectively. It should be noted that % complexation would depend on the total concentrations of levofloxacin.

[00464] The binding constants as well as the stoichiometry of complexation for the levofloxacin complexes with the metal cations were determined as follows:



Where M, A and MA_n represent the metal cation, levofloxacin and the complex, respectively. K_b would be the equilibrium binding constant. The above reaction assumes that 'n' moles of levofloxacin react with one mole of metal to yield one mole of complex.

$$K_b = [\text{MA}_n] / \{[\text{M}][\text{A}]^n\} \text{ (units } \text{M}^{-n}\text{)} \text{ ----- Eq.1}$$

$[\text{MA}_n]$ is the concentration of complex formed

$[\text{M}]$ and $[\text{A}]$ are the concentrations of the unbound metal and unbound levofloxacin, respectively.

Rearranging Eq.1,

$$[MA_n]/[A]^n = K_b * [M] \text{ ----- Eq.2}$$

$$[A] = [A]_{\text{Total}} - [A]_{\text{bound}} = [A]_{\text{Total}} - [\text{NaOH}]_{\text{used}}$$

$$[M] = [M]_{\text{Total}} - [M]_{\text{bound}} = [M]_{\text{Total}} - [\text{NaOH}]_{\text{used}}/n$$

$$[MA_n] = [A]_{\text{bound}}/n = [\text{NaOH}]_{\text{used}}/n$$

[00465] Note: $[\text{NaOH}]_{\text{used}}$ is the concentration of sodium hydroxide used at any given point to neutralize the change in pH caused by the addition of metal cation (presumably due to complexation).

[00466] Eq.2 can be modified to obtain,

$$[A]_{\text{bound}}/[A]^n = nK_b * [M] \text{ ----- Eq.3}$$

It is inferred from Eq.3 that a plot of $[M]$ versus $[A]_{\text{bound}}/[A]^n$ would result in a straight line with a slope of nK_b when,

$$n = 1, \text{ for a 1:1 complex}$$

$$n = 2, \text{ for a 2:1 complex}$$

$$n = 3, \text{ for a 3:1 complex etc.}$$

[00467] Shown below in Figures 46-49 are these plots for Ca^{2+} , Mg^{2+} , Fe^{2+} and Zn^{2+} , respectively.

[00468] As shown in Figures 46-49, for each of the cations evaluated a plot of $[A]_{\text{bound}}/[A]^n$ versus $nK_b * [M]$ was linear when $n=2$ (for Ca^{2+} $n=2$ resulted in a better fit than $n=1$). These results suggest that levofloxacin complexes with Ca^{2+} , Mg^{2+} , Fe^{2+} and Zn^{2+} are formed with a stoichiometry of 2 moles of drug per mole of cation (2:1).

[00469] Using $n=2$, the binding constants for the above complexes can be determined from the slopes of the respective linear plots.

[00470] The binding constants for 2:1 complexes represented as $\log(K_b)$ are as follows: $\text{Ca}^{2+}=2.75$, $\text{Mg}^{2+}=3.69$, $\text{Zn}^{2+}=4.44$, $\text{Fe}^{2+}=4.54$.

Solubility

[00471] This method allows for a relatively simple way of determining the stoichiometry of complexation. The approach involved evaluation of solubility of the drug (levofloxacin) in the presence of increasing concentrations of complexation agent (a

given metal cation). The total solubility of the drug (complexed + uncomplexed) was expected to increase linearly owing to complexation and to reach a plateau corresponding to the saturation solubility of both the drug and the complex. Determination of the stoichiometry from such a solubility curve was explained in detail elsewhere (Physical Pharmacy: 4th Edition by Alfred Martin; pp 265).

Experimental Methodology

[00472] Excess quantities of levofloxacin (amounts were recorded) were agitated, in the presence of increasing concentrations of MgCl₂, with 25 mM MES buffer (pH 5.99) using a vortex mixer. The samples were then filtered and the filtrate was diluted appropriately and analyzed spectrophotometrically to determine levofloxacin concentrations (Figure 50).

Results

[00473] As shown in Figure 50, the solubility of levofloxacin did increase with increasing MgCl₂ concentrations. However, beyond the plateau solubility (~ 650mM levofloxacin), further increase in solubility was observed, which is not consistent with the expected profile. This was attributed to the effect of ionic strength on levofloxacin solubility. It is important to note that the final pH of all the solutions were constant, albeit greater than 5.99 (final pH ~7.0).

[00474] Subsequently, the experiment was repeated at a constant ionic strength of ~1.0M (adjusted with NaCl) and with 0.5M MES buffer (pH 5.99) to enhance the buffer capacity of the solution (Figure 51).

Spectrofluorometry

[00475] This approach was adopted to evaluate levofloxacin complexation based on existing literature evidence that the complexation process is associated with a change in the fluoroquinolone fluorescence properties. By monitoring the change in fluorescence emission of levofloxacin in the presence of different concentrations of a given metal cation it was possible to determine the binding constant of complexation as well as the stoichiometry.

Experimental Methodology

[00476] The fluorescence emission of levofloxacin was evaluated at excitation and emission wavelengths of 298 nm and 498 nm, respectively. Studies were conducted at two different pH values i.e. 5.0 (acetate) and 9.0 (histidine). A series of solutions containing a constant levofloxacin concentration but increasing concentrations of a given cation were analyzed for fluorescence emission due to levofloxacin. Metal salts studied included CaCl₂, MgCl₂, FeCl₂, ZnCl₂ and Al₂(SO₄)₃.

Results

[00477] As shown in Table 34, significant data were obtained only for Fe²⁺ and Zn²⁺. For the remaining cations, the relative concentrations of levofloxacin and the cation need to be further optimized to observe a specific trend in change in levofloxacin fluorescence.

[00478] The influence of increasing concentrations of Fe²⁺ and Zn²⁺ on levofloxacin fluorescence emission are shown in Figures 52 and 53, respective

[00479] As described above, both Fe²⁺ and Zn²⁺ appear to form 2:1 complexes with levofloxacin; however, their influence on levofloxacin fluorescence are dissimilar (Figures 52 and 53). The exact reason for this is unclear at this point.

Table 34. Fluorescence Characteristics of Levofloxacin in the Presence of Cations.

Cation	Fluorescence of levofloxacin		Results	Comments
	pH 5.0	pH 9.0		
Ca ²⁺	Change not significant	Change not significant	N/A	-
Mg ²⁺	Change not significant	Change not significant	N/A	-
Fe ²⁺	Decrease in emission with increasing Fe ²⁺	N/A	Figure 3.12 (pH 5.0)	FeCl ₂ insoluble at pH 9.0
Zn ²⁺	Change not significant	Increase in emission with increasing Zn ²⁺	Figure 3.13 (pH 9.0)	-
Al ³⁺	Change not significant	N/A	N/A	Al ₂ (SO ₄) ₃ insoluble at pH 9.0

Samples of Levofloxacin Complexes

[00480] Seven samples of levofloxacin complexes were evaluated *in vivo* for efficacy and pharmacokinetics. Details of the samples tested are shown in Table 35 below.

Table 35. Molar Ratios of Levofloxacin Complexes.

Sample identifier	Cation	Molar ratio used	Total Levofloxacin (mg/mL)	Final pH of the solution
NB-049-001-06-066A	Mg ²⁺	1:1	40.2	6.24
NB-049-001-06-066B	Fe ²⁺	1:1	40.1	6.30
NB-049-001-06-066C	Mg ²⁺	1:1	202	5.98
NB-049-001-06-081A	Ca ²⁺	1:1	40.1	6.53
NB-049-001-06-081B	Ca ²⁺	1:1	201	6.04
NB-049-001-06-081C	Zn ²⁺	1:1	40	6.33
NB-049-001-06-081D	Zn ²⁺	1:1	200	5.69

Conclusions and Next Steps

[00481] Results obtained from our dual titration studies suggest that levofloxacin forms 2:1 complexes with all the divalent metal cations. The binding constants (log K_b) for complexation with Ca²⁺, Mg²⁺, Fe²⁺ and Zn²⁺ are 2.75, 3.69, 4.44 and 4.54, respectively.

Example 11 – Levofloxacin and Gemifloxacin Formulations with Organic Acids.

Experimental Methodology

[00482] Levofloxacin solution was prepared by dissolving either 50 or 100 mg levofloxacin base in 15-20 ml water. The initial pH of levofloxacin solution in water was about 7.3. The pH of the solution was adjusted with about 10% solution of acid prepared in water. The following acids were used to adjust the pH of the levofloxacin solution: acetic acid, ascorbic acid, citric, lactic, tartaric and propionic acid. After making up the volume of the solution to approximately 90% of the final volume, the osmolality of the solution was measured and adjusted to 300 mOsm/ kg with about 20% solution of sodium chloride prepared in water. After pH and osmolality adjustment, the volume of the

solution was made up to about 25 ml with water and its surface tension measured. The pH and osmolality were measured after making up the volume and are reported in Table 36. (The exact quantities of levofloxacin weighed, acid required to adjust pH, sodium chloride to adjust osmolality and final volume of solutions are listed in Table 36). The content of levofloxacin in the solutions was determined by HPLC.

Results

[00483] The details about the levofloxacin formulations with organic acids are shown in Table 36. The results of HPLC are shown in Table 37.

[00484] When tartaric acid was used to adjust the pH of the 100 mg/ml levofloxacin solution, a precipitate was formed.

[00485] Note: Solutions with acetic acid, citric acid and ascorbic acid were remade for HPLC analysis and hence the theoretical concentration for these solutions in Table 36 and Table 37 are different.

Gemifloxacin Formulations with Organic Bases

Experimental Methodology and Results

Gemifloxacin Formulation with Sodium Ascorbate

[00486] 50.30 mg of Gemifloxacin mesylate (equivalent to 40.37 mg Gemifloxacin) was added to 1.5 ml water. The resulting solution was cloudy. It was filtered through a 0.45 micron filter. 1.3 ml of solution was obtained after filtering having a pH of 4.28. The pH of this solution was adjusted to 5.48 with 400 uL of a 10% solution of sodium ascorbate prepared in water (Quantity of base required to adjust pH = 0.04 g). The osmolality of this solution was 308 mOsm/kg, hence sodium chloride was not used to adjust osmolality. The final volume of the solution was 1.7 ml. *Theoretical concentration of gemifloxacin in this formulation would be 20.59 mg/ml.

Table 36. Formulations of Levofloxacin With Organic Acids.

Wt of Levo used (g)	9.94% acetic acid used (ml)	Acetic acid used (g)	19.7% NaCl used (ml)	NaCl used (g)	Measured Final Vol of solution (ml)	Levo conc (mg/ml)	Final osmolality (mOsm/kg)	Final pH	Surface tension (mN/m)
1.253	1.05	0.104	0.681	0.134	25.105	49.9	312	6.48	63.2
2.501	2.05	0.204	0.326	0.064	25.935	96.4	300	6.53	62.5

Wt of Levo used (g)	9.99 % ascorbic acid used (ml)	ascorbic acid used (g)	19.7% NaCl used (ml)	NaCl used (g)	Measured Final Vol of solution (ml)	Levo conc (mg/ml)	Final osmolality (mOsm/kg)	Final pH	Surface tension (mN/m)
1.253	3.400	0.339	0.550	0.108	25.135	49.8	297	6.40	64.4
2.505	7.400	0.739	0.300	0.059	25.135	99.7	298	6.47	62.5

Wt of Levo used (g)	10.05% citric acid used (ml)	citric acid used (g)	21.54 % NaCl used (ml)	NaCl used (g)	Measured Final Vol of solution (ml)	Levo conc (mg/ml)	Final osmolality (mOsm/kg)	Final pH	Surface tension (mN/m)
1.251	1.25	0.126	1.005	0.216	25.12	49.8	299	6.54	61.5
2.498	2.6	0.261	0.918	0.198	25.82	96.7	301	6.53	61.4

Wt of Levo used (g)	10% lactic used (ml)	Lactic acid used (g)	21.54 % NaCl used (ml)	NaCl used (g)	Measured Final Vol of solution (ml)	Levo conc (mg/ml)	Final osmolality (mOsm/kg)	Final pH	Surface tension (mN/m)
1.258	2.1	0.21	0.745	0.160	25.135	50.1	297	6.54	59.4
2.497	4.2	0.42	0.392	0.084	25.605	97.5	301	6.63	57.5

Wt of Levo used (g)	10 % tartaric acid used (ml)	Tartaric acid used (g)	21.54% NaCl used (ml)	NaCl used (g)	Measured Final Vol of solution (ml)	Levo conc (mg/ml)	Final osmolality (mOsm)	Final pH	Surface tension (mN/m)
1.252	1.55	0.155	0.948	0.204	25.180	49.7	298	6.51	61.5

Wt of Levo used (g)	9.79% propionic acid used (ml)	propionic acid used (g)	21.53% NaCl used (ml)	NaCl used (g)	Measured Final Vol of solution (ml)	Levo conc (mg/ml)	Final osmolality (mOsm)	Final pH	Surface tension (mN/m)
1.25281	1.310	0.128	0.737	0.159	25.045	50.02	298	6.50	58.1
2.51342	2.610	0.256	0.310	0.067	25.030	100.42	297	6.57	52.0

[00487] * Theoretical concentration = Theoretical quantity of gemifloxacin in filtered solution (in this case 35 mg Gemifloxacin in filtered 1.3 mL)/ Final volume of solution (in this case 1.7 mL).

Table 37. Theoretical and Measured Concentrations of Levofloxacin in the Formulations.

Acid	Theoretical Conc. (mg/mL)	Measured Concentration (mg/ml) by HPLC
acetic acid	50.05	51.45
acetic acid	99.9	102.32
citric acid	49.91	50.31
citric acid	99.86	102.99
L-ascorbic acid	49.95	50.01
L-ascorbic acid	100	102.49
lactic acid	50.05	50.07
lactic acid	97.54	95.27
tartaric acid	49.74	51.07

[00488] Note: Solutions with acetic acid, citric acid and ascorbic acid were remade for HPLC analysis and hence the theoretical concentration for these solutions in Table 36 and Table 37 are different.

Gemifloxacin Formulation with Sodium Lactate

[00489] 50.05 mg of Gemifloxacin mesylate (equivalent to 40.17 mg Gemifloxacin) was added to 1.8 ml water. The resulting solution was cloudy. It was filtered through a 0.45 micron filter. 1.52 ml of solution was obtained after filtering having a pH of 4.21. The pH of this solution was adjusted to 5.42 with 180 uL of a 20% solution of sodium lactate prepared in water (Quantity of base required to adjust pH = 0.036 g). The osmolality of this solution was 478 mOsm/kg. The final volume of the solution was 1.7 ml. Theoretical concentration of gemifloxacin in this formulation would be 19.95 mg/ml.

Gemifloxacin Formulation with Sodium Acetate

[00490] 50.47 mg of Gemifloxacin mesylate (equivalent to 40.50 mg Gemifloxacin) was added to 2.0 ml water. The resulting solution was cloudy. It was filtered through a 0.45 micron filter. 1.77 ml of solution was obtained after filtering having a pH of 4.40. The pH of this solution was adjusted to 5.40 with 50 uL of a 10% solution of sodium acetate prepared in water (Quantity of base required to adjust pH =

0.005 g). The osmolality of this solution was 192 mOsm/kg. The osmolality of this solution was adjusted to 295 mOsm/ kg using 28 uL of 20 % solution of sodium chloride prepared in water.

Gemifloxacin Formulation with Sodium Propionate

[00491] 50.00 mg of Gemifloxacin mesylate (equivalent to 40.13 mg Gemifloxacin) was added to 1.9 ml water. The resulting solution was cloudy. It was filtered through a 0.45 micron filter. 1.39 ml of solution was obtained after filtering having a pH of 4.32. The pH of this solution was adjusted to 5.50 with 30 uL of a 20% solution of sodium propionate prepared in water (Quantity of base required to adjust pH = 0.006 g). The osmolality of this solution was 183 mOsm/kg. The osmolality of this solution was adjusted to 296 mOsm/ kg using 25 uL of 22 % solution of sodium chloride prepared in water. This solution was remade with osmolality adjustment to 237mOsm/Kg.

Gemifloxacin Formulation with Sodium Citrate

[00492] 49.92 mg of Gemifloxacin mesylate (equivalent to 40.06 mg Gemifloxacin) was added to 1.9 ml water. The resulting solution was cloudy. It was filtered through a 0.45 micron filter. 1.63 ml of solution was obtained after filtering having a pH of 4.20. The pH of this solution was adjusted to 5.39 with 15uL of a 20% solution of sodium citrate prepared in water (Quantity of base required to adjust pH = 0.003 g).

Example 12 – Microspheres of Levofloxacin.

[00493] The goal of this study was to prepare various microsphere forms of levofloxacin that may gain taste-masking, and AUC shape-enhancing properties through decreased solubility and/or dissolution. These benefits may enhance the pharmacodynamic properties of levofloxacin following pulmonary administration using either nanoparticle suspension or dry powder inhalation. These formulations are being optimized to prolong the release of levofloxacin from decreased solubility or dissolution forms. These properties may also be incorporated into other fluoroquinolone antibiotics including, without limitation gemifloxacin, gatifloxacin, norfloxacin, tosufloxacin,

sitafloxacin sarafloxacin, prulifloxacin, and pazufloxacin. Studies are underway to characterize microspheres of gemifloxacin for taste masking, AUC shape-enhancement, nanoparticle suspension and dry powder inhalation administration. Other approaches for dry powder administration currently being investigated include spray-dry and in situ micronization techniques.

Preformulation for Levofloxacin

Preformulation Studies

[00494] Preformulation studies were performed to determine the solubility of levofloxacin and polymers in various solvents which are expected to be used during processing.

Preparation of Microspheres

[00495] A spray drying technique is being used to formulate polymer microparticles loaded with levofloxacin. Formulation of microspheres will typically involve dissolving the drug and polymer in a suitable solvent. The solution is being spray dried using a spray dryer to evaporate the solvent, thereby entrapping the drug in a polymer matrix. Optimization of formulation parameters (drug: polymer ratio, polymer solution concentration and production parameters) is being performed to achieve a desired micro particle size, optimum drug loading and *in vitro* drug release.

Characterization of Microspheres

[00496] The microparticles will be characterized for their morphology using SEM while microscopy or a suitable technique (Laser diffraction) will be used for estimating their size.

[00497] Drug loading will be determined by extracting the drug from the microspheres in a suitable solvent and analyzing the extract by UV/ HPLC.

[00498] Drug release from the microspheres will be carried out using a USP dissolution apparatus.

Example 13 – Inhalation Toxicology in Rats.

[00499] In a 4 day non-GLP ascending dose study of aerosolized levofloxacin in

male and female Sprague-Dawley rats, a 25 mg/ml solution of levofloxacin was administered for one hour on day one and a 50 mg/ml solution of levofloxacin was administered for two hours per day on days 2 thru 4. No clinical signs of toxicity were observed during the treatment period. Necropsy 24 hours after administration of the last dose did not show any findings.

[00500] In a GLP study of aerosolized levofloxacin in male and female Sprague-Dawley rats, aerosolized levofloxacin was administered daily with an average dose of 6.92 mg/kg/day to the males and 10.04 mg/kg/day for the females over 4 days using a nose-only aerosol delivery device. Total exposures were 29 and 42 mg/kg for males and females, respectively over the study period. Each dose was delivered over 2 hours daily. The dose for this study was chosen based on the maximum solubility of levofloxacin that could be administered in the device over 2 hours. No clinical signs of toxicity were observed, and all animals survived during the 4 day treatment period. Necropsy of animals after administration of the last dose did not show any findings.

[00501] In a 28-day GLP study in Sprague-Dawley rats, animals were randomized to 3 dose levels of aerosolized levofloxacin or saline. Additional recovery groups using the vehicle control and the highest dose were also treated and observed for a 14 day recovery period following the last dose. Average aerosolized levofloxacin doses were 1.49, 3.63, and 7.29 mg/kg/day for male rats, and 2.20, 5.35, and 11.01 mg/kg/day in female rats. The total exposures over the 28-day treatment period ranged between 41.7 and 204.1 mg/kg for males and 61.6 and 308.3 mg/kg for females. Each dose was delivered over 2 hours daily. No dose related clinical signs of toxicity were observed, and all animals survived during the 28 day treatment period. Necropsy of animals after administration of the last dose showed a dose related squamous cell hyperplasia of the larynx which declined in severity during a 14 day recovery period.

WHAT IS CLAIMED IS:

1. A pharmaceutical composition, comprising a solution of levofloxacin or ofloxacin and a divalent or trivalent cation.
2. The composition of claim 1, wherein the divalent or trivalent cation is magnesium.
3. The composition of claim 1, wherein the divalent or trivalent cation is selected from one or more of calcium, aluminum, zinc, and iron.
4. The composition of claim 1, 2 or 3, wherein the solution has a concentration of levofloxacin or ofloxacin greater than about 10 mg/ml.
5. The composition of claim 1, 2 or 3, wherein the solution has a concentration of levofloxacin or ofloxacin greater than about 25 mg/ml.
6. The composition of claim 1, 2 or 3, wherein the solution has a concentration of levofloxacin or ofloxacin greater than about 35 mg/ml.
7. The composition of claim 1, 2 or 3, wherein the solution has a concentration of levofloxacin or ofloxacin greater than about 40 mg/ml.
8. The composition of claim 1, 2 or 3, wherein the solution has a concentration of levofloxacin or ofloxacin greater than about 50 mg/ml.
9. The composition of claim 1, 2 or 3, wherein the solution has a concentration of levofloxacin or ofloxacin of 100 mg/ml.
10. The composition of any one of claims 1 to 9, wherein the solution has an osmolality from about 200 mOsmol/kg to about 1250 mOsmol/kg.

11. The composition of any one of claims 1 to 9, wherein the solution an osmolality from about 250 mOsmol/kg to about 1050 mOsmol/kg.
12. The composition of any one of claims 1 to 9, wherein the solution an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg.
13. The composition of any one of claims 1 to 12, wherein the solution has a pH from about 4.5 to about 7.5.
14. The composition of any one of claims 1 to 12, wherein the solution has a pH from about 5 to about 6.5.
15. The composition of any one of claims 1 to 12, wherein the solution has a pH from about 5.5 to about 6.5.
16. The composition of any one of claims 1 to 15, comprising a sweetener.
17. The composition of claim 16, wherein the sweetener is aspartame or sucrose.
18. The composition of claim 16, wherein the sweetener is a mono- or di-saccharide.
19. The composition of claim 16, wherein the sweetener is lactose, sucrose, dextrose, or glucose.
20. The composition of any one of claims 1 to 19, comprising one or more of dornase alpha, a hypertonic formulation, mannitol, and sodium chloride.
21. The composition of claim 1, wherein the solution comprises levofloxacin and magnesium, has a concentration greater than about 50 mg/ml, has an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg, and has a pH from about 5.5 to about 6.5.
22. The composition of any one of claims 1 to 20, comprising another antimicrobial.
23. The composition of claim 22, wherein the other antimicrobial is an aminoglycoside.

24. The composition of claim 23, wherein the aminoglycoside is tobramycin.
25. The composition of claim 22, wherein the other antimicrobial is a polymyxin.
26. The composition of claim 25, wherein the polymyxin is colistin.
27. The composition of claim 22, wherein the other antimicrobial is a monobactam.
28. The composition of claim 27, wherein the monobactam is aztreonam.
29. The composition of claim 22, wherein the other antimicrobial is a macrolide or ketolide.
30. The composition of claim 22, wherein the other antimicrobial is a glycopeptide.
31. The composition of claim 30, wherein the glycopeptide is vancomycin.
32. The composition of claim 22, wherein the other antimicrobial is a fluoroquinolone.
33. The composition of claim 32, wherein the fluoroquinolone is selected from the group consisting of lomefloxacin, pefloxacin, ciprofloxacin, gatifloxacin, gemifloxacin, moxifloxacin, tosufloxacin, pazufloxacin, rufloxacin, fleroxacin, balofloxacin, sparfloxacin, trovafloxacin, enoxacin, norfloxacin, clinafloxacin, grepafloxacin, sitafloxacin, temafloxacin, marbofloxacin, orbifloxacin, sarafloxacin, danofloxacin, difloxacin, enrofloxacin, garenoxacin, prulifloxacin, olamufloxacin, DX-619, TG-873870 and DW-276.
34. A sterile, single use container, comprising the solution of any one of claims 1 to 33.
35. The container of claim 34, comprising from about 1 ml to about 5 ml of the solution.
36. A kit comprising:
the container of claim 34 or 35; and

a nebulizer adapted to aerosolize the solution for delivery to the lower respiratory tract through oral inhalation.

37. A kit comprising: a sterile, single use container, that comprises the solution of any one of claims 1 to 21; a second container that comprises another antimicrobial; and a nebulizer adapted to aerosolize the solution for delivery to the lower respiratory tract through oral inhalation.

38. The kit of claim 37, wherein the other antimicrobial is an aminoglycoside.

39. The kit of claim 38, wherein the aminoglycoside is tobramycin.

40. The kit of claim 37, wherein the other antimicrobial is a polymyxin.

41. The kit of claim 40, wherein the polymyxin is colistin.

42. The kit of claim 40, wherein the other antimicrobial is a monobactam.

43. The kit of claim 45, wherein the monobactam is aztreonam.

44. The kit of claim 40, wherein the other antimicrobial is a macrolide or ketolide.

45. The kit of claim 40, wherein the other antimicrobial is a glycopeptides.

46. The kit of claim 48, wherein the glycopeptide is vancomycin.

47. The kit of claim 40, wherein the other antimicrobial is a fluoroquinolone.

48. The kit of claim 50, wherein the fluoroquinolone is selected from the group consisting of lomefloxacin, pefloxacin, ciprofloxacin, gatifloxacin, gemifloxacin, moxifloxacin, tosufloxacin, pazufloxacin, rufloxacin, fleroxacin, balofloxacin, sparfloxacin, trovafloxacin, enoxacin, norfloxacin, clinafloxacin, grepafloxacin, sitafloxacin, temafloxacin, marbofloxacin, orbifloxacin, sarafloxacin, danofloxacin, difloxacin, enrofloxacin, garenoxacin, prulifloxacin, olamufloxacin, DX-619, TG-873870 and DW-276.

49. The kit of any one of claims 37 to 48, wherein the second container further comprises one or more of dornase alpha, a hypertonic formulation, mannitol, and sodium chloride.
50. The kit of any one of claims 36 to 49, wherein the nebulizer operates by ultrasonic atomization.
51. The kit of any one of claims 36 to 49, wherein the nebulizer operates by hydraulic atomization.
52. The kit of any one of claims 36 to 49, wherein the nebulizer operates by a vibrating mesh.
53. Use of levofloxacin or ofloxacin in combination with a divalent or trivalent cation in the preparation of an aerosol for treating a lung infection, a chronic obstructive pulmonary disease, cystic fibrosis, or sinusitis.
54. Use of levofloxacin or ofloxacin in combination with a divalent or trivalent cation for treating a lung infection, a chronic obstructive pulmonary disease, cystic fibrosis, or sinusitis.
55. The use of claim 53 or 54, wherein the lung infection is caused by one or more of the following bacteria: *Pseudomonas aeruginosa*, *Pseudomonas fluorescens*, *Pseudomonas acidovorans*, *Pseudomonas alcaligenes*, *Pseudomonas putida*, *Stenotrophomonas maltophilia*, *Aeromonas hydrophilia*, *Escherichia coli*, *Citrobacter freundii*, *Salmonella typhimurium*, *Salmonella typhi*, *Salmonella paratyphi*, *Salmonella enteritidis*, *Shigella dysenteriae*, *Shigella flexneri*, *Shigella sonnei*, *Enterobacter cloacae*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Klebsiella oxytoca*, *Serratia marcescens*, *Morganella morganii*, *Proteus mirabilis*, *Proteus vulgaris*, *Providencia alcalifaciens*, *Providencia rettgeri*, *Providencia stuartii*, *Acinetobacter calcoaceticus*, *Acinetobacter haemolyticus*, *Yersinia enterocolitica*, *Yersinia pestis*, *Yersinia pseudotuberculosis*, *Yersinia intermedia*, *Bordetella pertussis*, *Bordetella parapertussis*, *Bordetella bronchiseptica*, *Haemophilus influenzae*, *Haemophilus*

parainfluenzae, Haemophilus haemolyticus, Haemophilus parahaemolyticus, Haemophilus ducreyi, Pasteurella multocida, Pasteurella haemolytica, Helicobacter pylori, Campylobacter fetus, Campylobacter jejuni, Campylobacter coli, Borrelia burgdorferi, Vibrio cholera, Vibrio parahaemolyticus, Legionella pneumophila, Listeria monocytogenes, Neisseria gonorrhoeae, Neisseria meningitidis, Burkholderia cepacia, Francisella tularensis, Kingella, and Moraxella.

56. The use of claim 53 or 54, wherein the lung infection is pneumonia.

57. The use of claim 53 or 54, wherein the lung infection is caused by a gram-negative anaerobic bacteria.

58. The use of claim 53 or 54, wherein the lung infection is caused by one or more of the bacteria selected from the group consisting of *Bacteroides fragilis, Bacteroides distasonis, Bacteroides 3452A homology group, Bacteroides vulgatus, Bacteroides ovalus, Bacteroides thetaiotaomicron, Bacteroides uniformis, Bacteroides eggerthii, and Bacteroides splanchnicus.*

59. The use of claim 53 or 54, wherein the lung infection is caused by a gram-positive bacteria.

60. The use of claim 53 or 54, wherein the lung infection is caused by one or more of the bacteria selected from the group consisting of *Corynebacterium diphtheriae, Corynebacterium ulcerans, Streptococcus pneumoniae, Streptococcus agalactiae, Streptococcus pyogenes, Streptococcus milleri, Streptococcus (Group G), Streptococcus (Group C/F), Enterococcus faecalis, Enterococcus faecium, Staphylococcus aureus, Staphylococcus epidermidis, Staphylococcus saprophyticus, Staphylococcus intermedius,*

Staphylococcus hyicus subsp. hyicus, *Staphylococcus haemolyticus*, *Staphylococcus hominis*, and *Staphylococcus saccharolyticus*.

61. The use of claim 53 or 54, wherein the lung infection is caused by a gram-positive anaerobic bacteria.

62. The use of claim 53 or 54, wherein the lung infection is caused by one or more bacteria selected from the group consisting of *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetini*, and *Clostridium botulinum*.

63. The use of claim 53 or 54, wherein the lung infection is caused by an acid-fast bacteria.

64. The use of claim 53 or 54, wherein the lung infection is caused by one or more bacteria selected from the group consisting of *Mycobacterium tuberculosis*, *Mycobacterium avium*, *Mycobacterium intracellulare*, and *Mycobacterium leprae*.

65. The use of claim 53 or 54, wherein the lung infection is caused by an atypical bacteria.

66. The use of claim 53 or 54, wherein the lung infection is caused by one or more bacteria selected from the group consisting of *Chlamydia pneumoniae* and *Mycoplasma pneumoniae*.

67. The use of any one of claims 53 to 66, wherein the divalent or trivalent cation is selected from one or more of magnesium, calcium, aluminum, zinc, and iron.

68. The use of any one of claims 53 to 67, wherein the levofloxacin or ofloxacin and divalent or trivalent cation are in a solution.

69. The use of claim 68, wherein the solution comprises levofloxacin and magnesium, has a concentration greater than about 50 mg/ml, has an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg, and has a pH from about 5.5 to about 6.5.

70. The use of any one of claims 53 to 67, wherein the levofloxacin or ofloxacin and divalent or trivalent cation are in a dry powder.

71. A method of taste masking a fluoroquinolone, comprising combining the fluoroquinolone with a divalent or trivalent cation.

72. The method of claim 71, wherein the divalent or trivalent cation are selected from one or more of magnesium, calcium, aluminum, zinc, and iron.

73. The method of claim 71 or 72, wherein the fluoroquinolone is levofloxacin or ofloxacin.

74. The method of claim 71 or 72, wherein the fluoroquinolone is selected from the group consisting of ciprofloxacin, gatifloxacin, gemifloxacin, norfloxacin, moxifloxacin, marbofloxacin, orbifloxacin, sparfloxacin, tosufloxacin, trovafloxacin, pefloxacin, lomefloxacin, clinafloxacin, prulifloxacin, sarafloxacin, sitafloxacin, and pazufloxacin.

75. The method of any one of claims 71 to 74, wherein the fluoroquinolone and divalent or trivalent cation are combined in a single solution.

76. The method of any one of claims 71 to 75, further comprising aerosolizing the combination.

77. Use of a fluoroquinolone in combination with a divalent or trivalent cation in the preparation of an aerosol for treating a lung infection, a chronic obstructive pulmonary disease, cystic fibrosis, or sinusitis.

78. Use of a fluoroquinolone in combination with a divalent or trivalent cation for treating a lung infection, a chronic obstructive pulmonary disease, cystic fibrosis, or sinusitis.

79. The use of claim 77 or 78, wherein the fluoroquinolone is levofloxacin or ofloxacin.

80. The pharmaceutical composition of claim 1, wherein the solution is suitable for inhalation into a lung.

81. The composition of claim 80, wherein the concentration of the divalent or trivalent cation is sufficient to substantially complex the levofloxacin or ofloxacin.

82. The composition of claim 80 or 81, wherein the solution has a concentration of levofloxacin or ofloxacin greater than about 10 mg/ml.

83. The composition of claim 80 or 81, wherein the solution has a concentration of levofloxacin or ofloxacin greater than about 25 mg/ml.

84. The composition of claim 80 or 81, wherein the solution has a concentration of levofloxacin or ofloxacin greater than about 35 mg/ml.

85. The composition of claim 80 or 81, wherein the solution has a concentration of levofloxacin or ofloxacin greater than about 40 mg/ml.

86. The composition of claim 80 or 88, wherein the solution has a concentration of the levofloxacin or the ofloxacin greater than about 50 mg/ml.

87. The composition of claim 80 or 81, wherein the solution has a concentration of levofloxacin from about 100 mM to about 625 mM.

88. The composition of claim 80 or 81, wherein the concentration of the levofloxacin or the ofloxacin is about 100 mg/ml.

89. The composition of any one of claims 80 to 88, wherein the solution has an osmolality from about 200 mOsmol/kg to about 1250 mOsmol/kg.

90. The composition of any one of claims 80 to 88, wherein the solution has an osmolality from about 250 mOsmol/kg to about 1050 mOsmol/kg.

91. The composition of any one of claims 80 to 88, wherein the solution has an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg.

92. The composition of any one of claims 80 to 91, wherein the solution has a pH from about 4.5 to about 7.5.

93. The composition of any one of claims 80 to 91, wherein the solution has a pH from about 5 to about 6.5

94. The composition of any one of claims 80 to 91, wherein the solution has a pH from about 5.5 to about 6.5.

95. The composition of claim 80 or 81, consisting essentially of an aqueous solution of greater than about 50 mg/ml levofloxacin and the divalent or trivalent cation, wherein the solution has a pH from about 5.5 to about 6.5 and an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg.

96. The composition of any one of claims 80 to 95, wherein the divalent or trivalent cation is selected from one or more of calcium, aluminum, zinc, and iron.

97. The composition of any one of claims 80 to 95, wherein the solution consists essentially of levofloxacin and magnesium chloride.

98. The composition of any one of claims 80 to 95, wherein the divalent or trivalent cation is magnesium.

99. The composition of claim 98, wherein the solution comprises magnesium chloride.

100. The composition of claim 97 or 99, wherein concentration of the chloride is from about 25 mM to about 400 mM.

101. The composition of claim 97 or 99, wherein the magnesium chloride has a concentration from about 100 mM to about 250 mM.

102. The composition of claim 97 or 99, wherein the magnesium chloride has a concentration from about 125 mM to about 250 mM.

103. A sterile, single use container, comprising the composition of any one of claims 80 to 102.

104. The container of claim 103, comprising from about 20 mg to about 400 mg of levofloxacin.

105. The container of claim 103, comprising from about 28 mg to about 280 mg of levofloxacin.

106. The container of claim 103, comprising from about 1 ml to about 5 ml of the composition.

107. The container of claim 103, comprising at least about 100 mg of levofloxacin.

108. The container of claim 103, comprising at least about 400 mg of levofloxacin.

109. An aerosol of a solution comprising levofloxacin or ofloxacin and a divalent or trivalent cation.

110. The aerosol of claim 109, wherein the divalent or trivalent cation is magnesium.

111. The aerosol of claim 110, wherein the solution comprises magnesium chloride.

112. The aerosol of claim 111, wherein concentration of the chloride is from about 25 mM to about 400 mM.

113. The aerosol of claim 111, wherein the magnesium chloride has a concentration from about 100 mM to about 250 mM.

114. The aerosol of claim 111, wherein the magnesium chloride has a concentration from about 125 mM to about 250 mM.

115. The aerosol of claim 109, wherein the divalent or trivalent cation is selected from one or more of calcium, aluminum, zinc, and iron.

116. The aerosol of any one of claims 109 to 115, wherein the solution has a concentration of the levofloxacin or ofloxacin greater than about 10 mg/ml.

117. The aerosol of any one of claims 109 to 115, wherein the solution has a concentration of the levofloxacin or ofloxacin greater than about 25 mg/ml.

118. The aerosol of any one of claims 109 to 115, wherein the solution has a concentration of the levofloxacin or ofloxacin greater than about 35 mg/ml.

119. The aerosol of any one of claims 109 to 115, wherein the solution has a concentration of the levofloxacin or ofloxacin greater than about 40 mg/ml.

120. The aerosol of any one of claims 109 to 115, wherein the solution has a concentration of the levofloxacin or ofloxacin greater than about 50 mg/ml.

121. The aerosol of any one of claims 109 to 115, wherein the solution has a concentration of the levofloxacin or ofloxacin of 100 mg/ml.

122. The aerosol of any one of claims 109 to 115, wherein the solution has a concentration of levofloxacin from about 100 mM to about 625 mM.

123. The aerosol of any one of claims 109 to 122, wherein the solution has an osmolality from about 200 mOsmol/kg to about 1250 mOsmol/kg.

124. The aerosol of any one of claims 109 to 122, wherein the solution has an osmolality from about 250 mOsmol/kg to about 1050 mOsmol/kg.

125. The aerosol of any one of claims 109 to 122, wherein the solution has an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg.

126. The aerosol of any one of claims 109 to 125, wherein the solution has a pH from about 4.5 to about 7.5.

127. The aerosol of any one of claims 109 to 125, wherein the solution has a pH from about 5 to about 6.5.

128. The aerosol of any one of claims 109 to 125, wherein the solution has a pH from about 5.5 to about 6.5.

129. The aerosol of claim 109, wherein the solution comprises greater than about 50 mg/ml levofloxacin and magnesium chloride, has a pH from about 5.5 to about 6.5, and an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg.

130. The aerosol of claim 129, wherein the solution comprises the levofloxacin at a concentration from about 100 mM to about 625 mM and the magnesium chloride at a concentration from about 100 mM to about 250 mM.

131. The aerosol of any one of claims 109 to 129, comprising a mass median aerodynamic diameter from about 2 microns to about 5 microns with a geometric standard deviation less than or equal to about 2.5 microns.

132. The aerosol of any one of claims 109 to 129, comprising a mass median aerodynamic diameter from about 2.8 microns to about 4.3 microns with a geometric standard deviation less than or equal to about 2 microns.

133. The aerosol of any one of claims 109 to 129, comprising a mass median aerodynamic diameter from about 2.5 microns to about 4.5 microns with a geometric standard deviation less than or equal to about 1.8 microns.

134. An aerosol dose of a levofloxacin and magnesium solution comprised of:
a concentration of levofloxacin greater than 50 mg/ml and a taste-masking concentration of magnesium wherein the aerosol is comprised of a mist having a mean particle size of between 2 and 5 microns or a particle size geometric standard deviation of less than or equal to about 2 microns.

135. The aerosol dose of claim 134, wherein the magnesium concentration is from about 100 mM to about 250 mM.

136. The aerosol dose of claim 134 or 135, wherein the aerosol has a pH of from about 5.5 to about 7.5.

137. The aerosol dose of claim 134, 135 or 136, wherein a quantity of the mist contains at least 1 mg of levofloxacin.

138. The aerosol dose of any one of claims 134 to 137, wherein the levofloxacin and magnesium solution has an osmolarity from about 350 to about 750 mOsmol/kg.

139. The aerosol dose of any one of claims 134 to 138, having a permeant ion concentration of about 400 mM.

140. A kit comprising:

the container of any one of claims 103 to 108, and
a nebulizer adapted to receive solution from the vial and to aerosolize the
solution for delivery to the lung through oral inhalation.

141. The kit of claim 140, wherein the nebulizer operates by ultrasonic atomization.
142. The kit of claim 140, wherein the nebulizer operates by hydraulic atomization.
143. The kit of claim 140, wherein the nebulizer operates by a vibrating mesh.
144. The kit of claim 143, wherein the vibrating mesh nebulizer is a PARI E-FLOW®
nebulizer.

145. The kit of any one of claims 140 to 144, wherein the nebulizer is adapted to
produce particles having a mass median aerodynamic diameter from about 2 microns to about 5
microns with a geometric standard deviation less than or equal to about 2.5 microns.

146. The kit of any one of claims 140 to 144, wherein the nebulizer is adapted to
produce particles having a mass median aerodynamic diameter from about 2.8 microns to about
4.3 microns with a geometric standard deviation less than or equal to about 2 microns.

147. The kit of any one of claims 140 to 144, wherein the nebulizer is adapted to
produce particles having a mass median aerodynamic diameter from about 2.5 microns to about
4.5 microns with a geometric standard deviation less than or equal to about 1.8 microns.

148. The kit of any one of claims 140 to 144, wherein the pharmaceutical
composition in the container comprises levofloxacin at a concentration from about 100 mM to
about 625 mM and magnesium chloride at a concentration from about 125 mM to about 250
mM, has a pH from about 5.5 to about 6.5, and an osmolality from about 350 mOsmol/kg to
about 750 mOsmol/kg.

149. A system, comprising:
a reservoir comprising the pharmaceutical composition of any one of claims 84 to 106; and
a nebulizer configured to aerosolize the solution for delivery to the lung through oral inhalation.
150. The system of claim 149, wherein the nebulizer operates by ultrasonic atomization.
151. The system of claim 149, wherein the nebulizer operates by hydraulic atomization.
152. The system of claim 149, wherein the nebulizer operates by a vibrating mesh.
153. The system of claim 152, wherein the vibrating mesh nebulizer is a PARI E-FLOW® nebulizer.
154. The system of any one of claims 149 to 153, wherein the nebulizer is adapted to produce particles having a mass median aerodynamic diameter from about 2 microns to about 5 microns with a geometric standard deviation less than or equal to about 2.5 microns.
155. The system of any one of claims 149 to 153, wherein the nebulizer is adapted to produce particles having a mass median aerodynamic diameter from about 2.8 microns to about 4.3 microns with a geometric standard deviation less than or equal to about 2 microns.
156. The system of any one of claims 149 to 153, wherein the nebulizer is adapted to produce particles having a mass median aerodynamic diameter from about 2.5 microns to about 4.5 microns with a geometric standard deviation less than or equal to about 1.8 microns.
157. The system of any one of claims 149 to 153, wherein the pharmaceutical composition comprises levofloxacin at a concentration from about 100 mM to about 625 mM

and magnesium chloride at a concentration from about 125 mM to about 250 mM, has a pH from about 5.5 to about 6.5, and an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg.

158. The use of claim 53, wherein the aerosol comprises a mass median aerodynamic diameter from about 2 microns to about 5 microns with a geometric standard deviation less than or equal to about 2.5 microns.

159. The use of claim 53, wherein the aerosol comprises a mass median aerodynamic diameter from about 2.5 microns to about 4.5 microns with a geometric standard deviation less than or equal to about 1.8 microns.

160. The use of claim 53, wherein the aerosol comprises a mass median aerodynamic diameter from about 2.8 microns to about 4.3 microns with a geometric standard deviation less than or equal to about 2 microns.

161. The use of any one of claims 53 and 158 to 160, wherein the aerosol is produced with a vibrating mesh nebulizer.

162. The use of claim 161, wherein the vibrating mesh nebulizer is a PARI E-FLOW® nebulizer.

163. The use of claim 161 or 162, wherein the nebulizer is configured to administer at least about 20 mg of the levofloxacin or ofloxacin to the lung.

164. The use of claim 161 or 162, wherein the nebulizer is configured to administer at least about 100 mg or more of the levofloxacin or ofloxacin to the lung.

165. The use of claim 161 or 162, wherein the nebulizer is configured to administer at least about 125 mg of levofloxacin or ofloxacin to the lung.

166. The use of claim 161 or 162, wherein the nebulizer is configured to administer at least about 150 mg of levofloxacin or ofloxacin to the lung.

167. The use of any one of claims 161 to 166, wherein the nebulizer is configured to administer the aerosol to the lung in less than about 10 minutes.

168. The use of any one of claims 161 to 166, wherein the nebulizer is configured to administer the aerosol to the lung in less than about 5 minutes.

169. The use of any one of claims 161 to 166, wherein the nebulizer is configured to administer the aerosol to the lung in less than about 3 minutes.

170. The use of any one of claims 161 to 166, wherein the nebulizer is configured to administer the aerosol to the lung in less than about 2 minutes.

171. The use of any one of claims 53 and 158 to 170, wherein the aerosol is prepared from a solution comprising levofloxacin at a concentration from about 100 mM to about 625 mM and magnesium chloride at a concentration from about 125 mM to about 250 mM, having a pH from about 5.5 to about 6.5, and an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg.

172. The use of any one of claims 53 and 158 to 171, wherein the aerosol is for use in alternation with a second inhaled antimicrobial.

173. The use of claim 172, wherein the second inhaled antimicrobial is an aminoglycoside.

174. The use of claim 173, wherein the aminoglycoside is tobramycin.

175. The use of claim 172, wherein the second inhaled antimicrobial is a polymyxin.

176. The use of claim 175, wherein the polymyxin is colistin.

177. The use of claim 172, wherein the second inhaled antimicrobial is a monobactam.

178. The use of claim 177, wherein the monobactam is aztreonam.

179. Use of the composition of any one of claims 80 to 102 in preparation of an aerosol to provide a concentration in said lung of at least 32 $\mu\text{g/ml}$ of the levofloxacin or ofloxacin, for treating or preventing a microbial infection in a patient.

180. The use of claim 179, wherein the lung concentration is at least 128 $\mu\text{g/ml}$ of the levofloxacin or ofloxacin.

181. The use of claim 179, wherein the lung concentration is at least 512 $\mu\text{g/ml}$ of the levofloxacin or ofloxacin.

182. The use of claim 179, wherein the lung concentration is from 800 $\mu\text{g/mL}$ to 1600 $\mu\text{g/mL}$ of the levofloxacin or ofloxacin.

183. The use of any one of claims 179 to 182, wherein the aerosol comprises greater than about 50 mg/ml levofloxacin and magnesium chloride, has a pH from about 5.5 to about 6.5, and an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg.

184. A method of making a taste-masked pharmaceutical composition comprising forming a solution of levofloxacin or ofloxacin and a divalent or trivalent cation having a pH from about 5.5 to about 6.5 and an osmolality from about 350 mOsmol/kg to about 750 mOsmol/kg.

185. The method of claim 184, wherein the solution comprises greater than about 50 mg/ml levofloxacin.

186. The method of claim 184 or 185, wherein the divalent or trivalent cation is provided by magnesium chloride.

1/61




FIG. 1




FIG. 2

Serum or Sputum vs Time Profile of Ciprofloxacin after oral Administration of 750mg to CF Patients




FIG. 3

Time-Kill studies Using Logarithmic Cells of PAM1020




FIG. 4A

3/61




FIG. 4B




FIG. 5A




FIG. 5B

5/61

PAM1020 Regrowth After 10-minute Treatment With Levofloxacin




FIG. 6A

6/61




FIG. 6B

7/61




FIG. 6C




FIG. 6D

9/61

Time-Kill Studies Using Lete-Log Cells of PAM1020 Grown Under Oxygen Limiting Conditions




FIG. 7A

10/61

Time-Kill Studies Using Late-Log Cells of PAM1032 Grown Under Oxygen Limiting Conditions




FIG. 7B

Killing Kinetics of Levofloxacin against PAM1032 in MHB




FIG. 8A

11/61

Killing Kinetics of Levofloxacin against PAM1032 in Sputum




FIG. 8B

PAM1032 Biofilm Treatment with Levofloxacin




FIG. 9

12/61




FIG. 10




FIG. 11

13/61




FIG. 12

Sample: 049-001-01-029-C
Size: 1.0030 mg
Method: Heat/Cool/Heat
Comment: Micronized Levo 2.5 um (Sample 1)

DSC

File: C:\...\DSC\049-001\049-001-01-029-C.001
Operator: RSU
Run Date: 27-Sep-05 23:08
Instrument: DSC Q1000 V6.2 Build 204




FIG. 13

Sample: 049-001-01-029-B
Size: 1.4100 mg
Method: Heat/Cool/Heat
Comment: Unmicronized Levo (Sample 2)

DSC

File: C:\DSC\049-001\049-001-01-029-B.001
Operator: RSU
Run Date: 27-Sep-05 20:21
Instrument: DSC Q1000 V6.2 Build 204





FIG. 13 continued


16/61



Title: Unmicronized
Comment:

Date: 09-28-2005 Time: 10:38
Filename: TEMP.TIF

FIG. 14A



Title: MICRONIZED2.5
Comment:

Date: 09-28-2005 Time: 11:08
Filename: TEMP.TIF

FIG. 14B

17/61





FIG. 15

18/61




FIG. 16




FIG. 17

19/61




FIG. 18




FIG. 19

20/61




FIG. 20




FIG. 21

b) Levofloxacin base
mp = 239.2°C

Sample: 049-001-01-007
Size: 2.8910 mg
Method: Ramp
Comment: levofloxacin 50 psi

File: C:\...\049-001-01-007-levofloxacin.001
Operator: SJB
Run Date: 13-Oct-05 17:09
Instrument: DSC Q1000 V6.2 Build 204

DSC




FIG. 22 continued

c) Levofloxacin pamoic acid co-crystals
mp = 210.6°C

Sample: 049-001-03-011 dry
Size: 2.3430 mg
Method: Ramp
Comment: levofloxacin pamoate "salt"

File: C:\...\049-001-03-011 dry.001
Operator: SJB
Run Date: 13-Oct-05 15:50
Instrument: DSC Q1000 V6.2 Build 204

DSC




FIG. 22 continued

d) 1:1 levo-pamoate mixture

Sample: Levo + Pamoic mix
Size: 3.1900 mg
Method: Ramp
Comment: Levofloxacin + Pamoic acid mixture

File: C:\...\Salt Selection\Levo-Pamoic Mix.001
Operator: ABJ
Run Date: 04-Oct-05 16:35
Instrument: DSC Q1000 V6.2 Build 204

DSC




FIG. 22 continued




FIG. 23




FIG. 23 continued




FIG. 23 continued




FIG. 23 continued

29/61




FIG. 24

File: C:\...Levo Xinafoate\049-001-01-081.001
Operator: SJB
Run Date: 28-Oct-05 14:57
Instrument: DSC Q1000 V6.2 Build 204

Sample: 049-001-01-081
Size: 2.3710 mg
Method: Ramp
Comment: Levo xinafoate after drying @ 10 deg C/min
DSC




FIG. 24 continued




FIG. 25




FIG. 25 continued

Sample: Stearic Acid
Size: 2.9090 mg
Method: Ramp
Comment: Stearic Acid CH-B-02112

DSC

File: C:\... \Levo Stearic\Stearic Acid.001
Operator: SJB
Run Date: 04-Nov-05 13:18
Instrument: DSC Q1000 V6.2 Build 204




FIG. 26

File: ...049-001-01-123 @ 10 C per minute.001
Operator: SJB
Run Date: 17-Nov-05 16:49
Instrument: DSC Q1000 V6.2 Build 204

Sample: 049-001-01-123
Size: 2.2540 mg
Method: Ramp
Comment: Levo Stearic salt @ 10 C per minute




FIG. 26 continued

Sample: 049-001-01-114
Size: 2.7130 mg
Method: Ramp
Comment: Physical Mixture Stearic Acid + Levofloxacin Salt

File: ...049-001-01-114@ 10 C per minute.001
Operator: SJB
Run Date: 15-Nov-05 12:46
Instrument: DSC Q1000 V6.2 Build 204

DSC




FIG. 26 continued




FIG. 27




FIG. 27 continued




FIG. 27 continued

File: C:\... \levo oleate salt\oleic acid.002
Operator: xlw
Run Date: 30-Nov-05 17:06
Instrument: DSC Q1000 V6.2 Build 204

Sample: oleic acid
Size: 8.5840 mg
Method: Ramp
DSC




FIG. 28

Sample: 049-001-02-079-1
Size: 3.8820 mg
Method: Ramp
Comment: levofloxacin oleate salt @ 40oC

File: C:\...\1oC per min\049-001-02-079-1.001
Operator: xlw
Run Date: 12-Dec-05 15:57
Instrument: DSC Q1000 V6.2 Build 204

DSC




FIG. 28 continued

File: C:\... \1oC per min\049-001-02-083-1.001
Operator: xlw
Run Date: 13-Dec-05 04:07
Instrument: DSC Q1000 V6.2 Build 204

Sample: 049-001-02-083-1
Size: 3.8690 mg
Method: Ramp
Comment: levofloxacin oleic acid physical mixture

DSC




FIG. 28 continued

Sample: 049-001-02-082-3
Size: 8.0810 mg
Method: Ramp
Comment: levofloxacin oleic acid mixture(1-9)
DSC
File: C:\...049-001-02-082-3 mixture(1-9).001
Operator: xlw
Run Date: 02-Dec-05 14:55
Instrument: DSC Q1000 V6.2 Build 204





FIG. 28 continued

Sample: 049-001-02-82-4
Size: 4.3430 mg
Method: Ramp
Comment: Levofloxacin oleic acid mixture(9:1)
DSC
File: C:\... \049-001-02-082-4 mixture(9-1).001
Operator: xlw
Run Date: 02-Dec-05 12:04
Instrument: DSC Q1000 V6.2 Build 204



Universal V3.4C TA Instruments

FIG. 28 continued




FIG. 29




FIG. 29 continued

46/61




FIG. 29 continued




FIG. 30

48/61




FIG. 31




FIG. 32

49/61

Dissolution profile of Levo xinafote from 10 to 30 mins




FIG. 33




FIG. 34

50/61

Average dissolution profile of levo pamoate




FIG. 35

Dissolution profile of levo pamoate from 2 - 10 minutes




FIG. 36

51/61

Dissolution profile of levo pamoate
from 10 - 60 mins




FIG. 37

Average dissolution profile of levo stearate




FIG. 38

52/61

Dissolution profile of levo stearate
from 2 - 10 mins




FIG. 39

Dissolution profile of levo stearate
from 10 - 30 mins




FIG. 40




FIG. 41

54/61

Levofloxacin complexation with Mg²⁺: Dual titration II




FIG. 42

Levofloxacin complexation with FeCl₂: Dual titration




FIG. 43

55/61


Levofloxacin complexation with CaCl₂: Dual titration

FIG. 44


Levofloxacin complexation with Zn²⁺: Dual titration

FIG. 45

56/61





FIG. 46

57/61





FIG. 47

58/61





FIG. 48

59/61







FIG. 49

60/61

**FIG. 50****FIG. 51**

61/61

Complexation: Spectrofluorometry




FIG. 52

Complexation: Spectrofluorometry




FIG. 53