



HU000030916T2

(19) **HU**(11) Lajstromszám: **E 030 916**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 11 781949**(51) Int. Cl.: **C07K 16/24**

(2006.01)

(22) A bejelentés napja: **2011. 11. 02.**

(86) A nemzetközi (PCT) bejelentési szám:

PCT/US 11/058869(96) Az európai bejelentés bejelentési száma:
EP 20110781949

(87) A nemzetközi közzétételi szám:

WO 12061448(97) Az európai bejelentés közzétételi adatai:
EP 2635601 A1 2012. 05. 10.(97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2635601 B1 2016. 07. 20.

(30) Elsőbbségi adatai:

201161448785 P	2011. 03. 03.	US
412594 P	2010. 11. 11.	US
411953 P	2010. 11. 10.	US
410158 P	2010. 11. 04.	US

(73) Jogosult(ak):

**Boehringer Ingelheim International GmbH,
55216 Ingelheim am Rhein (DE)**

(72) Feltaláló(k):

BARRETT, Rachel Rebecca, Ridgefield, Connecticut 06877-0368 (US)
CANADA, Keith, Ridgefield, Connecticut 06877-0368 (US)
CATRON, Katrina Mary, Ridgefield, CT 06877-0368 (US)
COPENHAVER, Robert, PortlandOR 97239 (US)
FREGO, Lee Edward, Ridgefield, Connecticut 06877-0368 (US)
RAYMOND, Ernest Lee, Ridgefield, Connecticut 06877-0368 (US)
SINGH, Sanjaya, Ridgefield, Connecticut 06877-0368 (US)
ZHU, Xiangyang, Ridgefield, Connecticut 06877-0368 (US)

(74) Képviselő:

SBGK Szabadalmi Ügyvivői Iroda, Budapest

(54)

Anti-IL-23 antitestek

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11)

EP 2 635 601 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

20.07.2016 Bulletin 2016/29

(21) Application number: **11781949.0**

(22) Date of filing: **02.11.2011**

(51) Int Cl.:

C07K 16/24 (2006.01)

(86) International application number:

PCT/US2011/058869

(87) International publication number:

WO 2012/061448 (10.05.2012 Gazette 2012/19)

(54) **ANTI-IL-23 ANTIBODIES**

ANTI-IL-23-ANTIKÖRPER

ANTICORPS ANTI-IL-23

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

Designated Extension States:

BA ME

(30) Priority: **03.03.2011 US 201161448785 P**

11.11.2010 US 412594 P

10.11.2010 US 411953 P

04.11.2010 US 410158 P

(43) Date of publication of application:

11.09.2013 Bulletin 2013/37

(73) Proprietor: **Boehringer Ingelheim International**

GmbH

55216 Ingelheim am Rhein (DE)

(72) Inventors:

- **BARRETT, Rachel Rebecca**
Ridgefield, Connecticut 06877-0368 (US)
- **CANADA, Keith**
Ridgefield, Connecticut 06877-0368 (US)
- **CATRON, Katrina Mary**
Ridgefield, CT 06877-0368 (US)
- **COPENHAVER, Robert**
Portland
OR 97239 (US)
- **FREGO, Lee Edward**
Ridgefield, Connecticut 06877-0368 (US)
- **RAYMOND, Ernest Lee**
Ridgefield, Connecticut 06877-0368 (US)
- **SINGH, Sanjaya**
Ridgefield, Connecticut 06877-0368 (US)

- **ZHU, Xiangyang**
Ridgefield, Connecticut 06877-0368 (US)

(74) Representative: **Simon, Elke Anna Maria et al**
Boehringer Ingelheim GmbH
Binger Strasse 173
55216 Ingelheim am Rhein (DE)

(56) References cited:

**WO-A2-2006/036922 WO-A2-2007/005955
WO-A2-2007/024846 WO-A2-2007/027714**

- **CHAN ANDREW C ET AL: "Therapeutic antibodies for autoimmunity and inflammation", NATURE REVIEWS. IMMUNOLOGY, NATURE PUBLISHING GROUP, GB, vol. 10, no. 5, 1 May 2010 (2010-05-01), pages 301-316, XP002665072, ISSN: 1474-1733, DOI: 10.1038/NRI2761**
- **ALEGRE M-L ET AL: "A NON-ACTIVATING HUMANIZED ANTI-CD3 MONOCLONAL ANTIBODY RETAINS IMMUNOSUPPRESSIVE PROPERTIES IN VIVO", TRANSPLANTATION, WILLIAMS AND WILKINS, BALTIMORE US, vol. 57, no. 11, 1 June 1994 (1994-06-01), pages 1537-1543, XP008036788, ISSN: 0041-1337**
- **WOODLE E STEVE ET AL: "Phase I trial of a humanized, Fc receptor nonbinding OKT3 antibody, huOKT3gamma1(Ala-Ala) in the treatment of acute renal allograft rejection", TRANSPLANTATION, WILLIAMS AND WILKINS, BALTIMORE US, vol. 68, no. 5, 15 September 1999 (1999-09-15), pages 608-616, XP009134473, ISSN: 0041-1337**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Remarks:

The file contains technical information submitted after
the application was filed and not included in this
specification

Description**Technical Field of the Invention**

5 [0001] This invention generally relates to anti-IL-23p19 antibodies for diagnostic and therapeutic use. More specifically, humanized anti-IL-23p19 antibodies and methods of use for the treatment of various diseases or disorders are disclosed. Pharmaceutical compositions and kits comprising such compounds are also disclosed.

Background of the Invention

10 [0002] Higher eukaryotes have evolved an intricate response to pathogens that is initiated by the innate immune response and followed by the adaptive immune response. Together these two mechanisms not only eradicate pathogens that infect the organism but also establish a long term immunological response against future exposures. Deficiencies in these responses can result in increased susceptibility to infections and/or alterations of the adaptive immune response 15 leading to chronic inflammation and autoimmunity. IL-12, a heterodimeric cytokine consisting of a p40 and a p35 protein subunit, has long been considered the hallmark cytokine of the innate immune response with major influence on adaptive immunity. However, data from investigation of this cytokine's biological role led to confusing results. For example, while p40-deficient mice were resistant to Collagen Induced Arthritis (CIA) and Experimental Autoimmune Encephalomyelitis (EAE), p35-deficient mice were susceptible to both and even displayed exacerbated disease. Such conundrums began 20 to be resolved with the discovery in the late 1990's of a new member of the IL-12 cytokine family with a distinct role in the immune response - IL-23.

[0003] IL-23 is composed of a common subunit (p40) with IL-12 and a unique p19 subunit. Despite this shared p40 subunit, the roles for IL-23 and IL-12 are quite different. IL-12 is important for Th1 responses via promotion of Th1 cell differentiation, proliferation and activation. In contrast, IL-23 supports the development and maintenance of a recently defined set of CD4⁺ T helper cells termed Th17 cells due to their ability to produce IL-17 and related cytokines. There is mounting evidence that IL-23 is involved in chronic autoimmune inflammation and the modulation of IL-23 activity could provide promising therapies against autoimmune diseases.

[0004] There is therefore a need for antagonist molecules against IL-23 with beneficial pharmacological properties, which can be used as therapeutic agents to treat diseases, in particular immunological and autoimmune diseases in 30 humans. Accordingly, one aim of the present invention is to provide anti-IL-23 antagonist molecules, in particular anti-IL-23 antagonist molecules which have high binding affinity to IL-23.

[0005] A further aim of the present invention is to provide anti-IL-23 antagonist molecules, which have high specificity for IL-23.

[0006] A further aim of the present invention is to provide anti-IL-23 antagonists, which have high blocking activity for 35 the association of IL-23 and its receptor.

[0007] A further aim of the present invention is to provide anti-IL-23 antagonists, which have potent cellular activity.

[0008] A further aim of the present invention is to provide anti-IL-23 antagonists, which have a favorable bioavailability.

[0009] A further aim of the present invention is to provide anti-IL-23 antagonists, which have favorable biophysical properties.

40 [0010] Further aims of the present invention include combinations of any of the aims set forth above.

[0011] Antibodies against IL-23p19 are known from for example WO 2007/024846.

Summary of the Invention

45 [0012] The present invention addresses the above needs and provides antibodies that bind to the p19 subunit of the IL-23 protein. The invention is as set out in the claims. In one aspect, an antibody of the present disclosure binds to human IL-23 with high affinity. In another aspect, an antibody of the present disclosure inhibits the IL-23 stimulated production of IL-17 from mouse splenocytes. In another aspect, an antibody of the present disclosure does not bind to nor antagonize IL-12, which is a closely related family member to IL-23.

50 [0013] In one embodiment, the present disclosure provides anti-IL-23p19 antibodies that are derived from mouse hybridomas, for example monoclonal antibodies. In one embodiment, the present disclosure provides full length anti-IL-23p19 antibodies. In another embodiment, the present disclosure provides anti-IL-23p19 humanized antibodies, for example humanized monoclonal anti-IL-23p19 antibodies, for example full length humanized monoclonal anti-IL-23p19 antibodies. In one aspect, a humanized antibody of the present disclosure binds to human IL-23 with high affinity. In another aspect, a humanized antibody of the present disclosure also binds to cynomolgus IL-23 with high affinity. In a further aspect, a humanized antibody of the present disclosure inhibits IL-23-induced STAT3 phosphorylation in DB cells. In another aspect, a humanized antibody of the present disclosure antagonizes the action of IL-23 by binding to the p19 subunit of IL-23, for example as measured by the inhibition of cytokines such as IL-17 and IL-22, whose production

is stimulated by IL-23. In a further aspect, a humanized antibody of the present disclosure has a favorable pharmacokinetic (PK) profile. In a further aspect, a humanized antibody of the present invention has favorable biophysical properties, such as quality, stability or solubility, for example as defined by the percentage of antibody in monomer form.

[0014] Further embodiments encompass DNA molecules encoding antibodies of the present invention expression vectors and host cells comprising such DNA molecules, and methods of making antibodies of the present invention. The present invention further provides therapeutic uses for the antibodies of the present invention, in particular against immunological and autoimmune diseases.

[0015] In one embodiment, the present disclosure further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising a light chain CDR1 (L-CDR1) sequence of SEQ ID NO:1, 4, 6, 7, 8, 11, 15, 18, 19, 22, 27 or 30; a light chain CDR2 (L-CDR2) sequence of SEQ ID NO:2, 5, 9, 12, 16, 20, 23, 25, 28 or 31; a light chain CDR3 (L-CDR3) sequence of SEQ ID NO:3, 10, 13, 14, 17, 21, 24, 26, 29, or 32; a heavy chain CDR1 (H-CDR1) sequence of SEQ ID NO:33, 36, 38, 40, 43, 45, 48, 51, 54, 57, 60, 63, 66, 67, 68, 69, 77 or 80; a heavy chain CDR2 (H-CDR2) sequence of SEQ ID NO:34, 39, 41, 46, 49, 52, 55, 58, 61, 64, 70, 72, 73, 75, 78 or 81; and a heavy chain CDR3 (H-CDR3) sequence of SEQ ID NO:35, 37, 42, 44, 47, 50, 53, 56, 59, 62, 65, 71, 74, 76, 79 or 82. In one embodiment, the anti-IL-23p19 antibody or antigen-binding fragment thereof comprises a light chain variable region comprising a L-CDR1 listed above, a L-CDR2 listed above and a L-CDR3 listed above, and a heavy chain variable region comprising a H-CDR1 listed above, a H-CDR2 listed above and a H-CDR3 listed above.

[0016] In one embodiment, the present disclosure further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:1, 2, 3, 33, 34, and 35, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:4, 5, 3, 36, 34 and 37, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:1, 2, 3, 38, 39 and 35, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:6, 2, 3, 40, 41 and 42, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:7, 2, 3, 43, 41 and 44, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:8, 9, 10, 45, 46 and 47, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:8, 9, 10, 48, 49 and 50, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:11, 12, 13, 51, 52 and 53, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:7, 2, 14, 54, 55 and 56, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:15, 16, 17, 57, 58 and 59, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:18, 16, 17, 60, 61 and 62, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:19, 20, 21, 63, 66, 67 or 68, 64 and 65, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:22, 23, 24, 69, 70 and 71, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:22, 25, 26, 55, 72 and 71, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:8, 9, 10, 45, 73 and 74, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:27, 28, 29, 45, 75 and 76, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:8, 9, 10, 77, 78 and 79, respectively; or a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:30, 31, 32, 80, 81 and 82, respectively. In one embodiment, the anti-IL-23p19 antibody or antigen-binding fragment thereof comprises a light chain variable region comprising a L-CDR1, L-CDR2 and L-CDR3 combination listed above, and a heavy chain variable region comprising a H-CDR1, H-CDR2 and H-CDR3 combination listed above.

[0017] In one embodiment, the present disclosure further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:84 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:121; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:86 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:123; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:88 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:125; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:90 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:127; or a Light chain variable region comprising the amino acid sequence of SEQ ID NO:91 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:128; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:93 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:130; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:95 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:132; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:97 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:134; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:99 and a heavy chain variable region comprising the amino acid sequence SEQ

ID NO:136; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:101 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:138; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:103 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:140; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:105 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:142; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:107 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:144; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:109 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:146; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:111 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:148; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:113 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:150; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:115 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:152; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:117 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:154; or a light chain variable region comprising the amino acid sequence of SEQ ID NO:119 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:156.

[0018] In one embodiment, the present disclosure further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NO:158, 160, 162 and 164 and a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NO:166, 168 170 and 172.

[0019] In one embodiment, the present disclosure further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof has a K_D for IL-23 of less than 40pM, or a K_D for IL-23 of less than 20pM, or K_D for IL-23 of less than 10pM or K_D for IL-23 of less than 1pM.

[0020] In a further embodiment, the present disclosure provides an anti-IL-23p19 antibody or antigen-binding fragment thereof that binds to human IL-23p19 at an epitope consisting of amino acid residues 108 to 126 and amino acid residues 137 to 151 of SEQ ID NO: 181.

[0021] In a further embodiment, the present disclosure provides an anti-IL-23p19 antibody or antigen-binding fragment thereof that competitively binds to human IL-23p19 with an antibody of the present disclosure. In one embodiment, the present disclosure provides an anti-IL-23p19 antibody or antigen-binding fragment thereof that competitively binds to human IL-23p19 with a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:174 and a heavy chain comprising the amino acid sequence of SEQ ID NO:176. In one embodiment the present disclosure provides an anti-IL-23p19 antibody or antigen-binding fragment thereof that competitively binds to human IL-23p19 with a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:174 and a heavy chain comprising the amino acid sequence of SEQ ID NO:178. In one embodiment, the present disclosure provides an anti-IL-23p19 antibody or antigen-binding fragment thereof that competitively binds to human IL-23p19 with a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:180 and a heavy chain comprising the amino acid sequence of SEQ ID NO:176. In one embodiment, the present disclosure provides an anti-IL-23p19 antibody or antigen-binding fragment thereof that competitively binds to human IL-23p19 with a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO: 180 and a heavy chain comprising the amino acid sequence of SEQ ID NO: 178.

[0022] In the invention, the anti-IL-23p19 antibody is a humanized antibody. In one embodiment, the anti-IL-23p19 antibody is a monoclonal antibody. In one embodiment, the anti-IL-23p19 antibody is a full length antibody. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal antibody, for example a full length humanized monoclonal antibody. In one embodiment the antigen-binding fragment is a Fab, F(ab')₂, or single chain Fv fragment. In one embodiment, the antigen-binding fragment comprises a light chain variable region and a heavy chain variable region.

[0023] In one embodiment, the present disclosure further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises the amino acid sequence of SEQ ID NO:19 (CDR1-L); the amino acid sequence of SEQ ID NO:20 (CDR2-L); the amino acid sequence of (SEQ ID NO:21 (CDR3-L); the amino acid sequence of SEQ ID NO:63, 66, 67 or 68 (CDR1-H); the amino acid sequence of SEQ ID NO:64 (CDR2-H); and the amino acid sequence of SEQ ID NO:65 (CDR3-H).

[0024] In one embodiment, the present invention further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises the amino acid sequence of SEQ ID NO:19 (CDR1-L); the amino acid sequence of SEQ ID NO:20 (CDR2-L); the amino acid sequence of SEQ ID NO:21 (CDR3-L); the amino acid sequence of SEQ ID NO:66 (CDR1-H); the amino acid sequence of SEQ ID NO:64 (CDR2-H); and the amino acid sequence of SEQ ID NO:65 (CDR3-H).

[0025] In one embodiment, the present disclosure further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:19 (*CDR1-L*); the amino acid sequence of SEQ ID NO:20 (*CDR2-L*); and the amino acid sequence of SEQ ID NO:21 (*GDR3-L*); and a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:63, 66, 67 or 68 (*CDR1-H*); the amino acid sequence of SEQ ID NO:64 (*CDR2-H*); and the amino acid sequence of SEQ ID NO:65 (*CDR3-H*).

[0026] In one embodiment, the present invention further provides an anti-IL-23p19 or antigen-binding fragment thereof antibody, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:19 (*CDR1-L*); the amino acid sequence of SEQ ID NO:20 (*CDR2-L*); and the amino acid sequence of SEQ ID NO:21 (*CDR3-L*); and a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:66 (*CDR1-H*); the amino acid sequence of SEQ ID NO:64 (*CDR2-H*); and the amino acid sequence of SEQ ID NO:65 (*CDR3-H*).

[0027] In one embodiment, the present disclosure further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of any one of SEQ ID NO:158, 160, 162 or 164; and a heavy chain variable region comprising the amino acid sequence any one of SEQ ID NO:166, 168, 170 or 172.

[0028] In one embodiment, the present invention further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:160 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:166.

[0029] In one embodiment, the present invention further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:160 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:168.

[0030] In one embodiment, the present invention further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:158 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:166.

[0031] In one embodiment, the present invention further provides an anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:158 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:168.

[0032] In one embodiment, the anti-IL-23p19 antibody is a humanized antibody. In one embodiment, the anti-IL-23p19 antibody is a monoclonal antibody. In one embodiment, the anti-IL-23p19 antibody is a full length antibody. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal anti-IL-23p19 antibody, for example a full length humanized monoclonal antibody. In one embodiment, the antigen-binding fragment is a Fab, F(ab')₂, or single chain Fv fragment. In one embodiment, the antigen-binding fragment comprises a light chain variable region and a heavy chain variable region.

[0033] In one embodiment, the present disclosure further provides an antibody comprising the amino acid sequence SEQ ID NO:166 or 168 linked to a human IgG1, IgG2, IgG3, IgG4, IgM, IgA or IgE heavy chain constant region. An antibody comprising the amino acid sequence of SEQ ID NO: 166 or 168 linked to a human IgG1 heavy chain constant region. An antibody comprising the amino acid sequence of SEQ ID NO:158 or 160 linked to a human kappa or lambda light chain constant region. An antibody comprising the amino acid sequence of SEQ ID NO:158 or 160 linked to a human kappa light chain constant region.

[0034] In one embodiment, the present invention further provides an antibody comprising the amino acid sequence of SEQ ID NO:166 or 168 linked to a human IgG1 heavy chain constant region; and the amino acid sequence of SEQ ID NO:158 or 160 linked to a human kappa light chain constant region.

[0035] In one embodiment, the present disclosure further provides a humanized monoclonal anti-IL-23p19 antibody comprising a light chain variable region comprising the amino acid sequence selected from the group consisting of any one of SEQ ID NO:158, 160, 162 and 164 and a heavy chain variable region comprising the amino acid sequence selected from the group consisting of any one of SEQ ID NO:166, 168, 170 and 172.

[0036] In one embodiment, the present invention further provides a humanized monoclonal anti-IL-23p19 antibody comprising a light chain variable region comprising the amino acid sequence of SEQ ID NO:160 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:166.

[0037] In one embodiment, the present invention further provides a humanized monoclonal anti-IL-23p19 antibody comprising a light chain variable region comprising the amino acid sequence of SEQ ID NO:160 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:168.

[0038] In one embodiment, the present invention further provides a humanized monoclonal anti-IL-23p19 antibody

comprising a light chain variable region comprising the amino acid sequence of SEQ ID NO:158 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:166.

[0039] In one embodiment, the present invention further provides a humanized monoclonal anti-IL-23p19 antibody comprising a light chain variable region comprising the amino acid sequence of SEQ ID NO:158 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:168.

[0040] In one embodiment, the present invention further provides a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:174 or 180 and a heavy chain comprising the amino acid sequence of SEQ ID NO:176 or 178.

[0041] In one embodiment, the present invention further provides a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:174 and a heavy chain comprising the amino acid sequence of SEQ ID NO:176.

[0042] In one embodiment, the present invention further provides a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:174 and a heavy chain comprising the amino acid sequence of SEQ ID NO:178.

[0043] In one embodiment, the present invention further provides a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:180 and a heavy chain comprising the amino acid sequence of SEQ ID NO:176.

[0044] In one embodiment, the present invention further provides a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO: 180 and a heavy chain comprising the amino acid sequence of SEQ ID NO: 178.

[0045] In a further embodiment, the present invention relates to an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising a humanized light chain variable domain comprising the CDRs of SEQ ID NO:160 and framework regions having an amino acid sequence at least 90% identical to the amino acid sequence of the framework regions of the variable domain light chain amino acid sequence of SEQ ID NO:160 and a humanized heavy chain variable domain comprising the CDRs of SEQ ID NO:166 and framework regions having an amino acid sequence at least 90% identical to the amino acid sequence of the framework regions of the variable domain heavy chain amino acid sequence of SEQ ID NO:166. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal antibody, for example a full length humanized monoclonal antibody.

[0046] In a further embodiment, the present invention relates to an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising a humanized light chain variable domain comprising the CDRs of SEQ ID NO:160 and framework regions having an amino acid sequence at least 90% identical to the amino acid sequence of the framework regions of the variable domain light chain amino acid sequence of SEQ ID NO:160 and a humanized heavy chain variable domain comprising the CDRs of SEQ ID NO:168 and framework regions having an amino acid sequence at least 90% identical to the amino acid sequence of the framework regions of the variable domain heavy chain amino acid sequence of SEQ ID NO:168. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal antibody, for example a full length humanized monoclonal antibody.

[0047] In a further embodiment, the present invention relates to an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising a humanized light chain variable domain comprising the CDRs of SEQ ID NO:158 and framework regions having an amino acid sequence at least 90% identical to the amino acid sequence of the framework regions of the variable domain light chain amino acid sequence of SEQ ID NO:158 and a humanized heavy chain variable domain comprising the CDRs of SEQ ID NO:166 and framework regions having an amino acid sequence at least 90% identical to the amino acid sequence of the framework regions of the variable domain heavy chain amino acid sequence of SEQ ID NO:166. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal antibody, for example a full length humanized monoclonal antibody.

[0048] In a further embodiment, the present invention relates to an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising a humanized light chain variable domain comprising the CDRs of SEQ ID NO:158 and framework regions having an amino acid sequence at least 90% identical to the amino acid sequence of the framework regions of the variable domain light chain amino acid sequence of SEQ ID NO:158 and a humanized heavy chain variable domain comprising the CDRs of SEQ ID NO:168 and framework regions having an amino acid sequence at least 90% identical to the amino acid sequence of the framework regions of the variable domain heavy chain amino acid sequence of SEQ ID NO:168. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal antibody, for example a full length humanized monoclonal antibody.

[0049] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized by a K_D for human IL-23 equal or less than 1 pM. In one aspect, there is no shift in binding on-rate in 50% human serum.

[0050] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized in that it blocks IL-23 binding to human IL-23R/Fc in vitro.

[0051] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized in that it does not bind to human IL-12.

[0052] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized in that it inhibits human IL-23 induced IL-17 production in mouse splenocytes with IC₅₀'s equal or less than 20 pM.

[0053] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized in that it inhibits human IL-23 induced STAT3 phosphorylation in human DB cells with IC₅₀'s equal or less than 40 pM.

5 [0054] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized in that it has no predicted activity in ADCC/CDC.

[0055] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized in that it has a K_D equal or less than 1 pM for cynomolgus monkey IL-23.

10 [0056] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized in that it has no cross reactivity to mouse or rat IL-23.

[0057] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized in that it inhibits human IL-23 induced IL-17 and IL-22 production in a mouse ear at 80% or greater inhibition of both cytokines at 1 mg/kg.

15 [0058] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized by a melting temperature of 83°C as determined by differential scanning calorimetry.

[0059] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure may be further characterized by solubility equal or greater than 100 mg/ml, as measured by UV spectroscopy and monitored by turbidity.

20 [0060] In a further aspect, a humanized anti-IL-23p19 antibody of the present invention may be further characterized in that it is present in at least 90% monomer form, or in at least 92% monomer form, or in at least 95% monomer form in a buffer.

[0061] In a further aspect, a humanized anti-IL-23p19 antibody of the present invention may be further characterized in that it remains in at least 90% monomer form, or in at least 92% monomer form, or in at least 95% monomer form in a buffer for one month or for four months.

25 [0062] In one aspect, the humanized anti-IL-23p19 antibody is a humanized monoclonal antibody, for example a full length humanized monoclonal antibody.

[0063] Further embodiments encompass a DNA molecule encoding a variable light chain region above, a DNA molecule encoding a variable heavy chain region above, a DNA molecule encoding a light chain region above, or a DNA molecule encoding a heavy chain region above.

30 [0064] Further embodiments encompass an expression vector containing a DNA molecule above. In one embodiment, an expression vector comprises a DNA molecule encoding the constant heavy chain and/or the constant light chain, respectively, linked to the DNA molecule encoding the variable heavy chain and/or the variable light chain, respectively. Further embodiments encompass a host cell carrying one or more expression vectors above. In one embodiment, a host is a mammalian cell.

35 [0065] Further embodiments encompass a method for producing an antibody or antigen-binding fragment thereof above comprising transfecting a mammalian host cell with one or more of the vectors above, cultivating the host cell and recovering and purifying the antibody or antigen-binding fragment thereof.

40 [0066] Further embodiments encompass a method for producing an antibody or antigen-binding fragment thereof above comprising obtaining a mammalian host cell comprising one or more of the vectors above, and cultivating the host cell. In one embodiment, the method further comprises recovering and purifying the antibody or antigen-binding fragment thereof.

45 [0067] In one embodiment, the present disclosure further provides an antibody or antigen-binding fragment thereof above for use in medicine. In one embodiment, the use is the treatment of an inflammatory disease, of an autoimmune disease, of a respiratory disease, of a metabolic disorder or of cancer. In one embodiment, the use is for the treatment of psoriasis, inflammatory bowel disease (Crohn's disease, ulcerative colitis), psoriatic arthritis, multiple sclerosis, rheumatoid arthritis, or ankylosing spondylitis. In one embodiment, the use is for the treatment of psoriasis. In one embodiment, the use is for the treatment of inflammatory bowel disease.

50 [0068] In one embodiment, the present disclosure further provides a pharmaceutical composition comprising an antibody molecule or antigen-binding fragment above and a pharmaceutically acceptable carrier.

55 [0069] In one embodiment, the present disclosure further provides a method for treating an inflammatory disease, an autoimmune disease, a respiratory disease, a metabolic disorder or cancer comprising administering to a subject in need thereof, for example a patient, an effective amount of an anti-IL-23p19 antibody or antigen-binding fragment thereof above, or a pharmaceutical composition comprising the antibody or antigen-binding fragment thereof. In one embodiment, the antibody or antigen-binding fragment is administered by a parenteral route of administration; or is administered intravenously or subcutaneously. In one embodiment, the antibody or antigen-binding fragment is administered subcutaneously. In one embodiment, the disease is psoriasis, inflammatory bowel disease (Crohn's disease, ulcerative colitis), psoriatic arthritis, multiple sclerosis, rheumatoid arthritis, or ankylosing spondylitis. In one embodiment, the disease is psoriasis. In one embodiment, the disease is inflammatory bowel disease.

[0070] In one embodiment, the present disclosure further provides a method for inhibiting the binding of IL-23 to the

IL-23 receptor on a mammalian cell, comprising administering to the cell an antibody molecule or antigen-binding fragment above, whereby signaling mediated by the IL-23 receptor is inhibited.

[0071] In one embodiment, the present disclosure further provides a method for treating a subject having an IL-23-associated disorder, comprising administering to the subject an antibody or antigen-binding fragment above, which antibody or antigen-binding fragment binds to human IL-23.

[0072] In one embodiment, the present disclosure further provides a method for detecting and/or quantifying IL-23 levels in a biological sample by contacting the sample with an antibody or antigen binding fragment above and detecting binding of the antibody or fragment thereof with IL-23p19. This information can be used to diagnose an IL-23-associated disorder. Thus, methods are provided for diagnosing an IL-23-associated disorder or for determining if a subject has an increased risk of developing an IL-23-associated disorder, wherein the method comprises contacting a biological sample from a subject with an antibody or antigen binding fragment above and detecting binding of the antibody or antigen binding fragment to IL-23p19 to determine the expression or concentration of IL-23.

[0073] In one embodiment, the present disclosure further provides a method for inhibiting the binding of IL-23 to the IL-23 receptor on a cell, comprising administering to the cell or cellular environment an antibody or antigen-binding fragment above, whereby signaling mediated by the IL-23 receptor is inhibited.

Brief Description of the Figures

[0074]

Figure 1: Alignment of mouse and humanized variable regions. Figure 1a: Anti-IL-23p19 6B8 Engineered Vk regions. Figure 1b: Anti-IL-23p19 6B8 Engineered VH regions.

The numbering of the amino acids is by the standard Kabat numbering scheme. Regular font = Human; italic/unnderlined font = Murine; shadowed font = Synthetic; bold/italic/underlined = CDR.

Figure 2: Competition binding assay of human of IL-23 binding to IL-23R/Fc.

Detailed Description

[0075] The p19 subunit of IL-23 (also referred to herein as "IL-23p19" and "p19 subunit") is a 189 amino acid polypeptide containing a 21 aa leader sequence (Oppmann et al. Immunity 13:715 (2000), SEQ ID NO: 181). The biological activity of the molecule is only detected when it is partnered with the IL-12p40 subunit to form IL-23. IL-23 is predominantly expressed by activated dendritic cells (DCs) and phagocytic cells. The receptor for IL-23 was found to be composed of the IL-12R β 1 subunit of IL-12 receptor partnered with a unique subunit called IL-23R (Parham et al. J. Immunol. 168:5699 (2002)). Expression of the receptor is detected primarily on memory T cells and NK cells. Thus, expression of this cytokine:receptor pair appears to be restricted to specific populations of immune cells. While it was first thought that IL-12 and IL-23 would share many functions, the data has shown the picture to be different. Whereas IL-12 has a predominant role in the production of Th1 cells, IL-23 was found to be critically involved in the production and maintenance of a recently recognized Th cell subset termed Th17 (Kikly et al. Curr. Opin. Immunol. 18:670 (2006), Kastelein et al. Ann. Rev. Immunol. 25:221 (2007)). These cells produce IL-17A, IL-17F, IL-22 and other pro-inflammatory cytokines such as IL-6 and TNF- α . As described below, animal model studies on the role of these Th17 cells show their importance as a driving force in chronic inflammation and autoimmunity.

[0076] The present disclosure provides antibodies that bind to the p19 subunit of the IL-23, in particular human IL-23p19. The present disclosure also relates to humanized antibodies that recognize the p19 subunit of IL-23. In specific embodiments, the sequence of these humanized antibodies has been identified based on the sequences of certain lead mouse antibodies.

[0077] The lead mouse antibodies of the present disclosure were derived from mouse hybridomas. The immunization of the mice is carried out using different techniques. For example, antibodies that are specific for human IL-23p19 proteins or fragments thereof can be raised against an immunogenic antigen such as an isolated IL-23p19 protein, an isolated IL-23 protein, an isolated hybrid IL-23 protein, and/or a portion thereof of any of the above (including synthetic peptides). For example, a hybrid IL-23 protein comprising a mouse IL-23p40 subunit and a human IL-23p19 subunit is used to immunize mice. Preparation of immunogenic antigens and monoclonal antibody production can be performed using any suitable technique known in the art.

[0078] The lead mouse antibodies were selected based on their high affinity to human IL-23. Accordingly, in one aspect, the present disclosure provides an antibody that binds to human IL-23 with high affinity. Selected mouse antibodies were humanized to result in humanized antibodies. The humanized antibodies of the present invention bind to human IL-23 with high affinity. Accordingly, in another aspect, the present disclosure provides a humanized antibody that binds to human IL-23 with high affinity. Accordingly, in one embodiment, the present invention provides an anti-IL-23p19 antibody having a K_D of less than 40pM. In a further embodiment, the present disclosure provides an anti-IL-23p19

antibody having a K_D of less than 20pM. In a further embodiment, the present disclosure provides an anti-IL-23p19 antibody having a K_D less than 10pM. In a further embodiment, the present disclosure provides an anti-IL-23p19 antibody having a K_D less than 1pM.

[0079] In another aspect, an antibody of the present disclosure binds to IL-23p19 with high affinity in the absence of 5 human serum or in the presence of 50% human serum. In a further aspect, a humanized antibody of the present disclosure also binds to cynomolgus monkey IL-23 with high affinity.

[0080] In another aspect, an antibody of the present disclosure binds to IL-23, but does not bind to IL-12. In a further aspect, an antibody of the present invention does not interfere with the biological activity of IL-12, which is a closely related family member to IL-23.

[0081] In another aspect, an antibody of the present disclosure inhibits the IL-23 stimulated production of IL-17 from 10 mouse splenocytes.

[0082] In a further aspect, a humanized antibody of the present disclosure inhibits IL-23-induced STAT3 phosphorylation in DB cells.

[0083] In a further aspect, a humanized antibody of the present disclosure antagonizes the action of IL-23 by binding 15 to the p19 subunit of IL-23, as measured by the inhibition of cytokines such as IL-17 and IL-22, whose production is stimulated by IL-23, and detected by the reduction in the levels of these cytokines.

[0084] In a further aspect, a humanized antibody of the present disclosure has a favorable pharmacokinetic profile (PK) profile, as exemplified by in vivo half life in cynomolgus monkeys.

[0085] In a further aspect, a humanized monoclonal anti-IL-23p19 antibody of the present disclosure has favorable 20 biophysical properties, for example quality, stability, or solubility.

[0086] In one aspect, the anti-IL-23p19 antibody is a humanized antibody. In one aspect, the anti-IL-23p19 antibody is a monoclonal antibody. In one aspect, the anti-IL-23p19 antibody is a full length antibody. In one aspect, the anti-IL-23p19 antibody is a humanized monoclonal antibody, for example a full length humanized monoclonal antibody.

[0087] An antibody or antigen-binding fragment thereof of the present disclosure recognizes specific "IL-23p19 antigen epitope" or "IL-23p19 epitope". As used herein these terms refer to a molecule (e.g., a peptide) or a fragment of a molecule capable of immunoreactivity with an anti-IL-23p19 antibody and, for example, include an IL-23p19 antigenic determinant recognized by any of the antibodies having a light chain/heavy chain sequence combination of SEQ ID NO:84/121, 86/123, 88/125, 90/127, 91/128, 93/130, 95/132, 97/134, 99/136, 101/138, 103/140, 105/142, 107/144, 109/146, 111/148, 113/150, 115/152, 117/154, 119/156, 160/166, 160/168, 158/166 or 158/168. IL-23p19 antigen epitopes can be included in proteins, protein fragments, peptides or the like. The epitopes are most commonly proteins, 30 short oligopeptides, oligopeptide mimics (i.e., organic compounds that mimic antibody binding properties of the IL-23p19 antigen), or combinations thereof. The minimum size of a peptide or polypeptide epitope for an antibody is thought to be about four to five amino acids. Peptide or polypeptide epitopes contain for example at least seven amino acids or for example at least nine amino acids or for example between about 15 to about 20 amino acids. Since an antibody can 35 recognize an antigenic peptide or polypeptide in its tertiary form, the amino acids comprising an epitope need not be contiguous, and in some cases, may not even be on the same peptide chain. Epitopes may be determined by various techniques known in the art, such as X-ray crystallography, Hydrogen/Deuterium Exchange Mass Spectrometry (HXMS), site-directed mutagenesis, alanine scanning mutagenesis, and peptide screening methods.

[0088] The generalized structure of antibodies or immunoglobulin is well known to those of skill in the art. These 40 molecules are heterotetrameric glycoproteins, typically of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains and are typically referred to as full length antibodies. Each light chain is covalently linked to a heavy chain by one disulfide bond to form a heterodimer, and the heterotetrameric molecule is formed through a covalent disulfide linkage between the two identical heavy chains of the heterodimers. Although the light and heavy chains are linked together by one disulfide bond, the number of disulfide linkages between the two heavy 45 chains varies by immunoglobulin isotype. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at the amino terminus a variable domain (V_H), followed by three or four constant domains (C_{H1} , C_{H2} , C_{H3} , and C_{H4}), as well as a hinge region between C_{H1} and C_{H2} . Each light chain has two domains, an amino-terminal variable domain (V_L) and a carboxy-terminal constant domain (C_L). The V_L domain associates non-covalently with the V_H domain, whereas the C_L domain is commonly covalently linked to the C_{H1} domain via a disulfide bond. Particular amino acid residues are believed to form an interface between the light and heavy chain variable domains 50 (Chothia et al., 1985, J. Mol. Biol. 186:651-663). Variable domains are also referred herein as variable regions.

[0089] Certain domains within the variable domains differ extensively between different antibodies i.e., are "hypervariable." These hypervariable domains contain residues that are directly involved in the binding and specificity of each 55 particular antibody for its specific antigenic determinant. Hypervariability, both in the light chain and the heavy chain variable domains, is concentrated in three segments known as complementarity determining regions (CDRs) or hypervariable loops (HVLs). CDRs are defined by sequence comparison in Kabat et al., 1991, In: Sequences of Proteins of immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md., whereas HVLs (also referred herein as CDRs) are structurally defined according to the three-dimensional structure of the variable domain,

as described by Chothia and Lesk, 1987, J. Mol. Biol. 196: 901-917. These two methods result in slightly different identifications of a CDR. As defined by Kabat, CDR-L1 is positioned at about residues 24-34, CDR-L2, at about residues 50-56, and CDR-L3, at about residues 89-97 in the light chain variable domain; CDR-H1 is positioned at about residues 31-35, CDR-H2 at about residues 50-65, and CDR-H3 at about residues 95-102 in the heavy chain variable domain.

5 The exact residue numbers that encompass a particular CDR will vary depending on the sequence and size of the CDR. Those skilled in the art can routinely determine which residues comprise a particular CDR given the variable region amino acid sequence of the antibody. The CDR1, CDR2, CDR3 of the heavy and light chains therefore define the unique and functional properties specific for a given antibody.

10 [0090] The three CDRs within each of the heavy and light chains are separated by framework regions (FR), which contain sequences that tend to be less variable. From the amino terminus to the carboxy terminus of the heavy and light chain variable domains, the FRs and CDRs are arranged in the order: FR1, CDR1, FR2, CDR2, FR3, CDR3, and FR4. The largely β -sheet configuration of the FRs brings the CDRs within each of the chains into close proximity to each other as well as to the CDRs from the other chain. The resulting conformation contributes to the antigen binding site (see Kabat et al., 1991, NIH Publ. No. 91-3242, Vol. I, pages 647-669), although not all CDR residues are necessarily directly involved in antigen binding.

15 [0091] FR residues and Ig constant domains are not directly involved in antigen binding, but contribute to antigen binding and/or mediate antibody effector function. Some FR residues are thought to have a significant effect on antigen binding in at least three ways: by noncovalently binding directly to an epitope, by interacting with one or more CDR residues, and by affecting the interface between the heavy and light chains. The constant domains are not directly involved in antigen binding but mediate various Ig effector functions, such as participation of the antibody in antibody dependent cellular cytotoxicity (ADCC); complement dependent cytotoxicity (CDC) and antibody dependent cellular phagocytosis (ADCP).

20 [0092] The light chains of vertebrate immunoglobulin are assigned to one of two clearly distinct classes, kappa (κ) and lambda (λ), based on the amino acid sequence of the constant domain. By comparison, the heavy chains of mammalian immunoglobulins are assigned to one of five major classes, according to the sequence of the constant domains: IgA, IgD, IgE, IgG, and IgM. IgG and IgA are further divided into subclasses (isotypes), e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂. The heavy chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of the classes of native immunoglobulins are well known.

25 [0093] The terms, "antibody", "anti-IL-23p19 antibody", "humanized anti-IL-23p19 antibody", "humanized anti-IL-23p19 epitope antibody", and "variant humanized anti-IL-23p19 epitope antibody" specifically encompass monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments such as variable domains and other portions of antibodies that exhibit a desired biological activity, e.g., IL-23p19 binding. The term "monoclonal antibody" (mAb) refers to an antibody that is highly specific, being directed against a single antigenic determinant, an "epitope". Therefore, the modifier "monoclonal" is indicative of antibodies directed to the identical epitope and is not to be construed as requiring production of the antibody by any particular method. It should be understood that monoclonal antibodies can be made by any technique or methodology known in the art; including e.g., the hybridoma method (Kohler et al., 1975, Nature 256:495), or recombinant DNA methods known in the art (see, e.g., U.S. Pat. No. 4,816,567), or methods of isolation of monoclonal recombinantly produced using 30 phage antibody libraries, using techniques described in Clackson et al., 1991, Nature 352: 624-628, and Marks et al., 1991, J. Mol. Biol. 222: 581-597.

35 [0094] The term "monomer" refers to a homogenous form of an antibody. For examples, for a full-length antibody, monomer means a monomeric antibody having two identical heavy chains and two identical light chains.

40 [0095] Chimeric antibodies consist of the heavy and light chain variable regions of an antibody from one species (e.g., a non-human mammal such as a mouse) and the heavy and light chain constant regions of another species (e.g., human) antibody and can be obtained by linking the DNA sequences encoding the variable regions of the antibody from the first species (e.g., mouse) to the DNA sequences for the constant regions of the antibody from the second (e.g. human) species and transforming a host with an expression vector containing the linked sequences to allow it to produce a chimeric antibody. Alternatively, the chimeric antibody also could be one in which one or more regions or domains of the heavy and/or light chain is identical with, homologous to, or a variant of the corresponding sequence in a monoclonal antibody from another immunoglobulin class or isotype, or from a consensus or germline sequence. Chimeric antibodies can include fragments of such antibodies, provided that the antibody fragment exhibits the desired biological activity of its parent antibody, for example binding to the same epitope (see, e.g., U.S. Pat. No. 4,816,567; and Morrison et al., 1984, Proc. Natl. Acad. Sci. USA 81: 6851-6855).

45 [0096] The terms, "antibody fragment", "anti-IL-23p19 antibody fragment", "anti-IL-23p19 epitope antibody fragment", "humanized anti-IL-23p19 antibody fragment", "humanized anti-IL-23p19 epitope antibody fragment", "variant humanized anti-IL-23p19 epitope antibody fragment" refer to a portion of a full length anti-IL-23p19 antibody, in which a variable region or a functional capability is retained, for example, specific IL-23p19 epitope binding. Examples of antibody frag-

ments include, but are not limited to, a Fab, Fab', F(ab')₂, Fd, Fv, scFv and scFv-Fc fragment, a diabody, a linear antibody, a single-chain antibody, a minibody, a diabody formed from antibody fragments, and multispecific antibodies formed from antibody fragments.

[0097] Full length antibodies can be treated with enzymes such as papain or pepsin to generate useful antibody fragments. Papain digestion is used to produce two identical antigen-binding antibody fragments called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment. The Fab fragment also contains the constant domain of the light chain and the C_{H1} domain of the heavy chain. Pepsin treatment yields a F(ab')₂ fragment that has two antigen-binding sites and is still capable of cross-linking antigen.

[0098] Fab' fragments differ from Fab fragments by the presence of additional residues including one or more cysteines from the antibody hinge region at the C-terminus of the C_{H1} domain. F(ab')₂ antibody fragments are pairs of Fab' fragments linked by cysteine residues in the hinge region. Other chemical couplings of antibody fragments are also known.

[0099] "Fv" fragment contains a complete antigen-recognition and binding site consisting of a dimer of one heavy and one light chain variable domain in tight, non-covalent association. In this configuration, the three CDRs of each variable domain interact to define an antigen-biding site on the surface of the V_H-V_L dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody.

[0100] A "single-chain Fv" or "scFv" antibody fragment is a single chain Fv variant comprising the V_H and V_L domains of an antibody where the domains are present in a single polypeptide chain. The single chain Fv is capable of recognizing and binding antigen. The scFv polypeptide may optionally also contain a polypeptide linker positioned between the V_H and V_L domains in order to facilitate formation of a desired three-dimensional structure for antigen binding by the scFv (see, e.g., Pluckthun, 1994, In The Pharmacology of monoclonal Antibodies, Vol. 113, Rosenburg and Moore eds., Springer-Verlag, New York, pp. 269-315).

[0101] A "diabody" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy chain variable domain (V.sub.H) connected to a light chain variable domain (V.sub.L) in the same polypeptide chain (V.sub.H-V.sub.L or V.sub.L-V.sub.H). Diabodies are described more fully in, e.g., Holliger et al. (1993) Proc. Natl. Acad. Sci. USA 90: 6444-6448.

[0102] Other recognized antibody fragments include those that comprise a pair of tandem Fd segments (V_H-C_{H1}-V_H-C_{H1}) to form a pair of antigen binding regions. These "linear antibodies" can be bispecific or monospecific as described in, for example, Zapata et al. 1995, Protein Eng. 8(10):1057-1062.

[0103] A "humanized antibody" or a "humanized antibody fragment" is a specific type of chimeric antibody which includes an immunoglobulin amino acid sequence variant, or fragment thereof, which is capable of binding to a predetermined antigen and which, comprises one or more FRs having substantially the amino acid sequence of a human immunoglobulin and one or more CDRs having substantially the amino acid sequence of a non-human Immunoglobulin. This non-human amino acid sequence often referred to as an "import" sequence is typically taken from an "import" antibody domain, particularly a variable domain. In general, a humanized antibody includes at least the CDRs or HVLS of a non-human antibody, inserted between the FRs of a human heavy or light chain variable domain. The present invention describes specific humanized anti-IL-23p19 antibodies which contain CDRs derived from the mouse monoclonal antibodies or humanized CDRs shown in Tables 3 and 4 inserted between the FRs of human germline sequence heavy and light chain variable domains. It will be understood that certain mouse FR residues may be important to the function of the humanized antibodies and therefore certain of the human germline sequence heavy and light chain variable domains residues are modified to be the same as those of the corresponding mouse sequence.

[0104] In another aspect, a humanized anti-IL-23p19 antibody comprises substantially all of at least one, and typically two, variable domains (such as contained, for example, in Fab, Fab', F(ab')₂, Fabc, and Fv fragments) in which all, or substantially all, of the CDRs correspond to those of a non-human immunoglobulin, and specifically herein, all of the CDRs are mouse or humanized sequences as detailed in Tables 1 through 4 herein below and all, or substantially all, of the FRs are those of a human Immunoglobulin consensus or germline sequence. In another aspect, a humanized anti- IL-23p19 antibody also includes at least a portion of an immunoglobulin Fc region, typically that of a human immunoglobulin. Ordinarily, the antibody will contain both the light chain as well as at least the variable domain of a heavy chain. The antibody also may include one or more of the C_{H1}, hinge, C_{H2}, C_{H3}, and/or C_{H4} regions of the heavy chain, as appropriate.

[0105] A humanized anti-IL-23p19 antibody can be selected from any class of immunoglobulins, including IgM, IgG, IgD, IgA and IgE, and any isotype, including IgG₁, IgG₂, IgG₃, IgG₄, IgA₁ and IgA₂. For example, the constant domain can be a complement fixing constant domain where it is desired that the humanized antibody exhibit cytotoxic activity, and the isotype is typically IgG₁. Where such cytotoxic activity is not desirable, the constant domain may be of another isotype, e.g., IgG₂. An alternative humanized anti-IL-23p19 antibody can comprise sequences from more than one immunoglobulin class or isotype, and selecting particular constant domains to optimize desired effector functions is within the ordinary skill in the art. In specific embodiments, the present invention provides antibodies that are IgG₁ antibodies and more particularly, are IgG₁ antibodies in which there is a knock-out of effector functions.

[0106] The FRs and CDRs, or HVLS, of a humanized anti-IL-23p19 antibody need not correspond precisely to the

parental sequences. For example, one or more residues in the import CDR, or HVL, or the consensus or germline FR sequence may be altered (e.g., mutagenized) by substitution, insertion or deletion such that the resulting amino acid residue is no longer identical to the original residue in the corresponding position in either parental sequence but the antibody nevertheless retains the function of binding to IL-23p19. Such alteration typically will not be extensive and will be conservative alterations. Usually, at least 75% of the humanized antibody residues will correspond to those of the parental consensus or germline FR and import CDR sequences, more often at least 90%, and most frequently greater than 95%, or greater than 98% or greater than 99%.

[0107] Immunoglobulin residues that affect the interface between heavy and light chain variable regions ("the V_L - V_H interface") are those that affect the proximity or orientation of the two chains with respect to one another. Certain residues that may be involved in interchain interactions include V_L residues 34, 36, 38, 44, 46, 87, 89, 91, 96, and 98 and V_H residues 35, 37, 39, 45, 47, 91, 93, 95, 100, and 103 (utilizing the numbering system set forth in Kabat et al., Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md., 1987)). U.S. Pat. No. 6,407,213 also discusses that residues such as V_L residues 43 and 85, and V_H residues 43 and 60 also may be involved in this interaction. While these residues are indicated for human IgG only, they are applicable across species. Important antibody residues that are reasonably expected to be involved in interchain interactions are selected for substitution into the consensus sequence.

[0108] The terms "consensus sequence" and "consensus antibody" refer to an amino acid sequence which comprises the most frequently occurring amino acid residue at each location in all immunoglobulins of any particular class, isotype, or subunit structure, e.g., a human immunoglobulin variable domain. The consensus sequence may be based on immunoglobulins of a particular species or of many species. A "consensus" sequence, structure, or antibody is understood to encompass a consensus human sequence as described in certain embodiments, and to refer to an amino acid sequence which comprises the most frequently occurring amino acid residues at each location in all human immunoglobulins of any particular class, isotype, or subunit structure. Thus, the consensus sequence contains an amino acid sequence having at each position an amino acid that is present in one or more known immunoglobulins, but which may not exactly duplicate the entire amino acid sequence of any single immunoglobulin. The variable region consensus sequence is not obtained from any naturally produced antibody or Immunoglobulin. Kabat et al., 1991, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md., and variants thereof. The FRs of heavy and light chain consensus sequences, and variants thereof, provide useful sequences for the preparation of humanized anti-IL-23p19 antibodies. See, for example, U.S. Pat. Nos. 6,037,454 and 6,054,297.

[0109] Human germline sequences are found naturally in the human population. A combination of those germline genes generates antibody diversity. Germline antibody sequences for the light chain of the antibody come from conserved human germline , kappa or lambda v-genes and j-genes. Similarly the heavy chain sequences come from germline v-, d- and j-genes (LeFranc, M-P, and LeFranc, G, "The Immunoglobulin Facts Book" Academic Press, 2001).

[0110] As used herein, "variant", "anti- IL-23p19 variant", "humanized anti-IL-23p19 variant", or "variant humanized anti- IL-23p19" each refers to a humanized anti-IL-23p19 antibody having at least a light chain variable murine CDR from any of the sequences as shown in Table 1 or a heavy chain murine CDR sequence derived from the murine monoclonal antibody as shown in Table 2. Variants include those having one or more amino acid changes in one or both light chain or heavy chain variable domains, provided that the amino acid change does not substantially impair binding of the antibody to IL-23p19. Exemplary humanized antibodies produced herein include those designated as Antibody A, Antibody B, Antibody C and Antibody D, and the various light chains and heavy chains of the same are shown in SEQ ID Nos:174 and 180, and SEQ ID Nos:176 and 178, respectively.

[0111] An "isolated" antibody is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of the antibody's natural environment are those materials that may interfere with diagnostic or therapeutic uses of the antibody, and can be enzymes, hormones, or other proteinaceous or nonproteinaceous solutes. In one aspect, the antibody will be purified to at least greater than 95% isolation by weight of antibody.

[0112] An isolated antibody includes an antibody in situ within recombinant cells in which it is produced, since at least one component of the antibody's natural environment will not be present. Ordinarily however, an isolated antibody will be prepared by at least one purification step in which the recombinant cellular material is removed.

[0113] The term "antibody performance" refers to factors that contribute to antibody recognition of antigen or the effectiveness of an antibody in vivo. Changes in the amino acid sequence of an antibody can affect antibody properties such as folding, and can influence physical factors such as initial rate of antibody binding to antigen (k_a), dissociation constant of the antibody from antigen (k_d), affinity constant of the antibody for the antigen (K_d), conformation of the antibody, protein stability, and half life of the antibody.

[0114] The term "epitope tagged" when used herein, refers to an anti-IL-23p19 antibody fused to an "epitope tag". An "epitope tag" is a polypeptide having a sufficient number of amino acids to provide an epitope for antibody production, yet is designed such that it does not interfere with the desired activity of the humanized anti-IL-23p19 antibody. The epitope tag is usually sufficiently unique such that an antibody raised against the epitope tag does not substantially

cross-react with other epitopes. Suitable tag polypeptides generally contain at least 6 amino acid residues and usually contain about 8 to 50 amino acid residues, or about 9 to 30 residues. Examples of epitope tags and the antibody that binds the epitope include the flu HA tag polypeptide and its antibody 12CA5 (Field et al., 1988 Mol. Cell. Biol. 8: 2159-2165; c-myc tag and 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto (Evan et al., 1985, Mol. Cell. Biol. 5(12):3610-3616; and Herpes simplex virus glycoprotein D (gD) tag and its antibody (Paborsky et al. 1990, Protein Engineering 3(6): 547-553). In certain embodiments, the epitope tag is a "salvage receptor binding epitope". As used herein, the term "salvage receptor binding epitopes" refers to an epitope of the Fc region of an IgG molecule (such as IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the in vivo serum half-life of the IgG molecule.

[0115] In some embodiments, the antibodies of the present invention may be conjugated to a cytotoxic agent. This is any substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (such as I¹³¹, I¹²⁵, Y⁹⁰, and Re¹⁸⁶), chemotherapeutic agents, and toxins such as enzymatically active toxins of bacterial, fungal, plant, or animal origin, and fragments thereof. Such cytotoxic agents can be coupled to the humanized antibodies of the present invention using standard procedures, and used, for example, to treat a patient indicated for therapy with the antibody.

[0116] A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. There are numerous examples of chemotherapeutic agents that could be conjugated with the therapeutic antibodies of the present invention. Examples of such chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan, and piposulfan; aziridines such as benzodopa, carboquone, metutedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide, and trimethylololomelamine; acetogenins (especially bullatacin and bullatacinone); camptothecin (including the synthetic analogue topotecan); bryostatin; calystatin; CC-1065 (including its adozelesin, carzelesin, and bizelesin synthetic analogues); cryptophycines (particularly" cryptophycin 1 and cryptophycin 8); dolastatin, auristatins, (including analogues monomethyl-auristatin E and monomethyl-auristatin F); duocarmycin (including the synthetic analogues, KW-2189 and CBI-TMI); eleutherobin; pancratistatin; sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlomaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine; trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as the enediyne antibiotics (e.g., calicheamicin, especially calicheamicin gamma1I and calicheamicin phi1, see for example, Agnew, Chem. Intl. Ed. Engl., 33:183-186; dynemicin, including dynemicin A; bisphosphonates, such as clodronate; esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromomophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabicin, caminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin (Adriamycin™) (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin, and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycine, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ube nimex, zinostatin, zorubicin; anti-metabolites such a methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thioguanine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauroidine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitiostanol, mepitiostane, testolactone; anti-adranals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as folinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestrebutil; bisantrene; edatraxate; defofamine; democlocine; diaziquone; elfomithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; Ioniadamine; maytansinoids such as maytansine and ansamitocins; mitoguazone, mitoxanthrone; moperidol; nitracrine; pentostatin; phenacetin; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK®; razoxane; rhizoxin; sizofuran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomyctine; mitabronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, e.g., paclitaxel (TAXOL®, Bristol-Myers Squibb Oncology, Princeton, N.J.) and doxetaxel (TAXOTERE®, Rhone-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine (Gemzar™); 6-thioguanine; mercaptapurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; vinorelbine Navelbine™); novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine; and pharmaceutically acceptable salts, acids, or derivatives of any of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including Nolvadex™), raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene; keoxifene, LY117018, onapristone, and toremifene (Fareston™); aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, megestrol acetate (Megace™), exemestane, formestane, fadrozole, vorozole (Riv-

isor™), letrozole (Femara™), and anastrozole (Arimidex™); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprorelin, and goserelin; and pharmaceutically acceptable salts, acids, or derivatives of any of the above. Any one or more of these agents may be conjugated to the humanized antibodies of the present invention to provide a useful therapeutic agent for the treatment of various disorders.

5 [0117] The antibodies also may be conjugated to prodrugs. A "prodrug" is a precursor or derivative form of a pharmaceutically active substance that is less cytotoxic to tumor cells compared to the parent drug and is capable of being enzymatically activated or converted into the more active form. See, for example, Wilman, 1986, "Prodrugs in Cancer Chemotherapy", In Biochemical Society Transactions, 14, pp. 375-382, 615th Meeting Belfast and Stella et al., 1985, "Prodrugs: A Chemical Approach to Targeted Drug Delivery, Borchardt et al., (ed.), pp. 247-267, Humana Press. Useful prodrugs include, but are not limited to, phosphate-containing prodrugs, thiophosphate-containing prodrugs, sulfate-containing prodrugs peptide-containing prodrugs, D-amino acid-modified prodrugs, glycosylated prodrugs, β-lactam-containing prodrugs, optionally substituted phenoxyacetamide-containing prodrugs, and optionally substituted phenylacetamide-containing prodrugs, 5-fluorocytosine and other 5-fluorouridine prodrugs that can be converted into the more active cytotoxic free drug. Examples of cytotoxic drugs that can be derivatized into a prodrug form include, but are not limited to, those chemotherapeutic agents described above.

10 [0118] For diagnostic as well as therapeutic monitoring purposes, the antibodies of the invention also may be conjugated to a label, either a label alone or a label and an additional second agent (prodrug, chemotherapeutic agent and the like). A label, as distinguished from the other second agents refers to an agent that is a detectable compound or composition and it may be conjugated directly or indirectly to a humanized antibody of the present invention. The label may itself be 15 detectable (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition that is detectable. Labeled humanized anti-IL-23p19 antibody can be prepared and used in various applications including in vitro and in vivo diagnostics.

20 [0119] The antibodies of the present invention may be formulated as part of a liposomal preparation in order to affect delivery thereof in vivo. A "liposome" is a small vesicle composed of various types of lipids, phospholipids, and/or 25 surfactant. Liposomes are useful for delivery to a mammal of a compound or formulation, such as a humanized anti-IL-23p19 antibody disclosed herein, optionally, coupled to or in combination with one or more pharmaceutically active agents and/or labels. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

30 [0120] Certain aspects of the present invention related to isolated nucleic acids that encode one or more domains of the humanized antibodies of the present invention. An "isolated" nucleic acid molecule is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the antibody nucleic acid. An isolated nucleic acid molecule is distinguished from the nucleic acid molecule as it exists in natural cells.

35 [0121] In various aspects of the present disclosure one or more domains of the humanized antibodies will be recombinantly expressed. Such recombinant expression may employ one or more control sequences, i.e., polynucleotide sequences necessary for expression of an operably linked coding sequence in a particular host organism. The control sequences suitable for use in prokaryotic cells include, for example, promoter, operator, and ribosome binding site sequences. Eukaryotic control sequences include, but are not limited to, promoters, polyadenylation signals, and enhancers. These control sequences can be utilized for expression and production of humanized anti-IL-23p19 antibody 40 in prokaryotic and eukaryotic host cells.

45 [0122] A nucleic acid sequence is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, a nucleic acid presequence or secretory leader is operably linked to a nucleic acid encoding a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. However, enhancers are optionally contiguous. Linking can be accomplished by ligation at convenient restriction sites. If such sites do not exist, synthetic oligonucleotide adaptors or linkers can be used.

50 [0123] As used herein, the expressions "cell", "cell line", and "cell culture" are used interchangeably and all such designations include the progeny thereof. Thus, "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers.

55 [0124] The term "mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domesticated and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, and the like. Preferably, the mammal is human.

[0125] A "disorder", as used herein, is any condition that would benefit from treatment with a humanized anti-IL-23p19 antibody described herein. This includes chronic and acute disorders or diseases including those pathological conditions that predispose the mammal to the disorder in question. Non-limiting examples of disorders to be treated herein include inflammatory, angiogenic, autoimmune and immunologic disorders, respiratory disorders, cancer, hematological malig-

nancies, benign and malignant tumors, leukemias and lymphoid malignancies.

[0126] The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include, but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia.

5 [0127] As used herein, the term "IL-23-associated disorder" or "IL-23-associated disease" refers to a condition in which IL-23 activity contributes to the disease and typically where IL-23 is abnormally expressed. An IL-23-associated disorder includes diseases and disorders of the immune system, such as autoimmune disorders and inflammatory disorders. Such conditions include, but are not limited to, rheumatoid arthritis (RA), systemic lupus erythematosus (SLE), scleroderma, Sjogren's syndrome, multiple sclerosis, psoriasis, psoriatic arthritis, inflammatory bowel disease (e.g., ulcerative colitis and Crohn's disease), pulmonary inflammation, asthma, idiopathic thrombocytopenic purpura (ITP) and ankylosing spondylitis;

10 [0128] The term "intravenous infusion" refers to introduction of an agent into the vein of an animal or human patient over a period of time greater than approximately 15 minutes, generally between approximately 30 to 90 minutes.

15 [0129] The term "intravenous bolus" or "intravenous push" refers to drug administration into a vein of an animal or human such that the body receives the drug in approximately 15 minutes or less, generally 5 minutes or less.

[0130] The term "subcutaneous administration" refers to introduction of an agent under the skin of an animal or human patient, preferable within a pocket between the skin and underlying tissue, by relatively slow, sustained delivery from a drug receptacle. Pinching or drawing the skin up and away from underlying tissue may create the pocket.

20 [0131] The term "subcutaneous infusion" refers to introduction of a drug under the skin of an animal or human patient, preferably within a pocket between the skin and underlying tissue, by relatively slow, sustained delivery from a drug receptacle for a period of time including, but not limited to, 30 minutes or less, or 90 minutes or less. Optionally, the infusion may be made by subcutaneous implantation of a drug delivery pump implanted under the skin of the animal or human patient, wherein the pump delivers a predetermined amount of drug for a predetermined period of time, such as 30 minutes, 90 minutes, or a time period spanning the length of the treatment regimen.

25 [0132] The term "subcutaneous bolus" refers to drug administration beneath the skin of an animal or human patient, where bolus drug delivery is less than approximately 15 minutes; in another aspect, less than 5 minutes, and in still another aspect, less than 60 seconds. In yet even another aspect, administration is within a pocket between the skin and underlying tissue, where the pocket may be created by pinching or drawing the skin up and away from underlying tissue.

30 [0133] The term "therapeutically effective amount" is used to refer to an amount of an active agent that relieves or ameliorates one or more of the symptoms of the disorder being treated. In another aspect, the therapeutically effective amount refers to a target serum concentration that has been shown to be effective in, for example, slowing disease progression. Efficacy can be measured in conventional ways, depending on the condition to be treated.

35 [0134] The terms "treatment" and "therapy" and the like, as used herein, are meant to include therapeutic as well as prophylactic, or suppressive measures for a disease or disorder leading to any clinically, desirable or beneficial effect, including but not limited to alleviation, or relief of one or more symptoms, regression, slowing or cessation of progression of the disease or disorder. Thus, for example, the term treatment includes the administration of an agent prior to or following the onset of a symptom of a disease or disorder thereby preventing or removing one or more signs of the disease or disorder. As another example, the term includes the administration of an agent after clinical manifestation of 40 the disease to combat the symptoms of the disease. Further, administration of an agent after onset and after clinical symptoms have developed where administration affects clinical parameters of the disease or disorder, such as the degree of tissue injury or the amount or extent of metastasis, whether or not the treatment leads to amelioration of the disease, comprises "treatment" or "therapy" as used herein. Moreover, as long as the compositions of the invention either alone or in combination with another therapeutic agent alleviate or ameliorate at least one symptom of a disorder 45 being treated as compared to that symptom in the absence of use of the humanized anti-IL-23p19 antibody composition, the result should be considered an effective treatment of the underlying disorder regardless of whether all the symptoms of the disorder are alleviated or not.

50 [0135] The term "package insert" is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, administration, contraindications and/or warnings concerning the use of such therapeutic products.

Antibodies

55 [0136] In one aspect, described and disclosed herein are anti-IL-23 antibodies, in particular humanized anti-IL-23p19 antibodies, and compositions and articles of manufacture comprising one or more anti-IL-23 antibody, in particular one or more humanized anti-IL-23p19 antibody of the present invention. Also described are binding agents that include an antigen-binding fragment of an anti-IL-23 antibody, in particular a humanized anti-IL-23p19 antibody. The humanized anti-IL-23p19 antibodies and binding agents can inhibit the production of Th17 associated cytokines, which contribute

to chronic autoimmune and inflammatory diseases. The humanized anti-IL-23p19 antibodies and binding agents can thus be used in the treatment of a variety of diseases or disorders. A humanized anti-IL-23p19 antibody and an IL-23p19 binding agent each includes at least a portion that specifically recognizes an IL-23p19 epitope (i.e., an antigen-binding fragment).

5 [0137] In the initial characterization mouse antibodies were selected based on IL-23p19 binding characterization.

[0138] Accordingly in one aspect, an antibody of the present disclosure has a K_D for IL-23, in particular human IL-23, of less than 100pM. In another aspect, an antibody of the present disclosure has a K_D of less than 40pM. In another aspect, an antibody of the present disclosure has a K_D of less than 20pM. In another aspect, an antibody of the present disclosure has a K_D of less than 10pm. In another aspect, a monoclonal antibody of the present disclosure has a K_D of less than 1pM.

10 [0139] The selected mouse antibodies have the following light chain variable regions and heavy chain variable regions as shown in Table 1 and 2:

Table 1: Anti-IL-23p19 Mouse Leads - VK Sequences

15 2D1vk	GACATTGTGCTGACCCAATCTCCAGGTTCTTGGCTGTGTCTCTAGGGCAG AGGGCCACCATACTGCAGAACAGTGAAAGTGTATAGTTATGCCAA AATTTATACACTGGTACCCAGCAGAACCCAGGACAGCCACCCAAACTCCTC ATCTATCGTGCATCCAACCTGGAATCTGGATCCCTGCCAGGTTCACTGGC AGTGGGTCTAGGACAGACTTCACCCTCACCATGAATCCTGTGGAGGCTGAT GATGTTGCAACCTATTACTGTCACTGCAAACTAATGAGGATCCGTACACGTT GGAGGGGGGACCAAGCTGGAAATAAGA (SEQ ID NO:83)
20	DIVLTQSPGLAVSLGQRATISCRTSESVYSYQNFIIHWYQQKPGQPPKLL IYRASNLESGIPARFSGGSRTDFLTLMNPVEADDVATYYCQQTNEDPYTF GGGTKLEIR (SEQ ID NO:84)
25 30 6B8VK	GACATTGTGATGACCCAGTCTCACAAATTCTTGTCCACATCAGTGGGAGAC AGGGTCACCATACTTGCAAGGCCAGTCGGATGTGGCTATTGCTGTAGCC TGGTATCAACAGAACCCAGGGCAATCTCCTAAACTACTTCTTTCTGGGCA TCCACCCGACACACTGGGTCCCTGATCGCTTCACAGGCAGTGGATCTCGG ACAGATTCACTCTCACCAATTAGCAATGTGCAGTCTGAAGACTTGGCAGAT TATTCTGTACCAATATAGCAGCTATCCATTACGTTCTGGCTCGGGACA AAGTTGGAAATAAG (SEQ ID NO:85)
35 40 45 50 55 9D12-Vk	DIVMTQSHKELSTSVGDRVITCKASRDVAIAVAWYQQKPGQSPKLLLFWA STRHTGVPDFRTGSGSRTDFLTLSNVQSEDLADYFCHQYSSYPFTFGSGT KLEIK (SEQ ID NO:86)
55 9D12-Vk	GACATTGCGCTGACCCAATCTCCAGCTTCTTGGCTGTGTCTCTGGGGCAG AGGGCCACCATACTGCAGAGCCAGTGAAACTATTAAATTCTATGGCACT AGTTTATGCACTGGTACCCAGCAGAACCCAGGACAGTCACCCAAACTCCTC ATCTATCGTGCATCCAACCTAGAATCTGGATCCCTGCCAGGTTCACTGGC AGTGGGTCTAGGACAGACTTCACCCTCACCATTAATCCTGTGGAGGCTGAT GATGTTGCAACCTATTACTGTCACTGCAAACTAATGAGGATCCGTACACGTT GGAGGGGGGACTAAGTTGGAAATAAAA (SEQ ID NO:87)

(continued)

	DIALTQSPASLAVSLGQRATISCRASETINFYGTSMHWYQQKPGQSPKLL IYRASNLESGIPARFSGSGSRDFTLTINPVEADDVATYYCQQTNEDPYTF GGGKLEIK (SEQ ID NO:88)
5	
10	15C11vk GATGTTGTGATGACCCAACTCCACTCTCCCTGCCTGTCAGTCTGGAGAT CAAGCCTCCATCTCTTGCGAGATCTAGTCAGAGCCTTGTACACAGTAATGGA AACACCTATTACATTGGTACCTGCAGAAGCCAGGCCAGTCTCCAAAGCTC CTGATCTACAAAGTTCCAACCGATTTCTGGGTCCCAGACAGGTTCACT
15	GGCAGTGGATCAGGGACAGATTCAACACTCAAGATCAACAGAGTGGAGGCT GAGGATCTGGGAGTTATTCTGCTCTCAAAGTACACATGTTCCGTACACG TTCGGAGGGGGACCCAGCTGGAAATAAAA (SEQ ID NO:89)
20	DVVMTQTPLSLPVSLGDQASISCRSSQSLVHSNGNTYLHWYLQKPGQSPKL LIYKVSNRFSGVPDRFSGSGSDFTLKIIRVEAEDLGVYFCQSOTHVPYT FGGGTQLEIK (SEQ ID NO:90)
25	15F1vk DIVMTQSPATLSVTPGDRVSLSCRASQSIISDYLHWYQQRSHESPRLLIKYA SQSISGIPSRTGSGSGSDFTLTINSVEPEDVGVYYCQNGHSFPFTFGSGT KLEIK (SEQ ID NO:91)
30	
35	18D3vk GACATTGTGATGACTCAGTCTCCAGCCACCCCTGTCAGTCCAGGAGAT AGAGTCTCTCTTCCCTGCAGGCCAGTCAGAGTATTAGCGACTACTTATAC TGGTATCAACAAAAATCACATGAGTCTCAAGGCTTCTCATCAAATTGCT TCCCAATCCATCTGGGATCCCTCCAGGTTCACTGGCAGTGGATCAGGG TCAGATTCACTCTCAGTATCGACAGTGTGGAACCTGATGATGTTGGAGTC TTTTCTGTCAAAATGGTCACAGCTTCCGTTACGTTGGAGGGGGACC AAGCTGGAAATAAAA (SEQ ID NO:92)
40	DIVMTQSPATLSVTPGDRVSLSCRASQSIISDYLYWYQQKSHESPRLLIKFA SQSISGIPSRTGSGSGSDFTLSIDSVEPDDVGVFFCQNGHSFPFTFGGGT KLEIK (SEQ ID NO:93)
45	18C4vk GACATTGTGATGACTCAGTCTCCAGCCACCCCTGTCAGTCCAGGAGAT AGAGTCTCTCTTCCCTGCAGGCCAGCCAGAGTATTAGCGAGTACTTACAC TGGTATCAACAAAAATCACATGAGTCTCAAGGCTTCTCATCAAATATGCT TCCCAATCCATCTGGGATCCCTCCAGGTTCACTGGCAGTGGATCAGGG TCAGATTCACTCTCAGTATCACAGTGTGGAACCTGAAAGATGTTGGAGTG TATTACTGTCAAAATGGTCACAGCTTCCATTACGTTGGCTCGGGACAC AAGTTGGAAATAAAA (SEQ ID NO:94)
50	
55	

(continued)

	DIVMTQSPATLSVTPGDRVSLSCRASQSISEYLNHWYQQKSHESPRLLIKYA SQSISGIPSRFSGSGSDFTLSINSVEPEDVGVYYCQNGHSFPFTFGSGT KLEIK (SEQ ID NO:95)
18E5vk	GACATTGTGATGACTCAGTCTCCAGCCACCCTGTCTGTGACTCCAGGAGAT AGAGTCTCTCTTCCTGCAGGCCAGCCAGAGTATTAGCGACTACTTACAC TGGTATCAACAAAAATCACATGAGTCTCCAAGGCTTCTCATCAAATTGCT TCCCAATCCATCTGGATCCCCTCCAGGTTCACTGGCAGTGGATCAGGG TCAGATTCACTCTCAGTATCGACAGTGTGGAACCTGATGATGTTGGAGTC TTTCTGTCAAAATGGTCACAGCTTCCGTTACGTTGGAGGGGGACC AAGCTGGAAATAAAA (SEQ ID NO:96)
	DIVMTQSPATLSVTPGDRVSLSCRASQSISDYLYWYQQKSHESPRLLIKFA SQSISGIPSRTGSGSDFTLSIDSVEPDVGVFFCQNGHSFPFTFGGGT KLEIK (SEQ ID NO:97)
20E8vk	GACATTGTGATGACTCAGTCTCCAGCCACCCTGTCTGTGACTCCAGGAGAT AGAGTCTCTCTTCCTGCAGGCCAGCCAGAGTATTAGCGAGTATTACAC TGGTATCAACAAAAATCACATGAGTCTCCAAGGCTTCTCATCAAATATGCT TCCCAATCCATCTGGATCCCCTCCAGGTTCACTGGCAGTGGATCAGGG TCAGATTCACTCTCAGTATCACAGTGTGGAACCTGAAGATGTTGGAGTT TATTACTGTCAAAATGGTCACAGCTTCCATTACGTTGGCTCGGGACA AAGTTGGAAATAAAA (SEQ ID NO:98)
	DIVMTQSPATLSVTPGDRVSLSCRASQSISEYLNHWYQQKSHESPRLLIKYA SQSISGIPSRFSGSGSDFTLSINSVEPEDVGVYYCQNGHSFPFTFGSGT KLEIK (SEQ ID NO:99)
22E2vk	GACATTGTGATGACTCAGTCTCCAGCCACCCTGTCTGTGACTCCAGGAGAT AGAGTCTCTCTCCCTGCAGGCCAGCCAGAGTATTAGCGTCTACTTACAC TGGTATCAACAAAAATCACCTGAGTCTCCAAGGCTTCTCATCAAATATGCT TCCCAATCCATCTGGATCCCCTCCAGGTTCACTGGCAGTGGATCAGGG TCAGATTCACTCTCAGTATCACAGTGTGGAACCTGAAGATGTTGGAGTT TATTACTGTCAAAATGGTCACAGCTTCCATTACGTTGGCTCGGGACA AAGTTGGAAATAAAA (SEQ ID NO:100)
	DIVMTQSPATLSVTPGDRVSLSCRASQSISVYLHWYQQKSPESPRLLIKYA SQSISGIPSRFSGSGSDFTLSINSVEPEDVGVYYCQNGHSFPFTFGSGT KLEIK (SEQ ID NO:101)

(continued)

5	24A54vk	GACATTGTGATGACTCAGTCTCCAGCCACCCTGTCAGTGACTCCAGGAAAT AGAGTCTCTCTTCCTGCAGGCCAGCCAGAGTATTAGCGACTACTACAC TGGTATCAACAAAAATCACATGAGTCTCCAAGGCTCTCATCAAATATGCT TCCCAATCCATCTCTGGATCCCCTCCAGGTTAGTGGCAGTGGATCAGGG TCAAATTCACTCTCAGTATCAACAGTGTGGAACCTGAAGATGTGGAGTG TATTATTGTCAAAATGGTCACAGCTTCCATTACGTTGGCTCGGGACAA AAGTTGGAAATAAAA (SEQ ID NO:102)
10		DIVMTQSPATLSVTPGNRVSLSCRASQSISDYLHWYQQKSHESPRLLIKYA SQSISGIPSRSFGSGGSNFTLSINSVEPEDVGVYYCQNGHSFPFTFGSGT KLEIK (SEQ ID NO:103)
15	26F7V^k	GACATTGTGCTGACACAGTCTCCTGCTTCCTTAGCTGTTCTCTGGGGCAG AGGGCCACCATCTCATGCAGGCCAGCAAAAGTGTAGATTCTCTGACTAT TTTTATATGCACTGGTACCAACAGAAACCAGGACAGCCACCCAAACTCCTC ATCTACCTTGCATCCAACCTAGAATCTGGGGTCCCTGCCAGGTTAGTGGC
20		AGTGGGTCTGGGACAGACTTCACCCCTAACATCCATCCTGTGGAGGAGGAG GATGCTGCAACCTATTACTGTCAAGACAGTAGGGAGCTCCGTACACGTT GGAGGGGGGACCAAGCTGGAGATAAAA (SEQ ID NO:104)
25		DIVLTQSPASLAVSLGQRATISCRASKSVRFSDYFYMHWYQQKPGQPPKLL IYLASNLESVPARFSGSGSGTDFTLNIHPVEEDAATYYCQNSREL PYTF GGGTKLEIK (SEQ ID NO:105)
30	27G8v^k	GACATTGTGTTGACACAGTCTCCTGCTTCCTTAGCTGTATCTCTGGGGCAG AGGGCCACCATCTCATGCAGGCCAGCAAAAGTGTAGTACATCTGGCTAT AGTTATATACACTGGTACCAACAGAAACCAGGACAGCCACCCAAATTCTC ATCTATCTTGCATCCAACCTAGATTCTGGGGTCCCTGCCAGGTTAGTGGC AGTGGGTCTGGGACAGACTTCACCCCTAACATCCATCCTGTGGAGGAGGAG GATGCTGCAACCTATTACTGTCAAGACAGTAGGGAGCTCCGTACACGTT GGAGGGGGGACCAAGCTGGAAATAAAA (SEQ ID NO:106)
35		DIVLTQSPASLAVSLGQRATISCRASKSVSTSGYSYIHWYQQKPGQPPKFL IYLASNLDGVPARFSGSGSGTDFTLNIHPVEEDAATYYCQHSREL PYTF GGGTKLEIK (SEQ ID NO:107)
40		
45		

(continued)

5	31H9vk	GACATTGTGATGACTCAGTCTCCAGCCACCCTGTCTGTGACTCCAGGAGAT AGAGTCTCTCTTCCTGCAGGCCAGCCAGAGTATTAGCGACTACTAACAC TGGTATCAACAAAAATCACATGAGTCTCCAAGGCTCTCATCAAATATGCT TCCCAATCCATCTCTGGATCCCCTCCAGGTTAGTGGCAGTGGATCAGGG TCAGATTTCACCTCTCAGTATCAACAGTGTGAAACCTGAAGATGTGGAGTG TATTACTGTCAAAATGGTCACAGCTTCCGTACACGTTGGAGGGGGGACC AAGCTGGAAATAAAA (SEQ ID NO:108)
10		DIVMTQSPATLSVTPGDRVSLSCRASQSISDYLHWYQQKSHESPRLLIKYA SQSISGIPSRFSGSGSDFTLSINSVEPEDVGVYYCQNGHSFPYTFGGGT KLEIK (SEQ ID NO:109)
15	34G3V_k	GATGTTGTGATGACCCAAACTCCACTCTCCCTGCCTGTCAGTCTGGAGAT CAAGCCTCCATCTCTTGCAGATCTAGTCAGAGCCTTGTACACAGTAATGGA AACACCTATTACATTGGTACCTGCAGAAGCCAGGCCAGTCTCCAAAGCTC CTGATCTACAAAGTTCCAACCGATTTCTGGGTCCCCGACAGGTTAGT GGCAGTGGATCAGGGACAGATTACACTCAAGATCAGCAGAGTGGAGGCT GAGGATCTGGGAGTTATTCCTGCTCTCAAAGTACACATGTTCCGTACACG TTCGGAGGGGGGACCAAGCTGGAAATAAT (SEQ ID NO:110)
20		DVVMTQTPLSLPVSLGDQASISCRSSQSLVHSNGNTYLHWYLQKPGQSPKL LIYKVSNRFSGVPDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQSTHVPYT FGGGTKLEIN (SEQ ID NO:111)
25	34D9V_k	GACATTATGATGACCCAGTCTCACAAATTGTCACATGTCACATCAGTAGGAGAC AGGGTCAGCATCACCTGCAAGGCCAGTCAGGATGTGGTAATGCTGTGGTC TGGTATCAACAAAAACCAGGGCAATCTCCTAAACTACTGATTTACTGGGCA TCCACCCGGCACATTGGAGTCCCTGATCGCTTCACAGGCAGTGGATCTGGG ACAGATTTCACTCTCACCATTACCAATGTGCAGTCTGAAGACTTGGCAGAT TATTCTGTAGCAATATAGCAGCTATCTCACGTTGGCTGGGACCAAG CTGGAGCTGAAA (SEQ ID NO:112)
30		DIMMTQSHKFMSTSVDRLVSITCKASQDVGNAVVWYQQKPGQSPKLLIYWA STRHIGVPDRFTGSGSGTDFTLTIITNVQSEDLADYFCQQYSSYLTFGAGTK LELK (SEQ ID NO:113)
35	34D9V_k	
40		
45		

(continued)

5	43F5vk	GATGTTGTGATGACCAATCTCCACTCTCCCTGCCTGTCAGTCTGGAGAT CAAGCCTCCATCTCTTGAGATCTAGTCAGAGCCTGTACACAGTAATGGA AACACCTATCTACATTGGTACCTGCTGAAGCCAGGCCAGTCTCAAAGCTC CTGATCTACAAAGTTCCAACCGATTTCTGGGTCCCAGACAGGTTAGT GGCAGTGGATCAGGGACAGATTACACTAAGATCAGCAGAGTGGAGGCT GAGGATCTGGGAGTTATTCTGCTCTCAAAGTACACATGTTCCGTACACG TTCGGAGGGGGGACCAAGCTGGAAATAAAA (SEQ ID NO:114)
10		DVVMTQSPLSLPVSLGDQASISCRSSQSLVHSNGNTYLHWYLLKPGQSPKL LIYKVSNRFSGVPDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQSTHVPYT FGGGTKLEIK (SEQ ID NO:115)
15	73H10Vk	GACATCCAGATGACTCAGTCTCCAGTTTCTGTCTGCATCTGTGGAGAA ACTGTCACCACATGTCGAGCAAGTGAGAATATTGACAGTTATTAGCA TGGTATCAGCAGAACAGGGAAAATCTCCTCAGCTCCTGGTCTTGCTGCA CGAAACTTAGCAGATGGTGTGCCATCAAGGTTAGTGGCAGTGGATCAGGC ACACAGTATTCTCTCAAGATCAACAGAACATGCACTGAAGATGTTGCGAGA TACTACTGTCAACATTATTATAGTACTCCATTCACGTTGGCTCGGGACA AAGTTGGAAATAGAA (SEQ ID NO:116)
20		DIQMTQSPVFLSASVGETVTITCRASENIDSYLAWYQQKQGKSPQLLVFAA RNLADGVPSRFSGSGSGTQYSLKINRMQSEDVARYYCQHYYSTPFTFGSGT KLEIE (SEQ ID NO:117)
25	74H3Vk	GACATCCAGATGACTCAGTCGCCAGCTCCCTGTCTGCATCTGTGGAGAA ACTGTCATCTCACATGTCGAGCAAGTGAGAATATTGACAGTTATTAGCA TGGTATCAGCAGAACAGGGAAAATCTCCTCAGCTCCTGGTCTATGCTGCA ACAAACTTAGCAGATGGTGTGCCATCAAGGTTAGTGGCAGTGGATCAGGC ACACAGTATTCTCTCAAGATCAACAGCCTGCAGTCTGAAGATGTTGCGAGA TATTACTGTCTACATTATTATAGTACTCCATTCACGTTGGCTCGGGACA GAGTTGGAAATAAAA (SEQ ID NO:118)
30		DIQMTQSPASLSASVGETVIFTCRASENIDSYLAWYQQKQGKSPQLLVYAA TNLADGVPSRFSGSGSGTQYSLKINSLOSEDVARYYCLHYYSTPFTFGSGT ELEIK (SEQ ID NO:119)
35		
40		
45		

50

55

Table 2: Anti-IL-23p19 Mouse Leads - VH Sequences

	CAGGTGCAGCTGAAGGAGTCAGGACCTGGCCTGGTGGGCCCTCACAGAGC CTGTCATCACATGCACTGTCTCTGGTTCTCATTAACCACCTATGCTATA AGCTGGTTGCCAGTCACCAGGAAGGGCTGGAGTGGCTGGAGTCATA TGGACTGGTGGAGGCACAAAATATAATTCAAGCTCTCAAATCCAGACTGAGC ATCAGCAAAGACAACCTCAAGAGTCAAGTTCTAAAAATGAACAGTCTG CAAACGTGATGACACAGCCAGGTACTACTGTGCCAGAAAGGACTATAATTAC GGGGTGCTATGGACTACTGGGTCAAGGAACCTCAGTCACCGTCTCCTCA (SEQ ID NO:120)
5	QVQLKESGPGLVAPSQSLISITVSGFSLTTYAISWVRQSPGKGLEWLGV WTGGGTYNSALKSRLSISKDN SKSQVFLKMNSLQTDDTARYYCARKDNY GGAMDYWGQGTSVTVSS (SEQ ID NO:121)
10	
15	
20	CAGGTTCAGCTGCAACAGTCTGACGCTGAGTTGGTAAACCTGGCACTTCA GTGAAGACATCCTGCAAAATTCTGGCAACACCTTCACTGACCAAACATT CACTGGATGAAGCAGAGGCCTGAACAGGGCTGGAATGGATTGGATATATT TATCCTAGAGATGATAGTCTTAAGTACAATGAGAACTTCAAGGGCAAGGCC ACATTGACTGCAGACAAATCCTCCAGCACAGCCTACATGCAGCTAACAGT CTGACATCTGAGGA C TGCAGTCTATTCTGTCAATCCCAGACAGGTCA GGCTACGCC TGGTTATTACTGGGCCAAGGGACTCTGGTCACTGTCTCT TCA (SEQ ID NO:122)
25	
30	
35	QVQLQQSDAELVKPGTSVKTSCKISGNFTDQTIHWMKQRPEQGLEWIGYI YPRDDSPKYNENFKGKATLTADKSSSTAYMQLNSLTSEDSAVYFCAI PDRS GYAWFIYWGQGTLTVSS (SEQ ID NO:123)
40	
45	
50	CAGGTGCAGCTGAAGGAGTCAGGACCTGTCTGGTGGGCCCTCACAGAGC CTGTCATCACATGCACTGTCTCTGGTTCTCATTAACCAACTTGTATA AGTTGGTTCGTCAGCCACCAGGAAGGGCTGGAGTGGCTGGAGCAATA TGGACTGGTGGAGGCACAAATTATAATTCAAGCTCTCAAATCCAGACTGAGC ATCAGCAAAGACAACCTCAAGAGTCAAGTTCTAAAAATGAACAGTCTG CAAACGTGATGACACAGCCAGGTATTATTGTGTCAAGAAAGGACTATAGTTAC GGGGTGCTATGGACTACTGGGTCAAGGAACCTCAGTCACCGTCTCCTCA (SEQ ID NO:124)
55	QVQLKESGPVLVAPSQSLISITVSGFSLNNFAISWVRQPPGKGLELGAI WTGGGTYNSALKSRLSISKDN SKSQVFLKMNSLQTDDTARYYCVRKDYSY GGAMDYWGQGTSVTVSS (SEQ ID NO:125)

(continued)

	15C11vh	GAGGTCCAGCTGCAACAGTCTGGACCTGTGCTGGTAAGCCTGGGCTTCA GTGAAGATGTCCTGTAAGGCTCTGGATAACACATTCACTGACTACTATAG AACTGGGTGAAGCAGAGCCATGGAAAGAGCCTTGAGTGGATTGGAGTTATT ATTCCTAACACGGTGGTACTAGCTACAACCAGAAGTTCAAGGGCAAGGCC ACATTGACTGTTGACAAGTCCTCCAGCACAGCCTACATGGAGCTCAACAGC CTGACATCTGAGGACTCTGCAGTCTATTACTGTGCACGAGATGGTCACCGC TGGTACTTCGATGTCGGGACACAGGGACCACGGTACCGTCTCCTCA (SEQ ID NO:126)
5		EVQLQQSGPVLVKPGASVKMSCKASGYTFDYYMNWVKQSHGKSLEWIGVI IPYNGGTSYNQFKKGATLTVDKSSSTAYMELNSLTSEDSAVYYCARDGHR WYFDVWGTGTTVTVSS (SEQ ID NO:127)
10	15F1vh	EVQLQQSGPELVKPGASVKMSCKASGYFTCCIMHWVKQKPGQGLEWIGYI NPYNDGTYNEKFKGATLSDKSSSTAYMELSSLTSEDSAVYYCARRWDE AYWGQGTLTVSA (SEQ ID NO:128)
15		
20	18D3vh	GAGGTCCAGCTGCAGCAGTCTGGACCTGAGCTGGTAAGCCTGGGCTTCA GTGAAGATGTCCTGCAAGGCTCTGGATAACACATTCACTCGTATCTTATT CACTGGGTGAAACAGAACAGCCTGGCAGGGCCTTGAGTGGATTGGATATATT AATCCTAACATGATGGTACTAAATACAATGAGAAGTTCAAAGGCAGGCC ACACTGACTTCAGACAAATCCTCCAGCACAGCCTACATGGAGCTCAGCAGC CTGACCTCTGAGGACTCTGCAGTCTATTACTGTACCTCTAACACTGGACCTC GACTACTGGGCAAGGCACCACTCTCACAGTCTCCTCA (SEQ ID NO:129)
25		
30		
35		
40		EVQLQQSGPELVKPGASVKMSCKASGYTFTRYLIHWVKQKPGQGLEWIGYI NPYNDGTYNEKFKGATLSDKSSSTAYMELSSLTSEDSAVYYCTSNWDL DYWGQGTTLVSS (SEQ ID NO:130)
45	18C4vh	GAGGTCCAGCTGCAGCAGTCTGGACCTGAAGTGGTAAAGCCTGGGCTTCA GTGAAGATGTCCTGCAAGGCCTCTGGATAACACATTCACTAGTTCTGTTATA CACTGGGTGAAGCAGAACAGCCTGGCAGGGCCTTGAGTGGATTGGATATATC AATCCTATAATGATGGTACTAAAGTACAATGAGAAGTTCAAAGGCAGGCC ACACTGACTTCAGACAGATCCTCCAGCACAGCCTACATGGAGCTCAGCAGC CTGACCTCTGAGGACTCTGCAGTCTATTACTGTACAAAGACGGTTGGACGAG GCTTACTGGGCAAGGGACTCTGGTCACTGTCTCTGCA (SEQ ID NO:131)
50		
55		EVQLQQSGPEVVKPGASVKMSCKASGYTFSSVIHWVKQKAGQGLEWIGYI NPYNDGTYNEKFKGATLSDRSTSAYMELSSLTSEDSAVYYCTRRLDE AYWGQGTLTVSA (SEQ ID NO:132)

(continued)

	18E5vh	GAGGTCCAGCTGCAGCAGTCTGGACCTGAGCTGGTAAAGCCTGGGCTGCA GTGAAGATGTCCTGCAAGGCTCTGGATAACACATTCACTCGCTATCTTATT CACTGGGTGAAGCAGAAGCCTGGCAGGGCCTTGAGTGGATTGGATATATT AATCCTACAATGATGGTACTAAATATAATGAGAAGTTCAAAGGCAAGGCC ACACTGACTTCAGACAAATCCTCCAGCACAGCCTACATGGAGCTCAGCAGC CTGACCTCTGAGGACTCTGCGGTCTATTACTGTACCTCTAATTGGACCTC GACTACTGGGCCAAGGCACCCTCACAGTCTCCTCA (SEQ ID NO:133)
5		EVQLQQSGPELVKPGAAVKMSCKASGYTFTRYLIHWVKQKPGQGLEWIGYI NPYNDGTKYNEKFKGKATLTSKSSSTAYMELSSLTSEDSAVYYCTSNWDL DYWGQGTLTVSS (SEQ ID NO:134)
10	20E8vh	GAGGTCCAGCTGCAGCAGTCTGGACCTGAACTGGTAAAGCCTGGGCTTCA GTGAAGATGTCCTGCAAGGCCTCTGGATAACACATTCACTAGTTCTGTTATG CACTGGGTGAAGCAGAAGGCTGGCAGGGCCTTGAGTGGATTGGATATATC AATCCTATAATGATGGTACTCAGTACAATGAGAAGTTCAAAGGCAAGGCC ACACTGACTTCAGACAAATTCAGCACAGCCTACATGGAGCTCAGCAGC CTGACCTCTGAGGACTCTGCGGTCTATTACTGTACAAGACGGTTGGACGAG GCTTACTGGGCCAAGGGACTCTGGTCACTGTCTTGCA (SEQ ID NO:135)
15		EVQLQQSGPELVKPGASVVKMSCKASGYFTSSVMHWVKQKAGQGLEWIGYI NPYNDGTQYNEKFKGKATLTSKFSSTAYMELSSLTSEDSAVYYCTRRLDE AYWGQGTLTVSA (SEQ ID NO:136)
20	22E2vh	GAGGTCCAGCTGCAGCAGTCTGGACCTGAGCTGGTAAAGCCTGGGCTTCA GTGAAGATGTCCTGCAAGGCTCTGGATAACACATTCACTAGCTCTATTATT CACTGGGTGAAGCAGAGGCCTGGCAGGGCCTTGAGTGGATTGGATATATT AATCCTACGATGATGTTACTAAGTACAATGAGAAGTTCAAAGGCAAGGCC ACACTGACTTCAGACAAATCCTCCAGCACAGCCTACATGGAGCTCAGCAGC CTGACCTCTGAGGACTCTGCGGTCTATTACTGTGCAAGACGGTGGACGAG TCTTACTGGGCCAAGGGACTCTGGTCACTGTCTTGCA (SEQ ID NO:137)
25		EVQLQQSGPELVKPGASVVKMSCKASGYFTSSIIHWVKQRPGQGLEWIGYI NPYDDVTKYNEKFKGKATLTSKSSSTAYMELSSLTSEDSAVYYCARRWDE SYWGQGTLTVSA (SEQ ID NO:138)
30		
35		
40		
45		
50		

(continued)

5	24A5vh	GAGGTCCAGCTGCAGCAGTCTGGACCTGAGCTGGTAAAGCCTGGGCTTCA GTGAAGATGTCCTGCAAGGCTCTGGATACTTCACTACCTCTATTATG CACTGGGTGAAAACAGAACGCTGGCAGGGCCTTGAGTGGATTGGATATATT AATCCTTACGATGATGTTACTAAGTACAATGAAAAGTTCAAAGGCCAAGGCC
10		ACATTGACTTCAGACAAATCCTCCAGCACAGCCTACATGGAGCTCAGCAGC CTGACCTCTGAGGACTCTGCAGTCTATTACTGTGTAAGACGGTGGACGAG GCTTACTGGGCCAAGGGACTCTGGTCACTGTCTTGCA (SEQ ID NO:139)
15		EVQLQQSGPELVKPGASVKMSCKASGYTF TTSIMHWVKQKPGQGLEWIGYI NPYDDVTKYNEKFKGKATLSDKSSSTAYMELSSLTSEDSAVYYCVRRWDE AYWGQGTLTVSA (SEQ ID NO:140)
20	26F7VH	GAGGTCCAGCTGCAACAAATCTGGACCTGAGCTGGTGAAGCCTGGGCTTCA GTGAAGATATCCTGTAAGGCTCTGGATACACGTTACTGACTACTACATG AACTGGGTGAGGCAGAGCCATGGAGAGAGCCTTGAGTGGATTGGAGATT AATCATAACAATGATGTTATTACTTACAACCCGAAGTTCAAGGGCAAGGTC ACATGACTGTAGAGAAGTCTCCACCACAGCCTACATGGAGCTCCGCAGC CTGICATCTGAGGACTCTGCAGTCTATTACTGTGCAAGGGGGCTACGAGGC TACTATGCTATGGACTACTGGGTCAAGGAACCTCAGTCACCGTCTCCTCA (SEQ ID NO:141)
25		EVQLQQSGPELVKPGASVKISCKASGYTF TDYYMNWVRQSHGESLEWIGDF NHNNNDVITYNPKFKGKVLTVEKSSTAYMELRSLSED SAVYYCARGLRG YYAMDYWGQGTSVTVSS (SEQ ID NO:142)
30	27G8vh	CAGGTTCAGCTGCAACAGTCTGACGCTGAGTTGGTGAACACCTGGAGCTTCA GTGAAGATATCCTGCAAGGTTCTGGCTACACCTTCACTGACCATACTATT CACTGGATGAAGCAGAGGCCTGAACAGGGCCTGGAATGGATTGGATATATT TATCCTAGAGATGGTTATCCTAACGTTCAATGAGAAGTTCAAGGGCAAGGCC ACATTGACTGCAGACAAATCCTCCAGCACAGCCTACATGCAGCTAACAGC CTGACATCTGAGGACTCTGCAGTCTATTCTGTGCAAGACGGCCCCCTTAC TATGCTATGGACTACTGGGTCAAGGAACCTCAGTCGCCGTCTCCTCA (SEQ ID NO:143)
35		QVQLQQSDAELVKPGASVKISCKVSGYTF DHTIHWMKQRPEQGLEWIGYI YPRDGYPKFNEKFKGKATLADKSSSTAYMQLNSLTSEDSAVYFCARRPPY YAMDYWGQGTSVAVSS (SEQ ID NO:144)
40		
45		
50		

(continued)

5	31H9vh	GAGGTCCAAC TGCAGCAGTCTGGACCTGAGCTGGTAAAGCCTGGGCTTCA GTGAAGATGT CCGTCAAGGCTCTGGATAACACATTCACTAGGTATCTTATG CACTGGGTGAAGCAGAAGCCTGGGCAGGGCTTGAGTGGATTGGTTATATT AATCCTTACAATGATGGTACTAATTACAATGAGAAGTTCAAAGGCAAGGCC ACACTGACTTCAGACAAATCCTCCAGCACAGCCTACATGGAGCTCAGCAGC CTGACCTCTGAGGACTCTGCAGTCTATTACTGTTCCCTTAAC TGGGACTAT GCTTACTGGGCCAAGGGACTCTGGTCACTGTCTTGCA (SEQ ID NO:145)
10		EVQLQQSGPELVKPGASVKMSCKASGYTFTRYLMHWVKQKPGQGLEWIGYI NPYNDGTNYNEKFKGKATLTSDKSSSTAYMELSSLTSEDSAVYYCSLNWDY AYWGQGTLTVSA (SEQ ID NO:146)
15	34G3VH	GAGTCCAGCTGCAGCAGTCTGGACCTGAGCTGGTAAAGCCTGGCCTTCA GTGAAGATATCCTGCAAGGCTCTGGTTACTCATTCACTGACTACAACATG AACTGGGTGAAGCAGAGCAAAGGAAAGAGCCTTGAGTGGATTGGAGTAATT ATTCTTAACATATGGTTTACTAGCTACAATCAGAAC TTCAAGGGCAAGGCC ACTTGACTGTAGACCAGTCTCCAGCACAGCCCACATGCAGCTAACAGT GTGACATCTGAGGACTCTGCAGTCTATTACTGTGTAAGAGATGGGGAATA CTCCTCTGGTATCTCGATGTCTGGGCACAGGGACCACGGTCACCGTCTCC TCA (SEQ ID NO:147)
20		EFQLQQSGPELVKPGASVKISCKASGYSFTDYNMNWVKQSKGKSLEWIGVI IPNYGFTSYNQNFKGKATLTV DQSSSTAHMQLNSVTSEDSAVYYCVRDGGI LLWYLDVWGTGTTTVSS (SEQ ID NO:148)
25	34D9VH7	GAGGTCCAGCTGCAACAGTCTGGACCTGAGCTGGTAAAGCCTGGGCTTCA GTGAAGATACCCCTGCAAGGCTCTGGATAACACATTCACTGACTACAACATG GACTGGGTGAAGAAGAGCCATGGAAAGAGCCTTGAGTGGATTGGAGATATC AATCCTCACAAATGGTGGTACTATCTACAACCAGAAC TTCAAGGGCAAGGCC ACATTGACTGTAGACAAGTCCCTCCAGCACAGCCCACATGGAGCTCCGAGC CTGACATCTGAGGACACTGCAGTCTATTACTGTGCAAGAAATTACTACGGT AGTAGTTACGGCTGGTACTTCGATGTCTGGGCACAGGGACCACGGTCACC GTCTCGTCA (SEQ ID NO:149)
30		EVQLQQSGPELVKPGASVKIPCKASGYTFTDYNMDWVKKSHGKSLEWIGDI NPHNGGTIYNQKFKGKATLTVDKSSSTAHMELRSLTSEDTAVYYCARNYYG SSYGWYFDVWGTGTTTVSS (SEQ ID NO:150)
35		
40		
45		
50		
55		

(continued)

5	43F5vh	GAGGTCCAGCTGCAGCAGTCTGGACCTGAGCTGGTGAAGCCTGGGGCTTCA GTGAAGATTCCTGCAGGGCTCTGGTTACTCATTCACTGGCTACTACATG AACTGGGTGAAGCAAAGTCTGAAAAGAGCCTTGAGTGGATTGGAGAGATT ATT CCTTACCACTGGTGGTACTTCCTACAACCAGAAGTTCAAGGCCAAGGCC ACATTGACTGTAGACAAATCCTCCAGCACAGCCTACATGCAGCTCAAGAGC CTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGAGAGAGCGGTGGG TTCTACTGGTACTTCGATGTCTGGGCACAGGGACCACGGTCACCGTCTCC TCA (SEQ ID NO:151)
10		EVQLQQSGPELVKPGASVKISCRASGYSFTGYYMNWVKQSPEKSLEWIGEI IPTTGGTSYNQKFKAATLTVDKSSSTAYMQLKSLTSEDSAVYYCARESGG FYWYFDVWGTGTTVTVSS (SEQ ID NO:152)
15	73H10VH	GAGGTCCAGCTGCAGCAGTCTGGACCTGAGCTGGTAAAGCCTGGGGCTTCA GTGAAGATGTCTGCAAGGCTCTGGATAACACATTCACTAGGTATGTTATG CACTGGGTGAAGCAGAACAGCCTGGCAGGGCCTTGAGTGGATTGGATATATT AATCCTACAATGATGTTACTAAGTACAATGAGAAGTTCAAAGGCCAAGGCC ACACTGACTTCAGACAGATCCTCCAGCACAGCCTACATGAAACTCAGCAGC CTGACCTCTGAGGACTCTGCAGTCTATTATTGTGCAAGAAACTGGGACGTT CCTTACTGGGCCAAGGGACTCTGATCACTGTCTCTGCA (SEQ ID NO:153)
20		EVQLQQSGPELVKPGASVKMSCKASGYTFTRYVMHWVKQKPGQGLEWIGYI NPYNDVTKYNEFKKGKATLTSRSSLTSEDSAVYYCARNWDV PYWGQGTLITVSA (SEQ ID NO:154)
25	74H3VH	GAGGTCCAGCTGCAGCAGTCTGGACCTGAGCTGGTAAAGCCTGGGGCTTCA GTGAAGATGTCTGCAAGGCTCTGGATAACACATTCACTAGGTATCTTATG CACTGGGTGAAGCAGAACAGCCTGGACAGGGCCTTGAGTGGATTGGATATATT AATCCTACAATGATGGTACTAAGTACAATGAGAGGTTCAAAGGCCAAGGCC ACACTGACTTCAGACAAATCCTCCAGCACAGCCTACATGGAGCTCAGCAGC CTGACCTCTGAGGACTCTGCAGTCTATTACTGTGCAAGAAACTGGGACGTA CCTTACTGGGCCAAGGGACTCTGGTCACTGTCTCTGCA (SEQ ID NO:155)
30		EVQLQQSGPELVKPGASVKMSCKASGYTFTRYLMHWVKQKPGQGLEWIGYI NPYNDGTVKYNERFKKGKATLTDKSSSTAYMELSSLTSEDSAVYYCARNWDV PYWGQGTLITVSA (SEQ ID NO:156)
35		
40		
45		
50		

[0140] Human framework sequences were selected for each of the mouse leads based on the framework homology, CDR structure, conserved canonical residues, conserved interface packing residues and other parameters.

[0141] The mouse light chain and heavy chain CDRs of the various mouse antibodies are shown in Table 3 and Table 4, respectively. Table 4 also shows three heavy chains CDRs derived from the mouse antibody 6B8 through the hu-

manization process.

Table 3: LIGHT CHAIN CDR sequences

	L-CDR1	L-CDR2	L-CDR3
5	18C4 RASQSISEYLH (SEQ ID NO:1)	YASQSIS (SEQ ID NO:2)	QNGHSFPFT (SEQ ID NO:3)
18E5 RASQSISDYLY (SEQ ID NO:4)	FASQSIS (SEQ ID NO:5)	QNGHSFPFT (SEQ ID NO:3)	
18D3 RASQSISDYLY (SEQ ID NO:4)	FASQSIS (SEQ ID NO:5)	QNGHSFPFT (SEQ ID NO:3)	
10 20E8 RASQSISEYLH (SEQ ID NO:1)	YASQSIS (SEQ ID NO:2)	QNGHSFPFT (SEQ ID NO:3)	
22E2 RASQSISVYLN (SEQ ID NO:6)	YASQSIS (SEQ ID NO:2)	QNGHSFPFT (SEQ ID NO:3)	
24A5 RASQSISDYLN (SEQ ID NO:7)	YASQSIS (SEQ ID NO:2)	QNGHSFPFT (SEQ ID NO:3)	
15 15C11 RSSQLVHSNGNTYLH (SEQ ID NO:8)	KVSNRFS (SEQ ID NO:9)	SQSTHVPYT (SEQ ID NO:10)	
43F5 RSSQLVHSNGNTYLH (SEQ ID NO:8)	KVSNRFS (SEQ ID NO:9)	SQSTHVPYT (SEQ ID NO:10)	
27G8 RASKSVSTSGYSYIH (SEQ ID NO:11)	LASNLDs (SEQ ID NO:12)	QHSRELPYT (SEQ ID NO:13)	
31H9 RASQSISDYLN (SEQ ID NO:7)	YASQSIS (SEQ ID NO:2)	QNGHSFPYT (SEQ ID NO:14)	
20 2D1 RTSESVSYGQNFIH (SEQ ID NO:15)	RASNLES (SEQ ID NO:16)	QQTNEDPYT (SEQ ID NO:17)	
9D12 RASETINFYGTSFMH (SEQ ID NO:18)	RASNLES (SEQ ID NO:16)	QQTNEDPYT (SEQ ID NO:17)	
6B8 KASRDVAIAAVA (SEQ ID NO:19)	WASTRHT (SEQ ID NO:20)	HQYSSYPFT (SEQ ID NO:21)	
25 73H10 RASENIDSYLA (SEQ ID NO:22)	AARNLAD (SEQ ID NO:23)	QHYYSTPFT (SEQ ID NO:24)	
74H3 RASENIDSYLA (SEQ ID NO:22)	AATNLAD (SEQ ID NO:25)	LHYYSTPFT (SEQ ID NO:26)	
35H8 RSSQLVHSNGNTYLH (SEQ ID NO:8)	KVSNRFS (SEQ ID NO:9)	SQSTHVPYT (SEQ ID NO:10)	
30 26F7 RASKSVRFSDYFYMH (SEQ ID NO:27)	LASNLES (SEQ ID NO:28)	QNSRELPYT (SEQ ID NO:29)	
34G3 RSSQLVHSNGNTYLH (SEQ ID NO:8)	KVSNRFS (SEQ ID NO:9)	SQSTHVPYT (SEQ ID NO:10)	
34D9 KASQDVGNNAV (SEQ ID NO:30)	WASTRH (SEQ ID NO:31)	QQYSSYLT (SEQ ID NO:32)	

Table 4: HEAVY CHAIN CDR sequences

	H-CDR1	H-CDR2	H-CDR3
40 18C4 GYTFTSSVIH (SEQ ID NO:33)	YINPYNDGTKYNEKFKG (SEQ ID NO:34)	RLDEAY (SEQ ID NO:35)	
18E5 GYTFTTRYLIH (SEQ ID NO:36)	YINPYNDGTKYNEKFKG (SEQ ID NO:34)	NWDLDY (SEQ ID NO:37)	
45 18D3 GYTFTTRYLIH (SEQ ID NO:36)	YINPYNDGTKYNEKFKG (SEQ ID NO:34)	NWDLDY (SEQ ID NO:37)	
20E8 GYTFTSSVMH (SEQ ID NO:38)	YINPYNDGTQYNEKFKG (SEQ ID NO:39)	RLDEAY (SEQ ID NO:35)	
50 22E2 GYTFTSSIIH (SEQ ID NO:40)	YINPYDDVTKYNEKFKG (SEQ ID NO:41)	RWDESY (SEQ ID NO:42)	
24A5 GYTFTTSIMH (SEQ ID NO:43)	YINPYDDVTKYNEKFKG (SEQ ID NO:41)	RWDEAY (SEQ ID NO:44)	
55 15C11 GYTFTDYYMN (SEQ ID NO:45)	VIIIPYNGGTSYNQKFKG (SEQ ID NO:46)	DGHRWYFDV (SEQ ID NO:47)	
43F5 GYSFTGYYMN (SEQ ID NO:48)	EIIPPTGGTSYNQKFKA (SEQ ID NO:49)	ESGGFYWYFDV (SEQ ID NO:50)	

(continued)

	H-CDR1	H-CDR2	H-CDR3
5	27G8 GYTFTDHTIH (SEQ ID NO:51)	YIYPRDGYPKFNEKFKG (SEQ ID NO:52)	RPPYYAMDY (SEQ ID NO:53)
10	31H9 GYTFTRYLMH (SEQ ID NO:54)	YINPYNDGTYNEKFKG (SEQ ID NO:55)	NWDYAY (SEQ ID NO:56)
15	2D1 GFSLTTYAIS (SEQ ID NO:57)	VIWTGGGTKYNSALKS (SEQ ID NO:58)	KDNYGGAMDY (SEQ ID NO:59)
20	9D12 GFSLNNFAIS (SEQ ID NO:60)	AIWTGGGTNYNSALKS (SEQ ID NO:61)	KDYSYGGAMDY (SEQ ID NO:62)
25	6B8 GNTFTDQTIH (SEQ ID NO:63)	YIYPRDDSPKYNENFKG (SEQ ID NO:64)	PDRSGYAWFIY (SEQ ID NO:65)
30	Hu_6B8-2 GYTFTDQTIH (SEQ ID NO:66)	YIYPRDDSPKYNENFKG (SEQ ID NO:64)	PDRSGYAWFIY (SEQ ID NO:65)
35	Hu_6B8-5 Hu_6B8 -36/65 GYTFTDQTIH (SEQ ID NO:67)	YIYPRDDSPKYNENFKG (SEQ ID NO:64)	PDRSGYAWFIY (SEQ ID NO:65)
40	73H10 GYTFTRYVMH (SEQ ID NO:69)	YINPYNDVTKYNEKFKG (SEQ ID NO:70)	NWDVPY (SEQ ID NO:71)
45	74H3 GYTFTRYLMH (SEQ ID NO:54)	YINPYNDGTYNERFKG (SEQ ID NO:72)	NWDVPY (SEQ ID NO:71)
50	35H8 GYTFTDYYMN (SEQ ID NO:45)	VIIPNYNGGISYNQKFKG (SEQ ID NO:73)	NDYDWYFDV (SEQ ID NO:74)
55	26F7 GYTFTDYYMN (SEQ ID NO:45)	DFNHNNNDVITYNPKFKG (SEQ ID NO:75)	GLRGYYAMDY (SEQ ID NO:76)
	34G3 GYSFTDYNMN (SEQ ID NO:77)	VIIPNYGFTSYNQNFKG (SEQ ID NO:78)	DGGILLWYLDV (SEQ ID NO:79)
	34D9 GYTFTDYNMD (SEQ ID NO:80)	DINPHNGGTIYN4KFKG (SEQ ID NO:81)	NYYGSSYGWYFDV (SEQ ID NO:82)

[0142] The CDRs listed above in Tables 3 and 4 are defined using the Chothia lumbering system (Al-Lazikani et al., (1997) JMB 273,927-948).

[0143] Fab that showed better or equal binding as compared to the chimeric parent Fab were selected for conversion to IgG. 6B8 was converted to an IgG1KO format. IgG1KO (knock-out of effector functions) has two mutations in the Fc region, Leu234Ala and Leu235Ala, which reduce effector function such as Fc γ R and complement binding. The IgG format is described in the literature (see for example Hezareh et al. (2001) Journal of Virology 75: 12161-12168). Example 1 describes the humanization process in further detail. The results of such humanization resulted in humanized-antibody sequences. A representative number of humanized light chain and heavy chain variable regions derived from mouse antibody 6B8 are provided and shown in Tables 5 and 6. An alignment between the humanized light chain and heavy chain variable regions derived from mouse antibody 6B8 and the chain and heavy chain, variable regions from mouse antibody 6B8 is shown in Figure 1.

[0144] Selected combination of humanized light chain and heavy chain variable regions derived from mouse antibody 6B8 resulted in Antibodies A, B, C and D:

Antibody A: 6B8-IgG1KO-2 with IgK-66 (heavy chain variable region 6B8CVH-02 and light chain variable region 6B8CVK-66);

Antibody B: 6B8-IgG1KO-5 with IgK-66 (heavy chain variable region 6B8CVH-05 and light chain variable region 6B8CVK-66);

Antibody C: 6B8-IgG1KO-2 with IgK-65 (heavy chain variable region 6B8CVH-02 and light chain variable region

6B8CVK-65);

Antibody D: 6B8-IgG1KO-5 with IgK-65 (heavy chain variable region 6B8CVH-05 and light chain variable region 6B8CVK-65).

- 5 [0145] Antibodies A, B, C and D have the heavy and light chain sequences shown in Table 7.

Table 5: Humanized 6B8-VK Sequences

	6B8CVK-65	Gacatccagatgacccagagccccaaaggcgcggctgagcgccagcgtgg cgaccgcgtgaccatcacctgcaaggccagccgcacgtggccatcg ccgtggcttgttaccaggcagaagccaggcaaggtgccaaagctgctg ctgttctggccagcaccgcacccggccacaccggctgcccagaccgcctcag cggcagcggcagcggcaccgacttcaccctgaccatcagcagcctgc agccagaggacgtggccactactactgcccaccagtacagcagctac ccattcaccttcggccagggcaccaagctggagatcaag (SEQ ID NO:157)
10		DIQMTQSPSSLSASVGDRVITCKASRDVAIAVAWYQQKPGKVPKLL LFWAISTRHTGVPDFSGSGSTDFLTISLQPEDLADYYCHQYSSY PFTFGQGTKEIK (SEQ ID NO:158)
15	6B8CVK-66	Gacatccagatgacccagagccccaaaggcgcggctgagcgccagcgtgg cgaccgcgtgaccatcacctgcaaggccagccgcacgtggccatcg ccgtggcttgttaccaggcagaagccaggcaaggtgccaaagctgctg atctactggccagcaccgcacccggccacaccggctgcccagccgcctcag cggcagcggcagccgcaccgacttcaccctgaccatcagcagcctgc agccagaggacgtggccactactactgcccaccagtacagcagctac ccattcaccttcggccagggcaccaagctggagatcaag (SEQ ID NO:159)
20		DIQMTQSPSSLSASVGDRVITCKASRDVAIAVAWYQQKPGKVPKLL IYWASTRHTGVPSRFSGSGSRTDFLTISLQPEDVADYFCHQYSSY PFTFGSGTKEIK (SEQ ID NO:160)
25	6B8CVK-67	Gacatccagatgacccagagccccaaaggcgcggctgagcgccagcgtgg cgaccgcgtgaccatcacctgcaaggccagccgcacgtggccatcg ccgtggcttgttaccaggcagaagccaggcaaggtgccaaagctgctg ctgtactggccagcaccgcacccggccacaccggctgcccagccgcctcag cggcagcggcagccgcaccgacttcaccctgaccatcagcagcctgc agccagaggacgtggccacctactactgcccaccagtacagcagctac ccattcaccttcggccagggcaccaagctggagatcaag (SEQ ID NO:161)
30		
35		
40		
45		
50		
55		

(continued)

5

10

15

20

25

	DIQMTQSPSSLSASVGDRVTITCKASRDVAIAAVAWYQQKPGKVPKLL LYWASTRHTGVPSRESGSGSRTDFLTISLQPEDVATYYCHQYSSY PFTFGSGTKLEIK (SEQ ID NO:162)
6B8CVK-78	Gacatccagatgaccaggcccagaagcagcctgagcgccagcgtgg cgaccgcgtgaccatcacctgcaaggccagccgcacgtggccatcg ccgtggcctggtaccaggcagaagccaggcaaggtgccaaagctgctg ctgttctggccagcaccgcacccggcgtgccagaccgcttcag cggcagcggcagccgcaccgacttcaccctgaccatcagcagcctgc agccagaggacacctggccactactactgcccaccagtacagcagctac ccattcaccttcggcagcggcaccaagctggagatcaag (SEQ ID NO:163)
	DIQMTQSPSSLSASVGDRVTITCKASRDVAIAAVAWYQQKPGKVPKLL LFWASTRHTGVPDFSGSGSRTDFLTISLQPEDLADYYCHQYSSY PFTFGSGTKLEIK (SEQ ID NO:164)

Table 6: Humanized 6B8-VH Sequence

30

35

40

45

50

55

6B8CVH-02	Caggtgcagctggtgcagagcggcgccgaggtgaagaagccaggcag cagcgtgaagggtgagctgcaaggccagcggctacaccttaccgacc agaccatccactggatgcgccaggccccaggccagggcctggagtgg atcggctacatctacccacgcgacgcacagccaaagtacaacgagaa cttcaaggcaaggtcaccatcacgcgacaagagcaccagcaccg cctacatggagctgagcagcctgcgcagcggacaccgcgtgtac tactgcgccatcccagaccgcagcggctacgcctggttcatctactg ggccagggcaccctggtaccgtgagcagc (SEQ ID NO:165)
	QVQLVQSGAEVKPGSSVKVSCKASGYTFDQTIHWMRQAPGQGLEW IGYIYPRDDSPKYNENFKGVITADKSTSTAYMELSSLRSEDTAVY YCAIPDRSGYAWFIYWGQGTLTVSS (SEQ ID NO:166)
6B8CVH-05	Caggtgcagctggtgcagagcggcgccgaggtgaagaagccaggcag cagcgtgaagggtgagctgcaaggccagcggcttacacccatccgacc agaccatccactggatgcgccaggccccaggccagggcctggagtgg atcggctacatctacccacgcgacgcacagccaaagtacaacgagaa cttcaaggcaaggtcaccctgaccgcgacaccgcgtgtac cctacatggagctgagcagcctgcgcagcggacaccgcgtgtac tactgcgccatcccagaccgcagcggctacgcctggttcatctactg ggccagggcaccctggtaccgtgagcagc (SEQ ID NO:167)

(continued)

5		QVQLVQSGAEVKPGSSVKVSCKASGGFTTDQTIHWVRQAPGQGLEW MGYIYPRDDSPKYNENFKGVTLTADKSTSTAYMELSSLRSEDTAVY YCAIPDRSGYAWFIYWGQGTLTVSS (SEQ ID NO:168)
10	6B8CVH-36	Caggtgcagctggtgcagagcggcgccgaggtgaagaaggccaggcag cagcgtgaagaccagactgcaaggccagcggcgacacctcacccgacc agaccatccactgggtgcgccagcggcccaggccaggccctggagtgg atgggctacatctacccacgcgacacagccaaagtacaacgagaa cttcaaggccgcgtcaccatcaccgcgacaagagcaccaggcaccg cctacatggagctgagcagcctgcgcagcggacaccgcgtgtac tactgcgccatccagaccgcagccgtacgcctggttcatctactg ggccagggcaccctggtgaccgtgagc (SEQ ID NO:169)
15		QVQLVQSGAEVKPGSSVKTSCKASGGFTTDQTIHWVRQRPGQGLEW MGYIYPRDDSPKYNENFKGRVTITADKSTSTAYMELSSLRSEDTAVY YCAIPDRSGYAWFIYWGQGTLTVSS (SEQ ID NO:170)
20	6B8CVH-65	Caggtgcagctggtgcagagcggcgccgaggtgaagaaggccaggcag cagcgtgaaggtgagctgcaaggccagcggcgacacctcacccgacc agaccatccactgggtgcgccaggccccaggccaggccctggagtgg atgggctacatctacccacgcgacacagccaaagtacaacgagaa tttcaaggccgcgtcaccctgaccgcgacaagagcaccaggcaccg cctacatggagctgagcagcctgcgcagcggacaccgcgtgtac ttctgcgcggccagaccgcagccgtacgcctggttcatctactg ggccagggcaccctggtgaccgtgagc (SEQ ID NO:171)
25		QVQLVQSGAEVKPGSSVKVSCKASGGFTTDQTIHWVRQAPGQGLEW MGYIYPRDDSPKYNENFKGRVTLTADKSTSTAYMELSSLRSEDTAVY FCARPDRSGYAWFIYWGQGTLTVSS (SEQ ID NO:172)
30		
35		
40		
45		
50		
55		

Table 7: Heavy and Light Chain DNA and Amino Acid Sequences for Antibodies A, B, C, and D

Antibody A	IgK light Chain #66	<p><u>Gacatccagatgaccqaqgcccagaqcgcctgagcq</u> <u>ccaqcgtqqcqaccqcgatccatcacctqcaaqgc</u> <u>caqccqcgacqgtggccatcqccgtqgcctqgtaccqaq</u> <u>caqaaqccaaqqcaagqtqccaaagqctqctqatctact</u> <u>qqaccaggcaccqccacaccqgcgtgccaqcccgtt</u> <u>caqcgqcaqcqgcaqccgaccqacttcaccctgacc</u> <u>atcagcagcctgcaqccagaggacgtqgcccqactact</u> <u>tctqccaccaggtaqcaqcaqctaccattcacccattcgq</u> <u>cagcggcaccaaqctqgagatcaagcgtactgtggct</u> <u>gcaccatctgtcttcatcttccgccatctgatgagc</u> <u>agttgaaatctggaaactgcctctgttgtgtgcctgct</u> <u>gaataacttctatcccagagagggccaaagtacagtgg</u> <u>aaggtggataacgcccctccaatcggttaactcccagg</u> <u>agagtgtcacagagcaggacagcaaggacagcaccta</u> <u>cagcctcagcagcaccctgacgctgagcaaagcagac</u> <u>tacgagaaacacaaaagtctacgcctgcaagtaccc</u> <u>atcagggcctgagctcgccgtcacaaagagcttcaa</u> <u>caggggagagtgt (SEQ ID NO:173)</u></p>
		<p><u>DIQMTQSPSSLSASVGDRVTITCKASRDVAIAVAWYQ</u> <u>QKPGKVPKLLIYWASTRHTGVPSRFSGSGSRTDFTLT</u> <u>ISSLQPEDVADYFCHOYSSYPFTFGSGTKLEIKRTVA</u> <u>APSVFIFPPSDQLKSGTASVVCLNNFYPREAKVQW</u> <u>KVDNALQSGNSQESVTEQDSKDSTYSLSSTLTSKAD</u> <u>YEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID</u> <u>NO:174)</u></p>
	IgG1 KO Heavy Chain #2	<p><u>Caggtaqctqgtqcaqaaqcggcgccqaaqgtqaaqa</u> <u>aqccaggcaqcaqcgtaqgtqagctqcaaggccaaq</u> <u>cqgctacaccctcaccqaccqaccatccactqgatq</u> <u>cqccaaqqcccccaqqccaaqqccctqqaqtggatcqgct</u> <u>acatctacccacqcgacgacagccccaaqtaacaacqa</u> <u>gaactcaaggcaaggtaaccatcaccqccgacaag</u></p>

(continued)

5		<u>aqcacc</u> <u>aggcacc</u> <u>ggcctacat</u> <u>ggagct</u> <u>gagcag</u> <u>cctgc</u> <u>gcagcg</u> <u>aggacacc</u> <u>ggccgt</u> <u>gtactact</u> <u>qcgccatccc</u> <u>agaccg</u> <u>cagcgg</u> <u>ctacgc</u> <u>cctgg</u> <u>ttcatctact</u> <u>ggggc</u> <u>caggq</u> <u>cacc</u> <u>ctgg</u> <u>gtgacc</u> <u>gtgagg</u> <u>tcac</u> <u>actt</u> <u>ccctcc</u> <u>agggccc</u> <u>atcggt</u> <u>tttcccc</u> <u>ctggc</u> <u>accct</u> <u>cctccaa</u> <u>gagcac</u> <u>cttgggg</u> <u>cacagc</u> <u>ggcc</u> <u>ctgg</u> <u>gtgc</u> <u>ctgt</u> <u>gtcaagg</u> <u>actactt</u> <u>cccc</u> <u>gaacc</u> <u>gggt</u> <u>gacgg</u> <u>gtgt</u> <u>ggaact</u> <u>caggc</u> <u>ggcc</u> <u>ctgacc</u> <u>agcgg</u> <u>gtgc</u> <u>acac</u> <u>ctt</u> <u>ccggct</u> <u>gtcct</u> <u>acagt</u> <u>cctc</u> <u>aggact</u> <u>ctact</u> <u>ccctc</u> <u>agcagc</u> <u>gtgg</u> <u>tgacc</u> <u>gtgc</u> <u>ccct</u> <u>ccagc</u> <u>agctt</u> <u>ggca</u> <u>cccagac</u> <u>ctacat</u> <u>ctgca</u> <u>acgt</u> <u>gaat</u> <u>caca</u> <u>agccc</u> <u>ag</u> <u>caacacca</u> <u>agg</u> <u>tcgaca</u> <u>agagag</u> <u>gtt</u> <u>gagcc</u> <u>aaat</u> <u>tct</u> <u>tgtgacaaa</u> <u>actcac</u> <u>acat</u> <u>gccc</u> <u>accgt</u> <u>gccc</u> <u>agcac</u> <u>cagaag</u> <u>ctgct</u> <u>gggg</u> <u>accgt</u> <u>cagt</u> <u>ttcct</u> <u>cttccc</u> <u>cccaaa</u> <u>accca</u> <u>agg</u> <u>gacacc</u> <u>ctcat</u> <u>gat</u> <u>ctccc</u> <u>ggacc</u> <u>cctgaggt</u> <u>cacat</u> <u>gcgt</u> <u>gtgg</u> <u>acgt</u> <u>gagcc</u> <u>acg</u> <u>aagacc</u> <u>ctgagg</u> <u>tcaagt</u> <u>tcaact</u> <u>ggta</u> <u>cgtgg</u> <u>acgg</u> <u>cgtggagg</u> <u>gtcata</u> <u>atgcca</u> <u>agaca</u> <u>aaagcc</u> <u>gggg</u> <u>ag</u> <u>gagcagt</u> <u>taca</u> <u>acac</u> <u>acgt</u> <u>accgt</u> <u>gtgg</u> <u>tacgt</u> <u>tcaccgt</u> <u>cctgc</u> <u>accagg</u> <u>actgg</u> <u>ctga</u> <u>atgg</u> <u>caagg</u> <u>gtaca</u> <u>agtgc</u> <u>aggt</u> <u>ctcca</u> <u>acaa</u> <u>agcc</u> <u>ctccc</u> <u>agcc</u> <u>cccat</u> <u>cgagaaa</u> <u>accat</u> <u>ctccaa</u> <u>agccaa</u> <u>agg</u> <u>ggc</u> <u>agc</u> <u>ccc</u> <u>gagaacc</u> <u>acagg</u> <u>gttacacc</u> <u>ctgcccc</u> <u>atccc</u> <u>gg</u> <u>gaggagat</u> <u>gacca</u> <u>agaacc</u> <u>agg</u> <u>tcgc</u> <u>ctgac</u> <u>ctgc</u> <u>ctgg</u> <u>tcaa</u> <u>agg</u> <u>cttct</u> <u>atccc</u> <u>agc</u> <u>gacat</u> <u>cgccgt</u> <u>gg</u> <u>agtgg</u> <u>gag</u> <u>gca</u> <u>atgg</u> <u>gagcc</u> <u>gg</u> <u>gaga</u> <u>acaact</u> <u>aca</u> <u>gaccac</u> <u>gctcc</u> <u>ctgt</u> <u>ggact</u> <u>ccg</u> <u>acgg</u> <u>cttc</u> <u>ttcct</u> <u>ctac</u> <u>agca</u> <u>agct</u> <u>accgt</u> <u>gtgg</u> <u>aca</u> <u>agag</u> <u>cagg</u> <u>ggc</u> <u>agc</u> <u>agg</u> <u>gaa</u> <u>acgt</u> <u>ttct</u> <u>cat</u> <u>gtcc</u> <u>gtat</u> <u>gca</u> <u>tgagg</u> <u>ctct</u> <u>gcaca</u> <u>accact</u> <u>acac</u> <u>gcaga</u> <u>aa</u> <u>gagc</u> <u>ctc</u> <u>tccct</u> <u>gtct</u> <u>ccgg</u> <u>gt</u> (SEQ ID NO:175)
50		<u>QVOLVQSGAEVKPGSSVKVSCKASGYTFIDQTIHWM</u> <u>ROAPGOGLEWIGYIYPRDDSPKYNENFKGKVTTADK</u> <u>STSTAYMELSSLRSEDTAVYYCAIPDRSGYAWFIYWG</u> <u>QGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCL</u>

(continued)

5		VKDYFPEPVTVSWNSGALTSGVHTFPALQSSGLYSL SSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKS CDKTHTCPPCPAPEAAGGPSVFLFPPPKDLMISRT PEVTCVVVDVSHEDEVKFNWYVDGVEVHNNAKTKPRE EQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTC LVKGFYPSDIAVEWE SNGQPENNYKTPPVLDSDGSF FLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL SLSPG (SEQ ID NO:176)
10	Antibody B	IgK light Chain #66 (SEQ ID NO:173)
15		(SEQ ID NO:174)
20	IgG1KO Heavy Chain #5	<u>caggtgcagctqqtgcagagcqggccggatgtaaqa</u> <u>aqccaggcagcaqctqaaqgtqagctqcaaggccag</u> <u>cggttacacctaccgaccagaccatccactggta</u> <u>cgccaggcccccaaggccaggccctggatggatggct</u> <u>acatctaccacccacqgqacqacaqcccuaagtacaacqa</u> <u>gaacttcaaggcaaggtaaccctgaccgcccacaaq</u> <u>agcaccacgaccqccctacatggagctgagcagcctqc</u> <u>gcagcqaggacaccqccgtgtactactgcqccatccc</u> <u>agaccgcagcggctacgcctggatcatctactgggc</u> <u>cagggcaccctggatggcaggctgacgcggccatccca</u> <u>agggccatcggtttccctggcaccctctccaa</u> <u>gagcacctctggggcacagcggccctggctgcctg</u> <u>gtcaaggactacttccccgaaccggtgacggtgtcgt</u> <u>ggaactcaggcgccctgaccagcggcgtgcacacctt</u> <u>cccggctgtcctacagtccctcaggactctactccctc</u> <u>agcagcgtggtgaccgtgcccctccagcagcttggca</u> <u>cccagacctacatctgcaacgtgaatcacaagcccag</u>
25		
30		
35		
40		
45		

(continued)

		<pre> 5 caacaccaaggcgacaagagagttgagccaaatct tgtgacaaaactcacacatgccaccgtgccagcac cagaagctgctggggaccgtcagtcttccttccc cccaaaacccaaggacaccctcatgatctcccgacc cctgaggtcacatgcgtcgtggacgtgagccacg aagaccctgaggtcaagttcaactggtacgtggacgg cgtggaggtgcataatgccaagacaaagccgcggag gagcagtacaacacgacgtaccgtgtggtcagcgtcc tcaccgtcctgcaccaggactggctgaatggcaagga gtacaagtgcaggtctccaacaaagccctcccagcc cccatcgagaaaaccatctccaaagccaaagggcagc cccgagaaccacaggtgtacaccctgccccatcccg ggaggagatgaccaagaaccaggtcagcctgacctgc ctggtcaaaggctcttatccagcgcacatcgcgtgg agtggagagcaatggcagccggagaacaactacaa gaccacgcctcccgctggactccgacggctccttc ttcctctacagcaagctcaccgtggacaaggcaggt ggcagcagggaaacgtcttctatgctccgtatgca tgaggctctgcacaaccactacacgcagaagagcctc tccctgtctccgggt (SEQ ID NO:177) </pre>
		<pre> 35 <u>OVLVOSGAEVKKPGSSVKVSCKASGFTETD</u><u>OTIHWV</u> <u>ROAPGOGLEWMGYIYPRDDSPKYNENFKGKVTLTADK</u> <u>STSTAYMELSSLRSEDTAVYYCAIPDRSGYAWFIYWG</u> <u>QGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAA</u><u>LGCL</u> VKDYFPEPVTVSWNSGALTSGVHTFPALQSSGLYSL SSVTVPSLGTQTYICNVNHPKSNTKVDKRVEPKS CDKTHTCPPCPAPEAAGGPSVFLFPPPKDLMISRT PEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKP EQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTC LVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSE FLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL SLSPG (SEQ ID NO:178) </pre>

(continued)

5	Antibody C	IgK light Chain #65	<u>Gacatccagatgaccagagcccaaqcagcctgagcq</u> <u>ccagcgtggcgaccgcgtgaccatcacctgcaaggc</u> <u>cagccgcacgtggccatcgccgtggcctggttccat</u> <u>cagaagccaggcaagggtccaaagctgctgctgtttct</u> <u>gggccaqcccccccacaccggcgtgccagaccgctt</u> <u>cagcggcagcggcagcggcaccgacttcaccctgacc</u> <u>atcagcagcctgacagccaggacactggccactact</u> <u>actgccaccagtacagcaqctaccattcacccattcgg</u> <u>ccaggqcaccaaqctggagatcaagcgtactgtggct</u> <u>gcaccatctgtcttcatcttccgcacatctgatgagc</u> <u>agttgaaatctggactgcctctgttgtgcctgct</u> <u>gaataacttctatcccagagaggccaaagtacagtgg</u> <u>aagggtggataacgcctccaatcggttaactcccagg</u> <u>agagtgtcacagagcaggacagcaaggacaccta</u> <u>cagcctcagcagcaccctgacgctgagcaaagcagac</u> <u>tacgagaaacacaaaagtctacgcctgcaagtccaccc</u> <u>atcagggctgagctgccccgtcacaaagagcttc</u> aa caggggagagtgt (SEQ ID NO:179)
30			<u>DIQMTQSPSSLSASVGDRVTITCKASRDVAIAVAWYQ</u> <u>OKPGKVPKLLLFWASTRHTGVPDRFSGSGSGTDFTLT</u> <u>ISSLOPEDLADYYCHOYSSYPFTFGQGTKLEIKRTVA</u> <u>APSVFIFPPSDEQLKSGTASVVCLNNFYPREAKVQW</u> <u>KVDNALQSGNSQESVTEQDSKDSTYSLSSTTLSKAD</u> <u>YEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID</u> NO:180)
35		IgG1KO Heavy Chain #2	(SEQ ID NO:175)
40			(SEQ ID NO:176)
45	Antibody D	IgK light Chain #65	(SEQ ID NO:179)
			(SEQ ID NO:180)
50		IgG1KO Heavy Chain #5	(SEQ ID NO:177)
			(SEQ ID NO:178)

[0146] Light chains and heavy chain variable regions of Antibodies A, B, C, and D are underlined in Table 7 above.

[0147] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure has at least one of the properties below. In a further aspect, a humanized anti-IL-23p19 antibody of the present disclosure has any combination of at least two, or at least 3, 4, 5, 6, 7, 8, 9, 10, or 11 of the properties below. In a further aspect, a humanized anti-IL-23p19 antibody of the present disclosure has all the properties below.

- K_D for human IL-23 ≤ 1 pM (no shift in binding on-rate in 50% human serum)
- Blocks IL-23 binding to human IL-23R/Fc in vitro
- No binding to human IL-12
- Inhibits human IL-23 induced IL-17 production in mouse splenocytes with IC_{50} 's ≤ 20 pM
- 5 • Inhibits human IL-23 induced STAT3 phosphorylation in human DB cells with IC_{50} 's ≤ 40 pM
- No predicted activity in ADCC/CDC
- $K_D \leq 1$ pM for cynomolgus monkey IL-23
- No cross reactivity to mouse or rat IL-23
- Inhibits human IL-23 induced IL-17 and IL-22 production in mouse ear ($\geq 80\%$ inhibition of both cytokines at 1 mg/kg)
- 10 • Stability 83 °C (melting temperature 83°C as determined by differential scanning calorimetry)
- Solubility ≥ 100 mg/ml (as measured by UV spectroscopy and monitored by turbidity)
- Subcutaneous administration of 1.0 mg/kg in three cynomolgus monkeys shows sustained ≥ 10 nM exposure for approximately 28 days with a bioavailability of approximately 70%.

15 [0148] By no predicted activity in ADCC/DC, it is meant herein that a humanized anti-IL-23p19 antibody of the present invention has reduced affinity to Fc receptor and therefore is predicted not to have activity in ADCC/CDC.

[0149] In one aspect, a humanized anti-IL-23p19 antibody of the present disclosure has at least one of the properties below. In a further aspect, a humanized anti-IL-23p19 antibody of the present disclosure has any combination of at least two, or at least 3, 4, 5, 6, 7, 8, 9, or 10 of the properties below. In a further aspect, a humanized anti-IL-23p19 antibody of the present disclosure has all the properties below.

- K_D for human IL-23 ≤ 1 pM (no shift in binding on-rate in 50% human serum)
- Blocks IL-23 binding to human IL-23R/Fc in vitro
- No binding to human IL-12
- 25 • Inhibits human IL-23 induced IL-17 production in mouse splenocytes with IC_{50} 's ≤ 20 pM
- Inhibits human IL-23 induced STAT3 phosphorylation in human DB cells with IC_{50} 's ≤ 40 pM
- No predicted activity in ADCC/CDC
- $K_D \leq 1$ pM for cynomolgus monkey IL-23
- No cross reactivity to mouse or rat IL-23
- 30 • Inhibits human IL-23 induced IL-17 and IL-22 production in mouse ear ($\geq 80\%$ inhibition of both cytokines at 1 mg/kg)
- Stability 83°C (melting temperature 83°C as determined by differential scanning calorimetry)
- Solubility ≥ 100 mg/ml (as measured by UV spectroscopy and monitored by turbidity).

[0150] In a further aspect, a humanized antibody of the present disclosure has at least one of the following binding properties (properties A). In a further aspect, a humanized anti-IL-23p19 antibody of the present disclosure has any combination of at least two, or at least 3, of the properties below. In a further aspect, a humanized anti-IL-23p19 antibody of the present disclosure has all the properties below.

- K_D for human IL-23 ≤ 1 pM (no shift in binding on-rate in 50% human serum)
- 40 • No binding to human IL-12
- $K_D \leq 1$ pM for cynomolgus monkey IL-23
- 45 • No cross reactivity to mouse or rat IL-23.

[0151] In particular, a humanized antibody of the present disclosure has a K_D for human IL-23 ≤ 1 pM (no shift in binding on-rate in 50% human serum) and no binding to human IL-12.

[0152] In a further aspect, a humanized antibody of the present disclosure has at least one of the following functional properties (properties B). In a further aspect, a humanized anti-IL-23p19 antibody of the present disclosure has any combination of at least two, or at least 3, of the properties below. In a further aspect, a humanized anti-IL-23p19 antibody of the present disclosure has all the properties below.

- Blocks IL-23 binding to human IL-23R/Fc in vitro
- 55 • Inhibits human IL-23 induced IL-17 production in mouse splenocytes with IC_{50} 's ≤ 20 pM
- Inhibits human IL-23 induced STAT3 phosphorylation in human DB cells with IC_{56} 's ≤ 40 pM
- Inhibits human IL-23 induced IL-17 and IL-22 production in mouse ear ($\geq 80\%$ inhibition of both cytokines at 1 mg/kg).

[0153] In a further aspect, a humanized antibody of the present disclosure has at least one of the following properties (properties C). In a further aspect, a humanized anti-IL-23p19 antibody of the present disclosure has any combination of at least two, or at least 3, of the properties below. In a further aspect, a humanized Anti-IL-23019 antibody of the present invention has all the properties below.

- 5 • No predicted activity in ADCC/CDC
- Stability 83°C (melting temperature 83°C as determined by differential scanning calorimetry)
- Solubility ≥ 100 mg/ml (as measured by UV spectroscopy and monitored by turbidity)
- Subcutaneous administration of 1.0 mg/kg in three cynomolgus monkeys shows sustained ≥ 10 nM exposure for approximately 28 days with a bioavailability of approximately 70%.

[0154] In a further aspect, a humanized antibody of the present disclosure has at least one of the following properties (properties C). In a further aspect, a humanized anti-IL-23p19 antibody of the present disclosure has any combination of at least two of the properties below. In a further aspect, a humanized anti-IL-23p19 antibody of the present invention has all the properties below.

- 10 • No predicted activity in ADCC/CDC
- Stability 83°C (melting temperature 83°C as determined by differential scanning calorimetry)
- Solubility ≥ 100 mg/ml (as measured by UV spectroscopy and monitored by turbidity).

[0155] In a further aspect, a humanized antibody of the present disclosure has at least one property A, at least one property B and at least one property C. In a further aspect, a humanized anti-IL-23p19 antibody of the present invention has any combination of at least two, or at least 3, of the properties A, B, and C.

[0156] In some aspects, the humanized antibody displays blocking activity, whereby it decreases the binding of IL-23 to IL-23 receptor by at least 45%, by at least 50%; by at least 55%, by at least 60%, by at least 65%, by at least 70%, by at least 75%, by at least 80%, by at least 85%, by at least 90%, or by at least 95%. The ability of an antibody to block binding of IL-23 to the IL-23 receptor can be measured using competitive binding assays known in the art. Alternatively, the blocking activity of an antibody can be measured by assessing the biological effects of IL-23, such as the production of IL-17 and IL-22 to determine if signaling mediated by the IL-23 receptor is inhibited.

[0157] In a further aspect, the present invention provides a humanized anti-IL-23p19 antibody having favorable biophysical properties. In one aspect, a humanized anti-IL-23p19 antibody of the present invention is present in at least 90% monomer form, or in at least 92% monomer form, or in at least 95% monomer form in a buffer. In a further aspect, a humanized anti-IL-23p19 antibody of the present invention remains in at least 90% monomer form, or in at least 92% monomer form, or in at least 95% monomer form in a buffer for one month or for four months.

[0158] In one aspect, a humanized antibody of the present invention is Antibody A, Antibody B, Antibody C or Antibody D. Accordingly, in one embodiment, a humanized antibody of the present invention comprises the light chain sequence of SEQ ID NO:174 and the heavy chain sequence of SEQ ID NO:176 (Antibody A). In another embodiment, a humanized antibody of the present invention comprises the light chain sequence of SEQ ID NO:174 and the heavy chain sequence of SEQ ID NO:178 (Antibody B). In another embodiment, a humanized antibody of the present invention comprises the light chain sequence of SEQ ID NO:180 and the heavy chain sequence of SEQ ID NO:176 (Antibody C). In another embodiment, a humanized antibody of the present invention comprises the light chain sequence of SEQ ID NO:180 and the heavy chain sequence of SEQ ID NO:178 (Antibody D).

[0159] In a further embodiment, a humanized antibody of the present invention consists of the light chain sequence of SEQ ID NO:174 and the heavy chain sequence of SEQ ID NO:176 (Antibody A). In a further embodiment, a humanized antibody of the present invention consists of the light chain sequence of SEQ ID NO:174 and the heavy chain sequence of SEQ ID NO:178 (Antibody B). In a further embodiment, a humanized antibody of the present invention consists of the light chain sequence of SEQ ID NO:180 and the heavy chain sequence of SEQ ID NO:176 (Antibody C). In a further embodiment, a humanized antibody of the present invention consists of the light chain sequence of SEQ ID NO:180 and the heavy chain sequence of SEQ ID NO:178 (Antibody D).

[0160] In some embodiments, the humanized anti-IL-23p19 antibodies, including antigen-binding fragments thereof, such as heavy and light chain variable regions, comprise an amino acid sequence of the residues derived from Antibody A (light chain sequence = SEQ ID NO:174; heavy chain sequence = SEQ ID NO:176), Antibody B (light chain sequence = SEQ ID NO:174; heavy chain sequence = SEQ ID NO:178), Antibody C (light chain sequence = SEQ ID NO:180; heavy chain sequence = (SEQ ID NO:176) or Antibody D (light chain sequence = SEQ ID NO:180; heavy chain sequence = SEQ ID NO:178).

[0161] In a further embodiment, the present disclosure provides an anti-IL-23p19 antibody or an antigen-binding fragment thereof that binds to human IL-23p19 at an epitope consisting of amino acid residues 108 to 126 and amino acid residues 137 to 151 of SEQ ID NO: 181.

[0162] In a further embodiment, the present disclosure provides an anti-IL-23p19 antibody or antigen-binding fragment thereof that competitively binds to human IL-23p19 with an antibody of the present invention, for example Antibody A, Antibody B, Antibody C or Antibody D described herein. The ability of an antibody or antigen-binding fragment to competitively bind to IL-23p19 can be measured using competitive binding assays known in the art.

5 [0163] The humanized anti-IL-23p19 antibodies optionally include specific amino acid substitutions in the consensus or germline framework regions. The specific substitution of amino acid residues in these framework positions can improve various aspects of antibody performance including binding affinity and/or stability, over that demonstrated in humanized antibodies formed by "direct swap" of CDRs or HVLs into the human germline framework regions.

10 [0164] In some embodiments, the present disclosure describes other monoclonal antibodies with a light chain variable region having the amino acid sequence set forth in of SEQ ID NO: 84, 86, 88, 90, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117 or 119. In some embodiments, the present invention describes other monoclonal antibodies with a heavy chain variable region having the amino acid sequence set forth in of SEQ ID NO: 121, 123, 125, 127, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154 or 156 (see Tables 1 and 2 above). The CDR sequence of these mouse antibodies are shown in Tables 3 and 4. Placing such CDRs into FRs of the human consensus heavy and light chain variable domains will yield useful humanized antibodies of the present invention.

15 [0165] In particular, the present disclosure provides monoclonal antibodies with the combinations of light chain variable and heavy chain variable regions of SEQ ID NO:84/121, 86/123, 88/125, 90/127, 91/128, 93/130, 95/132, 97/134, 99/136, 101/138, 103/140, 105/142, 107/144, 109/146, 111/148, 113/150, 115/152, 117/154 or 119/156. Such variable regions can be combined with human constant regions.

20 [0166] In some embodiments, the present disclosure describes other humanized antibodies with light chain variable region sequences having the amino acid sequence set forth in of SEQ ID NO:158, 160, 162 or 164. In some embodiments, the present disclosure describes other humanized antibodies with heavy chain variable region sequences having the amino acid sequence set forth in of SEQ ID NO:166, 168, 170 or 172 (see Tables 5 and 6 above). The CDR sequences of these antibodies are shown in Tables 3 and 4. In particular, the present invention provides monoclonal antibodies with the combinations of light chain variable and heavy chain variable regions of SEQ ID NO: 160/166, 160/168, 158/166 or 158/168. Such variable regions can be combined with human constant regions.

25 [0167] In a further embodiment, the present invention relates to an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising a humanized light chain variable domain comprising the CDRs of SEQ ID NO:160 and framework regions having an amino acid sequence at least 90% identical, at least 93% identical or at least 95% identical to the amino acid sequence of the framework regions of the variable domain light chain amino acid sequence of SEQ ID NO:160 and a humanized heavy chain variable domain comprising the CDRs of SEQ ID NO:166 and framework regions having an amino acid sequence at least 90% identical, at least 93% identical or at least 95% identical to the amino acid sequence of the framework regions of the variable domain heavy chain amino acid sequence of SEQ ID NO:166. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal antibody.

30 [0168] In a further embodiment, the present invention relates to an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising a humanized light chain variable domain comprising the CDRs of SEQ ID NO:160 and framework regions having an amino acid sequence at least 90% identical, at least 93% identical or at least 95% identical to the amino acid sequence of the framework regions of the variable domain light chain amino acid sequence of SEQ ID NO:160 and a humanized heavy chain variable domain comprising the CDRs of SEQ ID NO:168 and framework regions having an amino acid sequence at least 90% identical, at least 93% identical or at least 95% identical to the amino acid sequence of the framework regions of the variable domain heavy chain amino acid sequence of SEQ ID NO:168. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal antibody.

35 [0169] In a further embodiment, the present invention relates to an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising a humanized light chain variable domain comprising the CDRs of SEQ ID NO:158 and framework regions having an amino acid sequence at least 90% identical, at least 93% identical or at least 95% identical to the amino acid sequence of the framework regions of the variable domain light chain amino acid sequence of SEQ ID NO:158 and a humanized heavy chain variable domain comprising the CDRs of SEQ ID NO:166 and framework regions having an amino acid sequence at least 90% identical, at least 93% identical or at least 95% identical to the amino acid sequence of the framework regions of the variable domain heavy chain amino acid sequence of SEQ ID NO:166. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal antibody.

40 [0170] In a further embodiment, the present invention relates to an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising a humanized light chain variable domain comprising the CDRs of SEQ ID NO:158 and framework regions having an amino acid sequence at least 90% identical, at least 93% identical or at least 95% identical to the amino acid sequence of the framework regions of the variable domain light chain amino acid sequence of SEQ ID NO:158 and a humanized heavy chain variable domain comprising the CDRs of SEQ ID NO:168 and framework regions having an amino acid sequence at least 90% identical, at least 93% identical or at least 95% identical to the amino acid sequence of the framework regions of the variable domain heavy chain amino acid sequence of SEQ ID NO:168. In one embodiment, the anti-IL-23p19 antibody is a humanized monoclonal antibody.

[0171] In some specific embodiments, the humanized anti-IL-23p19 antibodies disclosed herein comprise at least a heavy or a light chain variable domain comprising the CDRs or HVLS of the murine monoclonal antibodies or humanized antibodies as shown in Tables 1 through 6 above and the FRs of the human germline heavy and light chain variable domains.

[0172] The CDRs of these sequences are shown in Tables 3 and 4. Accordingly, in one aspect, the present disclosure provides an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising a light chain CDR1 (L-CDR1) sequence of SEQ ID NO:1, 4, 6, 7, 8, 11, 15, 18, 19, 22, 27 or 30; a light chain CDR2 (L-CDR2) sequence of SEQ ID NO:2, 5, 9, 12, 16, 20, 23, 25, 28 or 31; a light chain CDR3 (L-CDR3) sequence of SEQ ID NO:3, 10, 13, 14, 17, 21, 24, 26, 29, or 32; a heavy chain CDR1 (H-CDR1) sequence of SEQ ID NO:33, 36, 38, 40, 43, 45, 48, 51, 54, 57, 60, 63, 66, 67, 68, 69, 77 or 80; a heavy chain CDR2 (H-CDR2) sequence of SEQ ID NO:34, 39, 41, 46, 49, 52, 55, 58, 61, 64, 70, 72, 73, 75, 78 or 81; and a heavy chain CDR3 (H-CDR3) sequence of SEQ ID NO:35, 37, 42, 44, 47, 50, 53, 56, 59, 62, 65, 71, 74, 76, 79 or 82. In one aspect, the anti-IL-23p19 antibody or antigen-binding fragment thereof comprises a light chain variable region comprising a L-CDR1 listed above, a L-CDR2 listed above and a L-CDR₃ listed above, and a heavy chain variable region comprising a H-CDR1 listed above, a H-CDR2 listed above and a H-CDR3 listed above.

[0173] In a further aspect, the present disclosure provides an anti-IL-23p19 antibody or antigen-binding fragment thereof comprising:

- a) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:1, 2, 3, 33, 34, and 35, respectively; or
- b) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:4, 5, 3, 36, 34 and 37, respectively; or
- c) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:1, 2, 3, 38, 39 and 35, respectively; or
- d) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:6, 2, 3, 40, 41 and 42, respectively; or
- e) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:7, 2, 3, 43, 41 and 44, respectively; or
- f) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:8, 9, 10, 45, 46 and 47, respectively; or
- g) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:8, 9, 10, 48, 49 and 50, respectively; or,
- h) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:11, 12, 13, 51, 52 and 53, respectively; or
- i) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:7, 2, 14, 54, 55 and 56, respectively; or
- j) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:15, 16, 17, 57, 58 and 59, respectively; or
- k) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:18, 16, 17, 60, 61 and 62, respectively; or
- l) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:19, 20, 21, 63, 66, 67 or 68, 64 and 65, respectively;
- or
- m) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:22, 23, 24, 69, 70 and 71, respectively; or
- n) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:22, 25, 26, 55, 72 and 71, respectively; or
- o) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:8, 9, 10, 45, 73 and 74, respectively; or
- p) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:27, 28, 29, 45, 75 and 76, respectively; or
- q) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:8, 9, 10, 77, 78 and 79, respectively; or
- r) a L-CDR1, a L-CDR2, a L-CDR3, a H-CDR1, a H-CDR2 and a H-CDR3 sequence of SEQ ID NO:30, 31, 32, 80, 81 and 82, respectively.

[0174] In one aspect, the anti-IL-23p19 antibody or antigen-binding fragment thereof comprises a light chain variable region comprising a L-CDR1, L-CDR2 and L-CDR3 combination listed above, and a heavy chain variable region comprising a H-CDR1, H-CDR2 and H-CDR3 combination listed above.

[0175] In specific embodiments, it is contemplated that chimeric antibodies with switched CDR regions (i.e., for example switching one or two CDRs of one of the mouse antibodies or humanized antibody derived therefrom with the analogous CDR from another mouse antibody or humanized antibody derived therefrom) between these exemplary immunoglobulins may yield useful antibodies.

5 **[0176]** In certain embodiments, the humanized anti-IL-23p19 antibody is an antibody fragment. Various antibody fragments have been generally discussed above and there are techniques that have been developed for the production of antibody fragments. Fragments can be derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto et al., 1992, Journal of Biochemical and Biophysical Methods 24:107-117; and Brennan et al., 1985, Science 229:81). Alternatively, the fragments can be produced directly in recombinant host cells. For example, Fab'-SH fragments can be directly recovered from E. coli and chemically coupled to form F(ab')₂ fragments (see, e.g., Carter et al., 1992, Bio/Technology 10:163-167). By another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. Accordingly, in one aspect, the present invention provides antibody fragments comprising the CDRs described herein, in particular one of the combinations of L-CDR1, L-CDR2, L-CDR3, H-CDR1, H-CDR2 and H-CDR3 described herein. In a further aspect, the present disclosure provides antibody fragments comprising the variable regions described herein, for example one of the combinations of light chain variable regions and heavy chain variable regions described herein.

10 **[0177]** Certain embodiments include an F(ab')₂ fragment of a humanized anti-IL-23p19 antibody comprise a light chain sequence of any of SEQ ID NO: 174 or 180 in combination with a heavy chain sequence of SEQ ID NO: 176 or 178. Such embodiments can include an intact antibody comprising such an F(ab')₂.

15 **[0178]** In some embodiments, the antibody or antibody fragment includes a constant region that mediates effector function. The constant region can provide antibody-dependent cellular cytotoxicity (ADCC), antibody-dependent cellular phagocytosis (ADCP) and/or complement-dependent cytotoxicity (CDC) responses against an IL-23 expressing target cell. The effector domain(s) can be, for example, an Fc region of an Ig molecule.

20 **[0179]** The effector domain of an antibody can be from any suitable vertebrate animal species and isotypes. The isotypes from different animal species differ in the abilities to mediate effector functions. For example, the ability of human immunoglobulin to mediate CDC and ADCC/ADCP is generally in the order of IgM≈IgG₁≈IgG₃>IgG₂>IgG₄ and IgG₁≈IgG₃>IgG₂/IgM/IgG₄, respectively. Murine immunoglobulins mediate CDC and ADCC/ADCP generally in the order of murine IgM≈IgG₃>>IgG_{2b}>IgG_{2a}>>IgG₁ and IgG_{2b}>IgG_{2a}>IgG₁>>IgG₃, respectively. In another example, murine IgG_{2a} mediates ADCC while both murine IgG_{2a} and IgM mediate CDC.

25 **[0180]** The humanized anti-IL-23p19 antibodies and agents can include modifications of the humanized anti-IL-23p19 antibody or antigen-binding fragment thereof. For example, it may be desirable to modify the antibody with respect to effector function, so as to enhance the effectiveness of the antibody in treating cancer. One such modification is the introduction of cysteine residue(s) into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated can have improved internalization capability and/or increased complement-mediated cell killing and/or antibody-dependent cellular cytotoxicity (ADCC). See, for example, Caron et al., 1992, J. Exp Med. 176:1191-1195; and Shopes, 1992, J. Immunol. 148:2918-2922. Homodimeric antibodies having enhanced anti-tumor activity can also be prepared using heterobifunctional cross-linkers as described in Wolff et al., 1993, Cancer Research 53: 2560-2565. Alternatively, an antibody can be engineered to contain dual Fc regions, enhancing complement lysis and ADCC capabilities of the antibody. See Stevenson et al., 1989, Anti-Cancer Drug Design 3: 219-230.

30 **[0181]** Antibodies with improved ability to support ADCC have been generated by modifying the glycosylation pattern of their Fc region. This is possible, since antibody glycosylation at the asparagine residue, N297, in the C_{H2} domain is involved in the interaction between IgG and Fc_Y receptors prerequisite to ADCC. Host cell lines have been engineered to express antibodies with altered glycosylation, such as increased bisecting N-acetylglucosamine or reduced fucose. Fucose reduction provides greater enhancement to ADCC activity than does increasing the presence of bisecting N-acetylglucosamine. Moreover, enhancement of ADCC by low fucose antibodies is independent of the Fc_YR_{IIIA} V/F polymorphism.

Antibody Modifications

35 **[0182]** Modifying the amino acid sequence of the Fc region of antibodies is an alternative to glycosylation engineering to enhance ADCC. The binding site on human IgG₁ for Fc_Y receptors has been determined by extensive mutational analysis. This led to the generation of humanized IgG₁ antibodies with Fc mutations that increase the binding affinity for Fc_YR_{IIIA} and enhance ADCC in vitro. Additionally, Fc variants have been obtained with many different permutations of binding properties, e.g., improved binding to specific Fc_YR receptors with unchanged or diminished binding to other Fc_YR receptors.

40 **[0183]** Another aspect includes immunoconjugates comprising the humanized antibody or fragments thereof conjugated to a cytotoxic agent such as a chemotherapeutic agent, a toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

[0184] Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used to form useful immunoconjugates include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolaca americana proteins (PAPI, PAPII, and PAP-S), Momordica charantia inhibitor, curcin, crotin, Sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, the tricothecenes, and the like. A variety of radionuclides are available for the production of radioconjugated humanized anti-IL-23p19 antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

[0185] Conjugates of the humanized anti-IL-23p19 antibody and cytotoxic or chemotherapeutic agent can be made by known methods, using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis-(p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., 1987, *Science* 238:1098. Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. Conjugates also can be formed with a cleavable linker.

[0186] The humanized anti-IL-23p19 antibodies disclosed herein can also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., 1985, *Proc. Natl. Acad. Sci. USA* 82:3688; Hwang et al., 1980, *Proc. Natl. Acad. Sci. USA* 77:4030; and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes having enhanced circulation time are disclosed, for example, in U.S. Pat. No. 5,013,556.

[0187] Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of an antibody disclosed herein can be conjugated to the liposomes as described in Martin et al., 1982, *J. Biol. Chem.* 257:286-288 via a disulfide interchange reaction. A chemotherapeutic agent (such as doxorubicin) is optionally contained within the liposome. See, e.g., Gabizon et al., 1989, *J. National Cancer Inst.* 81 (19):1484.

[0188] The antibodies described and disclosed herein can also be used in ADEPT (Antibody-Directed Enzyme Prodrug Therapy) procedures by conjugating the antibody to a prodrug-activating enzyme that converts a prodrug (e.g., a peptidyl chemotherapeutic agent), to an active anti-cancer drug. See, for example, WO 81/01145, WO 88/07378, and U.S. Pat. No. 4,975,278. The enzyme component of the immunoconjugate useful for ADEPT is an enzyme capable of acting on a prodrug in such a way so as to convert it into its more active, cytotoxic form. Specific enzymes that are useful in ADEPT include, but are not limited to, alkaline phosphatase for converting phosphate-containing prodrugs into free drugs; aryl-sulfatase for converting sulfate-containing prodrugs into free drugs; cytosine deaminase for converting non-toxic 5-fluorocytosine into the anti-cancer drug, 5-fluorouracil; proteases, such as serratia protease, thermolysin, subtilisin, carboxypeptidases, and cathepsins (such as cathepsins B and L), for converting peptide-containing prodrugs into free drugs; D-alanylcarboxypeptidases, for converting prodrugs containing D-amino acid substituents; carbohydrate-cleaving enzymes such as β -galactosidase and neuraminidase for converting glycosylated prodrugs into free drugs; β -lactamase for converting drugs derivatized with β -lactams into free drugs; and penicillin amidases, such as penicillin V amidase or penicillin G amidase, for converting drugs derivatized at their amine nitrogens with phenoxyacetyl or phenylacetyl groups, respectively, into free drugs. Alternatively, antibodies having enzymatic activity ("abzymes") can be used to convert the prodrugs into free active drugs (see, for example, Massey, 1987, *Nature* 328: 457-458). Antibody-abzyme conjugates can be prepared by known methods for delivery of the abzyme to a tumor cell population, for example, by covalently binding the enzyme to the humanized anti-IL-23p19 antibody/heterobifunctional crosslinking reagents discussed above. Alternatively, fusion proteins comprising at least the antigen binding region of an antibody disclosed herein linked to at least a functionally active portion of an enzyme as described above can be constructed using recombinant DNA techniques (see, e.g., Neuberger et al., 1984, *Nature* 312:604-608).

[0189] In certain embodiments, it may be desirable to use a humanized anti-IL-23p19 antibody fragment, rather than an intact antibody, to increase tissue penetration, for example. It may be desirable to modify the antibody fragment in order to increase its serum half life. This can be achieved, for example, by incorporation of a salvage receptor binding epitope into the antibody fragment. In one method, the appropriate region of the antibody fragment can be altered (e.g., mutated), or the epitope can be incorporated into a peptide tag that is then fused to the antibody fragment at either end or in the middle, for example, by DNA or peptide synthesis. See, e.g., WO 96/32478.

[0190] In other embodiments, covalent modifications of the humanized anti-IL-23p19 antibody are also included. Covalent modifications include modification of cysteinyl residues, histidyl residues, lysinyl and amino-terminal residues, arginyl residues, tyrosyl residues, carboxyl side groups (aspartyl or glutamyl), glutaminyl and asparaginyl residues, or seryl, or threonyl residues. Another type of covalent modification involves chemically or enzymatically coupling glycosides to the antibody. Such modifications may be made by chemical synthesis or by enzymatic or chemical cleavage of the

antibody, if applicable Other types of covalent modifications of the antibody can be introduced into the molecule by reacting targeted amino acid residues of the antibody with an organic derivatizing agent that is capable of reacting with selected side chains or the amino- or carboxy-terminal residues.

[0191] Removal of any carbohydrate moieties present on the antibody can be accomplished chemically or enzymatically. Chemical deglycosylation is described by Hakimuddin et al., 1987, Arch. Biochem. Biophys. 259:52 and by Edge et al., 1981, Anal. Biochem., 118:131. Enzymatic cleavage of carbohydrate moieties on antibodies can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., 1987, Meth. Enzymol 138:350.

[0192] Another type of useful covalent modification comprises linking the antibody to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, or polyoxyalkylenes, in the manner set forth in one or more of U.S. Pat. No. 4,640,835, U.S. Pat. No. 4,496,689, U.S. Pat. No. 4,301,144, U.S. Pat. No. 4,670,417, U.S. Pat. No. 4,791,192 and U.S. Pat. No. 4,179,337.

Humanization and Amino Acid Sequence Variants

[0193] Amino acid sequence variants of the anti-IL-23p19 antibody can be prepared by introducing appropriate nucleotide changes into the anti-IL-23p19 antibody DNA, or by peptide synthesis. Such variants include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the amino acid sequences of the anti-IL-23p19 antibodies of the examples herein. Any combination of deletions, insertions, and substitutions is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the humanized or variant anti-IL-23p19 antibody, such as changing the number or position of glycosylation sites.

[0194] A useful method for identification of certain residues or regions of the anti-IL-23p19 antibody that are preferred locations for mutagenesis is called "alanine scanning mutagenesis," as described by Cunningham and Wells (Science, 244:1081-1085 (1989)). Here, a residue or group of target residues are identified (e.g., charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (typically alanine) to affect the interaction of the amino acids with IL-23p19 antigen. Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation per se need not be predetermined. For example, to analyze the performance of a mutation at a given site, alanine scanning or random mutagenesis is conducted at the target codon or region and the expressed anti-IL-23p19 antibody variants are screened for the desired activity.

[0195] Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides-containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an anti-IL-23p19 antibody fused to an epitope tag. Other insertional variants of the anti-IL-23p19 antibody molecule include a fusion to the N- or C-terminus of the anti-IL-23p19 antibody of an enzyme or a polypeptide which increases the serum half-life of the antibody.

[0196] Another type of variant is an amino acid substitution variant. These variants have at least one amino acid residue in the anti-IL-23p19 antibody molecule removed and a different residue inserted in its place. The sites of greatest interest for substitutional mutagenesis include the hypervariable regions, but FR alterations are also contemplated. Conservative substitutions are shown in Table 5 under the heading of "preferred substitutions". If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions", or as further described below in reference to amino acid classes, may be introduced and the products screened.

TABLE 8:

	Original Residue	Exemplary Substitutions	Preferred Substitutions
45	Ala (A)		val; leu; ile val
	Arg (R)		lys; gln; asn lys
	Asn (N)		gln; his; asp, lys; arg
		gln	
50	Asp (D)		glu; asn glu
	Cys (C)		ser; ala ser
	Gln (Q)		asn; glu asn
	Glu (E)		asp; gln asp
55	Gly (G)		ala ala
	His (H)		arg; asn; gln; lys;
		arg	

(continued)

	Original Residue	Exemplary Substitutions	Preferred Substitutions
5	Ile (I) phe; norleucine leu		leu; val; met; ala;
	Leu (L) met; ala; phe	ile	ile; norleucine; val;
10	Lys (K)		arg; gln; asn arg
	Met (M)		leu; phe; ile leu
	Phe (F)		tyr; leu; val; ile; ala;
		tyr	
15	Pro (P)		ala ala
	Ser (S)		thr thr
	Thr (T)		ser ser
	Trp (W)		tyr; phe tyr
	Tyr (Y)		phe; trp; thr; ser
		phe	
20	Val (V)		leu; ile; met; phe
	ala; norleucine;	leu	

[0197] In protein chemistry, it is generally accepted that the biological properties of the antibody can be accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

[0198] Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

[0199] Any cysteine residue not involved in maintaining the proper conformation of the humanized or variant anti-IL-23p19 antibody also may be substituted, generally with serine, to improve the oxidative stability of the molecule, prevent aberrant crosslinking, or provide for established points of conjugation to a cytotoxic or cytostatic compound. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

[0200] A type of substitutional variant involves, substituting one or more hypervariable region residues of a parent antibody (e.g., a humanized or human antibody). Generally, the resulting variant(s) selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants is affinity maturation using phage display. Briefly, several hypervariable region sites (e.g., 6-7 sites) are mutated to generate all possible amino substitutions at each site. The antibody variants thus generated are displayed in a monovalent fashion from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (e.g., binding affinity). In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or in addition, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the antibody and human IL-23p19. Such contact residues and neighboring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected

to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

[0201] Another type of amino acid variant of the antibody alters the original glycosylation pattern of the antibody. By "altering" is meant deleting one or more carbohydrate moieties found in the antibody, and/or adding one or more glycosylation sites that are not present in the antibody.

[0202] In some embodiments, it may be desirable to modify the antibodies of the invention to add glycosylation sites. Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylglucosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used. Thus, in order to glycosylate a given protein, e.g., an antibody, the amino acid sequence of the protein is engineered to contain one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

[0203] Nucleic acid molecules encoding amino acid sequence variants of the anti-IL-23p19 antibody are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the anti-IL-23p19 antibody.

Polynucleotides, Vectors, Host Cells, and Recombinant Methods

[0204] Other embodiments encompass isolated polynucleotides that comprise a sequence encoding a humanized anti-IL-23p19 antibody, vectors, and host cells comprising the polynucleotides, and recombinant techniques for production of the humanized antibody. The isolated polynucleotides can encode any desired form of the anti-IL-23p19 antibody including, for example, full length monoclonal antibodies, Fab, Fab', F(ab')₂, and Fv fragments, diabodies, linear antibodies, single-chain antibody molecules, and multispecific antibodies formed from antibody fragments.

[0205] Some embodiments include isolated polynucleotides comprising sequences that encode the light chain variable region of an antibody or antibody fragment having the amino acid sequence of any of SEQ ID NO: SEQ ID NO:84, 86, 88, 90, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117 or 119. Exemplary polynucleotide sequences encoding such amino acid sequences are SEQ ID NO:83, 85, 87, 89, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116 and 118. Some embodiments include isolated polynucleotides comprising sequences that encode the heavy chain variable region of an antibody or antibody fragment having the amino acid sequence of SEQ ID NO:121, 123, 125, 127, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154 or 156. Exemplary polynucleotide sequences encoding such amino acid sequences are SEQ ID NO: 120, 122, 124, 126, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153 or 155.

[0206] Some embodiments include isolated polynucleotides comprising sequences that encode the light chain variable region of an antibody or antibody fragment having the amino acid sequence of any of SEQ ID NO:158, 160, 162 or 164. Exemplary polynucleotide sequences encoding such amino acid sequences are SEQ ID NO:157, 159, 161 or 163. Some embodiments include isolated polynucleotides comprising sequences that encode the heavy chain variable region of an antibody or antibody fragment having the amino acid sequence of SEQ ID NO: 166, 168, 170 or 172. Exemplary polynucleotide sequences encoding such amino acid sequences are SEQ ID NO: 165, 167, 169 or 171.

[0207] Some embodiments include isolated polynucleotides comprising sequences that encode the light chain of an antibody having the amino acid sequence of any of SEQ ID NO:174 or 180. Exemplary polynucleotide sequences encoding such amino acid sequences are SEQ ID NO:173 or 179. Some embodiments include isolated polynucleotides comprising sequences that encode the heavy chain of an antibody having the amino acid sequence of SEQ ID NO:176 or 178. Exemplary polynucleotide sequences encoding such amino acid sequences are SEQ ID NO:175 or 177.

[0208] In one aspect, the isolated polynucleotide sequence(s) encodes an antibody or antibody fragment having a light chain and a heavy chain variable region comprising the amino acid sequences of SEQ ID NO:174 and SEQ ID NO:176, respectively; SEQ ID NO:174 and SEQ ID NO:178, respectively; SEQ ID NO:180 and SEQ ID NO:176, respectively; SEQ ID NO:180 and SEQ ID NO:178, respectively. Exemplary polynucleotide sequences encoding such amino acid sequences are SEQ ID NO: 173 and 175, respectively, SEQ ID NO: 173 and 177, respectively, SEQ ID NO: 179 and 175, respectively, SEQ ID NO: 179 and 177, respectively.

[0209] The polynucleotide(s) that comprise a sequence encoding a humanized anti-IL-23p19 antibody or a fragment or chain thereof can be fused to one or more regulatory or control sequence, as known in the art, and can be contained in suitable expression vectors or host cell as known in the art. Each of the polynucleotide molecules encoding the heavy

or light chain variable domains can be independently fused to a polynucleotide sequence encoding a constant domain, such as a human constant domain, enabling the production of intact antibodies. Alternatively, polynucleotides, or portions thereof, can be fused together, providing a template for production of a single chain antibody.

[0210] For recombinant production, a polynucleotide encoding the antibody is inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Many suitable vectors for expressing the recombinant antibody are available. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

[0211] The humanized anti-IL-23p19 antibodies can also be produced as fusion polypeptides, in which the antibody is fused with a heterologous polypeptide, such as a signal sequence or other polypeptide having a specific cleavage site at the amino terminus of the mature protein or polypeptide. The heterologous signal sequence selected is typically one that is recognized and processed (i.e., cleaved by a signal peptidase) by the host cell. For prokaryotic host cells that do not recognize and process the humanized anti-IL-23p19 antibody signal sequence, the signal sequence can be substituted by a prokaryotic signal sequence. The signal sequence can be, for example, alkaline phosphatase, penicillinase, lipoprotein, heat-stable enterotoxin II leaders, and the like. For yeast secretion, the native signal sequence can be substituted, for example, with a leader sequence obtained from yeast invertase alpha-factor (including *Saccharomyces* and *Kluyveromyces* α -factor leaders), acid phosphatase, *C. albicans* glucoamylase, or the signal described in WO90/13646. In mammalian cells, mammalian signal sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, can be used. The DNA for such precursor region is ligated in reading frame to DNA encoding the humanized anti-IL-23p19 antibody.

[0212] Expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2- ν . plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV, and BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

[0213] Expression and cloning vectors may contain a gene that encodes a selectable marker to facilitate identification of expression. Typical selectable marker genes encode proteins that confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, or alternatively, are complement auxotrophic deficiencies, or in other alternatives supply specific nutrients that are not present in complex media, e.g., the gene encoding D-alanine racemase for Bacilli.

[0214] One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene produce a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin, mycophenolic acid, and hygromycin. Common selectable markers for mammalian cells are those that enable the identification of cells competent to take up a nucleic acid encoding a humanized anti-IL-23p19 antibody, such as DHFR (dihydrofolate reductase), thymidine kinase, metallothionein-I and -II (such as primate metallothionein genes), adenosine deaminase, ornithine decarboxylase, and the like. Cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity (e.g., DG44).

[0215] Alternatively, host cells (particularly wild-type hosts that contain endogenous DHFR) transformed or co-transformed with DNA sequences encoding anti-IL-23p19 antibody, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3'-phosphotransferase (APH), can be selected by cell growth in medium containing a selection agent for the selectable marker such as an aminoglycosidic antibiotic, e.g., kanamycin, neomycin, or G418. See, e.g., U.S. Pat. No. 4,965,199.

[0216] Where the recombinant production is performed in a yeast cell as a host cell, the TRP1 gene present in the yeast plasmid YRp7 (Stinchcomb et al., 1979, *Nature* 282: 39) can be used as a selectable marker. The TRP1 gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 (Jones, 1977, *Genetics* 85:12). The presence of the trp1 lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan. Similarly, Leu2p-deficient yeast strains such as ATCC 20,622 and 38,626 are complemented by known plasmids bearing the LEU2 gene.

[0217] In addition, vectors derived from the 1.6 μ m circular plasmid pKD1 can be used for transformation of *Kluyveromyces* yeasts. Alternatively, an expression system for large-scale production of recombinant calf chymosin was reported for *K. lactis* (Van den Berg, 1990, *Bio/Technology* 8:135). Stable multi-copy expression vectors for secretion of mature recombinant human serum albumin by industrial strains of *Kluyveromyces* have also been disclosed (Fleer et al., 1991, *Bio/Technology* 9:968-975).

[0218] Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the nucleic acid molecule encoding an anti-IL-23p19 antibody or polypeptide chain thereof. Promoters suitable for use with prokaryotic hosts include phoA promoter, β -lactamase and lactose promoter systems, alkaline phosphatase, tryptophan (trp) promoter system, and hybrid promoters such as the tac promoter. Other known bacterial promoters are also suitable. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the humanized anti-IL-23p19 antibody.

[0219] Many eukaryotic promoter sequences are known. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CNCAAT region where N may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

[0220] Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

[0221] Inducible promoters have the additional advantage of transcription controlled by growth conditions. These include yeast promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, derivative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657. Yeast enhancers also are advantageously used with yeast promoters.

[0222] Humanized anti-IL-23p19 antibody transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, or from heat-shock promoters, provided such promoters are compatible with the host cell systems.

[0223] The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. The immediate early promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment. A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is disclosed in U.S. Pat. No. 4,419,446. A modification of this system is described in U.S. Pat. No. 4,601,978. See also Reyes et al., 1982, Nature 297:598-601, disclosing expression of human p-interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus. Alternatively, the Rous sarcoma virus long terminal repeat can be used as the promoter.

[0224] Another useful element that can be used in a recombinant expression vector is an enhancer sequence, which is used to increase the transcription of a DNA encoding a humanized anti-IL-23p19 antibody by higher eukaryotes. Many enhancer sequences are now known from mammalian genes (e.g., globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, an enhancer from a eukaryotic cell virus is used. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, 1982, Nature 297:17-18 for a description of enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the humanized anti-IL-23p19 antibody-encoding sequence, but is preferably located at a site 5' from the promoter.

[0225] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) can also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding anti-IL-23p19 antibody. One useful transcription termination component is the bovine growth hormone polyadenylation region. See WO94/11026 and the expression vector disclosed therein. In some embodiments, humanized anti-IL-23p19 antibodies can be expressed using the CHEF system. (See, e.g., U.S. Pat. No. 5,888,809).

[0226] Suitable host cells for cloning or expressing the DNA in the vectors herein are the prokaryote, yeast, or higher eukaryote cells described above. Suitable prokaryotes for this purpose include eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as Escherichia, e.g., E. coli, Enterobacter, Erwinia, Klebsiella, Proteus, Salmonella, e.g., Salmonella typhimurium, Serratia, e.g., Serratia marcescans, and Shigella, as well as Bacilli such as B. subtilis and B. licheniformis (e.g., B. licheniformis 41 P disclosed in DD 266,710 published Apr. 12, 1989), Pseudomonas such as P. aeruginosa, and Streptomyces. One preferred E. coli cloning host is E. coli 294 (ATCC 31,446), although other strains such as E. coli B, E. coli X1776 (ATCC 31,537), and E. coli W31110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting.

[0227] In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or

expression hosts for humanized anti-IL-23p19 antibody-encoding vectors. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *Schizosaccharomyces pombe*; *Kluyveromyces* hosts such as, e.g., *K. lactis*, *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickeramii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilae* (ATCC 36,906), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070); *Candida*; *Trichoderma reesii* (EP 244,234); *Neurospora crassa*; *Schwanniomyces* such as *Schwanniomyces occidentalis* and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus* hosts such as *A. nidulans* and *A. niger*.

[0228] Suitable host cells for the expression of glycosylated humanized anti-IL-23p19 antibody are derived from multicellular organisms. Examples of invertebrate cells include plant and insect cells, including, e.g., numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* (silk worm). A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used, particularly for transfection of *Spodoptera frugiperda* cells.

[0229] Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can also be utilized as hosts.

[0230] In another aspect, expression of humanized anti-IL-23p19 is carried out in vertebrate cells. The propagation of vertebrate cells in culture (tissue culture) has become routine procedure and techniques are widely available. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651), human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, (Graham et al., 1977, J. Gen Virol. 36: 59), baby hamster kidney cells (BHK, ATCC CCL 10), Chinese hamster ovary cells/-DHFR1 (CHO, Urlaub et al., 1980, Proc. Natl. Acad. Sci. USA 77: 4216; e.g., DG44), mouse sertoli cells (TM4, Mather, 1980, Biol. Reprod. 23:243-251), monkey kidney cells (CV1 ATCC CCL 70), African green monkey kidney cells (VERO-76, ATCC CRL-1587), human cervical carcinoma cells (HELA, ATCC CCL 2), canine kidney cells (MDCK, ATCC CCL 34), buffalo rat liver cells (BRL 3A, ATCC CRL 1442), human lung cells (W138, ATCC CCL 75), human liver cells (Hep G2, HB 8065), mouse mammary tumor (MMT 060562, ATCC CCL51), TR1 cells (Mather et al., 1982, Annals N.Y. Acad. Sci. 383: 44-68), MRC 5 cells, FS4 cells, and human hepatoma line (Hep G2).

[0231] Host cells are transformed with the above-described expression or cloning vectors for humanized anti-IL-23p19 antibody production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

[0232] The host cells used to produce a humanized anti-IL-23p19 antibody described herein may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma-Aldrich Co., St. Louis, Mo.), Minimal Essential Medium ((MEM), (Sigma-Aldrich Co.), RPMI-1640 (Sigma-Aldrich Co.), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma-Aldrich. Co.) are suitable for culturing the host cells. In addition, any of the media described in one or more of Ham et al., 1979, Meth. Enz. 58: 44, Barnes et al., 1980, Anal. Biochem. 102: 255, U.S. Pat. No. 4,767,704, U.S. Pat. No. 4,657,866; U.S. Pat. No. 4,927,762, U.S. Pat. No. 4,560,655, U.S. Pat. No. 5,122,469, WO 90/103430, and WO 87/00195 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as gentamicin), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Other supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

[0233] When using recombinant techniques, the antibody can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the antibody is produced intracellularly, the cells may be disrupted to release protein as a first step. Particulate debris, either host cells or lysed fragments, can be removed, for example, by centrifugation or ultrafiltration. Carter et al., 1992, Bio/Technology 10:163-167 describes a procedure for isolating antibodies that are secreted to the periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 minutes. Cell debris can be removed by centrifugation. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants. A variety of methods can be used to isolate the antibody from the host cell.

[0234] The antibody composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being a typical purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglob-

ulin Fc domain that is present in the antibody. Protein A can be used to purify antibodies that are based on human gamma1, gamma2, or gamma4 heavy chains (see, e.g., Lindmark et al., 1983 J. Immunol. Meth. 62:1-13). Protein G is recommended for all mouse isotypes and for human gamma3 (see, e.g., Guss et al., 1986 EMBO J. 5:1567-1575). A matrix to which an affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrene/divinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a C_H3 domain, the Bakerbond ABX™ resin (J. T. Baker, Phillipsburg, N.J.) is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, reverse phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™ chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered.

[0235] Following any preliminary purification step(s), the mixture comprising the antibody of interest and contaminants may be subjected to low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, typically performed at low salt concentrations (e.g., from about 0-0.25M salt).

[0236] Also included are nucleic acids that hybridize under low, moderate, and high stringency conditions, as defined herein, to all or a portion (e.g., the portion encoding the variable region) of the nucleotide sequence represented by isolated polynucleotide sequence(s) that encode an antibody or antibody fragment of the present invention. The hybridizing portion of the hybridizing nucleic acid is typically at least 15 (e.g., 20, 25, 30 or 50) nucleotides in length. The hybridizing portion of the hybridizing nucleic acid is at least 80%, e.g., at least 90%, at least 95%, or at least 98%, identical to the sequence of a portion or all of a nucleic acid encoding an anti-IL-23p19 polypeptide (e.g., a heavy chain or light chain variable region), or its complement. Hybridizing nucleic acids of the type described herein can be used, for example, as a cloning probe, a primer, e.g., a PCR primer, or a diagnostic probe.

[0237] Some embodiments include isolated polynucleotides including sequences that encode an antibody or antibody fragment having the amino acid sequence of any one of SEQ ID NO: 84, 86, 88, 90, 91, 93, 95, 97; 99, 101, 103, 105, 107, 109, 111, 113, 115, 117 or 119, and that is at least 80%, at least 90%, at least 95%, at least 98%, or at least 99% identical to the polynucleotide sequences of SEQ ID NO:83, 85, 87, 89, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116 or 118.

[0238] Some embodiments include isolated polynucleotides including sequences that encode an antibody or antibody fragment having the amino acid sequence of any one of SEQ ID NO: 158, 160, 162 or 164, and that is at least 80%, at least 90%, at least 95%, at least 98%, or at least 99% identical to the polynucleotide sequences of SEQ ID NO:157, 159, 161 or 163.

[0239] Some embodiments include isolated polynucleotides including sequences that encode an antibody or antibody fragment having the amino acid sequence of any one of SEQ ID NO: 121, 123, 125, 127, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154 or 156, and that is at least 80%, at least 90%, at least 95%, at least 98%, or at least 99% identical to the polynucleotide sequences of SEQ ID NO: 120, 122, 124, 126, 129, 131, 133, 135, 137, 139, 141, 143, 145, 147, 149, 151, 153 or 155.

[0240] Some embodiments include isolated polynucleotides including sequences that encode an antibody or antibody fragment having the amino acid sequence of any one of SEQ ID NO: 166, 168, 170 or 172, and that is at least 80%, at least 90%, at least 95%, at least 98%, or at least 99% identical to the polynucleotide sequences of SEQ ID NO: 165, 167, 169 or 171.

[0241] Some embodiments include isolated polynucleotides including sequences that encode an antibody or antibody fragment having the light chain variable region amino acid sequence that is at least 80%, at least 90%, at least 95%, at least 98%, or at least 99% identical to the amino acid sequence of any one of SEQ ID NO: 84, 86, 88, 90, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117 or 119. Some embodiments include isolated polynucleotides including sequences that encode an antibody or antibody fragment having the light chain variable region amino acid sequence that is at least 80%, at least 90%, at least 95%, at least 98%, or at least 99% identical to the amino acid sequence of any one of SEQ ID NO:158; 160, 162 or 164. Some embodiments include isolated polynucleotides including sequences that encode an antibody or antibody fragment having the heavy chain variable region amino acid sequence that is at least 80%, at least 90%, at least 95%, at least 98%, or at least 99% identical to the amino acid sequence of any one of SEQ ID NO:121, 123, 125, 127, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154 or 156. Some embodiments include isolated polynucleotides including sequences that encode an antibody or antibody fragment having the heavy chain variable region amino acid sequence that is at least 80%, at least 90%, at least 95%, at least 98%, or at least 99% identical to the amino acid sequence of any one of SEQ ID NO:166, 168, 170 or 172. As used herein, the terms "identical" or "percent identity," in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned for maximum correspondence. To determine the percent identity, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The

amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity=# of identical positions/total # of positions (e.g., overlapping positions)x100). In some embodiments, the two sequences that are compared are the same length after gaps are introduced within the sequences, as appropriate (e.g., excluding additional sequence extending beyond the sequences being compared). For example, when variable region sequences are compared, the leader and/or constant domain sequences are not considered. For sequence comparisons between two sequences, a "corresponding" CDR refers to a CDR in the same location in both sequences (e.g., CDR-H1 of each sequence).

[0242] The determination of percent identity or percent similarity between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. USA 87:2264-2268, modified as in Karlin and Altschul, 1993, Proc. Natl. Acad. Sci. USA 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al., 1990, J. Mol. Biol. 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score=100, wordlength=12, to obtain nucleotide sequences homologous to a nucleic acid encoding a protein of interest. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3, to obtain amino acid sequences homologous to protein of interest. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., 1997, Nucleic Acids Res. 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (Id.). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, CABIOS (1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Additional algorithms for sequence analysis are known in the art and include ADVANCE and ADAM as described in Torellis and Robotti, 1994, Comput. Appl. Biosci. 10:3-5; and FASTA described in Pearson and Lipman, 1988, Proc. Natl. Acad. Sci. USA 85:2444-8. Within FASTA, ktup is a control option that sets the sensitivity and speed of the search. If ktup=2, similar regions in the two sequences being compared are found by looking at pairs of aligned residues; if ktup=1, single aligned amino acids are examined. ktup can be set to 2 or 1 for protein sequences, or from 1 to 6 for DNA sequences. The default if ktup is not specified is 2 for proteins and 6 for DNA. Alternatively, protein sequence alignment may be carried out using the CLUSTAL W algorithm, as described by Higgins et al., 1996, Methods Enzymol. 266:383-402.

35 Non-Therapeutic Uses

[0243] The antibodies described herein are useful as affinity purification agents. In this process, the antibodies are immobilized on a solid phase such a Protein A resin, using methods well known in the art. The immobilized antibody is contacted with a sample containing the IL-23p19 protein (or fragment thereof) to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the IL-23p19 protein, which is bound to the immobilized antibody. Finally, the support is washed with another suitable solvent that will release the IL-23p19 protein from the antibody.

[0244] Anti-IL-23p19 antibodies, for example humanized anti-IL-23p19 antibodies, are also useful in diagnostic assays to detect and/or quantify IL-23 protein, for example, detecting IL-23 expression in specific cells, tissues, or serum. The anti-IL-23p19 antibodies can be used diagnostically to, for example, monitor the development or progression of a disease as part of a clinical testing procedure to, e.g., determine the efficacy of a given treatment and/or prevention regimen. Detection can be facilitated by coupling the anti-IL-23p19 antibody. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. See, for example, U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention.

[0245] The anti-IL-23p19 antibodies can be used in methods for diagnosing an IL-23-associated disorder (e.g., a disorder characterized by abnormal expression of IL-23) or to determine if a subject has an increased risk of developing an IL-23-associated disorder. Such methods include contacting a biological sample from a subject with an IL-23p19 antibody and detecting binding of the antibody to IL-23p19. By "biological sample" is intended any biological sample obtained from an individual, cell line, tissue culture, or other source of cells potentially expressing IL-23. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art.

[0246] In some embodiments, the method can further comprise comparing the level of IL-23 in a patient sample to a

control sample (e.g., a subject that does not have an IL-23-associated disorder) to determine if the patient has an IL-23-associated disorder or is at risk of developing an IL-23-associated disorder.

[0247] It will be advantageous in some embodiments, for example, for diagnostic purposes to label the antibody with a detectable moiety. Numerous detectable labels are available, including radioisotopes, fluorescent labels, enzyme substrate labels and the like. The label may be indirectly conjugated with the antibody using various known techniques.

[0248] For example, the antibody can be conjugated with biotin and any of the three broad categories of labels mentioned above can be conjugated with avidin, or vice versa. Biotin binds selectively to avidin and thus, the label can be conjugated with the antibody in this indirect manner. Alternatively, to achieve indirect conjugation of the label with the antibody, the antibody can be conjugated with a small hapten (such as digoxin) and one of the different types of labels mentioned above is conjugated with an anti-hapten antibody (e.g., anti-digoxin antibody). Thus, indirect conjugation of the label with the antibody can be achieved.

[0249] Exemplary radioisotopes labels include ^{35}S , ^{14}C , ^{125}I , ^3H , and ^{131}I . The antibody can be labeled with the radioisotope, using the techniques described in, for example, Current Protocols in Immunology, Volumes 1 and 2, 1991, Coligan et al., Ed. Wiley-Interscience, New York, N.Y., Pubs. Radioactivity can be measured, for example, by scintillation counting.

[0250] Exemplary fluorescent labels include labels derived from rare earth chelates (europium chelates) or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, Lissamine, phycoerythrin, and Texas Red are available. The fluorescent labels can be conjugated to the antibody via known techniques, such as those disclosed in Current Protocols in Immunology, for example. Fluorescence can be quantified using a fluorimeter.

[0251] There are various, well-characterized enzyme-substrate labels known in the art (see, e.g., U.S. Pat. No. 4,275,149 for a review). The enzyme generally catalyzes a chemical alteration of the chromogenic substrate that can be measured using various techniques. For example, alteration may be a color change in a substrate that can be measured spectrophotometrically. Alternatively, the enzyme may alter the fluorescence or chemiluminescence of the substrate. Techniques for quantifying a change in fluorescence are described above. The chemiluminescent substrate becomes electronically excited by a chemical reaction and may then emit light that can be measured, using a chemiluminometer, for example, or donates energy to a fluorescent acceptor.

[0252] Examples of enzymatic labels include luciferases such as firefly luciferase and bacterial luciferase (U.S. Pat. No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, malate dehydrogenase, urease, peroxidase such as horseradish peroxidase (HRPO), alkaline phosphatase, β -galactosidase, glucoamylase, lysozyme, saccharide oxidases (such as glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase), heterocyclic oxidases (such as uricase and xanthine oxidase), lactoperoxidase, microperoxidase, and the like. Techniques for conjugating enzymes to antibodies are described, for example, in O'Sullivan et al., 1981, Methods for the Preparation of Enzyme-Antibody Conjugates for use in Enzyme immunoassay, in Methods in Enzym. (J. Langone & H. Van Vunakis, eds.), Academic press, N.Y., 73: 147-166.

[0253] Examples of enzyme-substrate combinations include, for example: Horseradish peroxidase (HRPO) with hydrogen peroxidase as a substrate, wherein the hydrogen peroxidase oxidizes a dye precursor such as orthophenylenediamine (OPD) or 3,3',5,5'-tetramethyl benzidine hydrochloride (TMB); alkaline phosphatase (AP) with para-Nitrophenyl phosphate as chromogenic substrate; and β -D-galactosidase (β -D-Gal) with a chromogenic substrate such as p-nitrophenyl- β -D-galactosidase or fluorogenic substrate 4-methylumbelliferyl- β -D-galactosidase.

[0254] Numerous other enzyme-substrate combinations are available to those skilled in the art. For a general review of these, see U.S. Pat. No. 4,275,149 and U.S. Pat. No. 4,318,980.

[0255] In another embodiment, the humanized anti-IL-23p19 antibody is used unlabeled and detected with a labeled antibody that binds the humanized anti-IL-23p19 antibody.

[0256] The antibodies described herein may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. See, e.g., Zola, Monoclonal Antibodies: A Manual of Techniques, pp. 147-158 (CRC Press, Inc. 1987).

[0257] The anti-IL-23p19 antibody or antigen binding fragment thereof can be used to inhibit the binding of IL-23 to the IL-23 receptor. Such methods comprise administering an anti-IL-23p19 antibody or antigen binding fragment thereof to a cell (e.g., a mammalian cell) or cellular environment, whereby signaling mediated by the IL-23 receptor is inhibited. These methods can be performed *in vitro* or *in vivo*. By "cellular environment" is intended the tissue, medium, or extracellular matrix surrounding a cell. The anti-IL-23p19 antibody or antigen binding fragment thereof is administered to the cellular environment of a cell in such a manner that the antibody or fragment is capable of binding to IL-23 molecules outside of and surrounding the cell, therefore, preventing the binding of IL-23 to its receptor.

55 Diagnostic Kits

[0258] An anti-IL-23p19, antibody can be used in a diagnostic kit, i.e., a packaged combination of reagents in predetermined amounts with instructions for performing the diagnostic assay. Where the antibody is labeled with an enzyme,

the kit may include substrates and cofactors required by the enzyme such as a substrate precursor that provides the detectable chromophore or fluorophore. In addition, other additives may be included such as stabilizers, buffers (for example a block buffer or lysis buffer), and the like. The relative amounts of the various reagents may be varied widely to provide for concentrations in solution of the reagents that substantially optimize the sensitivity of the assay. The reagents may be provided as dry powders, usually lyophilized, including excipients that on dissolution will provide a reagent solution having the appropriate concentration.

Therapeutic Uses

[0259] In another embodiment, a humanized anti-IL-23p19 antibody disclosed herein is useful in the treatment of various disorders associated with the expression of IL-23p19 as described herein. Methods for treating an IL-23 associated disorder comprise administering a therapeutically effective amount of a humanized anti-IL-23p19 antibody to a subject in need thereof.

[0260] The humanized anti-IL-23p19 antibody or agent is administered by any suitable means, including parenteral, subcutaneous, intraperitoneal, intrapulmonary, and intranasal, and, if desired for local immunosuppressive treatment, intralesional administration (including perfusing or otherwise contacting the graft with the antibody before transplantation). The humanized anti-IL-23p19 antibody or agent can be administered, for example, as an infusion or as a bolus. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. In addition, the humanized anti-IL-23p19 antibody is suitably administered by pulse infusion, particularly with declining doses of the antibody. In one aspect, the dosing is given by injections, most preferably intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic.

[0261] For the prevention or treatment of disease, the appropriate dosage of antibody will depend on a variety of factors such as the type of disease to be treated, as defined above, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments.

[0262] Depending on the type and severity of the disease, about 1 µg/kg to 20 mg/kg (e.g., 0.1-15 mg/kg) of antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily dosage might range from about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays. An exemplary dosing regimen is that disclosed in WO 94/04188.

[0263] The term "suppression" is used herein in the same context as "amelioration" and "alleviation" to mean a lessening of one or more characteristics of the disease.

[0264] The antibody composition will be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The "therapeutically effective amount" of the antibody to be administered will be governed by such considerations, and is the minimum amount necessary to prevent, ameliorate, or treat the disorder associated with IL-23 expression.

[0265] The antibody need not be, but is optionally, formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends on the amount of humanized anti-IL-23p19 antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as used hereinbefore or about from 1 to 99% of the heretofore employed dosages.

IL-23-Associated Disorders

[0266] The anti-IL-23p19 antibodies or agents are useful for treating or preventing an immunological disorder characterized by abnormal expression of IL-23, e.g., by inappropriate activation of immune cells (e.g., lymphocytes or dendritic cells). Such abnormal expression of IL-23 can be due to, for example, increased IL-23 protein levels. The anti-IL-23p19 antibodies or antigen binding fragments thereof also find use in the treatment or prevention of respiratory disorders, metabolic disorders, for example diabetes mellitus, and certain cancers. Treatment or prevention of the immunological disorder, respiratory disorder, metabolic disorder or cancer, according to the methods described herein, is achieved by administering to a subject in need of such treatment or prevention an effective amount of the anti-IL-23p19 antibody or agent, whereby the antibody decreases the activity of IL-23 associated with the disease state.

[0267] immunological diseases that are characterized by inappropriate activation of immune cells and that can be

treated or prevented by the methods described herein can be classified, for example, by the type(s) of hypersensitivity reaction(s) that underlie the disorder. These reactions are typically classified into four types: anaphylactic reactions, cytotoxic (cytolytic) reactions, immune complex reactions, or cell-mediated immunity (CMI) reactions (also referred to as delayed-type hypersensitivity (DTH) reactions). (See, e.g., Fundamental Immunology (William E. Paul ed., Raven Press, N.Y., 3rd ed. 1993).) Immunological diseases include inflammatory diseases and autoimmune diseases.

[0268] Specific examples of such immunological diseases include the following: rheumatoid arthritis, autoimmune demyelinative diseases (e.g., multiple sclerosis, allergic encephalomyelitis), endocrine ophthalmopathy, uveoretinitis, systemic lupus erythematosus, myasthenia gravis, Grave's disease, glomerulonephritis, autoimmune hepatological disorder, inflammatory bowel disease (e.g., Crohn's disease or ulcerative colitis), anaphylaxis, allergic reaction, Sjogren's syndrome, type I diabetes mellitus, primary biliary cirrhosis, Wegener's granulomatosis, fibromyalgia, polymyositis, dermatomyositis, inflammatory myositis, multiple endocrine failure, Schmidt's syndrome, autoimmune uveitis, Addison's disease, adrehalitis, thyroiditis, Hashimoto's thyroiditis, autoimmune thyroid disease, pernicious anemia, gastric atrophy, chronic hepatitis, lupoid hepatitis, atherosclerosis, subacute cutaneous lupus erythematosus, hypoparathyroidism, Dressler's syndrome, autoimmune thrombocytopenia, idiopathic thrombocytopenic purpura, hemolytic anemia, pemphigus vulgaris, pemphigus, dermatitis herpetiformis, alopecia arcata, pemphigoid, scleroderma, progressive systemic sclerosis, CREST syndrome (calcinosis, Raynaud's phenomenon, esophageal dysmotility, sclerodactyl, and telangiectasia), male and female autoimmune infertility, ankylosing spondylitis, ulcerative colitis, mixed connective tissue disease, polyarteritis nodosa, systemic necrotizing vasculitis, atopic dermatitis, atopic rhinitis, Goodpasture's syndrome, Chagas' disease, sarcoidosis, rheumatic fever, asthma, recurrent abortion, anti-phospholipid syndrome, farmer's lung, erythema multiforme, post cardiotomy syndrome, Cushing's syndrome, autoimmune chronic active hepatitis, bird-fancier's lung, toxic epidermal necrolysis, Alport's syndrome, alveolitis, allergic alveolitis, fibrosing, alveolitis, interstitial lung disease, erythema nodosum, pyoderma gangrenosum, transfusion reaction, Takayasu's arteritis, polymyalgia rheumatica, temporal arteritis, schistosomiasis, giant cell arteritis, ascariasis, aspergillosis, Sampter's syndrome, eczema, lymphomatoid granulomatosis, Behcet's disease, Caplan's syndrome, Kawasaki's disease, dengue, encephalomyelitis, endocarditis, endomyocardial fibrosis, endophthalmitis, erythema elevatum et diutinum, psoriasis, psoriatic arthritis, erythroblastosis fetalis, eosinophilic faciitis, Shulman's syndrome, Felty's syndrome, filariasis, cyclitis, chronic cyclitis, heterochronic cyclitis, Fuch's cyclitis, IgA nephropathy, Henoch-Schonlein purpura, graft versus host disease, transplantation rejection, cardiomyopathy, Eaton-Lambert syndrome, relapsing polychondritis, cryoglobulinemia, Waldenstrom's macroglobulemia, Evan's syndrome, acute respiratory distress syndrome, pulmonary inflammation, osteoporosis, delayed type hypersensitivity and autoimmune gonadal failure.

[0269] In some embodiments, the immunological disorder is a T cell-mediated immunological disorder and accordingly, the anti-IL-23p19 antibodies and agents as described herein are also useful for treating or preventing T cell-mediated immunological disorders.

[0270] In one aspect, the anti-IL-23p19 antibodies or agents are useful for treating or preventing a respiratory disorder in which IL-23 is abnormally expressed. Treatment or prevention of the respiratory disorder, according to the methods described herein, is achieved by administering to a subject in need of such treatment or prevention an effective amount of the anti-IL-23p19 antibody or agent, whereby the antibody decreases the activity of IL-23 associated with the disease state. These include, but are not limited to: respiratory complaints, obstructive pulmonary diseases of various origins, pulmonary emphysema of various origins, restrictive pulmonary diseases, interstitial pulmonary diseases, interstitial lung disease, cystic fibrosis, bronchitis of various origins, bronchiectasis, ARDS (adult respiratory distress syndrome) and all forms, of pulmonary oedema; obstructive pulmonary diseases selected from among COPD (chronic obstructive pulmonary disease), asthma, bronchial asthma, paediatric asthma, severe asthma, acute asthma attacks and chronic bronchitis; pulmonary emphysema which has its origins in COPD (chronic obstructive pulmonary disease) or α 1-proteinase inhibitor deficiency; restrictive pulmonary diseases selected from among allergic alveolitis, restrictive pulmonary diseases triggered by work-related noxious substances, such as asbestosis or silicosis, and restriction caused by lung tumours, such as lymphangiosis carcinomatosa, bronchoalveolar carcinoma and lymphomas; pneumonia caused by infections, such as for example infection by viruses, bacteria, fungi, protozoa, helminths or other pathogens, pneumonitis caused by various factors, such as for example aspiration and left heart insufficiency, radiation-induced pneumonitis or fibrosis, collagenoses, such as for example lupus erythematosus, systemic scleroderma or sarcoidosis, granulomatoses, such as for example Boeck's disease, idiopathic interstitial pneumonia or idiopathic pulmonary fibrosis (IPF); mucoviscidosis, bronchitis caused by bacterial or viral infection, allergic bronchitis and toxic bronchitis; bronchiectasis; pulmonary oedema, for example, toxic pulmonary oedema after aspiration or inhalation of toxic substances and foreign substances; rhinitis, arthritis and related arthropathies, psoriasis, myeloid leukemia, multiple sclerosis, Alzheimer's disease, glomerulonephritis, and chronic atopic dermatitis.

[0271] In another aspect, the anti-IL-23p19 antibodies and agents as described herein are also useful for treating cancers, in which IL-23 is abnormally expressed.

[0272] IL-23-expressing cancers that can be treated by the methods described herein include, for example, leukemia, such as acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia (e.g., myeloblastic, promyelocytic,

myelomonocytic, monocytic, or erythroleukemia), chronic leukemia, chronic myelocytic (granulocytic) leukemia, or chronic lymphocytic leukemia; Polycythemia vera; Lymphoma (e.g., Hodgkin's disease or Non-Hodgkin's disease); multiple myeloma, Waldenstrom's macroglobulinemia; heavy chain disease; solid tumors such as sarcomas and carcinomas (e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, osteosarcoma, chordoma, angiiosarcoma, endothelioma, lymphangiosarcoma, lymphangioendothelioma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, colorectal carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, uterine cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, non small cell lung carcinoma, bladder carcinoma, epithelial, carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendrogioma, meningioma, melanoma, neuroblastoma, retinoblastoma, nasopharyngeal carcinoma, or esophageal carcinoma).

15 Pharmaceutical Compositions and Administration Thereof

[0273] A composition comprising an IL-23p19 binding agent (e.g., an anti-IL-23p19 antibody) can be administered to a subject having or at risk of having an immunological disorder, respiratory disorder or a cancer. The disclosure further provides for the use of a IL-23p19 binding agent (e.g., an anti-IL-23p19 antibody) in the manufacture of a medicament for prevention or treatment of a cancer, respiratory disorder or immunological disorder. The term "subject" as used herein means any mammalian patient to which an IL-23p19binding agent can be administered, including, e.g., humans and non-human mammals, such as primates, rodents, and dogs. Subjects specifically intended for treatment using the methods described herein include humans. The antibodies or agents can be administered either alone or in combination with other compositions in the prevention or treatment of the immunological disorder, respiratory disorder or cancer. Such compositions which can be administered in combination with the antibodies or agents include methotrexate (MTX) and immunomodulators, e.g. antibodies or small molecules.

[0274] Examples of antibodies for use in such pharmaceutical compositions are those that comprise a humanized antibody or antibody fragment having the light chain variable region amino acid sequence of any of SEQ ID NO: 84, 86, 88, 90, 91, 93, 95, 97, 99, 101, 103, 105, 107, 109, 111, 113, 115, 117 or 119. Examples of antibodies for use in such pharmaceutical compositions are also those that comprise a humanized antibody or antibody fragment having the heavy chain variable region amino acid sequence of any of SEQ ID NO: 121, 123, 125, 127, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, 154 or 156.

[0275] Further examples of antibodies for use in such pharmaceutical compositions are also those that comprise a humanized antibody or antibody fragment having the light chain variable region amino acid sequence of any of SEQ ID NO:158, 160, 162 or 164. Preferred antibodies for use in such pharmaceutical compositions are also those that comprise a humanized antibody or antibody fragment having the heavy chain variable region amino acid sequence of any of SEQ ID NO:166, 168, 170 or 172.

[0276] Further examples of antibodies for use in such pharmaceutical compositions are also those that comprise a humanized antibody or antibody fragment having the light chain variable region and heavy chain variable region of any of SEQ ID NO: 160 and 166, SEQ ID NO: 160 and 168, SEQ ID NO: 158 and 166 or SEQ ID NO: 158 and 168.

[0277] Further examples of antibodies for use in such pharmaceutical compositions are also those that comprise a humanized antibody having the light chain region amino acid sequence of any of SEQ ID NO:174 or 180. Preferred antibodies for use in such pharmaceutical compositions are also those that comprise humanized antibody having the heavy chain variable region amino acid sequence of any of SEQ ID NO:176 or 178.

[0278] Further examples of antibodies for use in such pharmaceutical compositions are also those that comprise Antibody A, Antibody B, Antibody C or Antibody D.

[0279] Various delivery systems are known and can be used to administer the IL-23p19 binding agent. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intra-nasal, epidural, and oral routes. The IL-23p19 binding agent can be administered, for example by infusion, bolus or injection, and can be administered together with other biologically active agents such as chemotherapeutic agents. Administration can be systemic or local. In preferred embodiments, the administration is by subcutaneous injection. Formulations for such injections may be prepared in for example prefilled syringes that may be administered once every other week.

[0280] In specific embodiments, the IL-23p19 binding agent composition is administered by injection, by means of a catheter, by means of a suppository, or by means of An implant, the implant being of a porous, non-porous, or gelatinous material, including a membrane, such as a sialastic membrane, or a fiber. Typically, when administering the composition, materials to which the anti-IL-23p19 antibody or agent does not absorb are used.

[0281] In other embodiments, the anti-IL-23p19 antibody or agent is delivered in a controlled release system. In one

embodiment, a pump may be used (see, e.g., Langer, 1990, Science 249:1527-1533; Sefton, 1989, CRC Crit. Ref. Biomed. End. 14:201; Buchwald et al., 1980, Surgery 88:507; Saudek et al., 1989, N. Engl. J. Med. 321:574). In another embodiment, polymeric materials can be used. (See, e.g., Medical Applications of Controlled Release (Langer and Wise eds., CRC Press, Boca Raton, Fla., 1974); Controlled Drug Bioavailability, Drug Product Design and Performance (Smolen and Ball eds., Wiley, New York, 1984); Ranger and Peppas, 1983, Macromol. Sci. Rev. Macromol. Chem. 23:61. See also Levy et al., 1985, Science 228:190; During et al., 1989, Ann. Neurol. 25:351; Howard et al., 1989, J. Neurosurg. 71:105.) Other controlled release systems are discussed, for example, in Langer, *supra*.

[0282] An IL-23p19 binding agent (e.g., an anti-IL-23p19 antibody) can be administered as pharmaceutical compositions comprising a therapeutically effective amount of the binding agent and one or more pharmaceutically compatible ingredients.

[0283] In typical embodiments, the pharmaceutical composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous or subcutaneous administration to human beings. Typically, compositions for administration by injection are solutions in sterile isotonic aqueous buffer. Where necessary, the pharmaceutical can also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette-indicating the quantity of active agent. Where the pharmaceutical is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the pharmaceutical is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients can be mixed prior to administration.

[0284] Further, the pharmaceutical composition can be provided as a pharmaceutical kit comprising (a) a container containing a IL-23p19 binding agent (e.g., an anti-IL-23p19 antibody) in lyophilized form and (b) a second container containing a pharmaceutically acceptable diluent (e.g., sterile water) for injection. The pharmaceutically acceptable diluent can be used for reconstitution or dilution of the lyophilized anti-IL-23p19 antibody or agent. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

[0285] The amount of the IL-23p19 binding agent (e.g., anti-IL-23p19 antibody) that is effective in the treatment or prevention of an immunological disorder or cancer can be determined by standard clinical techniques. In addition, in vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the stage of immunological disorder or cancer, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

[0286] Generally, the dosage of an anti-IL-23p19 antibody or IL-23p19 binding agent administered to a patient with an immunological disorder or IL-23p19-expressing cancer is typically about 0.1 mg/kg to about 100 mg/kg of the subject's body weight. The dosage administered to a subject is about 0.1 mg/kg to about 50 mg/kg, about mg/kg to about 30 mg/kg, about 1 mg/kg to about 20 mg/kg, about 1 mg/kg to about 15 mg/kg, or about 1 mg/kg to about 10 mg/kg of the subject's body weight.

[0287] Exemplary doses include, but are not limited to, from 1 ng/kg to 100 mg/kg. In some embodiments, a dose is about 0.5 mg/kg, about 1 mg/kg, about 2 mg/kg, about 3 mg/kg, about 4 mg/kg, about 5 mg/kg, about 6 mg/kg, about 7 mg/kg, about 8 mg/kg, about 9 mg/kg, about 10 mg/kg, about 11 mg/kg, about 12 mg/kg, about 13 mg/kg, about 14 mg/kg, about 15 mg/kg or about 16 mg/kg. The dose can be administered, for example, daily, once per week (weekly), twice per week; thrice per week, four times per week, five times per week, six times per week, biweekly or monthly, every two months, or every three months. In specific embodiments, the dose is about 0.5 mg/kg/week, about 1 mg/kg/week, about 2 mg/kg/week, about 3 mg/kg/week, about 4 mg/kg/week, about 5 mg/kg/week, about 6 mg/kg/week, about 7 mg/kg/week, about 8 mg/kg/week, about 9 mg/kg/week, about 10 mg/kg/week, about 11 mg/kg/week, about 12 mg/kg/week, about 13 mg/kg/week, about 14 mg/kg/week, about 15 mg/kg/week or about 16 mg/kg/week. In some embodiments, the dose ranges from about 1 mg/kg/week to about 15 mg/kg/week.

[0288] In some embodiments, the pharmaceutical compositions comprising the IL-23p19 binding agent can further comprise a therapeutic agent, either conjugated or unconjugated to the binding agent. The anti-IL-23p19 antibody or IL-23p19 binding agent can be co-administered in combination with one or more therapeutic agents for the treatment or prevention of immunological disorders or cancers.

[0289] Such combination therapy administration can have an additive or synergistic effect on disease parameters (e.g., severity of a symptom, the number of symptoms, or frequency of relapse).

[0290] With respect to therapeutic regimens for combinatorial administration, in a specific embodiment, an anti-IL-23p19 antibody or IL-23p19 binding agent is administered concurrently with a therapeutic agent. In another specific embodiment, the therapeutic agent is administered prior or subsequent to administration of the anti-IL-23p19 antibody or IL-23p19 binding agent, by at least an hour and up to several months, for example at least an hour, five hours, 12

hours, a day, a week, a month, or three months, prior or subsequent to administration of the anti-IL-23p19 antibody or IL-23p19 binding agent.

Articles of Manufacture

5 [0291] In another aspect, an article of manufacture containing materials useful for the treatment of the disorders described above is included. The article of manufacture comprises a container and a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition that is effective for treating the condition and may have a sterile access port. For example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle. The active agent in the composition is the humanized anti-IL-23p19 antibody. The label on or associated with the container indicates that the composition is used for treating the condition of choice. The article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution, and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

10 [0292] The invention is further described in the following examples, which are not intended to limit the scope of the invention.

15 20 **Examples**

Example 1: Production of Humanized Anti-IL-23p19 Antibodies

25 [0293] Mouse lead antibody 6B8 was converted to a chimeric antibody consisting of the mouse variable domain of 6B8 and a human constant IgG1KO domain. Mouse antibody 6B8 is shown in Tables 1 and 2 herein above. The IgG1KO (knock_out) has two replacement mutations (Leu234Ala and Leu235Ala) that eliminate ADCC and CDC activity by reducing effector functions such as Fc_yR and complement binding. The variable domains of the mouse and chimeric antibodies are identical. Chimeric antibodies are generated to confirm the function of the antibody and to ensure the correct sequence has been obtained. The variable region of the antibody is then humanized through a design and screening process. A library was made where human and mouse residues were varied in such a way that in any given position there could be either a human or mouse residue. Such a library was made for those amino acids that were different between human germline and mouse antibody. Only the clones that retain the function of the parent mouse antibody were selected. Representative humanized variable regions for antibody 6B8 are shown in Tables 5 and 6.

30 35 [0294] In this manner, Antibody A, Antibody B, Antibody C and Antibody D were humanized antibodies derived from mouse antibody 6B8 (cloned into a human IgG1-KO (KO=knock-out)/kappa backbone. Antibodies A, B, C and D are shown in Table 7.

Example 2: Binding of antibodies to recombinant IL-23 protein

40 45 [0295]

A) Kinetics and affinity of mouse anti-IL-23p19 antibodies binding to recombinant human IL-23 are shown below (Table 9). Kinetics and binding affinities were measured using the Fortebio Octet (ForteBio, Menlo Park, CA) using material generated from hybridoma following single column purification. Since the Octet is not a fluidics based technology, this method does not provide precise determination of off-rate. In some cases, only an estimate of affinity can be obtained.

50 55 Table 9

Antibody	ka(1/Ms)	Kd(1/s)	KD(pM)
18C4	3.84E+05	2.14E-06	5.57
18E 5	3.29E+05	2.61E-06	7.93
18D3	3.19E+05	2.16E-06	6.78
20 E8	4.21E+05	2.69E-04	638
22 E2	3.46E+05	3.53E-04	1024
24A5	2.02E+05	4.57E-06	22.6
15C11	4.11E+05	1.07E-05	26

(continued)

Antibody	ka(1/Ms)	Kd(1/s)	KD(pM)
43F5	1.72E+05	5.96E-06	34.6
27G8	1.57E+05	4.26E-06	27.2
31H9	2.99E+05	3.45E-06	11.5
2D1		< 1e-6	< 1
9D12		< 1e-6	< 1
6B8		< 1e-6	< 1
73H10	5.29E+04	5.24E-06	99.2
74H3	3.06E+04	2.09E-06	68.3
35H8			
26F7	4.76E+05	1.34E-05	28.1
34G3	9.18E+05	3.10E-05	32.8
34D9	3.44E+03	1.87E-06	544

B) Affinities were measured for humanized antibodies derived from mouse antibody 6B8. Kinetic binding data, measured using the ProteON XPR36 (Biorad, Hercules, CA) and globally fit to a 1:1 binding model, demonstrated the interactions with recombinant IL-23 either with or without a 21 amino acid linker covalently joining the p19 and p40 subunits to be of high affinity, in the range of 1 pM ~ 100 pM (Table 10). Antibody 6H12 (disclosed in WO 2007/027714), antibody QF20 (disclosed in WO 2007/024846) and antibody C1273 (disclosed in WO 2007/005955) were also tested.

Table 10

Antibody	Human IL-23 with linker			Human IL-23 no linker		
	ka (1/Ms)	kd (1/s)	KD (pM)	ka(1/Ms)	kd (1/s)	KD (pM)
Mouse 6B8				5.57E+05	1.38E-05	24.5
Antibody A	6.27E+05	< 1e-6	< 1	5.51 E+05	< 1e-6	< 1
antibody B	3.56E+05	< 1e-6	< 1	5.17E+05	< 1e-6	< 1
Antibody C	3.74E+05	1.19E-05	31.8	4.54E+05	1.65E-05	36.3
Antibody D	3.82E+05	4.07E-05	107	3.66E+05	4.93E-05	135
C-1273				3.60E+05	5.75E-06	15.8
6H12				4.99E+05	1.07E-04	214
QF20				2.03E+05	5.89E-06	2.91

C) Affinity and kinetic data for the anti-IL-23p19 antibodies binding to cynomolgous IL-23 were measured on the ProteON XPR36, and globally fit to a 1:1 binding model (Table 11). Antibody 6H12 (disclosed in WO 2007/027714), antibody QF20 (disclosed in WO 2007/024846) and antibody C1273 (disclosed in WO 2007/005955) were also tested.

Table 11

Antibody	K _D (pM)	k _a (1/Ms)	k _d (1/s)
Antibody A	< 1	2.95E+06	< 1e-6
Antibody B	< 1	2.99E+06	< 1e-6
Antibody C	2.9	3.23E+06	9.36E-06
Antibody D	15.9	2.07E+06	3.29E-05

(continued)

Antibody	K_D (pM)	k_a (1/Ms)	k_d (1/s)
C-1273	>5,000	n/a	n/a
6H12	157	9.91E+05	1.56E-04
QF20	1.2	3.90E+06	4.78E-06

10 D) Molecular Selectivity over human IL-12

The anti-IL-23p19 antibodies were also injected over a human IL-12 surface at a concentration of 100 nM. The binding signal for these antibodies measured using the Fortebio Octet is zero, which indicates that these antibodies selectively bind to human IL-23. The binding of the anti-IL-23p19 antibodies to IL-23 was also analyzed in the presence of 50% human serum and no significant effect of serum on binding on-rate was observed demonstrating high specificity.

Example 3: Competition binding assay of human IL-23 binding to human IL-23R/Fc

[0296] Human IL-23R-Fc was captured on the biosensor surface and 10 nM of human IL-23 was injected. The sensorgram indicates the specific binding between IL-23 and the IL-23 receptor (Figure 2, top trace). Antibodies were then co-injected with 10 nM human IL-23 to assess whether antibody binding to the IL-23 could inhibit the interaction between IL-23 and the IL-23 receptor. In this example, if the antibody binds to human IL-23 and is able to inhibit the interaction then reduced or no binding will be observed (Figure 2, bottom trace). In the example shown an equivalent molar concentration of Antibody A was co-injected with 10 nM recombinant human IL-23.

Example 4: Functional Cell Assays, Inhibition of IL-17 production from IL-23 stimulated mouse splenocytes

[0297] One functional cell assay for anti-IL-23p19 antibodies measures their ability to inhibit 1L-23 stimulated IL-17 production from mononuclear cells isolated from mouse spleens. Human recombinant IL-23 protein is capable of stimulating IL-17 release from mouse splenocytes. In addition, a natural source of human IL-23 found in the supernatant of activated human monocytic THP-1 cells can be used to stimulate IL-17 production from mouse mononuclear cells.

[0298] Human recombinant IL-23 or natural human IL-23 from activated THP-1 cells was preincubated with titrated anti-IL-23p19 antibodies. The IL-23/antibody combinations were then added to freshly isolated murine splenocytes. Recombinant IL-23 alone was used as a positive control. After two days in culture, cell supernatants were collected and assayed for IL-17 by ELISA (R&D Systems, Minneapolis, MN). Representative IC₅₀ values for anti-IL-23p19 antibodies are shown below. The tested antibodies are mouse antibodies derived from hybridomas (rows 1-19, see tables 1 and 2), chimeric antibodies (rows 20-23), and Antibodies A to D. Antibody 6H12 (disclosed in WO 2007/027714), antibody QF20 (disclosed in WO 2007/024846) and antibody C1273 (disclosed in WO 2007/005955) were also tested.

40 Table 12

Antibody	IC ₅₀ Values (pM), recombinant human IL-23	IC ₅₀ Values (pM), natural human IL-23
18C4	471	100, 413
18E5	not determined	9,9, 13
18D3	234	not determined
20E8	not determined	438,561
22 E2	61, 130	117, 35
24A5	22, 37	85, 31
15C11	126	232
43F5	250, 8000	8000
27G8	235	5000
31H9	960	2000
2D1	not determined	2336, 1911, 1597

(continued)

Antibody	IC₅₀ Values (pM), recombinant human IL-23	IC₅₀ Values (pM), natural human IL-23
9D12	59	281, 138
6B8	13	8, 2
73H10	1411	not determined
74H3	1352	not determined
36H8	not determined	not determined
26F7	27	2, 8
34G3	336	27, 25
34D9	510	456
Chimeric 18E5	31	8, 36, 10, 9
Chimeric 22E2	100	9, 178
Chimeric 24A5	404	95, 102
Chimeric 6B8	26, 37, 57	5, 2, 6, 3
Antibody A	5, 5, 5, 15	1, 1
Antibody B	13, 30, 54, 42	9, 8
Antibody C	53, 71, 162, 89	16, 32
Antibody D	236, 225, 614, 458	133, 125
6H12	1600, 806, 1300	957, 4400, 1013, 439
QF20	not determined	7, 12
C1273	not determined	93, 44

Example 5: Functional specificity testing against IL-12 in a human activated T cell assay

[0299] Anti-IL-23p19 antibodies were tested for functional inhibition of IL-12 in a human activated T cell assay. Human recombinant IL-12 (1 ng/ml) was preincubated with 5 µg/ml anti-IL-23p19 antibodies. The IL-12/antibody combinations were then added to human PHA-derived T cell blasts. Recombinant IL-12 alone was used as a positive control. An anti-IL-12p70 antibody (Bender MedSystems, Vienna, Austria) was used as a control inhibitory antibody. After two days in culture, cell supernatants were collected and assayed for IFN-γ by ELISA (R&D Systems). Samples were tested in triplicate and the average pg/ml of IFN-γ was determined. Results (with standard deviations) are shown in the table below.

Table 13

Antibody	Cytokine Stimulation	Average pg/ml IFN-γ +/- Standard Deviation
no antibody	None	87 +/- 7
no antibody	1 ng/ml IL-12	532 +/- 51
chimeric 18E5	1 ng/ml IL-12	511 +/- 3
chimeric 6B8	1 ng/ml IL-12	523 +/- 60
Antibody A	1 ng/ml IL-12	497 +/- 30
Antibody B	1 ng/ml IL-12	537 +/- 2
Antibody C	1 ng/ml IL-12	495 +/- 25
Antibody D	1 ng/ml IL-12	539 +/- 38
anti-IL-12p70 antibody	1 ng/ml IL-12	119 +/- 12

Example 6: Inhibition of IL-23 induced STAT3 phosphorylation in the human cell line DB

[0300] The human cell line DB (ATCC, Manassas, VA) responds to IL-23 stimulation through an endogenous IL-23Rcomplex (IL-23R and IL-12R β 1) and phosphorylates STAT3 in an IL-23 dose dependent manner. An assay was developed for testing anti-IL-23p19 antibody inhibition of IL-23 induced STAT3 phosphorylation. DB cells were plated at 1x10e6 cells/well in a 96 well plate. Antibodies to be tested were serially diluted and pre-incubated with recombinant human IL-23 (10 ng/ml) for 1 hour at room temperature. The antibody/IL-23 mixture was then added to the cells for 30 minutes at 37°C. Cells were harvested by centrifugation at 4°C for 10 minutes and then lysed in ice cold buffer (Cell Signaling Technology, Beverly, MA). A portion of the lysate was run in a phospho-STAT3 ELISA (Invitrogen). Antibody IC₅₀ values were calculated as percent inhibition of STAT3 phosphorylation compared to control wells without antibody. Representative IC₅₀ values are shown in the table below.

Table 14

Antibody	IC ₅₀ (pM)
Antibody A	25, 15, 38, 23, 13, 18
Antibody B	73, 84
Antibody C	132, 80
Antibody D	158
QF20	26, 26, 27
C-1273	163, 438

Example 7: In vivo model of IL-23 induced cytokine production in the mouse ear

[0301] An in vivo model in the mouse was used. Recombinant human IL-23 is injected into the skin of the mouse ear for 4 consecutive days resulting in epidermal thickening and up-regulation of IL-17 and IL-22 protein. Anti-IL-23p19 antibodies were evaluated in this model. A single intraperitoneal injection of 1 mg/kg or 5 mg/kg antibody was administered 1 hour prior to the initial IL-23 injection into the skin. Recombinant human IL-23 (with linker) was injected once daily for 3 additional days and tissue was collected for cytokine assessment. Inhibition of cytokine production was demonstrated for the antibodies. The results of three experiments are shown in the table bellow 1: rows 1-7, exp. 2: rows 8-10, exp. 3: rows 11-14).

Table 15

	Ear Tissue IL-17 pg/ml Mean +/- SEM	Ear Tissue IL-17 Percent Inhibition	Ear Tissue IL-22 pg/ml Mean +/- SEM	Ear Tissue IL-22 Percent Inhibition
0.1 % BSA + Citrate (Unstimulated Control)	3 +/- 1	NA	1 +/- 0	NA
0.3ug IL-23 + Citrate Buffer i.p. (Vehicle Control)	25 +/- 3	NA	274 +/- 30	NA
0.3ug IL-23 + 1 mg/kg Antibody 6B8	7 +/- 2	81	57 +/- 19	80
0.3ug IL-23 + 1 mg/kg Antibody A	2 +/- 1	101	17 +/- 3	94
0.3ug IL-23 + 1 mg/kg Antibody B	5 +/- 1	93	30 +/- 2	89
0.3ug IL-23 + 1 mg/kg Antibody C	11 +/- 1	66	108 +/- 12	61
0.3ug IL-23 + 1 mg/kg Antibody D	10 +/- 1	67	151 +/- 12	45

(continued)

	Ear Tissue IL-17 pg/ml Mean +/- SEM	Ear Tissue IL-17 Percent Inhibition	Ear Tissue IL-22 pg/ml Mean +/- SEM	Ear Tissue IL-22 Percent Inhibition	
5	0.1 % BSA + Vehicle (Unstimulated Control)	14 +/- 1	NA	1 +/- 1	NA
10	0.3ug IL-23 + Vehicle	31 +/- 4	NA	129 +/- 29	NA
15	0.3ug IL-23 + 5mg/kg 24A5	14 +/- 1	102	10 +/- 5	93
20	0.1% BSA + mlgG (Unstimulated Control)	17 +/- 1	NA	4 +/- 1	NA
25	0.3ug IL-23 + mlgG (Vehicle Control)	30 +/- 2	NA	208 +/- 40	
30	0.3ug IL-23 + 5mg/kg 24A5	16 +/- 0	109	28 +/- 5	88
35	0.3ug IL-23 + 5mg/kg 18E5	21 +/- 2	70	53 +/- 41	80

Example 8: Pharmacokinetic Studies in Cynomolgus Monkey

[0302] Humanized anti-IL-23p19 antibodies were administered by ten minute intravenous infusion at a dose of 1.0 mg/kg to three cynomolgus monkeys. Serum samples were collected over a 6 week time course and free antibody concentrations were measured using a specific ELISA. The serum concentration-time profiles for the antibodies and the corresponding pharmacokinetic parameters are summarized in the Table 16 below.

Table 16

Antibody	CL (ml/d/kg)	Vol (ml/kg)	AUC (nM•h/ml)	T _{1/2} (days)	MRT (days)
Antibody A	5.2	88	32262	12.1	17.2
Antibody B	6.0	87	27030	10.1	14.8
Antibody C	4.7	91	34642	14.1	19.6
Antibody D	3.4	67	47633	12.6	19.8

Example 9: Expression in NSO cells and biophysical data

[0303] Transfection of NS0 cells and generation of stable pools;

NS0 cells were grown in the presence of 1% FBS before transfection. 40x10e6 cells were collected and resuspended in 0.8ml in media containing 2% FBS with 20ug of linearized DNA (heavy chain and light chain expression vectors) and then cells were incubated on ice for approximately 15 min before electroporation of the cells at 750V/25uF (Bio-Rad Gene Pulser Xcell). Cells were recovered with 2% FBS for approximately 48 hours at 37°C and 5% CO₂ then plated in 96 well plates at 2x10e5 cells/ml containing G418 and mycophenolic acid for 14-21 days until formation of colonies.

[0304] Supernatant from 96 well plates with colonies were screened by ELISA. ELISA plates were coated with 1ug/ml of goat anti-kappa (Southern Biotech, Birmingham, AL) in PBS and diluted supernatant were incubated and then detected with goat anti-human IgG Fc-HRP (from Jackson ImmunoResearch Laboratories, West Grove, PA). The positive colonies were pooled for scale up. Titers for antibody production were determined by ForteBio using protein A tips according to manufacturer protocol. The titers for Antibody A and Antibody D were between 250-350 mg/L, with more than 80% recovery from protein purification, and more than 94% monomer after IEX purification. Proteins were resuspended in a final buffer containing 20mM Sodium Citrate and 115mM NaCl, pH 6.0 and are stable at 4°C for at least 4 months and with solubility up to 100 mg/ml in this buffer.

Table 17

	Protein A Column			IEX Column	
	Titer (mg/L)	Yield (mg/L)	Recovery	Yield (mg/L)	Recovery
Antibody A	345	275	80%	221	80%
Antibody D	248	225	90%	175	78%

Table 18

	Quality		Stability				Solubility
	AUC fresh (%M)	SEC fresh (%M)	AUC 1 month (%M)	SEC 1 month (%M)	AUC 4months (%M)	SEC 4months (%M)	AUC at 100 mg/ml (%M)
Antibody A	98	99	97	99	96	99	99
Antibody D	94	100	98	100	99	99	97

[0305] AUC: Analytical Ultracentrifugation as measured by the sedimentation Velocity method at concentrations of 0.5 -1 mg/ml; SEC: Size exclusion chromatography; %M: percent monomer.

Example 10: Epitope mapping

[0306] Hydrogen/Deuterium Exchange Mass Spectrometry (HXMS) was employed to map the epitope of Antibody A binding to human IL-23p19. This method determined the susceptibility of the amide backbone hydrogens of IL-23p19 to exchange with D₂O. The experiment was conducted with IL-23 alone and IL-23 with added Antibody A. Regions of the IL-23p19 sequence showing significant protection from exchange due to binding of Antibody A were thus identified. Resolution of the method is determined by the peptides produced by digestion with pepsin or Protease XVIII. These IL-23p19 derived peptides were identified by additional control experiments with unexchanged samples employing standard accurate mass and HPLC MS/MS technologies.

[0307] Recombinant human IL-23 was used. For the protein + antibody sample, 50ul of IL-23 (0.8mg/ml) was incubated with 10ul of Antibody A (12.7mg/ml) for 15 minutes at room temperature. The final molar ratio Antibody A/IL-23 was 1.2:1.

[0308] For the exchange 5ul of IL-23 protein was added to 50ul deuterated buffer (50mM PBS in D₂O) and incubated for 100 seconds at room temperature. 50ul of 2M Urea/0.5M TCEP was added and incubated for 60 seconds at room temperature. 5ul pepsin or Protease XVIII (4mg/ml in 0.1% formic acid) was added and the sample was immediately cooled to 4 °C.

[0309] After 5 minutes 50ul of sample was injected onto a Shimadzu HPLC system (SCL10A controller and two LC10AD pumps) under the following conditions:

Mobile Phase A = 99/1/0.1 (water/acetonitrile/formic acid).

Mobile Phase B = 95/5/0.1 (acetonitrile/water/formic acid).

Flow rate = 100ul/min.

Column = Phenomenex Jupiter C5, 5u, 50x1.0mm.

Mobile phase lines, column, injector loop are in ice baths.

Gradient = Time 0(3%B), Time 2.2(3%B), Time 10.1(90%B), Time 12.0(90%B), Time 12.1(3%B).

[0310] Mass Spectrometry was carried out as follows:

Mass Spec = Thermo Orbitrap Velos (0900865).

Methods:

A. Fragmentation (to ID peptides): 12 minute acquisition time (3 minute start delay), full-scan FTMS at 30,000 resolution, seven ion trap data dependant scans (CID).

B. MS Runs: 12 minute acquisition time (3 minute start delay), full-scan FTMS at 60,000 resolution.

5 [0311] Pepsin and Protease XVIII peptides were identified using fragmentation data and the program Proteome Discoverer (Thermoscientific, Waltham, MA). Identified peptides were visually compared (protein alone vs. protein with antibody present) using Xcalibur software (Thermoscientific). No significant shifts in exchange were observed for IL-23 alone vs. IL-23 with Antibody A outside of the IL-23p19 region. For the p19 portion of the protein, data was analyzed using the program PepMap (Thermoscientific). this program calculates the average mass for exchanged peptides. PepMap results were checked and those peptides that did not yield verified results were calculated with the aid of Microsoft Excel.

10 [0312] The regions of the IL-23 sequence showing significant protection from exchange due to binding of Antibody A were identified as amino acid resides 108 to 126 of SEQ ID NO:181 and amino acid residues 137 to 151 of SEQ ID NO:181.

SEQUENCE LISTING

15 [0313]

<110> Barrett, Rachel Rebecca

Canada, Keith

Catron, Katrina Mary

20 Copenhagen, Robert

Frego, Lee Edward

Raymond, Ernest Lee

Singh, Sanjaya

Zhu, Xiangyang

25 <120> Anti-IL-23 Antibodies

<130> 09-0561

30 <150> 61/410,158

<151> 2010-11-04

35 <150> 61/411,953

<151> 2010-11-10

40 <150> 61/412,594

<151> 2010-11-11

45 <150> 61/448,785

<151> 2011-03-03

50 <160> 181

45 <170> FastSEQ for Windows Version 4.0

55 <210> 1

<211> 11

<212> PRT

<213> Mouse

50 <400> 1

Arg	Ala	Ser	Gln	Ser	Ile	Ser	Glu	Tyr	Leu	His
1					5					10

55

<210> 2

<211> 7

<212> PRT

EP 2 635 601 B1

<213> Mouse

<400> 2

5 Tyr Ala Ser Gln Ser Ile Ser
 1 5

<210> 3

<211> 9

10 <212> PRT
 <213> Mouse

<400> 3

15 Gln Asn Gly His Ser Phe Pro Phe Thr
 1 5

<210> 4

<211> 11

20 <212> PRT
 <213> Mouse

<400> 4

25 Arg Ala Ser Gln Ser Ile Ser Asp Tyr Leu Tyr
 1 5 10

<210> 5

<211> 7

30 <212> PRT
 <213> Mouse

<400> 5

35 Phe Ala Ser Gln Ser Ile Ser
 1 5

<210> 6

<211> 11

40 <212> PRT
 <213> Mouse

<400> 6

45 Arg Ala Ser Gln Ser Ile Ser Val Tyr Leu His
 1 5 10

<210> 7

<211> 11

50 <212> PRT
 <213> Mouse

<400> 7

55 Arg Ala Ser Gln Ser Ile Ser Asp Tyr Leu His
 1 5 10

EP 2 635 601 B1

<210> 8
<211> 16
<212> PRT
<213> Mouse

5

<400> 8

Arg Ser Ser Gln Ser Leu Val His Ser Asn Gly Asn Thr Tyr Leu His
1 5 10 15

10

<210> 9
<211> 7
<212> PRT
<213> Mouse

15

<400> 9

Lys Val Ser Asn Arg Phe Ser
1 5

20

<210> 10
<211> 9
<212> PRT
<213> Mouse

25

<400> 10

Ser Gln Ser Thr His Val Pro Tyr Thr
1 5

30

<210> 11
<211> 15
<212> PRT
<213> Mouse

35

<400> 11

Arg Ala Ser Lys Ser Val Ser Thr Ser Gly Tyr Ser Tyr Ile His
1 5 10 15

40

<210> 12
<211> 7
<212> PRT
<213> Mouse

45

<400> 12

Leu Ala Ser Asn Leu Asp Ser
1 5

50

<210> 13
<211> 9
<212> PRT
<213> Mouse

55

<400> 13

Gln His Ser Arg Glu Leu Pro Tyr Thr
 1 5

5 <210> 14
 <211> 9
 <212> PRT
 <213> Mouse

10 <400> 14

Gln Asn Gly His Ser Phe Pro Tyr Thr
 1 5

15 <210> 15
 <211> 15
 <212> PRT
 <213> Mouse

20 <400> 15

Arg Thr Ser Glu Ser Val Tyr Ser Tyr Gly Gln Asn Phe Ile His
 1 5 10 15

25 <210> 16
 <211> 7
 <212> PRT
 <213> Mouse

30 <400> 16

Arg Ala Ser Asn Leu Glu Ser
 1 5

35 <210> 17
 <211> 9
 <212> PRT
 <213> Mouse

40 <400> 17

Gln Gln Thr Asn Glu Asp Pro Tyr Thr
 1 5

45 <210> 18
 <211> 15
 <212> PRT
 <213> Mouse

50 <900> 18

Arg Ala Ser Glu Thr Ile Asn Phe Tyr Gly Thr Ser Phe Met His
 1 5 10 15

55 <210> 19
 <211> 11
 <212> PRT
 <213> Mouse

<400> 19

Lys Ala Ser Arg Asp Val Ala Ile Ala Val Ala

1	5	10
---	---	----

5

<210> 20

<211> 7

<212> PRT

<213> Mouse

10

<400> 20

Trp Ala Ser Thr Arg His Thr

1	5
---	---

15

<210> 21

<211> 9

<212> PRT

<213> Mouse

20

<400> 21

His Gln Tyr Ser Ser Tyr Pro Phe Thr

1	5
---	---

25

<210> 22

<211> 11

<212> PRT

<213> Mouse

30

<400> 22

Arg Ala Ser Glu Asn Ile Asp Ser Tyr Leu Ala

1	5	10
---	---	----

35

<210> 23

<211> 7

<212> PRT

40

<213> Mouse

<400> 23

Ala Ala Arg Asn Leu Ala Asp

1	5
---	---

45

<210> 24

<211> 9

<212> PRT

50

<213> Mouse

<400> 24

Gln His Tyr Tyr Ser Thr Pro Phe Thr

1	5
---	---

55

<210> 25

<211> 7

<212> PRT
 <213> Mouse

<400> 25

5

Ala Ala Thr Asn Leu Ala Asp
 1 5

<210> 26

10

<211> 9
 <212> PRT
 <213> Mouse

<400> 26

15

Leu His Tyr Tyr Ser Thr Pro Phe Thr
 1 5

<210> 27

20

<211> 15
 <212> PRT
 <213> Mouse

<400> 27

25

Arg Ala Ser Lys Ser Val Arg Phe Ser Asp Tyr Phe Tyr Met His
 1 5 10 15

<210> 28

30

<211> 7
 <212> PRT
 <213> Mouse

<400> 28

35

Leu Ala Ser Asn Leu Glu Ser
 1 5

<210> 29

40

<211> 9
 <212> PRT
 <213> Mouse

<400> 29

45

Gln Asn Ser Arg Glu Leu Pro Tyr Thr
 1 5

<210> 30

50

<211> 11
 <212> PRT
 <213> Mouse

<400> 30

55

Lys Ala Ser Gln Asp Val Gly Asn Ala Val Val
 1 5 10

<210> 31
 <211> 7
 <212> PRT
 <213> Mouse

5

<400> 31

Trp Ala Ser Thr Arg His Ile
 1 5

10

<210> 32
 <211> 8
 <212> PRT
 <213> Mouse

15

<400> 32

Gln Gln Tyr Ser Ser Tyr Leu Thr
 1 5

20

<210> 33
 <211> 10
 <212> PRT
 <213> Mouse

25

<400> 33

Gly Tyr Thr Phe Thr Ser Ser Val Ile His
 1 5 10

30

<210> 34
 <211> 17
 <212> PRT
 <213> Mouse

35

<400> 34

Tyr Ile Asn Pro Tyr Asn Asp Gly Thr Lys Tyr Asn Glu Lys Phe Lys
 1 5 10 15
 Gly

40

<210> 35
 <211> 6
 <212> PRT
 <213> Mouse

45

<400> 35

Arg Leu Asp Glu Ala Tyr
 1 5

50

<210> 36
 <211> 10
 <212> PRT
 <213> Mouse

55

<400> 36

Gly Tyr Thr Phe Thr Arg Tyr Leu Ile His
 1 5 10

5 <210> 37
 <211> 6
 <212> PRT
 <213> Mouse

10 <400> 37

Asn Trp Asp Leu Asp Tyr
 1 5

15 <210> 38
 <211> 10
 <212> PRT
 <213> Mouse

20 <400> 38

Gly Tyr Thr Phe Thr Ser Ser Val Met His
 1 5 10

25 <210> 39
 <211> 17
 <212> PRT
 <213> Mouse

30 <400> 39

Tyr Ile Asn Pro Tyr Asn Asp Gly Thr Gln Tyr Asn Glu Lys Phe Lys
 1 5 10 15
 Gly

35 <210> 40
 <211> 10
 <212> PRT
 <213> Mouse

40 <400> 40

Gly Tyr Thr Phe Thr Ser Ser Ile Ile His
 1 5 10

45 <210> 41
 <211> 17
 <212> PRT
 <213> Mouse

50 <400> 41

Tyr Ile Asn Pro Tyr Asp Asp Val Thr Lys Tyr Asn Glu Lys Phe Lys
 1 5 10 15
 Gly

<210> 42

<211> 6
 <212> PRT
 <213> Mouse

5 <400> 42

Arg Trp Asp Glu Ser Tyr
 1 5

10 <210> 43
 <211> 10
 <212> PRT
 <213> Mouse

15 <400> 43

Gly Tyr Thr Phe Thr Thr Ser Ile Met His
 1 5 10

20 <210> 44
 <211> 6
 <212> PRT
 <213> Mouse

25 <400> 44

Arg Trp Asp Glu Ala Tyr
 1 5

30 <210> 45
 <211> 10
 <212> PRT
 <213> Mouse

35 <400> 45

Gly Tyr Thr Phe Thr Asp Tyr Tyr Met Asn
 1 5 10

40 <210> 46
 <211> 17
 <212> PRT
 <213> Mouse

45 <400> 46

Val Ile Ile Pro Tyr Asn Gly Gly Thr Ser Tyr Asn Gln Lys Phe Lys
 1 5 10 15
 Gly

50 <210> 47
 <211> 9
 <212> PRT
 <213> Mouse

55 <400> 47

Asp Gly His Arg Trp Tyr Phe Asp Val
 1 5

5 <210> 48
 <211> 10
 <212> PRT
 <213> Mouse

10 <400> 48
 10

Gly Tyr Ser Phe Thr Gly Tyr Tyr Met Asn
 1 5 10

15 <210> 49
 <211> 17
 <212> PRT
 <213> Mouse

20 <400> 49

Glu Ile Ile Pro Thr Thr Gly Gly Thr Ser Tyr Asn Gln Lys Phe Lys
 1 5 10 15
 Ala

25 <210> 50
 <211> 11
 <212> PRT
 <213> Mouse

30 <400> 50

Glu Ser Gly Gly Phe Tyr Trp Tyr Phe Asp Val
 1 5 10

35 <210> 51
 <211> 10
 <212> PRT
 <213> Mouse

40 <400> 51

Gly Tyr Thr Phe Thr Asp His Thr Ile His
 1 5 10

45 <210> 52
 <211> 17
 <212> PRT
 <213> Mouse

50 <400> 52

Tyr Ile Tyr Pro Arg Asp Gly Tyr Pro Lys Phe Asn Glu Lys Phe Lys
 1 5 10 15
 Gly

55 <210> 53

EP 2 635 601 B1

<211> 9
<212> PRT
<213> Mouse

5 <400> 53

Arg Pro Pro Tyr Tyr Ala Met Asp Tyr
1 5

10 <210> 54
<211> 10
<212> PRT
<213> Mouse

15 <400> 54

Gly Tyr Thr Phe Thr Arg Tyr Leu Met His
1 5 10

20 <210> 55
<211> 17
<212> PRT
<213> Mouse

25 <400> 55

Tyr Ile Asn Pro Tyr Asn Asp Gly Thr Asn Tyr Asn Glu Lys Phe Lys
1 5 10 15

30 Gly

<210> 56
<211> 6
35 <212> PRT
<213> Mouse

<400> 56

40 Asn Trp Asp Tyr Ala Tyr
1 5

45 <210> 57
<211> 10
<212> PRT
<213> Mouse

<400> 57

50 Gly Phe Ser Leu Thr Thr Tyr Ala Ile Ser
1 5 10

55 <210> 58
<211> 16
<212> PRT
<213> Mouse

<400> 58

Val Ile Trp Thr Gly Gly Gly Thr Lys Tyr Asn Ser Ala Leu Lys Ser
 1 5 10 15

5

<210> 59

<211> 11

<212> PRT

10 <213> Mouse

<400> 59

Lys Asp Tyr Asn Tyr Gly Gly Ala Met Asp Tyr
 1 5 10

15

<210> 60

<211> 10

<212> PRT

20 <213> Mouse

<400> 60

Gly Phe Ser Leu Asn Asn Phe Ala Ile Ser
 1 5 10

25

<210> 61

<211> 16

<212> PRT

30 <213> Mouse

<400> 61

Ala Ile Trp Thr Gly Gly Thr Asn Tyr Asn Ser Ala Leu Lys Ser
 1 5 10 15

35

<210> 62

<211> 11

<212> PRT

40 <213> Mouse

<400> 62

Lys Asp Tyr Ser Tyr Gly Gly Ala Met Asp Tyr
 1 5 10

45

<210> 63

<211> 10

<212> PRT

50 <213> Mouse

<400> 63

Gly Asn Thr Phe Thr Asp Gln Thr Ile His
 1 5 10

55

<210> 64

<211> 17

EP 2 635 601 B1

<212> PRT
<213> Mouse

5 <400> 64

Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe Lys
1 5 10 15
Gly

10 <210> 65
<211> 11
<212> PRT
<213> Mouse

15 <400> 65

Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr
1 5 10

20 <210> 66
<211> 10
<212> PRT
<213> Artificial Sequence

25 <220>
<223> Humanized sequence.

<400> 66

Gly Tyr Thr Phe Thr Asp Gln Thr Ile His
1 5 10

30 <210> 67
<211> 10
<212> PRT
<213> Artificial Sequence

35 <220>
<223> Humanized sequence.

40 <400> 67

Gly Phe Thr Phe Thr Asp Gln Thr Ile His
1 5 10

45 <210> 68
<211> 10
<212> PRT
<213> Artificial Sequence

50 <220>
<223> Humanized sequence.

<400> 68

Gly Gly Thr Phe Thr Asp Gln Thr Ile His
1 5 10

<210> 69
 <211> 10
 <212> PRT
 <213> Mouse

5

<400> 69

Gly Tyr Thr Phe Thr Arg Tyr Val Met His
 1 5 10

10

<210> 70
 <211> 17
 <212> PRT
 <213> Mouse

15

<400> 70

Tyr Ile Asn Pro Tyr Asn Asp Val Thr Lys Tyr Asn Glu Lys Phe Lys
 1 5 10 15

20

Gly

25

<210> 71
 <211> 6
 <212> PRT
 <213> Mouse

30

<400> 71

Asn Trp Asp Val Pro Tyr
 1 5

35

<210> 72
 <211> 17
 <212> PRT
 <213> Mouse

40

<400> 72

Tyr Ile Asn Pro Tyr Asn Asp Gly Thr Lys Tyr Asn Glu Arg Phe Lys
 1 5 10 15

45

Gly

50

<210> 73
 <211> 17
 <212> PRT
 <213> Mouse

<400> 73

55

Gly

Val Ile Ile Pro Tyr Asn Gly Gly Ile Ser Tyr Asn Gln Lys Phe Lys
 1 5 10 15

<210> 74
 <211> 9
 <212> PRT
 <213> Mouse

5

<400> 74

Asn	Asp	Tyr	Asp	Trp	Tyr	Phe	Asp	Val
1						5		

10

<210> 75
 <211> 17
 <212> PRT
 <213> Mouse

15

<400> 75

Asp	Phe	Asn	His	Asn	Asn	Asp	Val	Ile	Thr	Tyr	Asn	Pro	Lys	Phe	Lys
1										10				15	
Gly															

20

<210> 76
 <211> 10
 <212> PRT
 <213> Mouse

<400> 76

30

Gly	Leu	Arg	Gly	Tyr	Tyr	Ala	Met	Asp	Tyr
1						5			10

35

<210> 77
 <211> 10
 <212> PRT
 <213> Mouse

<400> 77

40

Gly	Tyr	Ser	Phe	Thr	Asp	Tyr	Asn	Met	Asn
1						5			10

45

<210> 78
 <211> 17
 <212> PRT
 <213> Mouse

<400> 78

50

Val	Ile	Ile	Pro	Asn	Tyr	Gly	Phe	Thr	Ser	Tyr	Asn	Gln	Asn	Phe	Lys
1							5				10				15
Gly															

55

<210> 79
 <211> 11
 <212> PRT
 <213> Mouse

EP 2 635 601 B1

<400> 79

Asp Gly Gly Ile Leu Leu Trp Tyr Leu Asp Val
1 5 10

5

<210> 80
<211> 10
<212> PRT
<213> Mouse

10

<400> 80

Gly Tyr Thr Phe Thr Asp Tyr Asn Met Asp
1 5 10

15

<210> 81
<211> 17
<212> PRT
<213> Mouse

20

<400> 81

Asp Ile Asn Pro His Asn Gly Gly Thr Ile Tyr Asn Gln Lys Phe Lys
1 5 10 15
Gly

25

<210> 82
<211> 13
<212> PRT
<213> Mouse

30

<400> 82

35

Asn Tyr Tyr Gly Ser Ser Tyr Gly Trp Tyr Phe Asp Val
1 5 10

40

<210> 83
<211> 333
<212> DNA
<213> Mouse

45

<210> 83
<211> 333
<212> DNA
<213> Mouse

gacattgtgc tgaccataatc tccaggttct ttggctgtgt ctctaggcca gagggccacc 60
atatacctgca gaaccagtga aagtgttat agttatggcc aaaattttat acactggtagc 120
cagcagaaac caggacagcc acccaaactc ctcatotatc gtgcattccaa cctgaaatct 180
gggatccctg ccaggttcag tggcagtggg tctaggacag acttcacccct caccatgaat 240
cctgtggagg ctgatgtatgt tgcaacctat tactgtcagc aaactaatga ggatccgtac 300
acgttcggag gggggaccaa gctggaaata aga 333

50

55

<210> 84
<211> 11
<212> PRT
<213> Mouse

<400> 84

Asp Ile Val Leu Thr Gln Ser Pro Gly Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Gln Arg Ala Thr Ile Ser Cys Arg Thr Ser Glu Ser Val Tyr Ser Tyr
 20 25 30
 Gly Gln Asn Phe Ile His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45
 Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Ile Pro Ala
 50 55 60
 Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Met Asn
 65 70 75 80
 Pro Val Glu Ala Asp Asp Val Ala Thr Tyr Tyr Cys Gln Gln Thr Asn
 85 90 95
 Glu Asp Pro Tyr Thr Phe Gly Gly Thr Lys Leu Glu Ile Arg
 100 105 110

15 <210> 85
 <211> 321
 <212> DNA
 <213> Mouse

20 <400> 85

gacattgtga tgacccagtc tcacaaattc ttgtccacat cagtggaga cagggtcacc 60
 atcacttgca aggccagtcg ggatgtggct attgctgttag cctggtatca acagaaaacca 120
 gggcaatctc ctaaactact tctttctgg gcatccaccc gacacactgg ggtccctgat 180
 cgcttcacag gcagtggatc tcgacagat ttcaacttca ccattagcaa tgtgcagtct 240
 gaagacttgg cagattattt ctgtcaccaa tatagcagct atccattcac gttcggtcg 300
 gggacaaagt tggaaataaa g 321

30 <210> 86
 <211> 107
 <212> PRT
 <213> Mouse

35 <400> 86

Asp Ile Val Met Thr Gln Ser His Phe Leu Ser Thr Ser Val Gly
 40
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu
 35 40 45
 Phe Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Thr Gly
 50 55 60
 Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Ser Asn Val Gln Ser
 65 70 75 80
 Glu Asp Leu Ala Asp Tyr Phe Cys His Gln Tyr Ser Ser Tyr Pro Phe
 85 90 95
 Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

55 <210> 87
 <211> 333
 <212> DNA
 <213> Mouse

<400> 87

5 gacattgcgc tgaccatac tccagttct ttggctgtgt ctatgggca gagggccacc 60
 atatctgca gagccagtga aactattaat ttttatggca ctagtttat gcactggcac 120
 cagcagaaac caggacagtc acccaaactc ctcatctatc gtgcacccaa cctagaatct 180
 gggatccctg ccaggttcag tggcagtggg tcttagacag acttcacccct caccattaat 240
 cctgtggagg ctgatgatgt tgcaacctat tactgtcagc aaactaatga ggatccgtac 300
 acgttcggag gggggactaa gttggaaata aaa 333

<210> 88

<211> 111

10 <212> PRT

<213> Mouse

<400> 88

15 Asp Ile Ala Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Thr Ile Asn Phe Tyr
 20 25 30
 Gly Thr Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro
 25 35 40 45
 Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Ile Pro Ala
 50 55 60
 Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asn
 65 70 75 80
 Pro Val Glu Ala Asp Asp Val Ala Thr Tyr Tyr Cys Gln Gln Thr Asn
 85 90 95
 Glu Asp Pro Tyr Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

<210> 89

30 <211> 336

<212> DNA

<213> Mouse

<400> 89

35

40 gatgttgtga tgaccataac tccactctcc ctgcctgtca gtcttggaga tcaaggctcc 60
 atctcttgcata gatctagtca gagccttgta cacagtaatg gaaacaccta tttacattgg 120
 tacctgcaga agccaggcca gtctccaaag ctcctgatct acaaagttc caaccgattt 180
 tctggggtcc cagacaggtt cagtggcagt ggatcaggga cagatttcac actcaagatc 240
 aacagagtgg aggctgagga tctgggagtt tatttctgct ctcaaagtac acatgttccg 300
 tacacgttcg gagggggac ccagctggaa ataaaa 336

<210> 90

<211> 112

45 <212> PRT

<213> Mouse

<400> 90

50

55

Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15
 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 5 Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 10 Asp Arg Phe Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Asn Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser
 85 90 95
 Thr His Val Pro Tyr Thr Phe Gly Gly Thr Gln Leu Glu Ile Lys
 100 105 110

15 <210> 91
 <211> 107
 <212> PRT
 <213> Mouse

20 <400> 91

Asp Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
 1 5 10 15
 Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Asp Tyr
 20 25 30
 25 Leu His Trp Tyr Gln Gln Arg Ser His Glu Ser Pro Arg Leu Ile
 35 40 45
 Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
 50 55 60
 30 Ser Gly Ser Gly Ser Asp Phe Thr Leu Thr Ile Asn Ser Val Glu Pro
 65 70 75 80
 Glu Asp Val Gly Val Tyr Tyr Cys Gln Asn Gly His Ser Phe Pro Phe
 85 90 95
 Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

35 <210> 92
 <211> 321
 <212> DNA
 40 <213> Mouse

<400> 92

45 gacattgtga tgactcagtc tccagccacc ctgtctgtga ctccaggaga tagagtctct 60
 ctttctgca gggccagtc gaggattagc gactacttat actggatataca aaaaaaatca 120
 catgagtctc caaggcttct catcaaattt gcttccaaat ccatctctgg gatccccctcc 180
 aggttcactg gcagtggtatc agggtcagat ttcaactctca gtatcgacag tgtggAACCT 240
 gatgatgttg gagttttt ctgtcaaaaat ggtcacagct ttccgttac gttcggaggg 300
 50 gggaccaagc tggaaaataaa a 321

55 <210> 93
 <211> 107
 <212> PRT
 <213> Mouse

<400> 93

Asp Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
 1 5 10 15
 Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Asp Tyr
 20 25 30
 5 Leu Tyr Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
 35 40 45
 Lys Phe Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Thr Gly
 50 55 60
 Ser Gly Ser Gly Ser Asp Phe Thr Leu Ser Ile Asp Ser Val Glu Pro
 65 70 75 80
 10 Asp Asp Val Gly Val Phe Phe Cys Gln Asn Gly His Ser Phe Pro Phe
 85 90 95
 Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

15 <210> 94
 <211> 321
 <212> DNA
 <213> Mouse

20 <400> 94

25 gacattgtga tgactca^gtc tccagccacc ctgtctgtga ctccaggaga tagagtctct 60
 ctttcc^ttgca gggccagcca gagtattagc gagta^ctta^cc actggtatca acaaaaatca 120
 catgagtctc caaggc^ttct catcaaata^tat gttcccaat cc^atctctgg gatccc^ttcc 180
 aggttc^tatgc gcagtg^gatgc aggtc^tatgc tt^cactctca gtatcaacag t^ttg^gaa^cct 240
 gaagatgtt^g gagt^tat^tat^tat^t ctgtcaaaat ggtcacagct ttccattcac gttcg^gctcg 300
 gggacaaaat^t tggaaataaaa a 321

30 <210> 95
 <211> 107
 <212> PRT
 <213> Mouse

35 <400> 95

35 Asp Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
 1 5 10 15
 Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Glu Tyr
 20 25 30
 40 Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
 35 40 45
 Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Ser Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Pro
 65 70 75 80
 45 Glu Asp Val Gly Val Tyr Tyr Cys Gln Asn Gly His Ser Phe Pro Phe
 85 90 95
 Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

50 <210> 96
 <211> 321
 <212> DNA
 <213> Mouse

55 <400> 96

5 gacattgtga tgactcagtc tccagccacc ctgtctgtga ctccaggaga tagagtctct 60
 ctttcctgca gggccagcca gagtattagc gactacttat actggtatca aaaaaatca 120
 catgagtctc caaggctct catcaaattt gttcccaat ccatctctgg gatcccctcc 180
 aggttcactg gcagtggatc aggctcagat ttcaactctca gtatcgacag tgtgaaacct 240
 gatgatgttg gagtctttt ctgtcaaaaat ggtcacagct ttccgttacac gttcggaggg 300
 gggaccaagc tggaaataaa a 321

10 <210> 97

<211> 107

<212> PRT

<213> Mouse

15 <400> 97

Asp Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
 1 5 10 15
 Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Asp Tyr
 20 25 30
 Leu Tyr Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
 35 40 45
 Lys Phe Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Thr Gly
 50 55 60
 Ser Gly Ser Gly Ser Asp Phe Thr Leu Ser Ile Asp Ser Val Glu Pro
 65 70 75 80
 Asp Asp Val Gly Val Phe Phe Cys Gln Asn Gly His Ser Phe Pro Phe
 85 90 95
 Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

30

<210> 98

<211> 321

<212> DNA

<213> Mouse

35

<400> 98

40 gacattgtga tgactcagtc tccagccacc ctgtctgtga ctccaggaga tagagtctct 60
 ctttcctgca gggccagcca gagtattttac gactatttatc actggtatca aaaaaatca 120
 catgagtctc caaggctct catcaaataat gttcccaat ccatctctgg gatcccctcc 180
 aggttcagtg gcagtggatc aggctcagat ttcaactctca gtatcaacag tgtgaaacct 240
 gaagatgttg gagtttatta ctgtcaaaaat ggtcacagct ttccatttacac gttcggctcg 300
 gggacaaagt tggaaataaa a 321

45

<210> 99

<211> 107

<212> PRT

<213> Mouse

50

<400> 99

55

Asp Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
 1 5 10 15
 Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Glu Tyr
 20 25 30
 Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
 35 40 45
 Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Ser Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Pro
 65 70 75 80
 Glu Asp Val Gly Val Tyr Tyr Cys Gln Asn Gly His Ser Phe Pro Phe
 85 90 95
 Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

15

<210> 100
 <211> 321
 <212> DNA
 <213> Mouse

20

<400> 100
 gacattgtga tgactcagtc tccagccacc ctgtctgtga ctccaggaga tagagtctct 60
 ctctctgtca gggccagcca gagtatttagc gtctacttac actggtatca aaaaaatca 120
 cctgagtctc caaggcttct catcaaataat gttccaaat ccattcttg gatccccctcc 180
 aggttcagtg gcagtggtatc aggttcagat ttcaacttca gtatcaacag tgtggAACCT 240
 gaagatgttg gagtttatta ctgtcaaaaat ggtcacagct ttccattcac gttcggtctcg 300
 gggacaaaat tggaaataaaa a 321

25

<210> 101
 <211> 107
 <212> PRT
 <213> Mouse

30

<400> 101

35

Asp Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
 1 5 10 15
 Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Val Tyr
 20 25 30
 Leu His Trp Tyr Gln Gln Lys Ser Pro Glu Ser Pro Arg Leu Leu Ile
 35 40 45
 Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Ser Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Pro
 65 70 75 80
 Glu Asp Val Gly Val Tyr Tyr Cys Gln Asn Gly His Ser Phe Pro Phe
 85 90 95
 Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

40

<210> 102
 <211> 321
 <212> DNA
 <213> Mouse

45

<400> 102

gacattgtga	tgactcagtc	tccagccacc	ctgtctgtga	ctccaggaaa	tagagtctct	60
ctttccctgca	gggccagcca	gagtattagc	gactacttac	actggtatca	acaaaaatca	120
catgagtctc	caaggcttct	catcaaataat	gcttcccaat	ccatctctgg	gatcccctcc	180
aggttcagtg	gcagtggatc	agggtcaaat	ttcactctca	gtatcaacag	tgtggAACCT	240
gaagatgttg	gagtgtatta	ttgtcaaaat	ggtcacagct	ttccatttcac	gttcggctcg	300
gggacaagaat	tggaaaataaa	a				321

<210> 103

<211> 107

10 <212> PRT

<213> Mouse

<400> 103

<210> 104

30 <211> 333

<212> DNA

<213> Mouse

<400> 104

35

```

gacattgtgc tgacacagtc tcctgcttcc ttagctgttt ctctggggca gagggccacc 60
atctcatgca gggccagcaa aagtgtcaga ttctctgact atttttatat gcactgtac 120
caacagaaaac caggacagcc acccaaactc ctcatactacc ttgcataccaa cctagaatct 180
gggttcctcg ccaggttcaag tggcagtggg tctgggacag acttcaccct caacatccat 240
cctgtggagg aggaggatgc tgcaacctat tactgtcaga acagtaggga gttccgtac 300
acgttccggag gggggaccaa gctggagata aaa 333

```

<210> 105

<211> 111

45 <212> PRT

<213> Mouse

<400> 105

50

55

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Lys Ser Val Arg Phe Ser
 20 25 30
 Asp Tyr Phe Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45
 Lys Leu Leu Ile Tyr Leu Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
 50 55 60
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His
 65 70 75 80
 Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Asn Ser Arg
 85 90 95
 Glu Leu Pro Tyr Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

15 <210> 106
 <211> 333
 <212> DNA
 <213> Mouse

20 <400> 106

gacattgtgt tgacacagtc tcctgcttcc ttagctgtat ctctggggca gagggccacc 60
 atctcatgca gggccagcaa aagtgtcagt acatctggct atagttatat acactggta 120
 caacagaaac cgggacagcc acccaaattc ctcatctatc ttgcataccaa cctagattct 180
 ggggtccctg ccagggttcag tggcagtggg tctggacag acttcacccat caacatccat 240
 cctgtggagg aggaggatgc tgcaacctat tactgtcagc acagtaggga gcttccgtac 300
 acgttcggag gggggaccaa gctggaaaata aaa 333

25 <210> 107
 <211> 111
 <212> PRT
 <213> Mouse

30 <400> 107

35 Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Lys Ser Val Ser Thr Ser
 20 25 30
 40 Gly Tyr Ser Tyr Ile His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45
 Lys Phe Leu Ile Tyr Leu Ala Ser Asn Leu Asp Ser Gly Val Pro Ala

45 50 55 60
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His
 65 70 75 80
 Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Ser Arg
 85 90 95
 50 Glu Leu Pro Tyr Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

55 <210> 108
 <211> 321
 <212> DNA
 <213> Mouse

<400> 108

5 gacattgtga tgactcagtc tccagccacc ctgtctgtga ctccaggaga tagagtctct 60
 ctttcctgca gggccagcca gagtattagc gactacttac actggtatca aaaaaaatca 120
 catgagtctc caaggcttct catcaaataat gttcccaat ccatctctgg gatcccctcc 180
 aggttcagtg gcagtggatc aggtcagat ttcaacttca gtatcaacag tgtgaaacct 240
 gaagatgttgc gagtgttatta ctgtcaaaaat ggtcacagct ttccgtacac gttcgaggg 300
 gggaccaagc tggaaataaa à 321

<210> 109

<211> 107

10 <212> PRT

<213> Mouse

<400> 109

15 Asp Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
 1 5 10 15
 Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Ser Ile Ser Asp Tyr
 20 25 30
 Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
 25 35 40 45
 Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Ser Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Pro
 65 70 75 80
 Glu Asp Val Gly Val Tyr Tyr Cys Gln Asn Gly His Ser Phe Pro Tyr
 75 85 90 95
 Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> 110

30 <211> 336

<212> DNA

<213> Mouse

<400> 110

35

gatgttgtga tgacccaaac tccactctcc ctgcctgtca gtcttgaga tcaaggctcc 60
 atctcttgc gatctagtca gagccttgta cacagtaatg gaaacaccta tttacattgg 120
 tacctgcaga agccaggcca gtctccaaag ctcctgatct acaaagttc caaccgattt 180
 tctggggtcc ccgacagggtt cagtggcagt ggatcaggga cagattcac actcaagatc 240
 agcagagtgg aggctgagga tctggagtt tatttctgct ctcaaagtac acatgttccg 300
 tacacgttgc gaggggggac caagctggaa ataaat 336

<210> 111

<211> 112

45 <212> PRT

<213> Mouse

<400> 111

50

55

Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15
 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 5 Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 10 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser
 85 90 95
 Thr His Val Pro Tyr Thr Phe Gly Gly Thr Lys Leu Glu Ile Asn
 100 105 110

15 <210> 112
 <211> 318
 <212> DNA
 <213> Mouse

20 <400> 112

25 gacattatga tgaccaggc tcacaaattc atgtccacat cagtaggaga cagggtcagc 60
 atcacctgca aggccagtca ggatgtgggt aatgctgtgg tctggtatca aaaaaacca 120
 gggcaatctc ctaaactact gatttactgg gcatccaccc gcacattgg agtccctgat 180
 cgcttcacag gcagtggatc tggacagat ttcactctca ccattaccaa tgtgcagtct 240
 gaagacttgg cagattattt ctgtcagcaa tatagcagct atctcacgtt cggtgctggg 300
 accaagctgg agctgaaa 318

30 <210> 113
 <211> 106
 <212> PRT
 <213> Mouse

35 <400> 113

Asp Ile Met Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Gly
 1 5 10 15
 Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Gly Asn Ala
 20 25 30
 40 Val Val Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile
 35 40 45
 Tyr Trp Ala Ser Thr Arg His Ile Gly Val Pro Asp Arg Phe Thr Gly
 50 55 60
 45 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Thr Asn Val Gln Ser
 65 70 75 80
 Glu Asp Leu Ala Asp Tyr Phe Cys Gln Gln Tyr Ser Ser Tyr Leu Thr
 85 90 95
 Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 100 105

50 <210> 114
 <211> 336
 <212> DNA
 <213> Mouse

55 <400> 114

5 gatgttgtga tgacccaatc tccactctcc ctgcctgtca gtcttgaga tcaaggctcc 60
 atctcttgca gatctagtca gagccttgc cacagtaatg gaaacaccta tctacattgg 120
 tacctgtca agccaggcc a gtc tccaaag ctc ctgatct acaaagtttc caaccgattt 180
 tctgggtcc cagacagggtt c agtggcagt ggatcaggga cagattcac actcaagatc 240
 agcagagtgg aggctgagga tctgggagtt tatttctgct ctcaaagtac acatgttccg 300
 tacacgttcg gagggggac caagctggaa ataaaa 336

10 <210> 115

<211> 112

<212> PRT

<213> Mouse

15 <400> 115

Asp	Val	Val	Met	Thr	Gln	Ser	Pro	Leu	Ser	Leu	Pro	Val	Ser	Leu	Gly	
1					5				10				15			
Asp	Gln	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Val	His	Ser	
								20				25			30	
Asn	Gly	Asn	Thr	Tyr	Leu	His	Trp	Tyr	Leu	Leu	Lys	Pro	Gly	Gln	Ser	
								35				40			45	
Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro	
								50			55			60		
Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile	
								65			70			75		80
Ser	Arg	Val	Glu	Ala	Glu	Asp	Leu	Gly	Val	Tyr	Phe	Cys	Ser	Gln	Ser	
								85			90			95		
Thr	His	Val	Pro	Tyr	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	
								100			105			110		

30 <210> 116

<211> 321

<212> DNA

<213> Mouse

35 <400> 116

40 gacatccaga tgactcagtc tccagtttc ctgtctgcat ctgtggaga aactgtcacc 60
 atcacatgtc gagcaagtga gaatattgac agttatggat catggatca gcagaaacag 120
 ggaaaatctc ctcagctcct ggtcttgct gcacgaaact tagcagatgg tgtgccatca 180
 aggtcagtg gcagtggatc aggacacag tattctctca agatcaacag aatgcagtct 240
 gaagatgtt cgagatacta ctgtcaacat tattatagta ctccattcac gttcggtcg 300
 gggacaaagt tggaaataga a 321

45 <210> 117

<211> 107

<212> PRT

<213> Mouse

50 <400> 117

Asp Ile Gln Met Thr Gln Ser Pro Val Phe Leu Ser Ala Ser Val Gly
 1 5 10 15
 Glu Thr Val Thr Ile Thr Cys Arg Ala Ser Glu Asn Ile Asp Ser Tyr
 20 25 30
 5 Leu Ala Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val
 35 40 45
 Phe Ala Ala Arg Asn Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Arg Met Gln Ser
 65 70 75 80
 10 Glu Asp Val Ala Arg Tyr Tyr Cys Gln His Tyr Tyr Ser Thr Pro Phe
 85 90 95
 Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Glu
 100 105

15 <210> 118
 <211> 321
 <212> DNA
 <213> Mouse

20 <400> 118
 gacatccaga tgactcagtc gccagcttcc ctgtctgcat ctgtgggaga aactgtcatc 60
 ttcacatgtc gagcaagtga gaatattgac agttatattag catggtatca gcagaaacag 120
 25 gaaaaatctc ctcagctect ggtctatgct gcaacaaact tagcagatgg tgtgccatca 180
 aggttcagtg gcagtggatc aggcacacag tattctctca agatcaacag cctgcagtct 240
 gaagatgttg cgagatatta ctgtctacat tattatagta ctccattcac gttcggctcg 300
 gggacagagt tggaaataaa a 321

30 <210> 119
 <211> 107
 <212> PRT
 <213> Mouse

<400> 119

35 Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Glu Thr Val Ile Phe Thr Cys Arg Ala Ser Glu Asn Ile Asp Ser Tyr
 20 25 30
 40 Leu Ala Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val
 35 40 45
 Tyr Ala Ala Thr Asn Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Ser
 65 70 75 80
 45 Glu Asp Val Ala Arg Tyr Tyr Cys Leu His Tyr Tyr Ser Thr Pro Phe
 85 90 95
 Thr Phe Gly Ser Gly Thr Glu Leu Glu Ile Lys
 100 105

50 <210> 120
 <211> 357
 <212> DNA
 <213> Mouse

<400> 120

5 caggtgcagc tgaaggagtc aggacctggc ctggggcgcc cctcacagag cctgtccatc 60
 acatgcactg tctctgggtt ctcatthaacc accttatgcta taagctgggt tcgcagtca 120
 ccaggaaagg gtctggagtg gcttggagtc atatggactg gtggaggcac aaaatataat 180
 tcagctctca aatccagact gagcatcagc aaagacaact ccaagagtca agtttctta 240
 aaaatgaaca gtctgcaaac tgatgacaca gccaggtact actgtgccag aaaggactat 300
 aattacgggg gtgctatgga ctactgggtt caaggaacct cagtcaccgt ctcctca 357

10 <210> 121

<211> 119

<212> PRT

<213> Mouse

15 <400> 121

Gln	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Ser	Gln
1															15
Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Tyr	
															30
20															
Ala	Ile	Ser	Trp	Val	Arg	Gln	Ser	Pro	Gly	Lys	Gly	Ieu	Glu	Trp	Leu
															45
25															
Gly	Val	Ile	Trp	Thr	Gly	Gly	Gly	Thr	Lys	Tyr	Asn	Ser	Ala	Leu	Lys
															50
50															
Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Asn	Ser	Lys	Ser	Gln	Val	Phe	Leu
															80
65															
Lys	Met	Asn	Ser	Leu	Gln	Thr	Asp	Asp	Thr	Ala	Arg	Tyr	Tyr	Cys	Ala
															95
Arg	Lys	Asp	Tyr	Asn	Tyr	Gly	Gly	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly
															100
															110

30 Thr Ser Val Thr Val Ser Ser
 115

<210> 122

<211> 360

35 <212> DNA

<213> Mouse

<400> 122

40 caggttcagc tgcaacagtc tgacgcttag ttgggtaaaac ctggcacttc agtgaagaca 60
 tcctgcaaaa ttctggcaa caccttact gaccaaacta ttcactggat gaägcagagg 120
 cctgaacagg gcctggaaatg gattggatat atttataacta gagatgatag tcctaagtac 180
 aatgagaact tcaaggcää ggccacattg actgcagaca aatccctccag cacagcctac 240
 atgcagacta acagtctgac atctgaggac tctgcagtct atttctgtgc aatcccagac 300
 45 aggtcaggct acgcctgggtt tatttactgg ggccaaggga ctctggcac tgtctttca 360

<210> 123

<211> 120

<212> PRT

50 <213> Mouse

<400> 123

Gln Val Gln Leu Gln Gln Ser Asp Ala Glu Leu Val Lys Pro Gly Thr
 1 5 10 15
 Ser Val Lys Thr Ser Cys Lys Ile Ser Gly Asn Thr Phe Thr Asp Gln
 20 25 30
 5 Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 10 Met Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
 85 90 95
 Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
 100 105 110
 Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 124

<211> 357

<212> DNA

20 <213> Mouse

<400> 124

25 caggtgcagc tgaaggagtc aggacctgtc ctggggcgcc cttcacagag cctgtccatc 60
 acatgcactg tctctgggtt ctcatataaac aactttgcta taagttgggt tcgtcagcca 120
 ccagggaaagg gtctggagtg gcttggagca atatggactg gtggaggcac aaattataat 180
 tcagctctca aatccagact gagcatcagc aaagacaact ccaagagtca agttttctta 240
 aaaatgaaca gtctgcaaac ttagtacaca gccaggattt atttgtgtcag aaaggactat 300
 agttacgggg gtgctatgga ctactgggtt caaggaacct cagtcaccgt ctccctca 357

30

<210> 125

<211> 119

<212> PRT

35 <213> Mouse

<400> 125

40 Gln Val Gln Leu Lys Glu Ser Gly Pro Val Leu Val Ala Pro Ser Gln
 1 5 10 15

Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Asn Asn Phe
 20 25 30
 Ala Ile Ser Trp Val Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Leu
 35 40 45
 45 Gly Ala Ile Trp Thr Gly Gly Thr Asn Tyr Asn Ser Ala Leu Lys
 50 55 60
 Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Phe Leu
 65 70 75 80
 Lys Met Asn Ser Leu Gln Thr Asp Asp Thr Ala Arg Tyr Tyr Cys Val
 85 90 95
 Arg Lys Asp Tyr Ser Tyr Gly Gly Ala Met Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Ser Val Thr Val Ser Ser
 115

55

<210> 126

<211> 354

<212> DNA

<213> Mouse

<400> 126

5 gaggccagc tgcaacagtc tggacctgtc ctggtaagg ctggggcttc agtgaagatg 60
 tcctgttaagg cttctggata cacattcaact gactactata tgaactgggt gaagcagagc 120
 catggaaaga gccttgagtg gattggagtt attattcctt acaacggtgg tactagctac 180
 aaccagaagt tcaaggccaa ggccacattg actgttgaca agtcctccag cacagcctac 240
 atggagctca acagecctgac atctgaggac tctgcagtctt attactgtgc acgagatgg 300
 10 cacccgttgtt acttcgatgt ctggggcaca gggaccacgg tcaccgtctc ctca 354

<210> 127

<211> 118

<212> PRT

15 <213> Mouse

<400> 127

20 Glu Val Gln Leu Gln Gln Ser Gly Pro Val Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 Tyr Met Asn Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
 35 40 45
 Gly Val Ile Ile Pro Tyr Asn Gly Gly Thr Ser Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asp Gly His Arg Trp Tyr Phe Asp Val Trp Gly Thr Gly Thr
 100 105 110
 Thr Val Thr Val Ser Ser
 115

35 <210> 128

<211> 115

<212> PRT

<213> Mouse

40 <400> 128

45 Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Cys Cys
 20 25 30

 Ile Met His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Asn Pro Tyr Asn Asp Gly Thr Lys Tyr Asn Glu Lys Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Arg Trp Asp Glu Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ala
 115

<210> 129
<211> 345
<212> DNA
<213> Mouse

5

<400> 129

gagggtccagc tgcagcagtc tggacctgag ctggtaaagc ctggggcttc agtgaagatg 60
tcctgcaagg cttctggata cacattcaact cgctatctta ttcactgggt gaaacagaag 120
cctgggcagg gccttgagtg gattggatat attaattcctt acaatgatgg tactaaatac 180
aatgagaagt tcaaaggcaa ggccacactg acttcagaca aatcctccag cacagcctac 240
atggagctca gcagcctgac ctctgaggac tctgcggctt attactgtac ctctaactgg 300
gacotcgact actggggcca aggaccact ctcacagtct cctca 345

15

<210> 130
<211> 115
<212> PRT
<213> Mouse

20

<400> 130

Glu	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala
1															15
Ser	Val	Lys	Met	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Arg	Tyr
															30
Leu	Ile	His	Trp	Val	Lys	Gln	Lys	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile
															45
Gly	Tyr	Ile	Asn	Pro	Tyr	Asn	Asp	Gly	Thr	Lys	Tyr	Asn	Glu	Lys	Phe
															50
Lys	Gly	Lys	Ala	Thr	Leu	Thr	Ser	Asp	Lys	Ser	Ser	Ser	Thr	Ala	Tyr
															65
Met	Glu	Leu	Ser	Ser	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys
															80
Thr	Ser	Asn	Trp	Asp	Leu	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Thr	Leu	Thr
															95
35	Val	Ser	Ser												100
															105
															110
															115

25

30

35

40

<210> 131
<211> 345
<212> DNA
<213> Mouse

<400> 131

45

gagggtccagc tgcagcagtc tggacctgaa gtggtaaagc ctggggcttc agtgaagatg 60
tcctgcaagg cctctggata cacattcaact agttctgtta tacactgggt gaagcagaag 120
gctgggcagg gccttgagtg gattggatat atcaatccct ataatgatgg tactaagtac 180
aatgagaagt tcaaaggcaa ggccacactg acttcagaca gatcctccag cacagcctac 240
atggagctca gcagcctgac ctctgaggac tctgcggctt attactgtac aagacggtg 300
gacgaggctt actggggcca aggactctg gtcactgtct ctgca 345

50

<210> 132
<211> 115
<212> PRT
<213> Mouse

<400> 132

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Val Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Ser
 20 25 30
 Val Ile His Trp Val Lys Gln Lys Ala Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Asn Pro Tyr Asn Asp Gly Thr Lys Tyr Asn Glu Lys Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Ser Asp Arg Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Thr Arg Arg Leu Asp Glu Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ala
 115

<210> 133

<211> 345

<212> DNA

20 <213> Mouse

<400> 133

25 gaggtccagc tgcagcagtc tggacctgag ctggtaaagc ctggggctgc agtgaagatg 60
 tcctgcaagg cttctggata cacattcaact cgctatctta ttcaactgggt gaagcagaag 120
 cctggcagg gccttgagtg gattggatat attaatcctt acaatgatgg tactaaatat 180
 aatgagaagt tcaaaggcaa ggccacactg acttcagaca aatcctccag cacagcctac 240
 atggagctca gcagcctgac ctctgaggac tctgcggtct attactgtac ctctaattgg 300
 gacctcgact actggggcca aggcaccact ctcacagtct cctca 345

30 <210> 134
 <211> 115
 <212> PRT
 <213> Mouse

35 <400> 134

40 Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ala Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30
 Leu Ile His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Asn Pro Tyr Asn Asp Gly Thr Lys Tyr Asn Glu Lys Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Thr Ser Asn Trp Asp Leu Asp Tyr Trp Gly Gln Gly Thr Thr Leu Thr
 100 105 110
 Val Ser Ser
 115

55 <210> 135

<211> 345

<212> DNA

<213> Mouse

<400> 135

5 gagggtccagc tgcagcagtc tggacctgaa ctggtaaagc ctggggcttc agtgaagatg 60
 tcctgcaagg cctctggata cacattcaact agttctgtta tgcactgggt gaagcagaag 120
 gctggcagg gccttgagtg gattggatat atcaatccct ataatgatgg tactcagtag 180
 aatgagaagt tcaaaggcaa ggccacactg acttcagaca aattttccag cacagcctac 240
 atggagctca gcagcctgac ctctgaggac tctgcgtct attactgtac aagacggttg 300
 gacgaggctt actggggcca agggactctg gtcactgtct ctgca 345

10 <210> 136

<211> 115

<212> PRT

<213> Mouse

15 <400> 136

Glu	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala
1															15
Ser	Val	Lys	Met	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser	Ser
															30
20															
Val	Met	His	Trp	Val	Lys	Gln	Lys	Ala	Gly	Gln	Gly	Leu	Glu	Trp	Ile
															45
35															
Gly	Tyr	Ile	Asn	Pro	Tyr	Asn	Asp	Gly	Thr	Gln	Tyr	Asn	Glu	Lys	Phe
															50
50															
Lys	Gly	Lys	Ala	Thr	Leu	Thr	Ser	Asp	Lys	Phe	Ser	Ser	Thr	Ala	Tyr
															80
25															
65															
Met	Glu	Leu	Ser	Ser	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys
															95
85															
Thr	Arg	Arg	Leu	Asp	Glu	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr
															100
30															
Val	Ser	Ala													
															110
115															

<210> 137

<211> 345

35 <212> DNA

<213> Mouse

<400> 137

40

gagggtccagc tgcagcagtc tggacctgag ctggtaaagc ctggggcttc agtgaagatg 60
 tcctgcaagg cttctggata cacattcaact agctctatta ttcaactgggt gaagcagagg 120
 cctggcagg gccttgagtg gattggatat attaattcctt acgtatgtgt tactaagtac 180
 aatgagaagt tcaaaggcaa ggccacactg acttcagaca aatcctccag cacagcctac 240
 atggagctca gcagcctgac ctctgaggac tctgcgtct attactgtgc aagacggttg 300
 gacgagtctt actggggcca agggactctg gtcactgtct ctgca 345

<210> 138

<211> 115

50 <212> PRT

<213> Mouse

<400> 138

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Ser
 20 25 30
 Ile Ile His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Asn Pro Tyr Asp Asp Val Thr Lys Tyr Asn Glu Lys Phe
 50 55 60

10 Lys Gly Lys Ala Thr Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Arg Trp Asp Glu Ser Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ala
 115

<210> 139

<211> 345

<212> DNA

<213> Mouse

<400> 139

gaggtccagc tgcagcagtc tggacctgag ctggtaaagc ctggggcttc agtgaagatg 60
 tcctgcagg cttctggata cacttcaact acctctatta tgcactgggt gaaacagaag 120
 cctgggcagg gccttgagtg gattggatat attaattcctt acgtatgtt tactaagtac 180
 30 aatgaaaagt tcaaaggcaa gcccacattt acttcagaca aatcctccag cacagcctac 240
 atggagctca gcagcctgac ctctgaggac tctgcaatgtt attactgtgt aagacgggtgg 300
 gacgaggctt actggggcca agggactctg gtcactgtct ctgca 345

<210> 140

<211> 115

<212> PRT

<213> Mouse

<400> 140

40 Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Thr Ser
 20 25 30
 Ile Met His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Asn Pro Tyr Asp Asp Val Thr Lys Tyr Asn Glu Lys Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Val Arg Arg Trp Asp Glu Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ala
 115

<210> 141

<211> 357

<212> DNA

<213> Mouse

<400> 141

5

gaggtccagc tgcaacaatc tggacctgag ctggtaagg ctggggcttc agtgaagata 60
 tcctgttaagg cttctggata cacgttact gactactaca tgaactgggt gaggcagagc 120
 catggagaga gccttgagtg gattggagat tttaatcata acaatgatgt tattactac 180
 aaccggaaat tcaagggcaa ggtcacattg actgttagaga agtcttccac cacagcctac 240
 atggagctcc gcagcctgtc atctgaggac tctgcagtct attactgtgc aagggggcta 300
 cgaggctact atgctatgga ctactgggt caaggaacct cagtcaccgt ctccctca 357

10

<210> 142

15

<211> 119

<212> PRT

<213> Mouse

<400> 142

20

Glu	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala
1									10					15	
Ser	Val	Lys	Ile	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Tyr
									20	25				30	
Tyr	Met	Asn	Trp	Val	Arg	Gln	Ser	His	Gly	Glu	Ser	Leu	Glu	Trp	Ile
									35	40				45	
Gly	Asp	Phe	Asn	His	Asn	Asn	Asp	Val	Ile	Thr	Tyr	Asn	Pro	Lys	Phe
									50	55				60	
Lys	Gly	Lys	Val	Thr	Leu	Thr	Val	Glu	Lys	Ser	Ser	Thr	Thr	Ala	Tyr
									65	70				75	80
Met	Glu	Leu	Arg	Ser	Leu	Ser	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys
									85	90				95	
Ala	Arg	Gly	Leu	Arg	Gly	Tyr	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly
									100	105				110	
Thr	Ser	Val	Thr	Val	Ser	Ser									
									115						

25

30

35

<210> 143

<211> 354

<212> DNA

40

<213> Mouse

<400> 143

caggttcagc tgcaacagtc tgacgctgag ttggtaaac ctggagcttc agtgaagata 60
 tcctgttaagg tttctggata caccttact gaccatacta ttcactggat gaagcagagg 120
 cctgaacagg gccttggatg gattggatat atttatacata gagatggta tcctaagttc 180
 aatgagaagt tcaagggcaa gcccacattg actgcagaca aatcctccag cacagcctac 240
 atgcagctca acagcctgac atctgaggac tctgcagtct atttctgtgc aagacggccc 300
 ctttactatg ctatggacta ctgggtcaa ggaacctcag tggccgttcc ctca 354

50

<210> 144

<211> 118

<212> PRT

55

<213> Mouse

<400> 144

Gln Val Gln Leu Gln Gln Ser Asp Ala Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Ile Ser Cys Lys Val Ser Gly Tyr Thr Phe Thr Asp His
 20 25 30
 5 Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Tyr Pro Arg Asp Gly Tyr Pro Lys Phe Asn Glu Lys Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 10 Met Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
 85 90 95
 Ala Arg Arg Pro Pro Tyr Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr
 100 105 110
 15 Ser Val Ala Val Ser Ser
 115

<210> 145

<211> 345

<212> DNA

20 <213> Mouse

<400> 145

25 gaggtccaac tgcagcagtc tggacctgag ctggtaaagg ctggggcttc agtgaagatg 60
 tcctgcaagg cttctggata cacattcaact aggtatctta tgcactgggt gaagcagaag 120
 cctgggcagg gccttgagtg gattggttat attaatcctt acaatgatgg tactaattac 180
 aatgagaagt tcaaaggcaa ggccacactg acttcagaca aatcctccag cacagcctac 240
 atggagctca gcagcctgac ctctgaggac tctgcggctt attactgttc ccttaactgg 300
 gactatgctt actggggcca agggactctg gtcactgtct ctgca 345

30 <210> 146
 <211> 115
 <212> PRT
 <213> Mouse

35 <400> 146

40 Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30
 Leu Met His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Asn Pro Tyr Asn Asp Gly Thr Asn Tyr Asn Glu Lys Phe
 50 55 60
 45 Lys Gly Lys Ala Thr Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 50 Ser Leu Asn Trp Asp Tyr Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ala
 115

55 <210> 147

<211> 360

<212> DNA

<213> Mouse

<400> 147

5 gagttccagc tgcagcagtc tggacctgag ctggtaaagc ctggcgcttc agtgaagata 60
 tcctgcaagg cttctggta ctcattcaact gactacaaca tgaactgggt gaagcagagc 120
 aaaggaaaaga gccttgagtg gattggagta attattccta actatggtt tactagctac 180
 aatcagaact tcaaggccaa gcccacattg actgttagacc agtctccag cacagccac 240
 atgcagctca acagtgtgac atctgaggac tctgcagtct attactgtgt aagagatggg 300
 ggaatactcc tctggtatct cgatgtctgg ggcacaggga ccacggcac cgtccctca 360

10

<210> 148

<211> 120

<212> PRT

<213> Mouse

15

<400> 148

20 Glu Phe Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
 20 25 30
 Asn Met Asn Trp Val Lys Gln Ser Lys Gly Lys Ser Leu Glu Trp Ile
 35 40 45
 Gly Val Ile Ile Pro Asn Tyr Gly Phe Thr Ser Tyr Asn Gln Asn Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Val Asp Gln Ser Ser Ser Thr Ala His
 65 70 75 80
 Met Gln Leu Asn Ser Val Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys

30

25 85 90 95
 Val Arg Asp Gly Gly Ile Leu Leu Trp Tyr Leu Asp Val Trp Gly Thr
 100 105 110
 Gly Thr Thr Val Thr Val Ser Ser
 115 120

35

<210> 149

<211> 366

<212> DNA

<213> Mouse

<400> 149

40 gagggtccagc tgcaacagtc tggacctgag ctggtaaagc ctggggcttc agtgaagata 60
 ccctgcaagg cttctggata cacattcaact gactacaaca tggactgggt gaagaagagc 120
 catggaaaaga gccttgagtg gattggagat atcaatcctc acaatggtt tactatctac 180
 aaccagaagt tcaaggccaa gcccacattg actgttagaca agtctccag cacagccac 240
 atggagctcc gcagcctgac atctgaggac actgcagtct attactgtgc aagaaaattac 300
 tacgttagta gttacggctg gtacttcgat gtctgggca cagggaccac ggtcaccgtc 360
 tcgtca 366

50

<210> 150

<211> 122

<212> PRT

<213> Mouse

<400> 150

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Ile Pro Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 5 Asn Met Asp Trp Val Lys Lys Ser His Gly Lys Ser Leu Glu Trp Ile
 35 40 45
 Gly Asp Ile Asn Pro His Asn Gly Gly Thr Ile Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala His
 65 70 75 80
 10 Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asn Tyr Tyr Gly Ser Ser Tyr Gly Trp Tyr Phe Asp Val Trp
 100 105 110
 Gly Thr Gly Thr Thr Val Thr Val Ser Ser
 115 120

<210> 151

<211> 360

<212> DNA

20 <213> Mouse

<400> 151

25 gaggtccagc tgcagcagtc tggaccttag ctggtaaggc ctggggcttc agtgaagatt 60
 tcctgcaggg cttctggta ctcattcaact ggctactaca tgaactgggt gaagcaaagt 120
 cctgaaaaga gccttgagtg gattggagag attattccta ccactgggt tacttcctac 180
 aaccagaagt tcaaggccaa ggccacattg actgttagaca aatcctccag cacagcctac 240
 atgcagctca agagcctgac atctgaggac tctgcagtctt attactgtgc aagagagagc 300
 ggtgggttctt actggtaactt cgatgtctgg ggcacaggga ccacggcac cgttcctca 360

30

<210> 152

<211> 120

<212> PRT

35 <213> Mouse

<400> 152

40 Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Ile Ser Cys Arg Ala Ser Gly Tyr Ser Phe Thr Gly Tyr
 20 25 30
 45 Tyr Met Asn Trp Val Lys Gln Ser Pro Glu Lys Ser Leu Glu Trp Ile
 35 40 45
 Gly Glu Ile Ile Pro Thr Thr Gly Gly Thr Ser Tyr Asn Gln Lys Phe
 50 55 60
 Lys Ala Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Gln Leu Lys Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Glu Ser Gly Gly Phe Tyr Trp Tyr Phe Asp Val Trp Gly Thr
 100 105 110
 Gly Thr Thr Val Thr Val Ser Ser
 115 120

55

<210> 153

<211> 345

<212> DNA

<213> Mouse

<400> 153

5 gaggccatc tgcagcgtc tggacctgag ctggtaaagc ctggggcttc agtgaagatg 60
 tcctcaagg cttctggata cacattcaact aggtatgtta tgcactgggt gaagcagaag 120
 cctggcagg gccttgagtg gattggatat attaatcctt acaatgatgt tactaagtac 180
 aatgagaagt tcaaaggcaa gcccacactg acttcagaca gatcctccag cacagctac 240
 10 atgaaactca gcagcctgac ctctgaggac tctgcggctt attattgtgc aagaaaactgg 300
 gacgttccctt actggggcca agggactctg atcactgtct ctgca 345

<210> 154

<211> 115

<212> PRT

<213> Mouse

<400> 154

20 Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30
 Val Met His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Asn Pro Tyr Asn Asp Val Thr Lys Tyr Asn Glu Lys Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Ser Asp Arg Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 25 Met Lys Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asn Trp Asp Val Pro Tyr Trp Gly Gln Gly Thr Leu Ile Thr
 100 105 110
 Val Ser Ala
 115

<210> 155

<211> 345

<212> DNA

<213> Mouse

<400> 155

35 gaggccatc tgcagcgtc tggacctgag ctggtaaagc ctggggcttc agtgaagatg 60
 tcctcaagg cttctggata cacattcaact aggtatgtta tgcactgggt gaagcagaag 120
 cctggcagg gccttgagtg gattggatat attaatcctt acaatgatgg tactaagtac 180
 aatgagaggt tcaaaggcaa gcccacactg acttcagaca aatcctccag cacagctac 240
 atggagctca gcagcctgac ctctgaggac tctgcggctt attactgtgc aagaaaactgg 300
 gacgttccctt actggggcca agggactctg gtcactgtct ctgca 345

<210> 156

<211> 115

<212> PRT

<213> Mouse

<400> 156

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30
 5 Leu Met His Trp Val Lys Gln Lys Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Asn Pro Tyr Asn Asp Gly Thr Lys Tyr Asn Glu Arg Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Ser Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 10 Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asn Trp Asp Val Pro Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 15 Val Ser Ala
 115

<210> 157

<211> 321

<212> DNA

20 <213> Artificial Sequence

<220>

<223> Humanized sequence.

25 <400> 157

gacatccaga tgacccagag cccaaaggcgc ctgagcgcca gcgtggcga ccgcgtgacc 60
 atcacctgca aggccagccg cgacgtggcc atcgccgtgg cctggtagcca gcagaagcca 120
 ggcaaggtgc caaagctgct gctgttctgg gccagcaccc gccacacccgg cgtgccagac 180
 cgcttcagcg gcagcggcag cggcaccgac ttccacctga ccatcagcag cctgcagcca 240
 gaggacctgg ccgactacta ctgccaccag tacagcagct acccattcac cttcggccag 300
 ggcaccaagc tggagatcaa g 321

35 <210> 158

<211> 107

<212> PRT

<213> Artificial Sequence

40 <220>

<223> Humanized sequence.

<400> 158

45 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala
 20 25 30
 50 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu
 35 40 45
 Phe Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Ser Gly

50	55	60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro		
65	70	75
Glu Asp Leu Ala Asp Tyr Tyr Cys His Gln Tyr Ser Ser Tyr Pro Phe		
	85	90
Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys		
	100	105

<210> 159

10 <211> 321

<212> DNA

<213> Artificial Sequence

<220>

15 <223> Humanized sequence.

<400> 159

20	gacatccaga	tgacccagag	ccaaaggcgc	ctgagcgcca	gcgtgggcga	ccgcgtgacc	60
	atcacctgca	aggccagccg	cgacgtggcc	atcgccgtgg	cctggtatcca	gcagaagccca	120
	ggcaaggtgc	caaagctgtct	gatctactgg	gccagcaccc	gccacacccgg	cgtgccaagc	180
	cgcttcagcg	gcagcggcag	ccgcacccgac	ttcacccctga	ccatcagcag	cctgcagccca	240
	gaggacgtgg	ccgactactt	ctgccaccag	tacagcagct	accattcac	cttcggcagc	300
	ggcaccaagc	tggagatcaa	g				321

<210> 160

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Hun

13

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Ile	Ser	Ala	Ser	Val	Gly
1					5				10					15	
Asp	Arg	Val	Thr	Ile	Thr	Cys	Lys	Ala	Ser	Arg	Asp	Val	Ala	Ile	Ala
							20		25					30	
Val	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Val	Pro	Lys	Leu	Leu	Ile
						35		40				45			
Tyr	Trp	Ala	Ser	Thr	Arg	His	Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
						50		55		60					
Ser	Gly	Ser	Arg	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
						65		70		75				80	
Glu	Asp	Val	Ala	Asp	Tyr	Phe	Cys	His	Gln	Tyr	Ser	Ser	Tyr	Pro	Phe
							85		90					95	
Thr	Phe	Gly	Ser	Gly	Thr	Lys	Leu	Glu	Ile	Lys					
						100		105							

<210> 161

<211> 321

<212> DNA

<213> Artificial Sequence

<220>

<223> Humanized sequence.

<400> 161

5 gacatccaga tgacctcagag cccaaaggcgc ctgagcgcca gcgtggcgca ccgcgtgacc 60
 atcacctgca aggccagccg cgacgtggcc atcgcctgtt cctggtagcc gcagaagcca 120
 ggcaaggtgc caaaagctgtt gatctactgg gccagcaccc gccacacccgg cgtgccaaagc 180
 cgcttcagcg gcagcggcag ccgcaccgac ttcaccctga ccatcagcag cctgcagcca 240

10 gaggacgtgg ccgactactt ctgccaccag tacagcagct acccattcac cttcggcagc 300
 ggcaccaagc tggagatcaa g 321

15 <210> 162
 <211> 107
 <212> PRT
 <213> Artificial Sequence

20 <220>
 <223> Humanized sequence.

<400> 162

25 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu
 35 40 45
 Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Val Ala Thr Tyr Tyr Cys His Gln Tyr Ser Ser Tyr Pro Phe
 85 90 95
 Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

30 <210> 163
 <211> 321
 <212> DNA
 <213> Artificial Sequence

35 <220>
 <223> Humanized sequence.

<400> 163

40 gacatccaga tgacctcagag cccaaaggcgc ctgagcgcca gcgtggcgca ccgcgtgacc 60
 atcacctgca aggccagccg cgacgtggcc atcgcctgtt cctggtagcc gcagaagcca 120
 ggcaaggtgc caaaagctgtt gctgttctgg gccagcaccc gccacacccgg cgtgccagac 180
 cgcttcagcg gcagcggcag ccgcaccgac ttcaccctga ccatcagcag cctgcagcca 240
 gaggacctgg ccgactacta ctgccaccag tacagcagct acccattcac cttcggcagc 300
 ggcaccaagc tggagatcaa g 321

45 <210> 164
 <211> 107
 <212> PRT
 <213> Artificial Sequence

50 <220>
 <223> Humanized sequence.

<400> 164

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Leu
 35 40 45
 Phe Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

Glu Asp Leu Ala Asp Tyr Tyr Cys His Gln Tyr Ser Ser Tyr Pro Phe
 85 90 95
 Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> 165

<211> 360

<212> DNA

<213> Artificial Sequence

<220>

<223> Humanized sequence.

<400> 165

caggtgcagc tgggtgcagag cggcgccgag gtgaagaagc caggcagcag cgtgaagg 60
 agctgcaagg ccagcggtta caccttacc gaccagacca tccactggat gcccaggcc 120
 ccaggccagg gcctggagtg gatcggtac atctacccac gcgacgacag cccaaagtac 180
 aacgagaact tcaaggggcaa ggtcaccatc accgccgaca agagcaccag caccgcctac 240
 atggagctga gcagcctgctg cagcgaggac accgcgtgt actactgcgc catccccac 300
 cgcaagcggtt acgcctgggtt catctactgg ggccagggca ccctggtgac cgtgagcagc 360

<210> 166

<211> 120

<212> PRT

<213> Artificial Sequence

<220>

<223> Humanized sequence.

<400> 166

45

50

55

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Gln
 20 25 30
 5 Thr Ile His Trp Met Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
 50 55 60
 10 Lys Gly Lys Val Thr Ile Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
 100 105 110
 15 Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 167

<211> 360

<212> DNA

20 <213> Artificial Sequence

<220>

<223> Humanized sequence.

25 <400> 167

caggtgcagg tgggtgcagag cggcgccgag gtgaagaagc cagggcagcag cgtgaagggtg 60
 agctgcagg ccagcggtt caccttcacc gaccagacca tccactgggt ggcggccaggcc 120
 30 ccaggccagg gcctggagtg gatgggctac atctacccac ggcacgacag cccaaagtac 180
 aacgagaact tcaagggcaa ggtcacccctg accgcggaca agagcaccag caccgcctac 240

atggagctga gcagcctgcg cagcgaggac accgcccgtgt actactgcgc catccccac 300
 35 cgcagcggct acgcctgggtt catctactgg ggccaggca ccctggtgac cgtgagcagc 360

<210> 168

<211> 120

<212> PRT

40 <213> Artificial Sequence

<220>

<223> Humanized sequence.

45 <400> 168

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Thr Phe Thr Asp Gln
 20 25 30
 5 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
 50 55 60
 10 Lys Gly Lys Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
 100 105 110
 15 Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 169

<211> 360

<212> DNA

20 <213> Artificial Sequence

<220>

<223> Humanized sequence.

25 <400> 169

caggtgcagc tgggtgcagag cggcgccgag gtgaagaagc caggcagcag cgtgaagacc 60
 agctgcagg ccagcggcgg caccttcacc gaccagacca tccactgggt ggcgcagcgc 120
 30 ccagggcagg gcctggagtg gatgggctac atctacccac ggcacgacag cccaaagtac 180
 aacgagaact tcaagggccg cgtcaccatc accgcccaca agagcaccag caccgcctac 240
 atggagctga gcagcctgcg cagcgaggac accgcctgtt actactgcgc catccccagac 300
 cgcagcggct acgcctggtt catctactgg ggccaggcga ccctgggtac cgtgagcagc 360

<210> 170

35 <211> 120

<212> PRT

<213> Artificial Sequence

<220>

40 <223> Humanized sequence.

<400> 170

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 45 1 5 10 15
 Ser Val Lys Thr Ser Cys Lys Ala Ser Gly Gly Thr Phe Thr Asp Gln
 20 25 30
 Thr Ile His Trp Val Arg Gln Arg Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

50

55

Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
 50 55 60
 Lys Gly Arg Val Thr Ile Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
 100 105 110
 Gly Thr Leu Val Thr Val Ser Ser
 115 120

5

10

<210> 171

<211> 360

<212> DNA

15

<213> Artificial Sequence

<220>

<223> Humanized sequence.

20

<900> 171

25

caggtgcagc tggcgcagag cggcgccgag gtgaagaagc caggcagcag cgtgaagggtg 60
 agctgcagg ccagcggcg cacccacc gaccagacca tccactgggt gcgcaggcc 120
 ccagccagg gcctggagtg gatgggctac atctaccac gcgacgacag cccaaagtac 180
 aacgagaatt tcaaggcccgc cgtcaccctg accgcccaca agagcaccag caccgcctac 240
 atggagctga gcagcctgcg cagcgaggac accgcccgtgt acttctgcgc cccgcccagac 300
 cgcagcggct acgcctggtt catctactgg ggccaggca ccctggtgac cgtgagcagc 360

30

<210> 172

<211> 120

<212> PRT

35

<213> Artificial Sequence

<220>

35

<223> Humanized sequence.

<400> 172

40

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Gly Thr Phe Thr Asp Gln
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
 50 55 60
 Lys Gly Arg Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Phe Cys
 85 90 95
 Ala Arg Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
 100 105 110
 Gly Thr Leu Val Thr Val Ser Ser
 115 120

50

55

<210> 173

<211> 642

<212> DNA

<213> Artificial Sequence

<220>

<223> Humanized sequence.

<400> 173

5

	gacatccaga	tgacccagag	cccaaggcagc	ctgagcgcca	gcgtggcgaa	ccgcgtgacc	60
	atcacctgca	aggccagccg	cgacgtggcc	atccgcgtgg	cctggtaacca	gcagaagcca	120
	ggcaagggtgc	caaagctgct	gatctactgg	gccagcaccc	gccacaccgg	cgtgccaagc	180
10	cgcttcagcg	gcagcggcag	cgcacccgac	ttcacccctga	ccatcagcag	cctgcagcca	240
	gaggacgtgg	ccgactactt	ctgccaccag	tacagcagct	accattcac	cttcggcagc	300
	ggcaccaagc	tggagatcaa	gcgtactgtg	gctgcaccat	ctgtcttcat	cttcccggca	360
	tctgtatgagc	agttgaaatc	tggaactgcc	tctgttgtgt	gcctgctgaa	taacttctat	420
	cccagagagg	ccaaagtaca	gtgaaagggtg	gataacgccc	tccaatcgaaa	taactcccaag	480
15	gagagtgtca	cagaggcagga	cagcaaggac	agcacctaca	gcctcagcag	caccctgacg	540
	ctgagcaaag	cagactacga	gaaacacaaa	gtctacgcct	gcgaagtcaac	ccatcagggc	600
	ctgagctcgc	ccgtcacaaa	gagctcaac	aggggagagt	gt		642

<210> 174

<211> 214

20

<212> PRT

<213> Artificial Sequence

<220>

<223> Humanized sequence.

25

<400> 174

	Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly						
	1	5	10	15			
30	Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Arg Asp Val Ala Ile Ala						
	20	25	30				
	Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Ile						
	35	40	45				
35	Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Ser Arg Phe Ser Gly						
	50	55	60				
	Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro						
	65	70	75	80			
	Glu Asp Val Ala Asp Tyr Phe Cys His Gln Tyr Ser Ser Tyr Pro Phe						
	85	90	95				
40	Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala						
	100	105	110				
	Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly						
	115	120	125				
	Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala						
	130	135	140				
45	Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln						
	145	150	155	160			
	Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser						
	165	170	175				
50	Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr						
	180	185	190				
	Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser						
	195	200	205				
	Phe Asn Arg Gly Glu Cys						
	210						

55

<210> 175

<211> 1347

<212> DNA

<213> Artificial Sequence

<220>

<223> Humanized sequence.

5

<400> 175

caggtgcagc tggcagag cggcgccgag gtgaagaagc caggcagcag cgtgaagg 60

10

agctgcaagg ccagcgcta cacccacc gaccagacca tccactggat ggcgcaggcc 120
 ccagggcagg gcctggagt gatcgctac atctacccac ggcacgacag cccaaagtac 180
 aacgagaact tcaaggccaa ggtcaccatc accggccaca agagcaccag caccgcctac 240
 atggagctga gcagcctgcg cagcgaggac accggccgtgt actactgcgc catcccagac 300
 cgcagcggtt acgcctgtt catctactgg ggccaggcga ccctggtgac cgtgagcagc 360
 gcctccacca agggcccatc ggttttcccc ctggcacccct cctccaaagag cacctctggg 420
 ggcacagcgg ccctgggctg cctggtcaag gactacttc cggaaaccgg gacgggtgtcg 480
 tggacttag cgcgcctgac cagcgccgtg cacacccctcc cggctgtctt acagtctca 540
 ggactctact ccctcagcag cgtggtgacc gtccctcca gcagcttggg caccaggacc 600
 tacatctgca acgtgaatca caagccccagg aacaccaagg tcgacaagag agttgagccc 660
 aaatcttggt gaaaaactca cacatccccca ccgtgccccag caccagaagc tgctggggga 720
 ccgtcagtct tccttcccc cccaaaaccc aaggacaccc tcatgatctc ccggacccct 780
 gaggtcacat gcgtcggtt ggacgtgago cacgaagacc ctgaggtcaa gttcaactgg 840
 tacgtggacg gcgtggaggt gcataatgcc aagacaaagg cgccgggagga gcagtacaac 900
 agcacgtacc gtgtggtcg cgttctcacc gtccctgcacc aggactggct gaatggcaag 960
 gagtacaagt gcaagggtctc caacaaaggcc ctcccaagccc ccattcgagaa aaccatctcc 1020
 aaagccaaag ggcagcccg agaaccacag gtgtacaccc tgccccccatc ccgggaggag 1080
 atgaccaaga accaggttcg cctgacctgc ctggtaaaag gcttctatcc cagcgacatc 1140
 gccgtggagt gggagagcaa tggcagccg gagaacaact acaagaccac gcctccctgt 1200
 ctggactccg acggctctt cttctctac agcaagctca ccgtggacaa gagcagggtgg 1260
 cagcagggga acgttcttc atgtccctgt atgcattgagg ctctgcacaa ccactacacg 1320
 cagaagagcc tctccctgtc tccgggt 1347

<210> 176

<211> 449

35

<212> PRT

<213> Artificial Sequence

<220>

<223> Humanized sequence.

40

<400> 176

45

50

55

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Gln
 20 25 30
 5 Thr Ile His Trp Met Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
 50 55 60
 10 Lys Gly Lys Val Thr Ile Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 15 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
 100 105 110
 Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
 115 120 125
 20 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
 130 135 140
 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
 145 150 155 160
 25 Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
 165 170 175
 Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
 180 185 190
 Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
 195 200 205
 Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
 210 215 220
 Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Ala Ala Gly Gly
 225 230 235 240
 30 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile

	245	250	255	
35	Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu			
	260	265	270	
	Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His			
	275	280	285	
	Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg			
	290	295	300	
40	Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys			
	305	310	315	320
	Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu			
	325	330	335	
	Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr			
	340	345	350	
45	Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu			
	355	360	365	
	Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp			
	370	375	380	
	Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val			
	385	390	395	400
50	Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp			
	405	410	415	
	Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His			
	420	425	430	
	Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro			
	435	440	445	
55	Gly			

<210> 177

<211> 1347

<212> DNA

<213> Artificial Sequence

<220>

5 <223> Humanized sequence.

<400> 177

10	cagggtgcagc	tggtgtcagag	cggcgcccag	gtgaagaagc	caggcagcag	cgtgaagggtg	60
	agctgcagg	ccagcggctt	cacccacc	gaccagacca	tccactgggt	gcgcaggcc	120
	ccagggcagg	gcctggagt	gatgggctac	atctaccac	gcgacgacag	cccaaagtac	180
	aacgagaact	tcaaggcua	ggtcacccctg	accgcggaca	agagcaccag	caccgcctac	240
	atggagctga	gcagcctgcg	cagcgaggac	accgcctgt	actactgcgc	catcccagac	300
15	cgcagcggct	acgcctgtt	catctactgg	ggccagggca	ccctgggtgac	cgtgagcagc	360
	gcctccacca	aggggccatc	ggtttcccc	ctggcaccc	cctccaagag	cacctctggg	420
	ggcacagcgg	ccctggctg	cctggtaaag	gactactcc	ccgaaccgt	gacggtgtcg	480
	tggactcag	gcgcctgtac	cagcggcgtg	cacaccc	cggtgtct	acagtctca	540
	ggactctact	ccctcagcag	cgtggtgacc	gtgccttcca	gcagcttggg	caccagacc	600
20	tacatctgca	acgtgaatca	caagcccagc	aacaccaagg	tgcacaagag	agttgagccc	660
	aaatcttgc	acaaaactca	cacatcccc	ccgtgcccag	caccagaagc	tgctggggga	720
	ccgtcagtct	tcctttccc	ccaaaaacc	aaggacaccc	tcatgatctc	ccggacccct	780
	gaggtcacat	gcgtcggtt	ggacgtgagc	cacgaagacc	ctgaggtcaa	gttcaactgg	840
25	tacgtggacg	gcgtggaggt	gcataatgcc	aagacaaaagc	cgcgggagga	gcagtacaac	900
	agcacgtacc	gtgtggtcag	cgtccctcacc	gtcctgcacc	aggactggct	gaatggcaag	960
	gagtacaagt	gcaagggtctc	caacaaagcc	ctcccaagccc	ccatcgagaa	aaccatctcc	1020
	aaagccaaag	ggcagcccc	agaaccacag	gtgtacaccc	tgccccccatc	ccgggaggag	1080
30	atgaccaaga	accaggctcag	cctgacactgc	ctggtaaaag	gtttctatcc	cagcgcacatc	1140
	gccgtggagt	gggagagcaa	tggcagccg	gagaacaact	acaagaccac	gcctccctg	1200
	ctggactccg	acggctcctt	cttcctctac	agcaagctca	ccgtggacaa	gagcagggtgg	1260
	cagcagggg	acgtttctc	atgcctcg	atgcatgagg	ctctgcacaa	ccactacacg	1320
35	cagaagagcc	tctccctgtc	tccgggt				1347

<210> 178

<211> 449

<212> PRT

35 <213> Artificial Sequence

<220>

<223> Humanized sequence.

40 <400> 178

45

50

55

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Thr Phe Thr Asp Gln
 20 25 30
 Thr Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Tyr Ile Tyr Pro Arg Asp Asp Ser Pro Lys Tyr Asn Glu Asn Phe
 50 55 60
 Lys Gly Lys Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ile Pro Asp Arg Ser Gly Tyr Ala Trp Phe Ile Tyr Trp Gly Gln
 100 105 110
 Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val
 115 120 125
 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala
 130 135 140
 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
 145 150 155 160
 Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
 165 170 175
 Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
 180 185 190
 Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys
 195 200 205
 Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp
 210 215 220
 Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Ala Ala Gly Gly
 225 230 235 240
 Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 245 250 255
 Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 260 265 270
 Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
 275 280 285
 Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
 290 295 300
 Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
 305 310 315 320
 Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
 325 330 335
 Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
 340 345 350
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu
 355 360 365
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
 370 375 380
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
 385 390 395 400
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
 405 410 415
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
 420 425 430
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro

55

435

Gly

440

445

<211> 642
 <212> DNA
 <213> Artificial Sequence

5 <220>
 <223> Humanized sequence.

<400> 179

10 gacatccaga tgacccagag cccaaggcgc ctgagcgcca gcgtggcga ccgcgtgacc 60
 atcacctgca aggccagccg cgacgtggcc atcgccgtgg cctggtaacca gcagaagcca 120
 ggcaagggtgc caaagctgct gctgttctgg gccagcaccc gccacacccgg cgtgccagac 180
 cgcttcagcg gcagcggcag cggcaccgac ttcacccctga ccatcagcag cctgcagcca 240
 gaggacctgg ccgactacta ctgcccaccag tacagcagct acccattcac cttccggccag 300
 ggcaccaagc tggagatcaa gcgtactgtg gctgcaccat ctgtcttcat cttccggcca 360
 tctgatgagc agttgaaatc tggaaactgcc tctgttgtgt gcctgctgaa taacttcttat 420
 cccagagagg ccaaagtaca gtggaaagggtg gataacgccc tccaatcggg taactcccgag 480
 gagagtgtca cagagcagga cagcaaggac agdactatac gcttcagcag caccctgacg 540
 ctgagcaagc cagactacga gaaacacaaa gtctacgcct gctgaagtac ccatcaggc 600
 ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gt 642

20
 <210> 180
 <211> 214
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Humanized sequence.

30 <400> 180

	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
	1				5					10					15	
35	Asp	Arg	Val	Thr	Ile	Thr	Cys	Lys	Ala	Ser	Arg	Asp	Val	Ala	Ile	Ala
					20				25					30		
	Val	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Val	Pro	Lys	Leu	Leu	Leu
					35				40				45			
40	Phe	Trp	Ala	Ser	Thr	Arg	His	Thr	Gly	Val	Pro	Asp	Arg	Phe	Ser	Gly
					50			55			60					
	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
					65			70			75				80	
45	Glu	Asp	Leu	Ala	Asp	Tyr	Tyr	Cys	His	Gln	Tyr	Ser	Ser	Tyr	Pro	Phe
					85				90					95		
	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala
					100				105				110			
50	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly
					115				120				125			
	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala
					130			135			140					
55	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln
					145			150			155				160	
	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser
					165				170				175			
	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr
					180				185				190			
	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser
					195			200					205			
	Phe	Asn	Arg	Gly	Glu	Cys										
					210											

<210> 181
 <211> 189
 <212> PRT
 <213> Homo sapiens

5

<400> 181

	Met	Leu	Gly	Ser	Arg	Ala	Val	Met	Leu	Leu	Leu	Leu	Leu	Pro	Trp	Thr
10	1					5			10						15	
	Ala	Gln	Gly	Arg	Ala	Val	Pro	Gly	Gly	Ser	Ser	Pro	Ala	Trp	Thr	Gln
						20			25						30	
	Cys	Gln	Gln	Leu	Ser	Gln	Lys	Leu	Cys	Thr	Leu	Ala	Trp	Ser	Ala	His
						35			40					45		
15	Pro	Leu	Val	Gly	His	Met	Asp	Leu	Arg	Glu	Glu	Gly	Asp	Glu	Glu	Thr
						50			55					60		
	Thr	Asn	Asp	Val	Pro	His	Ile	Gln	Cys	Gly	Asp	Gly	Cys	Asp	Pro	Gln
						65			70				75		80	
	Gly	Leu	Arg	Asp	Asn	Ser	Gln	Phe	Cys	Leu	Gln	Arg	Ile	His	Gln	Gly
						85			90				95			
20	Leu	Ile	Phe	Tyr	Glu	Lys	Leu	Leu	Gly	Ser	Asp	Ile	Phe	Thr	Gly	Glu
						100			105					110		
	Pro	Ser	Leu	Leu	Pro	Asp	Ser	Pro	Val	Gly	Gln	Leu	His	Ala	Ser	Leu
						115			120				125			
25	Leu	Gly	Leu	Ser	Gln	Leu	Leu	Gln	Pro	Glu	Gly	His	His	Trp	Glu	Thr
						130			135				140			
	Gln	Gln	Ile	Pro	Ser	Leu	Ser	Pro	Ser	Gln	Pro	Trp	Gln	Arg	Leu	Leu
						145			150				155		160	
	Leu	Arg	Phe	Lys	Ile	Leu	Arg	Ser	Leu	Gln	Ala	Phe	Val	Ala	Val	Ala
						165			170				175			
30	Ala	Arg	Val	Phe	Ala	His	Gly	Ala	Ala	Thr	Leu	Ser	Pro			
						180			185							

Claims

1. A humanized anti-IL-23p19 antibody or antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof comprises:
 - a) a light chain variable region comprising the amino acid sequence of SEQ ID NO:19 (*CDR1-L*); the amino acid sequence of SEQ ID NO:20 (*CDR2-L*); and the amino acid sequence of SEQ ID NO:21 (*CDR3-L*); and
 - b) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 66 or 67 (*CDR1-H*); the amino acid sequence of SEQ ID NO:64 (*CDR2-H*); and the amino acid sequence of SEQ ID NO:65 (*CDR3-H*).
2. The humanized anti-IL-23p19 antibody or antigen-binding fragment thereof according to claim 1, wherein the antibody or antigen-binding fragment thereof comprises:
 - a) a light chain variable region comprising the amino acid sequence of SEQ ID NO:19 (*CDR1-L*); the amino acid sequence of SEQ ID NO:20 (*CDR2-L*); and the amino acid sequence of SEQ ID NO:21 (*CDR3-L*); and
 - b) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:66 (*CDR1-H*); the amino acid sequence of SEQ ID NO:64 (*CDR2-H*); and the amino acid sequence of SEQ ID NO:65 (*CDR3-H*).
3. The humanized anti-IL-23p19 antibody or antigen-binding fragment thereof according to claim 1, wherein the antibody or antigen-binding fragment thereof comprises:
 - a) a light chain variable region comprising the amino acid sequence of SEQ ID NO:19 (*CDR1-L*); the amino acid sequence of SEQ ID NO:20 (*CDR2-L*); and the amino acid sequence of SEQ ID NO:21 (*CDR3-L*); and
 - b) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:67 (*CDR1-H*); the amino acid sequence of SEQ ID NO:64 (*CDR2-H*); and the amino acid sequence of SEQ ID NO:65 (*CDR3-H*).
4. The humanized anti-IL-23p19 antibody or antigen-binding fragment thereof according to claim 1, wherein the antibody

or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of any one of SEQ ID NO:158, 160, 162 or 164; and a heavy chain variable region comprising the amino acid sequence of any one of SEQ ID NO:166 or 168.

- 5 5. The humanized anti-IL-23p19 antibody or antigen-binding fragment thereof according to claim 4, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:160 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:166.
- 10 6. The humanized anti-IL-23p19 antibody or antigen-binding fragment thereof, according to claim 4, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:160 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:168.
- 15 7. The humanized anti-IL-23p19 antibody or antigen-binding fragment thereof according to claim 4, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO:158 and a heavy chain variable region comprising the amino acid sequence SEQ ID NO:166.
- 20 8. The humanized anti-IL-23p19 antibody or antigen-binding fragment thereof according to claim 4, wherein the antibody or antigen-binding fragment thereof comprises the amino acid sequence of SEQ ID NO:166 or 168 linked to a human IgG1 heavy chain constant region; and the amino acid sequence of SEQ ID NO:158 or 160 linked to a human kappa light chain constant region.
- 25 9. The humanized anti-IL-23p19 antibody or antigen-binding fragment thereof according to claim 1, wherein the antibody or antigen-binding fragment thereof comprises:
 - a) a humanized light chain variable domain comprising the CDRs of SEQ ID NO:158 or 160 and framework regions having an amino acid sequence at least 90% identical to the amino acid sequence of the framework regions of the variable domain light chain amino acid sequence of SEQ ID NO:158 or 160; and
 - b) a humanized heavy chain variable domain comprising the CDRs of SEQ ID NO:166 or 168 and framework regions having an amino acid sequence at least 90% identical to the amino acid sequence of the framework regions of the variable domain heavy chain amino acid sequence of SEQ ID NO:166 or 168.
- 30 10. The humanized anti-IL-23p19 antibody according to any one of claims 1 to 9, wherein said antibody is a monoclonal antibody.
- 35 11. The humanized anti-IL-23p19 antibody according to claim 1, wherein the antibody is a humanized monoclonal anti-IL-23p19 antibody comprising a light chain comprising the amino acid sequence of SEQ ID NO:174 or 180 and a heavy chain comprising the amino acid sequence of SEQ ID NO:176 or 178.
- 40 12. The humanized monoclonal anti-IL-23p19 antibody according to claim 11, wherein said antibody comprises a light chain comprising the amino acid sequence of SEQ ID NO:174 and a heavy chain comprising the amino acid sequence of SEQ ID NO:176.
- 45 13. The humanized monoclonal anti-IL-23p19 antibody according to claim 11, wherein said antibody comprises a light chain comprising the amino acid sequence of SEQ ID NO:174 and a heavy chain comprising the amino acid sequence of SEQ ID NO:178.
- 50 14. The humanized monoclonal anti-IL-23p19 antibody according to claim 11, wherein said antibody comprises a light chain comprising the amino acid sequence of SEQ ID NO:180 and a heavy chain comprising the amino acid sequence of SEQ ID NO:176.
- 55 15. An antibody or antigen-binding fragment according to any one of claims 1 to 14 for use in medicine.
16. The antibody or antigen-binding fragment according to claim 15, wherein the use is the treatment of an inflammatory disease, of an autoimmune disease, of a respiratory disease, of a metabolic disorder or of cancer.
17. The antibody or antigen-binding fragment according to claim 16, wherein the use is for the treatment of psoriasis, inflammatory bowel disease, psoriatic arthritis, multiple sclerosis, rheumatoid arthritis, Crohn's disease, ulcerative colitis or ankylosing spondylitis.

18. The antibody or antigen-binding fragment according to claim 16, wherein the use is for the treatment of asthma or COPD (chronic obstructive pulmonary disease).
- 5 19. A pharmaceutical composition comprising an antibody or antigen-binding fragment according to any one of claims 1 to 14 and a pharmaceutically acceptable carrier.
- 10 20. An isolated polynucleotide comprising a sequence encoding a light chain variable region of an antibody or antibody fragment having the amino acid sequence of SEQ ID NO:158, 160, 162 or 164, and encoding a heavy chain variable region of an antibody or antibody fragment having the amino acid sequence of SEQ ID NO: 166 or 168.
- 15 21. The isolated polynucleotide according to claim 20, wherein said polynucleotide comprises a sequence encoding a light chain of an antibody having the amino acid sequence of SEQ ID NO:174 or 180, and encoding a heavy chain of an antibody having the amino acid sequence of SEQ ID NO:176 or 178.
22. An expression vector comprising the polynucleotide of claim 20 or 21.
23. A host cell comprising:
- 20 a) an isolated polynucleotide comprising a sequence encoding a light chain variable region of an antibody or antibody fragment having the amino acid sequence of SEQ ID NO:158, 160, 162 or 164; and
b) an isolated polynucleotide comprising a sequence encoding a heavy chain variable region of an antibody or antibody fragment having the amino acid sequence of SEQ ID NO: 166 or 168.
24. A host cell comprising:
- 25 a) an isolated polynucleotide comprising a sequence encoding a light chain of an antibody having the amino acid sequence of SEQ ID NO:174 or 180; and
b) an isolated polynucleotide comprising a sequence encoding a heavy chain of an antibody having the amino acid sequence of SEQ ID NO:176 or 178.
- 30 25. A method for producing an antibody or antigen-binding fragment comprising:
- 35 a) obtaining a host cell according to claim 23 or 24; and
b) cultivating the host cell.
26. The method according to claim 25, further comprising recovering and purifying the antibody or antigen-binding fragment.

40 **Patentansprüche**

1. Humanisierter anti-IL-23p19-Antikörper oder Antigen-bindendes Fragment davon, wobei der Antikörper oder das Antigen-bindende Fragment davon:
- 45 a) eine variable Region der leichten Kette umfassend die Aminosäuresequenz von SEQ ID NO:19 (*CDR1-L*); die Aminosäuresequenz von SEQ ID NO:20 (*CDR2-L*); und die Aminosäuresequenz von SEQ ID NO:21 (*CDR3-L*); und
b) eine variable Region der schweren Kette umfassend die Aminosäuresequenz von SEQ ID NO: 66 oder 67 (*CDR1-H*); die Aminosäuresequenz of SEQ ID NO:64 (*CDR2-H*); und die Aminosäuresequenz von SEQ ID NO:65 (*CDR3-H*)
- 50 umfasst.
2. Humanisierter anti-IL-23p19-Antikörper oder Antigen-bindendes Fragment davon nach Anspruch 1, wobei der Antikörper oder das Antigen-bindende Fragment davon:
- 55 a) eine variable Region der leichten Kette umfassend die Aminosäuresequenz von SEQ ID NO:19 (*CDR1-L*); die Aminosäuresequenz von SEQ ID NO:20 (*CDR2-L*); und die Aminosäuresequenz von SEQ ID NO:21 (*CDR3-*

L); und

b) eine variable Region der schweren Kette umfassend die Aminosäuresequenz von SEQ ID NO:66 (*CDR1-H*); die Aminosäuresequenz von SEQ ID NO:64 (*CDR2-H*); und die Aminosäuresequenz von SEQ ID NO:65 (*CDR3-H*)

5

umfasst.

3. Humanisierter anti-IL-23p19-Antikörper oder Antigen-bindendes Fragment davon nach Anspruch 1, wobei der Antikörper oder das Antigen-bindende Fragment davon:

10

a) eine variable Region der leichten Kette umfassend die Aminosäuresequenz von SEQ ID NO:19 (*CDR1-L*); die Aminosäuresequenz von SEQ ID NO:20 (*CDR2-L*); und die Aminosäuresequenz von SEQ ID NO:21 (*CDR3-L*); und

15

b) eine variable Region der schweren Kette umfassend die Aminosäuresequenz von SEQ ID NO:67 (*CDR1-H*); die Aminosäuresequenz von SEQ ID NO:64 (*CDR2-H*); und die Aminosäuresequenz von SEQ ID NO:65 (*CDR3-H*) umfasst.

20

4. Humanisierter anti-IL-23p19-Antikörper oder Antigen-bindendes Fragment davon nach Anspruch 1, wobei der Antikörper oder das Antigen-bindende Fragment davon eine variable Region der leichten Kette umfassend die Aminosäuresequenz von irgendeinem von SEQ ID NO:158, 160, 162 oder 164; und eine variable Region der schweren Kette umfassend die Aminosäuresequenz von irgendeinem von SEQ ID NO:166 oder 168 umfasst.

25

5. Humanisierter anti-IL-23p19-Antikörper oder Antigen-bindendes Fragment davon nach Anspruch 4, wobei der Antikörper oder das Antigen-bindende Fragment davon eine variable Region der leichten Kette umfassend die Aminosäuresequenz von SEQ ID NO:160 und eine variable Region der schweren Kette umfassend die Aminosäuresequenz SEQ ID NO:166 umfasst.

30

6. Humanisierter anti-IL-23p19-Antikörper oder Antigen-bindendes Fragment davon nach Anspruch 4, wobei der Antikörper oder das Antigen-bindende Fragment davon eine variable Region der leichten Kette umfassend die Aminosäuresequenz von SEQ ID NO:160 und eine variable Region der schweren Kette umfassend die Aminosäuresequenz SEQ ID NO:168 umfasst.

35

7. Humanisierter anti-IL-23p19-Antikörper oder Antigen-bindendes Fragment davon nach Anspruch 4, wobei der Antikörper oder das Antigen-bindende Fragment davon eine variable Region der leichten Kette umfassend die Aminosäuresequenz von SEQ ID NO:158 und eine variable Region der schweren Kette umfassend die Aminosäuresequenz SEQ ID NO:166 umfasst.

40

8. Humanisierter anti-IL-23p19-Antikörper oder Antigen-bindendes Fragment davon nach Anspruch 4, wobei der Antikörper oder das Antigen-bindende Fragment davon die Aminosäuresequenz von SEQ ID NO:166 oder 168 verknüpft an eine konstante Region der schweren Kette von einem humanen IgG1; und die Aminosäuresequenz von SEQ ID NO:158 oder 160 verknüpft an eine konstante Region der humanen Kappa-Leichtkette umfasst.

45

9. Humanisierter anti-IL-23p19-Antikörper oder Antigen-bindendes Fragment davon nach Anspruch 1, wobei der Antikörper oder das Antigen-bindende Fragment davon:

50

a) eine humanisierte variable Leichtketten-Domäne umfassend die CDRs von SEQ ID NO:158 oder 160 und Gerüstregionen mit einer Aminosäuresequenz, die mindestens zu 90% identisch ist mit der Aminosäuresequenz der Gerüstregionen der Aminosäuresequenz der variablen Domäne der leichten Kette von SEQ ID NO:158 oder 160; und

55

b) eine humanisierte variable Schwerketten-Domäne umfassend die CDRs von SEQ ID NO:166 oder 168 und Gerüstregionen mit einer Aminosäuresequenz, die mindestens zu 90% identisch ist mit der Aminosäuresequenz der Gerüstregionen der Aminosäuresequenz der variablen Domäne der schweren Kette von SEQ ID NO:166 oder 168.

55

umfasst.

10. Humanisierter anti-IL-23p19-Antikörper nach irgendeinem der Ansprüche 1 bis 9, wobei der Antikörper ein monoklonaler Antikörper ist.

- 11.** Humanisierter anti-IL-23p19-Antikörper nach Anspruch 1, wobei der Antikörper ein humanisierter monoklonaler anti-IL-23p19-Antikörper umfassend eine leichte Kette umfassend die Aminosäuresequenz von SEQ ID NO:174 oder 180 und eine schwere Kette umfassend die Aminosäuresequenz von SEQ ID NO:176 oder 178 ist.
- 12.** Humanisierter monoklonaler anti-IL-23p19-Antikörper nach Anspruch 11, wobei der Antikörper eine leichte Kette umfassend die Aminosäuresequenz von SEQ ID NO:174 und eine schwere Kette umfassend die Aminosäuresequenz von SEQ ID NO:176 umfasst.
- 13.** Humanisierter monoklonaler anti-IL-23p19-Antikörper nach Anspruch 11, wobei der Antikörper eine leichte Kette umfassend die Aminosäuresequenz von SEQ ID NO:174 und eine schwere Kette umfassend die Aminosäuresequenz von SEQ ID NO:178 umfasst.
- 14.** Humanisierter monoklonaler anti-IL-23p19-Antikörper nach Anspruch 11, wobei der Antikörper eine leichte Kette umfassend die Aminosäuresequenz von SEQ ID NO:180 und eine schwere Kette umfassend die Aminosäuresequenz von SEQ ID NO:176 umfasst.
- 15.** Antikörper oder Antigen-bindendes Fragment nach irgendeinem der Ansprüche 1 bis 14 zur Verwendung in Medizin.
- 16.** Antikörper oder Antigen-bindendes Fragment nach Anspruch 15, wobei die Verwendung die Behandlung einer entzündlichen Erkrankung, einer Autoimmunerkrankung, einer Atemwegserkrankung, einer metabolischen Störung oder von Krebs ist.
- 17.** Antikörper oder Antigen-bindendes Fragment nach Anspruch 16, wobei die Verwendung für die Behandlung von Psoriasis, entzündlicher Darmerkrankung, psoriatischer Arthritis, multipler Sklerose, rheumatoider Arthritis, Morbus Crohn, Colitis ulcerosa oder Spondylitis ankylosans ist.
- 18.** Antikörper oder Antigen-bindendes Fragment nach Anspruch 16, wobei die Verwendung für die Behandlung von Asthma oder COPD (chronisch obstruktiver Lungenerkrankung) ist.
- 19.** Pharmazeutische Zusammensetzung umfassend einen Antikörper oder ein Antigen-bindendes Fragment nach irgendeinem der Ansprüche 1 bis 14 und einen pharmazeutisch akzeptablen Träger.
- 20.** Isoliertes Polynukleotid umfassend eine Sequenz kodierend eine variable Region einer leichten Kette von einem Antikörper oder Antikörperfragment mit der Aminosäuresequenz von SEQ ID NO:158, 160, 162 oder 164 und kodierend eine variable Region einer schweren Kette von einem Antikörper oder Antikörperfragment mit der Aminosäuresequenz von SEQ ID NO: 166 oder 168.
- 21.** Isoliertes Polynukleotid nach Anspruch 20, wobei das Polynukleotid eine Sequenz kodierend eine leichte Kette von einem Antikörper mit der Aminosäuresequenz von SEQ ID NO:174 oder 180 und kodierend eine schwere Kette von einem Antikörper mit der Aminosäuresequenz von SEQ ID NO:176 oder 178 umfasst.
- 22.** Expressionsvektor umfassend das Polynukleotid nach Anspruch 20 oder 21.
- 23.** Wirtszelle umfassend:
- a) ein isoliertes Polynukleotid umfassend eine Sequenz kodierend eine variable Region einer leichten Kette von einem Antikörper oder Antikörperfragment mit der Aminosäuresequenz von SEQ ID NO:158, 160, 162 oder 164; und
- b) ein isoliertes Polynukleotid umfassend eine Sequenz kodierend eine variable Region einer schweren Kette von einem Antikörper oder Antikörperfragment mit der Aminosäuresequenz von SEQ ID NO: 166 oder 168.
- 24.** Wirtszelle umfassend:
- a) ein isoliertes Polynukleotid umfassend eine Sequenz kodierend eine leichte Kette von einem Antikörper mit der Aminosäuresequenz von SEQ ID NO:174 oder 180; und
- b) ein isoliertes Polynukleotid umfassend eine Sequenz kodierend eine schwere Kette von einem Antikörper mit der Aminosäuresequenz von SEQ ID NO: 176 oder 178.

25. Verfahren zur Herstellung eines Antikörpers oder Antigen-bindenden Fragments umfassend:

- a) Erhalten einer Wirtszelle nach Anspruch 23 oder 24; und
- b) Kultivieren der Wirtszelle.

5

26. Verfahren nach Anspruch 25, ferner umfassend das Gewinnen und Reinigen des Antikörpers oder Antigen-bindenden Fragments.

10 **Revendications**

1. Anticorps humanisé anti-IL-23p19 ou fragment de liaison à l'antigène de cet anticorps, dans lequel l'anticorps ou le fragment de liaison à l'antigène de cet anticorps comprend :

- 15 a) une région variable de chaîne légère comprenant la séquence d'acides aminés de SEQ ID NO: 19 (*CDR1-L*) ; la séquence d'acides aminés de SEQ ID NO: 20 (*CDR2-L*) ; et la séquence d'acides aminés de SEQ ID NO: 21 (*CDR3-L*) ; et
- b) une région variable de chaîne lourde comprenant la séquence d'acides aminés de SEQ ID NO: 66 ou 67 (*CDR1-H*) ; la séquence d'acides aminés de SEQ ID NO: 64 (*CDR2-H*) ; et la séquence d'acides aminés de SEQ ID NO: 65 (*CDR3-H*).

20

2. Anticorps humanisé anti-IL-23p19 ou fragment de liaison à l'antigène de cet anticorps selon la revendication 1, dans lequel l'anticorps ou le fragment de liaison à l'antigène de cet anticorps comprend :

25

- a) une région variable de chaîne légère comprenant la séquence d'acides aminés de SEQ ID NO: 19 (*CDR1-L*) ; la séquence d'acides aminés de SEQ ID NO: 20 (*CDR2-L*) ; et la séquence d'acides aminés de SEQ ID NO: 21 (*CDR3-L*) ; et
- b) une région variable de chaîne lourde comprenant la séquence d'acides aminés de SEQ ID NO: 66 (*CDR1-H*) ; la séquence d'acides aminés de SEQ ID NO: 64 (*CDR2-H*) ; et la séquence d'acides aminés de SEQ ID NO: 65 (*CDR3-H*).

30

3. Anticorps humanisé anti-IL-23p19 ou fragment de liaison à l'antigène de cet anticorps selon la revendication 1, dans lequel l'anticorps ou le fragment de liaison à l'antigène de cet anticorps comprend :

35

- a) une région variable de chaîne légère comprenant la séquence d'acides aminés de SEQ ID NO: 19 (*CDR1-L*) ; la séquence d'acides aminés de SEQ ID NO: 20 (*CDR2-L*) ; et la séquence d'acides aminés de SEQ ID NO: 21 (*CDR3-L*) ; et
- b) une région variable de chaîne lourde comprenant la séquence d'acides aminés de SEQ ID NO: 67 (*CDR1-H*) ; la séquence d'acides aminés de SEQ ID NO: 64 (*CDR2-H*) ; et la séquence d'acides aminés de SEQ ID NO: 65 (*CDR3-H*).

40

4. Anticorps humanisé anti-IL-23p19 ou fragment de liaison à l'antigène de cet anticorps selon la revendication 1, dans lequel l'anticorps ou le fragment de liaison à l'antigène de cet anticorps comprend une région variable de chaîne légère comprenant la séquence d'acides aminés de l'une quelconque des séquences SEQ ID NO: 158, 160, 162 ou 164 ; et une région variable de chaîne lourde comprenant la séquence d'acides aminés de l'une quelconque des séquences SEQ ID NO: 166 ou 168.

45

5. Anticorps humanisé anti-IL-23p19 ou fragment de liaison à l'antigène de cet anticorps selon la revendication 4, dans lequel l'anticorps ou le fragment de liaison à l'antigène de cet anticorps comprend une région variable de chaîne légère comprenant la séquence d'acides aminés de SEQ ID NO: 160 et une région variable de chaîne lourde comprenant la séquence d'acides aminés SEQ ID NO: 166.

50

6. Anticorps humanisé anti-IL-23p19 ou fragment de liaison à l'antigène de cet anticorps selon la revendication 4, dans lequel l'anticorps ou le fragment de liaison à l'antigène de cet anticorps comprend une région variable de chaîne légère comprenant la séquence d'acides aminés de SEQ ID NO: 160 et une région variable de chaîne lourde comprenant la séquence d'acides aminés SEQ ID NO: 168.

55

7. Anticorps humanisé anti-IL-23p19 ou fragment de liaison à l'antigène de cet anticorps selon la revendication 4, dans

lequel l'anticorps ou le fragment de liaison à l'antigène de cet anticorps comprend une région variable de chaîne légère comprenant la séquence d'acides aminés de SEQ ID NO: 158 et une région variable de chaîne lourde comprenant la séquence d'acides aminés SEQ ID NO: 166.

- 5 **8.** Anticorps humanisé anti-IL-23p19 ou fragment de liaison à l'antigène de cet anticorps selon la revendication 4, dans lequel l'anticorps ou le fragment de liaison à l'antigène de cet anticorps comprend la séquence d'acides aminés de SEQ ID NO: 166 ou 168 reliée à une région constante de chaîne lourde d'une IgG1 humaine ; et la séquence d'acides aminés de SEQ ID NO: 158 ou 160 reliée à une région constante de chaîne légère kappa humaine.
- 10 **9.** Anticorps humanisé anti-IL-23p19 ou fragment de liaison à l'antigène de cet anticorps selon la revendication 1, dans lequel l'anticorps ou le fragment de liaison à l'antigène de cet anticorps comprend :
- 15 a) un domaine variable de chaîne légère humanisé comprenant les CDR de SEQ ID NO: 158 ou 160 et des régions charpentes ayant une séquence d'acides aminés présentant une identité d'au moins 90% avec la séquence d'acides aminés des régions charpentes de la séquence d'acides aminés de la chaîne légère du domaine variable de SEQ ID NO: 158 ou 160; et
- 20 b) un domaine variable de chaîne lourde humanisé comprenant les CDR de SEQ ID NO: 166 ou 168 et des régions charpentes ayant une séquence d'acides aminés présentant une identité d'au moins 90% avec la séquence d'acides aminés des régions charpentes de la séquence d'acides aminés de la chaîne lourde du domaine variable de SEQ ID NO: 166 ou 168.
- 25 **10.** Anticorps humanisé anti-IL-23p19 selon l'une quelconque des revendications 1 à 9, dans lequel ledit anticorps est un anticorps monoclonal.
- 30 **11.** Anticorps humanisé anti-IL-23p19 selon la revendication 1, dans lequel l'anticorps est un anticorps anti-IL-23p19 monoclonal humanisé comprenant une chaîne légère comprenant la séquence d'acides aminés de SEQ ID NO: 174 ou 180 et une chaîne lourde comprenant la séquence d'acides aminés de SEQ ID NO: 176 ou 178.
- 35 **12.** Anticorps anti-IL-23p19 monoclonal humanisé selon la revendication 11, dans lequel ledit anticorps comprend une chaîne légère comprenant la séquence d'acides aminés de SEQ ID NO: 174 et une chaîne lourde comprenant la séquence d'acides aminés de SEQ ID NO: 176.
- 40 **13.** Anticorps anti-IL-23p19 monoclonal humanisé selon la revendication 11, dans lequel ledit anticorps comprend une chaîne légère comprenant la séquence d'acides aminés de SEQ ID NO: 174 et une chaîne lourde comprenant la séquence d'acides aminés de SEQ ID NO: 178.
- 45 **14.** Anticorps anti-IL-23p19 monoclonal humanisé selon la revendication 11, dans lequel ledit anticorps comprend une chaîne légère comprenant la séquence d'acides aminés de SEQ ID NO: 180 et une chaîne lourde comprenant la séquence d'acides aminés de SEQ ID NO: 176.
- 50 **15.** Anticorps ou fragment de liaison à l'antigène selon l'une quelconque des revendications 1 à 14, pour une utilisation en médecine.
- 55 **16.** Anticorps ou fragment de liaison à l'antigène selon la revendication 15, dans lequel l'utilisation est le traitement d'une maladie inflammatoire, d'une maladie autoimmune, d'une maladie respiratoire, d'un trouble métabolique ou d'un cancer.
- 60 **17.** Anticorps ou fragment de liaison à l'antigène selon la revendication 16, dans lequel l'utilisation est destinée au traitement du psoriasis, d'une maladie inflammatoire des intestins, de l'arthrite psoriasique, de la sclérose en plaques, de la polyarthrite rhumatoïde, de la maladie de Crohn, de la rectocolite hémorragique ou de la spondylite ankylosante.
- 65 **18.** Anticorps ou fragment de liaison à l'antigène selon la revendication 16, dans lequel l'utilisation est destinée au traitement de l'asthme ou de la COPD (pneumopathologie obstructive chronique).
- 70 **19.** Composition pharmaceutique comprenant un anticorps ou fragment de liaison à l'antigène selon l'une quelconque des revendications 1 à 14 et un véhicule pharmaceutiquement acceptable.

20. Polynucléotide isolé comprenant une séquence codant pour une région variable de chaîne légère d'un anticorps ou fragment d'anticorps ayant la séquence d'acides aminés de SEQ ID NO: 158, 160, 162 ou 164, ou codant pour une région variable de chaîne lourde d'un anticorps ou fragment d'anticorps ayant la séquence d'acides aminés de SEQ ID NO: 166 ou 168.

5

21. Polynucléotide isolé selon la revendication 20, dans lequel ledit polynucléotide comprend une séquence codant pour une chaîne légère d'un anticorps ayant la séquence d'acides aminés de SEQ ID NO: 174 ou 180, ou codant pour une chaîne lourde d'un anticorps ayant la séquence d'acides aminés de SEQ ID NO: 176 ou 178.

10

22. Vecteur d'expression comprenant le polynucléotide selon la revendication 20 ou 21.

23. Cellule hôte comprenant :

15

a) un polynucléotide isolé comprenant une séquence codant pour une région variable de chaîne légère d'un anticorps ou d'un fragment d'anticorps ayant la séquence d'acides aminés de SEQ. ID N°158, 160, 162 ou 164 ; et
b) un polynucléotide isolé comprenant une séquence codant pour une région variable de chaîne lourde d'un anticorps ou d'un fragment d'anticorps ayant la séquence d'acides aminés de SEQ. ID N°166 ou 168.

24. Cellule hôte comprenant :

20

a) un polynucléotide isolé comprenant une séquence codant pour une région variable de chaîne légère d'un anticorps ou d'un fragment d'anticorps ayant la séquence d'acides aminés de SEQ. ID N°174 ou 180 ; et
b) un polynucléotide isolé comprenant une séquence codant pour une région variable de chaîne lourde d'un anticorps ou d'un fragment d'anticorps ayant la séquence d'acides aminés de SEQ. ID N°176 ou 178.

25

25. Méthode de production d'un anticorps ou d'un fragment de liaison à l'antigène comprenant :

30

a) l'obtention d'une cellule hôte selon la revendication 23 ou 24 ; et
b) la culture de la cellule hôte.

35

26. Méthode selon la revendication 25, comprenant en outre la récupération et la purification de l'anticorps ou du fragment de liaison à l'antigène.

40

45

50

55

Anti-IL-23p19 6B8 Engineered V_k regions

<pre>>6B8-VK >Hum VK#65 >Hum VK#66 >Hum VK#67 >Hum VK#78</pre>	Chothia --- Kabat --- IMGT --- DIVMTQSHKFKLSTSVGDRVITITCKASRDVAIAVAVYQQQKPGOSPKLLIFWASTRET DIQMTOQPSSLSASVGDRVITITCKASRDVAIAVAVYQQQKPGKVPKLLIFWASTRET DIQMTOQPSSLSASVGDRVITITCKASRDVAIAVAVYQQQKPGKVPKLLIFWASTRET DIQMTOQPSSLSASVGDRVITITCKASRDVAIAVAVYQQQKPGKVPKLLIFWASTRET DIQMTOQPSSLSASVGDRVITITCKASRDVAIAVAVYQQQKPGKVPKLLIFWASTRET	<pre>14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56</pre>
<pre>>6B8-VK >Hum VK#65 >Hum VK#66 >Hum VK#67 >Hum VK#78</pre>	Chothia --- Kabat --- IMGT --- GVPDRFTGSGSRTDFTLTISVQSEDLADYFECHQSSYPETFGSGTKLEIK GVPDRFTGSGSQTDFLFTLTISQPEDLADYFYCHQSSYSYTFGSGTKLEIK GVPDRFTGSGSRTDFTLTISQPEDVADYFCHQSSYSYTEGSGTKLEIK GVPDRFTGSGSRTDFTLTISQPEDVATYCHQSSYSYTEGSGTKLEIK GVPDRFTGSGSRTDFTLTISQPEDLADYFYCHQSSYSYTFGSGTKLEIK	<pre>57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107</pre>

Fig. 1a

Anti-IL-23p19 6B8 Engineered VH regions

Fig.




Fig. 2

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2007024846 A [0011] [0295] [0298]
- US 4816567 A [0093] [0095]
- US 6407213 B [0107]
- US 6037454 A [0108]
- US 6054297 A [0108]
- US 4485045 A [0186]
- US 4544545 A [0186]
- US 5013556 A [0186]
- WO 8101145 A [0188]
- WO 8807378 A [0188]
- US 4975278 A [0188]
- WO 9632478 A [0189]
- US 4640835 A [0192]
- US 4496689 A [0192]
- US 4301144 A [0192]
- US 4670417 A [0192]
- US 4791192 A [0192]
- US 4179337 A [0192]
- WO 9013646 A [0211]
- US 4965199 A [0215]
- EP 73657 A [0221]
- US 4419446 A [0223]
- US 4601978 A [0223]
- WO 9411026 A [0225]
- US 5888809 A [0225]
- DD 266710 [0226]
- EP 402226 A [0227]
- EP 183070 A [0227]
- EP 244234 A [0227]
- US 4767704 A [0232]
- US 4657866 A [0232]
- US 4927762 A [0232]
- US 4560655 A [0232]
- US 5122469 A [0232]
- WO 90103430 A [0232]
- WO 8700195 A [0232]
- US 4741900 A [0244]
- US 4275149 A [0251] [0254]
- US 4737456 A [0252]
- US 4318980 A [0254]
- WO 9404188 A [0262]
- WO 2007027714 A [0295] [0298]
- WO 2007005955 A [0295] [0298]
- WO 61410158 A [0313]
- WO 61411953 A [0313]
- WO 61412594 A [0313]
- WO 61448785 A [0313]

Non-patent literature cited in the description

- KIKLY et al. *Curr. Opin. Immunol.*, 2006, vol. 18, 670 [0075]
- CHOTHIA et al. *J. Mol. Biol.*, 1985, vol. 186, 651-663 [0088]
- KABAT et al. Sequences of Proteins of immunological Interest. Public Health Service, National Institutes of Health, 1991 [0089]
- CHOTHIA ; LESK. *J. Mol. Biol.*, 1987, vol. 196, 901-917 [0089]
- KABAT et al. *NIH*, 1991, vol. I, 647-669 [0090]
- KOHLER et al. *Nature*, 1975, vol. 256, 495 [0093]
- CLACKSON et al. *Nature*, 1991, vol. 352, 624-628 [0093]
- MARKS et al. *J. Mol. Biol.*, 1991, vol. 222, 581-597 [0093]
- MORRISON et al. *Proc. Natl. Acad. Sci. USA*, 1984, vol. 81, 6851-6855 [0095]
- PLUCKTHUN. The Pharmacology of monoclonal Antibodies. Springer-Verlag, 1994, vol. 113, 269-315 [0100]
- HOLLIGER et al. *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, 6444-6448 [0101]
- ZAPATA et al. *Protein Eng.*, 1995, vol. 8 (10), 1057-1062 [0102]
- KABAT et al. Sequences of Proteins of Immunological Interest. National Institutes of Health, 1987 [0107]
- KABAT et al. Sequences of Proteins of Immunological Interest. Public Health Service, National Institutes of Health, 1991 [0108]
- LEFRANC, M-P ; LEFRANC, G. The Immunoglobulin Facts Book. Academic Press, 2001 [0109]
- FIELD et al. *Mol. Cell. Biol.*, 1988, vol. 8, 2159-2165 [0114]
- EVAN et al. *Mol. Cell. Biol.*, 1985, vol. 5 (12), 3610-3616 [0114]
- PABORSKY et al. *Protein Engineering*, 1990, vol. 3 (6), 547-553 [0114]
- AGNEW. *Chem. Int. Ed. Engl.*, vol. 33, 183-186 [0116]

- **WILMAN.** Prodrugs in Cancer Chemotherapy. *Biochemical Society Transactions*, 1986, vol. 14, 375-382 [0117]
- Prodrugs: A Chemical Approach to Targeted Drug Delivery. **BELFAST ; STELLA et al.** Directed Drug Delivery. Humana Press, 1985, 247-267 [0117]
- **AL-LAZIKANI et al.** *JMB*, 1997, vol. 273, 927-948 [0142]
- **HEZAREH et al.** *Journal of Virology*, 2001, vol. 75, 12161-12168 [0143]
- **MORIMOTO et al.** *Journal of Biochemical and Biophysical Methods*, 1992, vol. 24, 107-117 [0176]
- **BRENNAN et al.** *Science*, 1985, vol. 229, 81 [0176]
- **CARTER et al.** *Bio/Technology*, 1992, vol. 10, 163-167 [0176] [0233]
- **CARON et al.** *J. Exp Med.*, 1992, vol. 176, 1191-1195 [0180]
- **SHOPES.** *J. Immunol.*, 1992, vol. 148, 2918-2922 [0180]
- **WOLFF et al.** *Cancer Research*, 1993, vol. 53, 2560-2565 [0180]
- **STEVENSON et al.** *Anti-Cancer Drug Design*, 1989, vol. 3, 219-230 [0180]
- **VITETTA et al.** *Science*, 1987, vol. 238, 1098 [0185]
- **EPSTEIN et al.** *Proc. Natl. Acad. Sci. USA*, 1985, vol. 82, 3688 [0186]
- **HWANG et al.** *Proc. Natl. Acad. Sci. USA*, 1980, vol. 77, 4030 [0186]
- **MARTIN et al.** *J. Biol. Chem.*, 1982, vol. 257, 286-288 [0187]
- **GABIZON et al.** *J. National Cancer Inst.*, 1989, vol. 81 (19), 1484 [0187]
- **MASSEY.** *Nature*, 1987, vol. 328, 457-458 [0188]
- **NEUBERGER et al.** *Nature*, 1984, vol. 312, 604-608 [0188]
- **HAKIMUDDIN et al.** *Arch. Biochem. Biophys.*, 1987, vol. 259, 52 [0191]
- **EDGE et al.** *Anal. Biochem.*, 1981, vol. 118, 131 [0191]
- **THOTAKURA et al.** *Meth. Enzymol.*, 1987, vol. 138, 350 [0191]
- **CUNNINGHAM ; WELLS.** *Science*, 1989, vol. 244, 1081-1085 [0194]
- **STINCHCOMB et al.** *Nature*, 1979, vol. 282, 39 [0216]
- **JONES.** *Genetics*, 1977, vol. 85, 12 [0216]
- **VAN DEN BERG.** *Bio/Technology*, 1990, vol. 8, 135 [0217]
- **FLEER et al.** *Bio/Technology*, 1991, vol. 9, 968-975 [0217]
- **REYES et al.** *Nature*, 1982, vol. 297, 598-601 [0223]
- **YANIV.** *Nature*, 1982, vol. 297, 17-18 [0224]
- **GRAHAM et al.** *J. Gen Virol.*, 1977, vol. 36, 59 [0230]
- **URLAUB et al.** *Proc. Natl. Acad. Sci. USA*, 1980, vol. 77, 4216 [0230]
- **MATHER.** *Biol. Reprod.*, 1980, vol. 23, 243-251 [0230]
- **MATHER et al.** *Annals N.Y. Acad. Sci.*, 1982, vol. 383, 44-68 [0230]
- **HAM et al.** *Meth. Enz.*, 1979, vol. 58, 44 [0232]
- **BARNES et al.** *Anal. Biochem.*, 1980, vol. 102, 255 [0232]
- **LINDMARK et al.** *J. Immunol. Meth.*, 1983, vol. 62, 1-13 [0234]
- **GUSS et al.** *EMBO J.*, 1986, vol. 5, 1567-1575 [0234]
- **KARLIN ; ALTSCHUL.** *Proc. Natl. Acad. Sci. USA*, 1990, vol. 87, 2264-2268 [0242]
- **KARLIN ; ALTSCHUL.** *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, 5873-5877 [0242]
- **ALTSCHUL et al.** *J. Mol. Biol.*, 1990, vol. 215, 403-410 [0242]
- **ALTSCHUL et al.** *Nucleic Acids Res.*, 1997, vol. 25, 3389-3402 [0242]
- **MYERS ; MILLER.** *CABIOS*, 1989 [0242]
- **TORELLIS ; ROBOTI.** *Comput. Appl. Biosci.*, 1994, vol. 10, 3-5 [0242]
- **PEARSON ; LIPMAN.** *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 2444-8 [0242]
- **HIGGINS et al.** *Methods Enzymol.*, 1996, vol. 266, 383-402 [0242]
- Current Protocols in Immunology. Wiley-Interscience, 1991, vol. 1 and 2 [0249]
- Methods for the Preparation of Enzyme-Antibody Conjugates for use in Enzyme immunoassay. **O'SULLIVAN et al.** Methods in Enzym. Academic press, 1981, vol. 73, 147-166 [0252]
- **ZOLA.** Monoclonal Antibodies: A Manual of Techniques. CRC Press, Inc, 1987, 147-158 [0256]
- Fundamental Immunology. Raven Press, 1993 [0267]
- **LANGER.** *Science*, 1990, vol. 249, 1527-1533 [0281]
- **SEFTON.** *Crit. Ref. Biomed. End.* CRC, 1989, vol. 14, 201 [0281]
- **BUCHWALD et al.** *Surgery*, 1980, vol. 88, 507 [0281]
- **SAUDEK et al.** *N. Engl. J. Med.*, 1989, vol. 321, 574 [0281]
- Medical Applications of Controlled Release. CRC Press, 1974 [0281]
- Controlled Drug Bioavailability, Drug Product Design and Performance. Wiley, 1984 [0281]
- **RANGER ; PEPPAS.** *Macromol. Sci. Rev. Macromol. Chem.*, 1983, vol. 23, 61 [0281]
- **LEVY et al.** *Science*, 1985, vol. 228, 190 [0281]
- **DURING et al.** *Ann. Neurol.*, 1989, vol. 25, 351 [0281]
- **HOWARD et al.** *J. Neurosurg.*, 1989, vol. 71, 105 [0281]

Anti-IL-23 antitestek

SZABADALMI IGÉNYPONTOK

1. Humanizált anti-IL-23p19 antitest vagy ennek antigén-kötő fragmentuma, ahol az antitest vagy ennek az antigén-kötő fragmentuma tartalmaz:

- a) egy könnyűlánc variábilis régiót, amely tartalmazza a SEQ ID NO: 19 (CDR1-L) aminosav-szekvenciát; a SEQ ID NO: 20 (CDR2-L) aminosav-szekvenciát; és a SEQ ID NO: 21 (CDR3-L) aminosav-szekvenciát; és
- b) egy nehézlánc variábilis régiót, amely tartalmazza a SEQ ID NO: 66 vagy 67 (CDR1-H) aminosav-szekvenciát; a SEQ ID NO: 64 (CDR2-H) aminosav-szekvenciát; és a SEQ ID NO: 65 (CDR3-H) aminosav-szekvenciát.

2. Az 1. igénypont szerinti humanizált anti-IL-23p19 antitest vagy ennek antigén-kötő fragmentuma,

ahol az antitest vagy ennek az antigén-kötő fragmentuma tartalmaz:

- a) egy könnyűlánc variábilis régiót, amely tartalmazza a SEQ ID NO: 19 (CDR1-L) aminosav-szekvenciát; a SEQ ID NO: 20 (CDR2-L) aminosav-szekvenciát; és a SEQ ID NO: 21 (CDR3-L) aminosav-szekvenciát; és
- b) egy nehézlánc variábilis régiót, amely tartalmazza a SEQ ID NO: 66 (CDR1-H) aminosav-szekvenciát; a SEQ ID NO: 64 (CDR2-H) aminosav-szekvenciát; és a SEQ ID NO: 65 (CDR3-H) aminosav-szekvenciát.

3. Az 1. igénypont szerinti humanizált anti-IL-23p19 antitest vagy ennek antigén-kötő fragmentuma,

ahol az antitest vagy ennek az antigén-kötő fragmentuma tartalmaz:

- a) egy könnyűlánc variábilis régiót, amely tartalmazza a SEQ ID NO: 19 (CDR1-L) aminosav-szekvenciát; a SEQ ID NO: 20 (CDR2-L) aminosav-szekvenciát; és a SEQ ID NO: 21 (CDR3-L) aminosav-szekvenciát; és
- b) egy nehézlánc variábilis régiót, amely tartalmazza a SEQ ID NO: 67 (CDR1-H) aminosav-szekvenciát; a SEQ ID NO: 64 (CDR2-H) aminosav-szekvenciát; és a SEQ ID NO: 65 (CDR3-H) aminosav-szekvenciát.

4. Az 1. igénypont szerinti humanizált anti-IL-23p19 antitest vagy ennek antigén-kötő fragmentuma, ahol az antitest vagy ennek az antigén-kötő fragmentuma tartalmaz egy könnyűlánc variábilis régiót, amely tartalmazza a SEQ ID NO: 158,

160, 162 vagy 164 aminosav-szekvenciák egyikét; és egy nehézlánc variabilis régiót, amely tartalmazza a SEQ ID NO: 166 vagy 168 aminosav-szekvenciák egyikét.

5. A 4. igénypont szerinti humanizált anti-IL-23p19 antitest vagy ennek antigén-kötő fragmentuma, ahol az antitest vagy ennek az antigén-kötő fragmentuma tartalmaz egy könnyűlánc variabilis régiót, amely tartalmazza a SEQ ID NO: 160 aminosav-szekvenciát; és egy nehézlánc variabilis régiót, amely tartalmazza a SEQ ID NO: 166 aminosav-szekvenciát.

6. A 4. igénypont szerinti humanizált anti-IL-23p19 antitest vagy ennek antigén-kötő fragmentuma, ahol az antitest vagy ennek az antigén-kötő fragmentuma tartalmaz egy könnyűlánc variabilis régiót, amely tartalmazza a SEQ ID NO: 160 aminosav-szekvenciát; és egy nehézlánc variabilis régiót, amely tartalmazza a SEQ ID NO: 168 aminosav-szekvenciát.

7. A 4. igénypont szerinti humanizált anti-IL-23p19 antitest vagy ennek antigén-kötő fragmentuma, ahol az antitest vagy ennek az antigén-kötő fragmentuma tartalmaz egy könnyűlánc variabilis régiót, amely tartalmazza a SEQ ID NO: 158 aminosav-szekvenciát; és egy nehézlánc variabilis régiót, amely tartalmazza a SEQ ID NO: 166 aminosav-szekvenciát.

8. A 4. igénypont szerinti humanizált anti-IL-23p19 antitest vagy ennek antigén-kötő fragmentuma, ahol az antitest vagy ennek az antigén-kötő fragmentuma tartalmazza a SEQ ID NO: 166 vagy 168 aminosav-szekvenciát egy humán IgG1 nehézlánc konstans régióhoz kötve; és a SEQ ID NO: 158 vagy 160 aminosav-szekvenciát egy humán kappa könnyűlánc konstans régióhoz kötve.

9. Az 1. igénypont szerinti humanizált anti-IL-23p19 antitest vagy ennek antigén-kötő fragmentuma, ahol az antitest vagy ennek az antigén-kötő fragmentuma tartalmaz:

- a) egy humanizált könnyűlánc variabilis domént, amely tartalmazza a SEQ ID NO: 158 vagy 160 CDR-eket és olyan váz régiókat, amelyeknek aminosav-szekvenciája legalább 90%-ban egyezik a SEQ ID NO: 158 vagy 160 variabilis domén könnyűlánc aminosav-szekvencia váz régiójának aminosav-szekvenciájával; és
- b) egy humanizált nehézlánc variabilis domént, amely tartalmazza a SEQ ID NO: 166 vagy 168 CDR-eket és olyan váz régiókat, amelyeknek aminosav-szekvenciája legalább 90%-ban egyezik a SEQ ID NO: 166 vagy 168 variabilis domén nehézlánc aminosav-szekvencia váz régiójának aminosav-szekvenciájával.

10. Az 1-9. igénypontok bármelyike szerinti humanizált anti-IL-23p19 antitest, ahol az antitest egy monoklonális antitest.

11. Az 1. igénypont szerinti humanizált anti-IL-23p19 antitest, ahol az antitest egy monoklonális anti-IL-23p19 antitest, amely tartalmaz egy, a SEQ ID NO: 174 vagy 180 aminosav-szekvenciát tartalmazó könnyűláncot és egy, a SEQ ID NO: 176 vagy 178 aminosav-szekvenciát tartalmazó nehézláncot.

12. A 11. igénypont szerinti humanizált monoklonális anti-IL-23p19 antitest, ahol az antitest tartalmaz egy, a SEQ ID NO: 174 aminosav-szekvenciát tartalmazó könnyűláncot és egy, a SEQ ID NO: 176 aminosav-szekvenciát tartalmazó nehézláncot.

13. A 11. igénypont szerinti humanizált monoklonális anti-IL-23p19 antitest, ahol az antitest tartalmaz egy, a SEQ ID NO: 174 aminosav-szekvenciát tartalmazó könnyűláncot és egy, a SEQ ID NO: 178 aminosav-szekvenciát tartalmazó nehézláncot.

14. A 11. igénypont szerinti humanizált monoklonális anti-IL-23p19 antitest, ahol az antitest tartalmaz egy, a SEQ ID NO: 180 aminosav-szekvenciát tartalmazó könnyűláncot és egy, a SEQ ID NO: 176 aminosav-szekvenciát tartalmazó nehézláncot.

15. Az 1-14. igénypontok bármelyike szerinti antitest vagy antigén-kötő fragmentum gyógyászatban való alkalmazásra.

16. A 15. igénypont szerinti antitest vagy antigén-kötő fragmentum, ahol az alkalmazás egy gyulladásos betegség, egy autoimmun betegség, egy légzőszervi betegség, egy metabolikus rendellenesség vagy rák kezelése.

17. A 16. igénypont szerinti antitest vagy antigén-kötő fragmentum, ahol az alkalmazás psoriasis, gyulladásos bélbetegség, psoriasisos arthritis, sclerosis multiplex, rheumatoid arthritis, Crohn-betegség, fekélyes kolitis vagy ankylosis spondylitis kezelésére szolgál.

18. A 16. igénypont szerinti antitest vagy antigén-kötő fragmentum, ahol az alkalmazás asztma vagy COPD (krónikus obstruktív tüdőbetegség) kezelésére szolgál.

19. Gyógyszerkészítmény, amely tartalmaz egy, az 1-14. igénypontok bármelyike szerinti antitestet vagy antigén-kötő fragmentumot és egy gyógyszerészeti leg elfogadható hordozót.

20. Izolált polinukleotid, amely tartalmaz egy, a SEQ ID NO: 158, 160, 162 vagy 164 aminosav-szekvenciával rendelkező antitest vagy antitest fragmentum könnyűlánc variabilis régióját kódoló, és egy, a SEQ ID NO: 166 vagy 168 aminosav-szekvenciával rendelkező antitest vagy antitest fragmentum nehézlánc variabilis régióját kódoló szekvenciát.

21. A 20. igénypont szerinti izolált polinukleotid, ahol a polinukleotid tartalmaz egy, a SEQ ID NO: 174 vagy 180 aminosav-szekvenciával rendelkező antitest könnyűláncát kódoló, és egy, a SEQ ID NO: 176 vagy 178 aminosav-szekvenciával rendelkező antitest nehézláncát kódoló szekvenciát.

22. Expressziós vektor, amely tartalmazza a 20. vagy 21. igénypont szerinti polinukleotidot.

23. Gazdasejt, amely tartalmaz:

- a) egy, a SEQ ID NO: 158, 160, 162 vagy 164 aminosav-szekvenciával rendelkező antitest vagy antitest fragmentum könnyűlánc variabilis régióját kódoló szekvenciát; és
- b) egy, a SEQ ID NO: 166 vagy 168 aminosav-szekvenciával rendelkező antitest vagy antitest fragmentum nehézlánc variabilis régióját kódoló szekvenciát.

24. Gazdasejt, amely tartalmaz:

- a) egy, a SEQ ID NO: 174 vagy 180 aminosav-szekvenciával rendelkező antitest könnyűláncát kódoló szekvenciát; és
- b) egy, a SEQ ID NO: 176 vagy 178 aminosav-szekvenciával rendelkező antitest nehézláncát kódoló szekvenciát.

25. Eljárás egy antitest vagy antitest-kötő fragmentum előállítására, amely magában foglalja:

- a) egy, a 23. vagy 24. igénypont szerinti gazdasejt nyerését; és
- b) a gazdasejt ténylezítését.

26. A 25. igénypont szerinti eljárás, amely továbbá magában foglalja az antitest vagy antigén-kötő fragmentum kinyerését és tisztítását.

A meghatalmazott:



Dr. Bánkás Árpád
szabadalmi igazgató
SSOK Szabadalmi Ügyelői Iroda
H-1055 Budapest, Andrásfalvi utca 113.
Telefon: +36-1-33006 Fax: +36-1-38999
E-mail: bankas@ssok.hu