

How do butterflies stick to branches during metamorphosis?

August 9 2024



Caterpillars spin silky carpets and seatbelt-like structures to protect themselves during metamorphosis. Credit: *ACS Biomaterials Science & Engineering* (2024). DOI: 10.1021/acsbiomaterials.4c00237

Most of us learned about butterfly metamorphosis as a kid—a wriggly



caterpillar molts its skin to form a tough chrysalis and emerges as a beautiful butterfly. But how exactly do chrysalises stay anchored as the butterfly brews within?

Research <u>published</u> in *ACS Biomaterials Science & Engineering* shows that, despite their silks being weak and thin on their own, caterpillars can expertly spin them into chrysalis support structures resembling hook-and-loop fasteners and multi-strand safety tethers.

Silk is a natural protein-based fiber that's commonly associated with silkworms, but it's also produced by other insects, including butterflies. In their caterpillar stage, butterflies use silk for spinning tethers to protect themselves from falls, tying up leaf bundles to hide in and weaving little silk carpets to latch onto while undergoing metamorphosis.

When forming a chrysalis, caterpillars use a special arm-like structure called a cremaster that can grab this silk carpet and secure its chrysalis to a branch. Some species of butterflies go a step further and spin a special seatbelt around their thorax for extra security.

But how strong is that silk, and what kind of structures make up these seatbelts and silk carpets? Qingyou Xia, Zhaoming Dong and colleagues raised some caterpillars and found out.

The researchers raised two species of butterflies—Danaus chrysippus (plain tiger) and Papilio polytes (Common mormon)—and two species of silkworms, which they allowed to form chrysalises or cocoons, respectively. The team collected the silk threads that attached the chrysalises to branches or formed the cocoons and measured the diameters and tensile strengths. They found that butterfly silk was thinner and weaker than silkworm silk.

When the primary structural proteins in the silks were analyzed, the



researchers saw that butterfly silk has fewer beta sheet structures compared to silkworm, and that likely contributes to its relative weakness.

Despite this, the team found that both species of butterflies adopt unique strategies to make the most of their silk. The cremaster appendage affixes itself to the silk carpet just like a hook-and-loop fastener and the anchor shape of the cremaster's hooks makes for a secure bond.

When forming their <u>silk</u> seatbelts, caterpillars spin together approximately 20 different strands into a rope-like form, which increases its strength eight-fold. The researchers say their results help explain the unique strategies that caterpillars use to stay safe and secure until they emerge in their final butterfly forms.

More information: Mengyao Lu et al, How Do Butterflies Use Silk to Attach their Pupae to Trees?, *ACS Biomaterials Science & Engineering* (2024). DOI: 10.1021/acsbiomaterials.4c00237

Provided by American Chemical Society

Citation: How do butterflies stick to branches during metamorphosis? (2024, August 9) retrieved 11 August 2024 from https://phys.org/news/2024-08-butterflies-metamorphosis.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.