

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use BYDUREON safely and effectively. See full prescribing information for BYDUREON.

BYDUREON® (exenatide) for extended-release injectable suspension, for subcutaneous use
Initial U.S. Approval: 2005

WARNING: RISK OF THYROID C-CELL TUMORS

See full prescribing information for complete boxed warning.

- Exenatide extended-release causes thyroid C-cell tumors at clinically relevant exposures in rats. It is unknown whether BYDUREON causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC) in humans, as the human relevance of exenatide extended-release-induced rodent thyroid C-cell tumors has not been determined (5.1, 13.1).
- BYDUREON is contraindicated in patients with a personal or family history of MTC or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). Counsel patients regarding the potential risk of MTC and the symptoms of thyroid tumors (4, 5.1).

RECENT MAJOR CHANGES

Warnings and Precautions, Acute Gallbladder Disease (5.10) 06/2022

INDICATIONS AND USAGE

BYDUREON is a glucagon-like peptide-1 (GLP-1) receptor agonist indicated as an adjunct to diet and exercise to improve glycemic control in adults and pediatric patients aged 10 years and older with type 2 diabetes mellitus. (1)

Limitations of Use:

- Not recommended as first-line therapy for patients who have inadequate glycemic control on diet and exercise. (1)
- Not indicated to treat type 1 diabetes mellitus. (1)
- BYDUREON is an extended-release formulation of exenatide and should not be used with other exenatide-containing products. (1)
- Has not been studied in patients with a history of pancreatitis. Consider other antidiabetic therapies in patients with a history of pancreatitis. (1, 5.2)

DOSAGE AND ADMINISTRATION

- Administer 2 mg by subcutaneous injection once every seven days (weekly), at any time of day and with or without meals. (2.1)
- Administer immediately after the dose is prepared. (2.2)
- See Full Prescribing Information for important preparation and administration instructions including required reconstitution prior to administration. (2.2)

DOSAGE FORMS AND STRENGTHS

For extended-release injectable suspension available as: (3)

- Single-dose pen containing 2 mg of exenatide.

CONTRAINDICATIONS

- Personal or family history of medullary thyroid carcinoma or in patients with Multiple Endocrine Neoplasia syndrome type 2. (4)
- Prior serious hypersensitivity reaction to exenatide or any of the product components. (4)

- History of drug-induced immune-mediated thrombocytopenia from exenatide products. (4)

WARNINGS AND PRECAUTIONS

- **Acute Pancreatitis:** Including fatal and non-fatal hemorrhagic or necrotizing pancreatitis has been reported. Discontinue promptly if pancreatitis is suspected. Do not restart if pancreatitis is confirmed. Consider other antidiabetic therapies if patient has history of pancreatitis. (5.2)
- **Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin:** Patients taking an insulin secretagogue or insulin may have an increased risk of hypoglycemia, including severe hypoglycemia. Reduction in the dose of insulin secretagogue or insulin may be necessary. (5.3)
- **Acute Kidney Injury:** May induce nausea and vomiting with transient hypovolemia and may worsen renal function. Postmarketing increased serum creatinine, renal impairment, worsened chronic renal failure and acute renal failure, sometimes requiring hemodialysis or kidney transplantation has been reported. Not recommended for use in patients with eGFR below 45 mL/min/1.73 m². (5.4, 8.6, 12.3)
- **Gastrointestinal Disease:** Not recommended in patients with severe gastrointestinal disease (e.g., gastroparesis). (5.5)
- **Immunogenicity-Associated Decreased Glycemic Control:** Patients may develop antibodies to exenatide. If there is worsening glycemic control or failure to achieve target glycemic control, consider alternative antidiabetic therapy. (5.6)
- **Hypersensitivity Reactions:** Serious hypersensitivity reactions (e.g., anaphylaxis and angioedema) have been reported. In such cases, patients are to discontinue BYDUREON and promptly seek medical advice. (5.7)
- **Drug-induced Immune-mediated Thrombocytopenia:** Serious bleeding which may be fatal has been reported. Discontinue BYDUREON promptly and avoid re-exposure to exenatide. (5.8)
- **Serious Injection-site Reactions:** Serious injection-site reactions with or without subcutaneous nodules have been reported. (5.9)
- **Acute Gallbladder Disease:** If cholelithiasis or cholecystitis are suspected, gallbladder studies are indicated. (5.10)

ADVERSE REACTIONS

Most common (≥5%) and occurring more frequently than comparator in clinical trials: nausea, diarrhea, headache, vomiting, constipation, injection-site pruritus, injection-site nodule, and dyspepsia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact AstraZeneca at 1-800-236-9933 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- May impact absorption of orally administered medications. (7)
- Warfarin: Postmarketing reports with exenatide of increased international normalized ratio (INR) sometimes associated with bleeding. Monitor INR frequently until stable upon initiation of BYDUREON therapy. (7)

USE IN SPECIFIC POPULATIONS

Pregnancy: Use during pregnancy only if the potential benefit justifies the risk to the fetus. (8.1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

Revised: 12/2022

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: RISK OF THYROID C-CELL TUMORS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosage
- 2.2 Preparation and Administration Instructions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Risk of Thyroid C-cell Tumors
- 5.2 Acute Pancreatitis
- 5.3 Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin
- 5.4 Acute Kidney Injury
- 5.5 Gastrointestinal Disease

5.6 Immunogenicity-Associated Decreased Glycemic Control

5.7 Hypersensitivity Reactions

5.8 Drug-Induced Thrombocytopenia

5.9 Serious Injection-Site Reactions

5.10 Acute Gallbladder Disease

6 ADVERSE REACTIONS

6.1 Clinical Trial Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6	Renal Impairment	14.2	Glycemic Control Trials in Adults with Type 2 Diabetes Mellitus - Monotherapy
10	OVERDOSAGE	14.3	Glycemic Control Trials in Adults with Type 2 Diabetes Mellitus - Combination Therapy
11	DESCRIPTION	14.4	EXSCEL Cardiovascular Outcomes Trial in Adult Patients with Type 2 Diabetes Mellitus
12	CLINICAL PHARMACOLOGY	14.5	Glycemic Control Trial in Pediatric Patients 10 Years of Age and Older with Type 2 Diabetes Mellitus
12.1	Mechanism of Action	16	HOW SUPPLIED/STORAGE AND HANDLING
12.2	Pharmacodynamics	17	PATIENT COUNSELING INFORMATION
12.3	Pharmacokinetics		
12.6	Immunogenicity		
13	NONCLINICAL TOXICOLOGY		
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility		
14	CLINICAL STUDIES		
14.1	Overview of Clinical Trials		

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: RISK OF THYROID C-CELL TUMORS

- Exenatide extended-release causes an increased incidence in thyroid C-cell tumors at clinically relevant exposures in rats compared to controls. It is unknown whether BYDUREON causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans, as the human relevance of exenatide extended-release-induced rodent thyroid C-cell tumors has not been determined [see *Warnings and Precautions (5.1)* and *Nonclinical Toxicology (13.1)*].
- BYDUREON is contraindicated in patients with a personal or family history of MTC and in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). Counsel patients regarding the potential risk for MTC with the use of BYDUREON and inform them of symptoms of thyroid tumors (e.g., mass in the neck, dysphagia, dyspnea, persistent hoarseness). Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for detection of MTC in patients treated with BYDUREON [see *Contraindications (4)* and *Warnings and Precautions (5.1)*].

1 INDICATIONS AND USAGE

BYDUREON is indicated as an adjunct to diet and exercise to improve glycemic control in adults and pediatric patients aged 10 years and older with type 2 diabetes mellitus [see *Clinical Studies (14)*].

Limitations of Use:

BYDUREON:

- Is not recommended as first-line therapy for patients who have inadequate glycemic control on diet and exercise because of the uncertain relevance of the rat thyroid C-cell tumor findings to humans [see *Warnings and Precautions (5.1)*].
- Is not indicated for use in patients with type 1 diabetes mellitus.
- Is an extended-release formulation of exenatide and should not be used with other exenatide-containing products.
- Has not been studied in patients with a history of pancreatitis. Consider other antidiabetic therapies in patients with a history of pancreatitis [see *Warnings and Precautions (5.2)* and *Adverse Reactions (6.2)*].

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

- The recommended dosage of BYDUREON is 2 mg subcutaneously once every 7 days (weekly). The dose can be administered at any time of day, with or without meals.
- Discontinue an immediate- or extended-release exenatide product prior to initiation of BYDUREON. Patients changing from immediate-release exenatide to BYDUREON may experience transient (approximately 2 to 4 weeks) elevations in blood glucose concentrations.
- The day of weekly administration can be changed, if necessary, as long as the last dose was administered 3 or more days before the new day of administration.

Recommendations Regarding Missed Dose

- If a dose is missed, administer the dose as soon as noticed, provided the next regularly scheduled dose is due at least 3 days later. Thereafter, administer subsequent doses once every 7 days (weekly).
- If a dose is missed and the next regularly scheduled dose is due 1 or 2 days later, do not administer the missed dose and instead resume BYDUREON with the next regularly scheduled dose.

2.2 Preparation and Administration Instructions

- BYDUREON requires reconstitution with the supplied diluent prior to use to obtain a final concentration of 2 mg of exenatide per 0.65 mL of suspension.
- Prior to initiation, train patients and caregivers on proper mixing and injection [*see Instructions for Use*]. Instruct caregivers to assist pediatric patients with mixing and administration.
- Inspect BYDUREON visually before use. The suspension should appear white to off-white and cloudy. Do not use if particulate matter is present or if discoloration is observed.
- Immediately after the dose is prepared, administer BYDUREON subcutaneously to the abdomen, thigh, or upper arm region. Instruct patients to use a different injection site each week when injecting in the same region.
- When using BYDUREON with insulin, always administer BYDUREON and insulin as separate injections and never mix the products. It is acceptable to inject BYDUREON and insulin in the same body region, but avoid injecting the products next to each other.

3 DOSAGE FORMS AND STRENGTHS

For extended-release injectable suspension available as a:

- Single-dose pen which contains 2 mg of exenatide white to off-white powder, 0.65 mL of diluent, and one needle. Each carton contains one spare needle.

4 CONTRAINDICATIONS

BYDUREON is contraindicated in patients with a:

- Personal or family history of medullary thyroid carcinoma (MTC) or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).
- Prior serious hypersensitivity reaction to exenatide or to any of the excipients in BYDUREON. Serious hypersensitivity reactions including anaphylaxis and angioedema have been reported with BYDUREON [see *Warnings and Precautions* (5.7)].
- History of drug-induced immune-mediated thrombocytopenia from exenatide products. Serious bleeding, which may be fatal, from drug-induced immune-mediated thrombocytopenia has been reported with exenatide use [see *Warnings and Precautions* (5.8)].

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Thyroid C-cell Tumors

In both genders of rats, exenatide extended-release caused a dose-related and treatment-duration–dependent increase in the incidence of thyroid C-cell tumors (adenomas and/or carcinomas) at clinically relevant exposures compared to controls [see *Nonclinical Toxicology* (13.1)]. A statistically significant increase in malignant thyroid C-cell carcinomas was observed in female rats receiving exenatide extended-release at 25-times clinical exposure compared to controls and higher incidences were noted in males above controls in all treated groups at ≥ 2 -times clinical exposure. The potential of exenatide extended-release to induce C-cell tumors in mice has not been evaluated. Other GLP-1 receptor agonists have also induced thyroid C-cell adenomas and carcinomas in male and female mice and rats at clinically relevant exposures. It is unknown whether BYDUREON will cause thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans as the human relevance of exenatide extended-release-induced rodent thyroid C-cell tumors has not been determined.

Cases of MTC in patients treated with liraglutide, another GLP-1 receptor agonist, have been reported in the postmarketing period; the data in these reports are insufficient to establish or exclude a causal relationship between MTC and GLP-1 receptor agonist use in humans.

BYDUREON is contraindicated in patients with a personal or family history of MTC or in patients with MEN 2. Counsel patients regarding the potential risk of MTC with the use of BYDUREON and inform them of symptoms of thyroid tumors (e.g., a mass in the neck, dysphagia, dyspnea, persistent hoarseness).

Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with BYDUREON. Such monitoring may increase the risk of unnecessary procedures, due to the low specificity of serum calcitonin testing for MTC and a high background incidence of thyroid disease. Significantly elevated serum calcitonin may indicate MTC and patients with MTC usually have values >50 ng/L. If serum calcitonin is measured and found to be elevated, the patient should be further evaluated. Patients with thyroid nodules noted on physical examination or neck imaging should also be further evaluated.

5.2 Acute Pancreatitis

Based on postmarketing data, exenatide has been associated with acute pancreatitis, including fatal and non-fatal hemorrhagic or necrotizing pancreatitis. After initiation of BYDUREON, observe patients carefully for signs and symptoms of pancreatitis (including persistent severe abdominal pain, sometimes radiating to the back, which may or may not be accompanied by vomiting). If pancreatitis is suspected, BYDUREON should promptly be discontinued and appropriate management should be initiated. If pancreatitis is confirmed, BYDUREON should not be restarted. Consider antidiabetic therapies other than BYDUREON in patients with a history of pancreatitis.

5.3 Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin

Patients receiving BYDUREON in combination with an insulin secretagogue (e.g., sulfonylurea) or insulin may have an increased risk of hypoglycemia, including severe hypoglycemia [see *Adverse Reactions (6.1) and Drug Interactions (7)*].

The risk of hypoglycemia may be lowered by a reduction in the dose of sulfonylurea (or other concomitantly administered insulin secretagogues) or insulin. Inform patients using these concomitant medications of the risk of hypoglycemia and educate them on the signs and symptoms of hypoglycemia.

5.4 Acute Kidney Injury

BYDUREON may induce nausea and vomiting with transient hypovolemia and may worsen renal function. There have been postmarketing reports of altered renal function with exenatide, including increased serum creatinine, renal impairment, worsened chronic renal failure and acute renal failure, sometimes requiring hemodialysis or kidney transplantation. Some of these events occurred in patients receiving one or more pharmacologic agents known to affect renal function or hydration status such as angiotensin converting enzyme inhibitors, nonsteroidal anti-inflammatory drugs, or diuretics. Some events occurred in patients who had been experiencing nausea, vomiting or diarrhea, with or without dehydration. Reversibility of altered renal function has been observed in many cases with supportive treatment and discontinuation of potentially causative agents, including BYDUREON. BYDUREON is not recommended for use in patients with an eGFR below 45 mL/min/1.73 m² [see *Use in Specific Populations (8.6)*].

5.5 Gastrointestinal Disease

Exenatide has not been studied in patients with severe gastrointestinal disease, including gastroparesis. Because exenatide is commonly associated with gastrointestinal adverse reactions, including nausea, vomiting, and diarrhea, the use of BYDUREON is not recommended in patients with severe gastrointestinal disease.

5.6 Immunogenicity-Associated Decreased Glycemic Control

In the 24- to 30-week treatment periods in Trials 1, 2, 3, 4 and 9 in adult patients with type 2 diabetes mellitus [see *Clinical Studies (14.2, 14.3) and Adverse Reactions (6.1)*], BYDUREON-treated patients who had high titer anti-exenatide antibodies (defined as ≥ 625) had a lower glycemic response compared

to BYDUREON-treated patients who did not have anti-exenatide antibodies [see *Clinical Pharmacology (12.6)*].

In the 24-week treatment period in Trial 8 in pediatric patients with type 2 diabetes mellitus [see *Clinical Studies (14.5)*], BYDUREON-treated patients who had high titer anti-exenatide antibodies (≥ 625) appeared to have had a lower glycemic response compared to BYDUREON-treated patients who did not have anti-exenatide antibodies. At Week 24, the mean change in HbA_{1c} from baseline in BYDUREON-treated patients was greater (-0.73%) in those with low titer anti-exenatide antibodies (< 625) compared to +0.07% in those with high titer anti-exenatide antibodies [see *Clinical Pharmacology (12.6)*].

If there is worsening glycemic control or failure to achieve targeted glycemic control, consider stopping BYDUREON and using alternative antidiabetic therapy.

5.7 Hypersensitivity Reactions

There have been postmarketing reports of serious hypersensitivity reactions (e.g., anaphylaxis and angioedema) in patients treated with exenatide. If a hypersensitivity reaction occurs, the patient should discontinue BYDUREON and promptly seek medical advice [see *Contraindications (4)* and *Adverse Reactions (6.2)*]. Inform and closely monitor patients with a history of anaphylaxis or angioedema with another GLP-1 receptor agonist for allergic reactions, because it is unknown whether such patients will be predisposed to anaphylaxis with BYDUREON.

5.8 Drug-Induced Thrombocytopenia

Serious bleeding, which may be fatal, from drug-induced immune-mediated thrombocytopenia has been reported in the postmarketing setting with exenatide use. Drug-induced thrombocytopenia is an immune-mediated reaction, with exenatide-dependent anti-platelet antibodies. In the presence of exenatide, these antibodies cause platelet destruction. If drug-induced thrombocytopenia is suspected, discontinue BYDUREON immediately and do not re-expose the patient to exenatide. Upon discontinuation, thrombocytopenia can persist due to the prolonged exenatide exposure from BYDUREON (about 10 weeks) [see *Adverse Reactions (6.2)*].

5.9 Serious Injection-Site Reactions

There have been postmarketing reports of serious injection-site reactions (e.g., abscess, cellulitis, and necrosis), with or without subcutaneous nodules, with the use of BYDUREON. Isolated cases required surgical intervention [see *Adverse Reactions (6.1)*].

5.10 Acute Gallbladder Disease

Acute events of gallbladder disease, such as cholelithiasis or cholecystitis, have been reported in GLP-1 receptor agonist trials and postmarketing. In the EXSCEL trial [see *Clinical Studies (14.4)*], 1.9% of BYDUREON-treated patients and 1.4% of placebo-treated patients reported an acute event of gallbladder disease, such as cholelithiasis or cholecystitis. If cholelithiasis is suspected, gallbladder studies and appropriate clinical follow-up are indicated.

6 ADVERSE REACTIONS

The following serious adverse reactions are described below or elsewhere in the Prescribing Information:

- Risk of Thyroid C-cell Tumors [see *Warnings and Precautions (5.1)*]
- Acute Pancreatitis [see *Warnings and Precautions (5.2)*]
- Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin [see *Warnings and Precautions (5.3)*]
- Acute Kidney Injury [see *Warnings and Precautions (5.4)*]
- Gastrointestinal Disease [see *Warnings and Precautions (5.5)*]
- Immunogenicity-Associated Decreased Glycemic Control [see *Warnings and Precautions (5.6)*]
- Hypersensitivity Reactions [see *Warnings and Precautions (5.7)*]
- Drug-Induced Thrombocytopenia [see *Warnings and Precautions (5.8)*]
- Serious Injection-Site Reactions [see *Warnings and Precautions (5.9)*]
- Acute Gallbladder Disease [see *Warnings and Precautions (5.10)*]

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety data presented below are derived from six comparator-controlled trials of BYDUREON in adult patients who entered the studies not achieving adequate glycemic control on their current therapy [see *Clinical Studies (14.2, 14.3)*].

- In a double-blind 26-week trial, patients on diet and exercise were treated with BYDUREON 2 mg given subcutaneously once every 7 days (weekly), sitagliptin 100 mg daily, pioglitazone 45 mg daily, or metformin 2000 mg daily (Trial 1).
- In a double-blind 26-week trial, patients on metformin were treated with BYDUREON 2 mg given subcutaneously once every 7 days (weekly), sitagliptin 100 mg daily, or pioglitazone 45 mg daily (Trial 2).
- In an open-label 26-week trial, patients on metformin or metformin plus sulfonylurea were treated with BYDUREON 2 mg given subcutaneously once every 7 days (weekly) or optimized insulin glargine (Trial 3).
- In two open-label 24- to 30-week studies, patients on diet and exercise or metformin, a sulfonylurea, a thiazolidinedione, or combination of oral agents were treated with BYDUREON 2 mg given subcutaneously once every 7 days (weekly) or BYETTA 10 mcg twice daily (Trials 4 and 9 [NCT00308139]).
- In an open-label 26-week trial, patients on metformin, a sulfonylurea, metformin plus a sulfonylurea, or metformin plus pioglitazone were treated with BYDUREON 2 mg given subcutaneously every 7 days (weekly) or liraglutide 1.8 mg once daily (Trial 5).

The safety of BYDUREON in pediatric patients age 10 to less than 18 years with type 2 diabetes was similar to that observed in adults (Trial 8) [see *Clinical Studies (14.5)*].

Common Adverse Reactions

Tables 1 and 2 summarize adverse reactions in adults with type 2 diabetes mellitus with an incidence $\geq 5\%$ reported in the six comparator-controlled 24- to 30-week trials of BYDUREON used as monotherapy or

as add-on to metformin, a sulfonylurea, a thiazolidinedione, or combination of these oral antidiabetic agents. Nausea was a common adverse reaction associated with initiation of treatment with BYDUREON and usually decreased over time.

Table 1: Adverse Reactions Reported in $\geq 5\%$ of BYDUREON-Treated Adult Patients with Type 2 Diabetes Mellitus in Monotherapy Trial (Trial 1)

26-Week Monotherapy Trial				
	BYDUREON 2 mg N = 248 %	Sitagliptin 100 mg N = 163 %	Pioglitazone 30-45 mg (mean dose 40 mg) N = 163 %	Metformin 1000-2500 mg (mean dose 2077 mg) N = 246 %
Nausea	11.3	3.7	4.3	6.9
Diarrhea	10.9	5.5	3.7	12.6
Injection-site nodule*	10.5	6.7	3.7	10.2
Constipation	8.5	2.5	1.8	3.3
Headache	8.1	9.2	8.0	12.2
Dyspepsia	7.3	1.8	4.9	3.3

N = number of intent-to-treat patients; Percentages are based on the number of intent-to-treat patients in each treatment group.

* Patients in the sitagliptin, pioglitazone, and metformin treatment groups received weekly placebo injections.

Table 2: Adverse Reactions Reported in $\geq 5\%$ of BYDUREON-Treated Patients with Type 2 Diabetes Mellitus in 24- to 30-Week Add-On Combination Therapy Trials

26-Week Add-On to Metformin Trial (Trial 2)			
	BYDUREON 2 mg (N = 160) %	Sitagliptin 100 mg (N = 166) %	Pioglitazone 45 mg (N = 165) %
Nausea	24.4	9.6	4.8
Diarrhea	20.0	9.6	7.3
Vomiting	11.3	2.4	3.0
Headache	9.4	9.0	5.5
Constipation	6.3	3.6	1.2
Fatigue	5.6	0.6	3.0
Dyspepsia	5.0	3.6	2.4
Decreased appetite	5.0	1.2	0.0
Injection-site pruritus*	5.0	4.8	1.2
26-Week Add-On to Metformin or Metformin + Sulfonylurea Trial (Trial 3)			
	BYDUREON 2 mg (N = 233) %	Insulin Glargine Titrated (N = 223) %	
Nausea	12.9	1.3	
Headache	9.9	7.6	
Diarrhea	9.4	4.0	
Injection-site nodule	6.0	0.0	

Table 2: Adverse Reactions Reported in ≥5% of BYDUREON-Treated Patients with Type 2 Diabetes Mellitus in 24- to 30-Week Add-On Combination Therapy Trials

30-Week Monotherapy or as Add-On to Metformin, a Sulfonylurea, a Thiazolidinedione, or Combination of Oral Agents Trial (Trial 9)		
	BYDUREON 2 mg once weekly (N = 148) %	BYETTA 10 mcg twice daily (N = 145) %
Nausea	27.0	33.8
Diarrhea	16.2	12.4
Vomiting	10.8	18.6
Injection-site pruritus	18.2	1.4
Constipation	10.1	6.2
Gastroenteritis viral	8.8	5.5
Gastroesophageal reflux disease	7.4	4.1
Dyspepsia	7.4	2.1
Injection-site erythema	7.4	0.0
Fatigue	6.1	3.4
Headache	6.1	4.8
Injection-site hematoma	5.4	11.0
24-Week Monotherapy or as Add-On to Metformin, a Sulfonylurea, a Thiazolidinedione, or Combination of Oral Agents Trial (Trial 4)		
	BYDUREON 2 mg (N = 129) %	BYETTA 10 mcg (N = 123) %
Nausea	14.0	35.0
Diarrhea	9.3	4.1
Injection-site erythema	5.4	2.4
26-Week Add-On to Metformin, a Sulfonylurea, Metformin + Sulfonylurea, or Metformin + Pioglitazone Trial (Trial 5)		
	BYDUREON 2 mg (N = 461) %	
Injection-site nodule	10.4	
Nausea	9.3	
Diarrhea	6.1	

N = number of intent-to-treat patients; Percentages are based on the number of intent-to-treat patients in each treatment group.

* Patients in the sitagliptin, pioglitazone, and metformin treatment groups received weekly placebo injections.

Adverse Reactions Leading to Study Withdrawal

The incidence of withdrawal due to adverse reactions in adults with type 2 diabetes mellitus was 4.1% (N=57) for BYDUREON-treated patients, 4.9% (N=13) for BYETTA-treated patients, and 2.9% (N=46) for other comparator-treated patients in the six comparator-controlled 24- to 30-week trials. The most common classes of adverse reactions (0.5%) leading to withdrawal for BYDUREON-treated patients were, Gastrointestinal Disorders 1.6% (N=22) versus 4.1% (N=11) for BYETTA and 1.9% (N=30) for

other comparators, and Administration Site Conditions 0.8% (N=11) versus 0.0% for BYETTA and 0.2% (N=3) for other comparators.

The most frequent adverse reactions within each of these respective classes were, nausea 0.4% (N=6) for BYDUREON versus 1.5% (N=4) for BYETTA and 0.8% (N=12) for other comparators, and injection-site nodule, 0.4% (N=6) for BYDUREON versus 0.0% for BYETTA and 0.0% for other comparators.

Hypoglycemia

Table 3 summarizes the incidence of hypoglycemia (defined as the presence of symptoms of hypoglycemia with a concomitant glucose <54 mg/dL and the ability to self-treat) in the six comparator-controlled 24- to 30-week trials of BYDUREON in adults with type 2 diabetes mellitus. BYDUREON was used as monotherapy or as add-on to metformin, a sulfonylurea, a thiazolidinedione, or combination of these oral antidiabetic agents.

Table 3: Incidence (% of Subjects) of Hypoglycemia* in Clinical Trials in Adult Patients with Type 2 Diabetes Mellitus

26-Week Monotherapy Trial (Trial 1)	
BYDUREON 2 mg (N = 248)	2.0%
Sitagliptin 100 mg (N = 163)	0.0%
Pioglitazone 30-45 (mean dose 40) mg (N = 163)	0.0%
Metformin 1000-2500 (mean dose 2077) mg (N = 246)	0.0%
26-Week Add-On to Metformin Trial (Trial 2)	
BYDUREON 2 mg (N = 160)	1.3%
Sitagliptin 100 mg (N = 166)	3.0%
Pioglitazone 45 mg (N = 165)	1.2%
26-Week Add-On to Metformin or Metformin + Sulfonylurea Trial (Trial 3)	
With Concomitant Sulfonylurea Use (N = 136)	
BYDUREON 2 mg (N = 70)	20.0%
Titrated Insulin Glargine (N = 66)	43.9%
Without Concomitant Sulfonylurea Use (N = 320)	
BYDUREON 2 mg (N = 163)	3.7%
Titrated Insulin Glargine† (N = 157)	19.1%
24-Week Monotherapy or Add-On to Metformin, a Sulfonylurea, a Thiazolidinedione, or Combination of Oral Agents Trial (Trial 4)	
With Concomitant Sulfonylurea Use (N = 74)	
BYDUREON 2 mg (N = 40)	12.5%
BYETTA 10 mcg (N = 34)	11.8%
Without Concomitant Sulfonylurea Use (N = 178)	
BYDUREON 2 mg (N = 89)	0.0%
BYETTA 10 mcg (N = 89)	0.0%
30-Week Monotherapy or Add-On to Metformin, a Sulfonylurea, a Thiazolidinedione, or Combination of Oral Agents Trial (Trial 9)	
With Concomitant Sulfonylurea Use (N = 107)	
BYDUREON 2 mg (N = 55)	14.5%
BYETTA 10 mcg (N = 52)	15.4%
Without Concomitant Sulfonylurea Use (N = 186)	
BYDUREON 2 mg (N = 93)	0.0%
BYETTA 10 mcg (N = 93)	1.1%
26-Week as Add-On to Metformin, a Sulfonylurea, Metformin + Sulfonylurea, or Metformin + Pioglitazone Trial (Trial 5)	
With Concomitant Sulfonylurea Use (N = 590)	
BYDUREON 2 mg (N = 294)	15.3%
Without Concomitant Sulfonylurea Use (N = 321)	
BYDUREON 2 mg (N = 167)	3.6%

N = number of intent-to-treat (ITT) patients; Percentages were based on the number of ITT patients in each treatment group.

* Reported event that has symptoms consistent with hypoglycemia with a concomitant glucose <54 mg/dL and the patient was able to self-treat.

† Insulin glargine was dosed to a target fasting glucose concentration of 72 to 100 mg/dL. The mean dose of insulin glargine was 10 units/day at baseline and 31 units/day at endpoint.

In the 24-week pediatric placebo-controlled clinical trial (Trial 8) [see *Clinical Studies (14.5)*], 2 (3.4%) of BYDUREON-treated patients with type 2 diabetes had hypoglycemia with a blood glucose <54 mg/dL with or without symptoms and 1 (1.7%) had severe hypoglycemia (defined as an episode with severe cognitive impairment requiring external assistance for recovery).

Injection-Site Adverse Reactions Including Nodules

In five comparator-controlled 24- to 30-week trials in adults, injection-site reactions were observed more frequently in BYDUREON-treated patients (17%) than in BYETTA-treated patients (13%), titrated insulin glargine-treated patients (1.8%), or placebo-treated patients (sitagliptin (11%), pioglitazone (6%), and metformin (13%) treatment groups). One percent of BYDUREON-treated patients withdrew due to injection-site adverse reactions (injection-site mass, injection-site nodule, injection-site pruritus, and injection-site reaction).

In a separate 15-week study of BYDUREON in adults in which information on nodules were collected and analyzed, 24 out of 31 subjects (77%) experienced at least 1 injection-site nodule during treatment; 2 subjects (7%) reported accompanying localized symptoms. The mean duration of events was 27 days.

Immunogenicity: Anti-Drug Antibody-Associated Adverse Reactions

Injection site reactions for BYDUREON-treated patients were more commonly observed in those who were antibody-positive (anti-exenatide antibodies) (14%) compared with those who were antibody-negative (3%) [see *Clinical Pharmacology (12.6)*].

Increase in Heart Rate

Increases in heart rate from baseline ranging from 1.5 to 4.5 beats per minute have been observed in BYDUREON-treated patients in comparator-controlled clinical trials in adults.

Cholelithiasis and cholecystitis

In the EXSCEL trial [see *Clinical Studies (14.4)*], 1.9% of BYDUREON-treated patients and 1.4% of placebo-treated patients reported an acute event of gallbladder disease, such as cholelithiasis or cholecystitis.

Other Adverse Reactions

The following adverse reactions were also reported in three 30-week controlled trials (in adults with type 2 diabetes mellitus) of BYETTA (N=963) add-on to metformin and/or sulfonylurea, with an incidence of $\geq 1\%$ and reported more frequently than with placebo: feeling jittery (9% BYETTA, 4% placebo), dizziness (9% BYETTA, 6% placebo), asthenia (4% BYETTA, 2% placebo), and hyperhidrosis (3% BYETTA, 1% placebo).

6.2 Postmarketing Experience

The following additional adverse reactions have been reported during post-approval use of BYDUREON or other formulations of exenatide. Because these events are reported voluntarily from a population of uncertain size, it is generally not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Allergy/Hypersensitivity: injection-site reactions (e.g., abscess, cellulitis, and necrosis, with or without subcutaneous nodules), generalized pruritus and/or urticaria, macular or papular rash, angioedema; anaphylactic reaction.

Blood and Lymphatic Systems: drug-induced thrombocytopenia.

Drug Interactions: increased international normalized ratio (INR), sometimes associated with bleeding, with concomitant warfarin use [*see Drug Interactions (7)*].

Gastrointestinal: nausea, vomiting, and/or diarrhea resulting in dehydration; abdominal distension, abdominal pain, eructation, constipation, flatulence, ileus, acute pancreatitis, hemorrhagic and necrotizing pancreatitis sometimes resulting in death [*see Indications and Usage (1)*].

Hepatobiliary: cholecystitis, cholelithiasis requiring cholecystectomy.

Neurologic: dysgeusia; somnolence.

Renal and Urinary Disorders: altered renal function, including increased serum creatinine, renal impairment, worsened chronic renal failure or acute renal failure (sometimes requiring hemodialysis), kidney transplant and kidney transplant dysfunction.

Skin and Subcutaneous Tissue Disorders: alopecia.

7 DRUG INTERACTIONS

Table 4: Clinically Relevant Interactions with BYDUREON

Concomitant Use of Insulin Secretagogues or Insulin	
Clinical Impact	Exenatide promotes insulin release from pancreatic beta-cells in the presence of elevated glucose concentrations. The risk of hypoglycemia is increased when exenatide is used in combination with insulin secretagogues (e.g., sulfonylureas) or insulin [see <i>Warnings and Precautions (5.3)</i> and <i>Adverse Reactions (6.1)</i>].
Intervention	When initiating BYDUREON, consider reducing the dose of concomitantly administered insulin secretagogue or insulin to reduce the risk of hypoglycemia.
Warfarin	
Clinical Impact	BYDUREON has not been studied with warfarin. However, in a drug interaction study, BYETTA did not have a significant effect on INR [see <i>Clinical Pharmacology (12.3)</i>]. There have been postmarketing reports for exenatide of increased INR with concomitant use of warfarin, sometimes associated with bleeding [see <i>Adverse Reactions (6.2)</i>].
Intervention	In patients taking warfarin, the INR should be monitored more frequently after initiating BYDUREON. Once a stable INR has been documented, the INR can be monitored at the intervals recommended for patients taking warfarin.
Orally Administered Drugs (e.g., acetaminophen)	
Clinical Impact	Exenatide slows gastric emptying. Therefore, BYDUREON has the potential to reduce the rate of absorption of orally administered drugs [see <i>Clinical Pharmacology (12.3)</i>].
Intervention	Use caution when administering oral medications with BYDUREON where a slower rate of oral absorption may be clinically meaningful.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Limited data with exenatide, the active ingredient in BYDUREON, in pregnant women are not sufficient to determine a drug-associated risk for major birth defects or miscarriage. There are risks to the mother and fetus associated with poorly controlled diabetes in pregnancy (see *Clinical Considerations*). Based on animal reproduction studies, there may be risks to the fetus from exposure to BYDUREON during pregnancy. BYDUREON should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Animal reproduction studies identified increased adverse fetal and neonatal outcomes from exposure to exenatide extended-release during pregnancy, or from exposure to exenatide, during pregnancy and lactation, in association with maternal effects. In rats, exenatide extended-release administered during the period of organogenesis reduced fetal growth and produced skeletal ossification deficits at doses that approximate clinical exposures at the maximum recommended human dose (MRHD) of 2 mg/week. In mice, exenatide administered during gestation and lactation caused increased neonatal deaths at doses that approximate clinical exposures at the MRHD (see *Data*). Based on animal data, advise pregnant women of the potential risk to a fetus.

The estimated background risk of major birth defects is 6-10% in women with pre-gestational diabetes with an HbA_{1c} >7 and has been reported to be as high as 20-25% in women with HbA_{1c} >10. The estimated background risk of miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Clinical Considerations

Disease-associated Maternal and/or Embryofetal Risk

Poorly controlled diabetes in pregnancy increases the maternal risk for diabetic ketoacidosis, preeclampsia, spontaneous abortions, preterm delivery and delivery complications. Poorly controlled diabetes increases the fetal risk for major birth defects, stillbirth, and macrosomia related morbidity.

Data

Animal Data

Pregnant rats given subcutaneous doses of 0.3, 1, or 3 mg/kg exenatide extended-release every 3 days during organogenesis had systemic exposures 3-, 7-, and 17-times human exposure, respectively, at the maximum recommended human dose (MRHD) of 2 mg/week BYDUREON based on plasma exenatide exposure (AUC) comparison. Reduced fetal growth at all doses and skeletal ossification deficits at 1 and 3 mg/kg occurred at doses that decreased maternal food intake and body weight gain.

In studies evaluating reproduction and development in pregnant mice and rabbits, maternal animals were administered exenatide, the active ingredient in BYDUREON, by subcutaneous injection twice a day. Differences in embryo-fetal developmental toxicity from subcutaneously injected exenatide extended-release and exenatide were not evaluated in mice, rats, or rabbits.

In pregnant mice given 6, 68, 460, or 760 mcg/kg/day exenatide during fetal organogenesis, skeletal variations associated with slowed fetal growth, including changes in number of rib pairs or vertebral ossification sites, and wavy ribs were observed at 760 mcg/kg/day, a dose that produced maternal toxicity and yielded systemic exposure 200 times the human exposure resulting from the MRHD of BYDUREON based on AUC comparison.

In pregnant rabbits given 0.2, 2, 22, 156, or 260 mcg/kg/day exenatide during fetal organogenesis, irregular fetal skeletal ossifications were observed at 2 mcg/kg/day, a dose yielding systemic exposure up to 4 times the human exposure from the MRHD of BYDUREON based on AUC comparison.

In maternal mice given 6, 68, or 760 mcg/kg/day exenatide from gestation day 6 through lactation day 20 (weaning), an increased number of neonatal deaths were observed on postpartum days 2 to 4 in dams given 6 mcg/kg/day, a dose yielding a systemic exposure equivalent to the human exposure from the MRHD of BYDUREON based on AUC comparison.

8.2 Lactation

Risk Summary

There is no information regarding the presence of exenatide in human milk, the effects of exenatide on the breastfed infant, or the effects of exenatide on milk production. Exenatide, the active ingredient in BYDUREON, was present in the milk of lactating mice. However, due to species-specific differences in lactation physiology, the clinical relevance of these data is not clear (*see Data*).

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for exenatide and any potential adverse effects on the breastfed child from exenatide or from the underlying maternal condition.

Data

In lactating mice subcutaneously injected twice a day with exenatide, the active ingredient in BYDUREON, the concentration of exenatide in milk was up to 2.5% of the concentration in maternal plasma.

8.4 Pediatric Use

The safety and effectiveness of BYDUREON as an adjunct to diet and exercise to improve glycemic control in type 2 diabetes mellitus have been established in pediatric patients aged 10 years and older. Use of BYDUREON for this indication is supported by a 24-week placebo-controlled trial with 28-week open-label uncontrolled extension (Trial 8) in 82 pediatric patients 10 years of age and older with type 2 diabetes, a pediatric pharmacokinetic study, and studies in adults with type 2 diabetes mellitus [*see Clinical Pharmacology (12.3) and Clinical Studies (14.2, 14.3, 14.5)*].

The safety and effectiveness of BYDUREON have not been established in pediatric patients less than 10 years of age.

8.5 Geriatric Use

In five comparator-controlled 24- to 30-week trials, BYDUREON was studied in 132 patients (17%) who were between 65 years of age and 75 years of age and 20 (2.5%) patients who were at least 75 years of age. In these studies, no meaningful differences in safety (N=152) and effectiveness (N=52) were observed between patients ≥ 65 years of age and younger adult patients, but these studies did not include sufficient numbers of patients ≥ 75 years of age to determine whether they respond differently from younger adult patients.

In a large cardiovascular outcomes trial [*see Clinical Studies (14.4)*], BYDUREON was studied in 2,959 patients (40%) who were at least 65 years old and of those, 605 patients (8%) were at least 75 years old.

Use caution when initiating BYDUREON in geriatric patients because they are more likely to have decreased kidney function [*see Use in Specific Populations (8.6)*].

8.6 Renal Impairment

Pharmacokinetic studies of adult patients with renal impairment who received BYDUREON indicate that there is an increase in exenatide exposure in those with mild and moderate renal impairment as compared

to patients with normal kidney function. BYDUREON may induce nausea and vomiting with transient hypovolemia and may worsen kidney function in patients with renal impairment.

Monitor patients with mild renal impairment for adverse reactions that may lead to hypovolemia. BYDUREON is not recommended for use in patients with eGFR below 45 mL/min/1.73 m² or end stage renal disease. If BYDUREON is used in patients with renal transplantation, closely monitor for adverse reactions that may lead to hypovolemia [see *Warnings and Precautions (5.4)* and *Clinical Pharmacology (12.3)*].

10 OVERDOSAGE

Effects of overdoses with BYETTA, another exenatide product, included severe nausea, severe vomiting, and rapidly declining blood glucose concentrations, including severe hypoglycemia requiring parenteral glucose administration. In the event of overdose, appropriate supportive treatment should be initiated according to the patient's clinical signs and symptoms.

11 DESCRIPTION

BYDUREON, (exenatide) for extended-release injectable suspension, is a GLP-1 receptor agonist supplied as a sterile powder to be suspended in diluent and administered by subcutaneous injection. Exenatide is a 39-amino acid synthetic peptide amide with an empirical formula of C₁₈₄H₂₈₂N₅₀O₆₀S and a molecular weight of 4186.6 Daltons. The amino acid sequence for exenatide is shown below.

H-His-Gly-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Leu-Ser-Lys-Gln-Met-Glu-Glu-Glu-Ala-Val-Arg-Leu-Phe-Ile-Glu-Trp-Leu-Lys-Asn-Gly-Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-NH₂

BYDUREON is a white to off-white powder that is available in a dosage strength of 2 mg exenatide per pen. Exenatide is incorporated in an extended-release microsphere formulation containing the 50:50 poly(D,L-lactide-co-glycolide) polymer (37.2 mg per dose) along with sucrose (0.8 mg per dose). The powder must be suspended in the diluent prior to injection.

The diluent (0.65 mL) for the BYDUREON Pen is contained within each single-dose pen. The diluent is a clear, colorless to pale-yellow solution composed of carboxymethylcellulose sodium (19 mg), polysorbate 20 (0.63 mg), sodium phosphate monobasic monohydrate (0.61 mg), sodium phosphate dibasic heptahydrate (0.51 mg), sodium chloride (4.1 mg), and water for injection. Sodium hydroxide may be added during manufacture of BYDUREON Pen for pH adjustment.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Incretins, such as glucagon-like peptide-1 (GLP-1), enhance glucose-dependent insulin secretion and exhibit other antihyperglycemic actions following their release into the circulation from the gut. BYDUREON is a GLP-1 receptor agonist that enhances glucose-dependent insulin secretion by the pancreatic beta-cell, suppresses inappropriately elevated glucagon secretion, and slows gastric emptying.

The amino acid sequence of exenatide partially overlaps that of human GLP-1. Exenatide is a GLP-1 receptor agonist that has been shown to bind and activate the human GLP-1 receptor *in vitro*. This leads

to an increase in both glucose-dependent synthesis of insulin and *in vivo* secretion of insulin from pancreatic beta cells, by mechanisms involving cyclic AMP and/or other intracellular signaling pathways. Exenatide promotes insulin release from pancreatic beta-cells in the presence of elevated glucose concentrations.


12.2 Pharmacodynamics

Exenatide improves glycemic control through the actions described below.

Glucose-Dependent Insulin Secretion

The effect of exenatide infusion on glucose-dependent insulin secretion rates (ISR) was investigated in 11 healthy subjects. In these healthy subjects, on average, the ISR response was glucose-dependent (Figure 1). Exenatide did not impair the normal glucagon response to hypoglycemia.

Figure 1: Mean (SE) Insulin Secretion Rates During Infusion of Exenatide or Placebo by Treatment, Time, and Glycemic Condition in Healthy Subjects



SE = standard error.

Notes: 5 mmol = 90 mg/dL, 4 mmol/L = 72 mg/dL, 3.2 mmol/L = 58 mg/dL; Study medication infusion was started at time = 0 minutes.

Statistical assessments were for the last 30 minutes of each glycemic step, during which the target glucose concentrations were maintained.

*p < 0.05, exenatide treatment relative to placebo.

Glucagon Secretion

In patients with type 2 diabetes, exenatide moderates glucagon secretion and lowers serum glucagon concentrations during periods of hyperglycemia.

Gastric Emptying

Exenatide slows gastric emptying, thereby reducing the rate at which postprandial glucose appears in the circulation.

Fasting and Postprandial Glucose

In a clinical study in adults with type 2 diabetes mellitus, treatment with once weekly BYDUREON resulted in mean reductions in fasting plasma glucose of -45 mg/dL and 2-hour PPG concentrations of -95 mg/dL.

Cardiac Electrophysiology

The effect of exenatide at therapeutic (253 pg/mL) and suprathreshold (627 pg/mL) concentrations, following an intravenous infusion on QTc interval was evaluated in a randomized, placebo- and active-controlled (moxifloxacin 400 mg) three-period crossover thorough QT study in 74 healthy subjects. The upper bound of the one-sided 95% confidence interval for the largest placebo adjusted, baseline-corrected QTc based on population correction method (QTcP) was below 10 ms. Therefore, exenatide was not associated with prolongation of the QTc interval at therapeutic and suprathreshold concentrations.

12.3 Pharmacokinetics

Absorption

Following a single dose of BYDUREON, exenatide is released from the microspheres over approximately 10 weeks. There is an initial period of release of surface-bound exenatide followed by a gradual release of exenatide from the microspheres, which results in two subsequent peaks of exenatide in plasma at around Week 2 and Week 6 to 7, respectively, representing the hydration and erosion of the microspheres.

Following initiation of once every 7 days (weekly) administration of 2 mg BYDUREON, a gradual increase in the plasma exenatide concentration is observed over 6 to 7 weeks. After 6 to 7 weeks, mean exenatide concentrations of approximately 300 pg/mL were maintained over once every 7 days (weekly) dosing intervals indicating that steady state was achieved.

Distribution

The mean apparent volume of distribution of exenatide following subcutaneous administration of a single dose of BYETTA is 28.3 L and is expected to remain unchanged for BYDUREON.

Metabolism

Elimination

Nonclinical studies have shown that exenatide is predominantly eliminated by glomerular filtration with subsequent proteolytic degradation. The mean apparent clearance of exenatide in humans is 9.1 L/hour and is independent of the dose. Approximately 10 weeks after discontinuation of BYDUREON therapy, plasma exenatide concentrations generally fall below the minimal detectable concentration of 10 pg/mL.

Specific Populations

Patients with Renal Impairment

BYDUREON has not been studied in patients with severe renal impairment (creatinine clearance <30 mL/min) or end-stage renal disease receiving dialysis. Population pharmacokinetic analysis of renally impaired patients receiving 2 mg BYDUREON indicate that there is a 62% and 33% increase in exposure in moderate (N=10) and mild (N=56) renally impaired patients, respectively, as compared to patients with normal renal function (N=84).

In a study of BYETTA in subjects with end-stage renal disease receiving dialysis, mean exenatide exposure increased by 3.4-fold compared to that of subjects with normal renal function [*see Use in Specific Populations (8.6)*].

Patients with Hepatic Impairment

BYDUREON has not been studied in patients with acute or chronic hepatic impairment.

Age, Male and Female Patients, Race, and Body Weight

Age, gender, race and body weight did not alter the pharmacokinetics of BYDUREON in population pharmacokinetic analyses.

Pediatric Patients

The clinical pharmacology of BYDUREON has been evaluated in the population pharmacokinetic study in adolescent patients with type 2 diabetes mellitus between age 11 and age less than 18 years old. The pharmacokinetic profile of BYDUREON in the pediatric population was consistent with that of the adults.

Drug Interaction Studies

Acetaminophen

When 1000 mg acetaminophen tablets were administered, either with or without a meal, following 14 weeks of BYDUREON therapy (2 mg weekly), no significant changes in acetaminophen AUC were observed compared to the control period. Acetaminophen C_{max} decreased by 16% (fasting) and 5% (fed) and T_{max} was increased from approximately 1 hour in the control period to 1.4 hours (fasting) and 1.3 hours (fed).

The following drug interactions have been studied using BYETTA. The potential for drug-drug interaction with BYDUREON is expected to be similar to that of BYETTA.

Warfarin

Administration of warfarin (25 mg) 35 minutes after repeated doses of BYETTA (5 mcg twice daily on days 1-2 and 10 mcg twice daily on days 3-9) in healthy volunteers delayed warfarin T_{max} by approximately 2 hours. No clinically relevant effects on C_{max} or AUC of *S*- and *R*-enantiomers of warfarin were observed. BYETTA did not significantly alter the pharmacodynamic properties (e.g., international normalized ratio) of warfarin [*see Drug Interactions (7)*].

Digoxin

Administration of repeated doses of BYETTA 30 minutes before oral digoxin (0.25 mg once daily) decreased the C_{\max} of digoxin by 17% and delayed the T_{\max} of digoxin by approximately 2.5 hours; however, the overall steady-state pharmacokinetic exposure (e.g., AUC) of digoxin was not changed.

Lovastatin

Administration of BYETTA (10 mcg twice daily) 30 minutes before a single oral dose of lovastatin (40 mg) decreased the AUC and C_{\max} of lovastatin by approximately 40% and 28%, respectively, and delayed the T_{\max} by about 4 hours compared with lovastatin administered alone. In the 30-week controlled clinical trials of BYETTA, the use of BYETTA in patients already receiving HMG CoA reductase inhibitors was not associated with consistent changes in lipid profiles compared to baseline.

Lisinopril

In patients with mild to moderate hypertension stabilized on lisinopril (5-20 mg/day), BYETTA (10 mcg twice daily) did not alter steady-state C_{\max} or AUC of lisinopril. Lisinopril steady-state T_{\max} was delayed by 2 hours. There were no changes in 24-hour mean systolic and diastolic blood pressure.

Oral Contraceptives

The effect of BYETTA (10 mcg twice daily) on single and on multiple doses of a combination oral contraceptive (30 mcg ethinyl estradiol plus 150 mcg levonorgestrel) was studied in healthy female subjects. Repeated daily doses of the oral contraceptive (OC) given 30 minutes after BYETTA administration decreased the C_{\max} of ethinyl estradiol and levonorgestrel by 45% and 27%, respectively, and delayed the T_{\max} of ethinyl estradiol and levonorgestrel by 3.0 hours and 3.5 hours, respectively, as compared to the oral contraceptive administered alone. Administration of repeated daily doses of the OC one hour prior to BYETTA administration decreased the mean C_{\max} of ethinyl estradiol by 15%, but the mean C_{\max} of levonorgestrel was not significantly changed as compared to when the OC was given alone. BYETTA did not alter the mean trough concentrations of levonorgestrel after repeated daily dosing of the oral contraceptive for both regimens. However, the mean trough concentration of ethinyl estradiol was increased by 20% when the OC was administered 30 minutes after BYETTA administration injection as compared to when the OC was given alone. The effect of BYETTA on OC pharmacokinetics is confounded by the possible food effect on OC in this study [see [Drug Interactions \(7\)](#)].

12.6 Immunogenicity

The observed incidence of anti-drug antibodies is highly dependent on the sensitivity and specificity of the assay. Differences in assay methods preclude meaningful comparisons of the incidence of anti-drug antibodies in the studies described below with the incidence of anti-drug antibodies in other studies, including those of BYDUREON or of other exenatide products.

Immunogenicity in Adult Patients

Anti-exenatide antibodies were measured at prespecified intervals (4-14 weeks) in BYDUREON-treated adult patients (N=918) in five active-controlled trials for the improvement of glycemic control in adults with type 2 diabetes mellitus (Trials 1, 2, 3, 4 and 9) [see [Clinical Studies \(14.2, 14.3\)](#) and [Adverse Reactions \(6.1\)](#)]. In these five trials, 107/918 (12%) of BYDUREON-treated patients had high titer anti-

exenatide antibodies (high titer was defined as ≥ 625) at endpoint. These patients had a lower glycemic response compared to BYDUREON-treated patients who did not have anti-exenatide antibodies at endpoint [see *Warnings and Precautions (5.6)*].

In a 30-week, randomized, open-label, controlled trial in adults with type 2 diabetes mellitus [see *Adverse Reactions (6.1)*], anti-exenatide antibody assessments were performed at baseline and at 4-week intervals from Week 6 to Week 30. The mean anti-exenatide antibody titer in the BYDUREON-treated patients peaked at Week 6.

Injection site reactions for BYDUREON-treated patients were more commonly observed in those who were antibody-positive (anti-exenatide antibodies) compared with those who were antibody-negative, with a greater incidence in those with higher titer anti-exenatide antibodies (defined as ≥ 625) [see *Adverse Reactions (6.1)*]. There is insufficient data to assess whether the observed anti-exenatide antibodies in adults had an effect on the severity of injection site reactions, or on the incidence or severity of hypersensitivity reactions.

A total of 32 adult patients with anti-exenatide antibodies in BYDUREON Trial 9 were tested for the presence of cross-reactive antibodies to GLP-1 and/or glucagon and no treatment-emergent cross-reactive antibodies were observed.

Immunogenicity in Pediatric Patients

In a 52-week trial (24-week controlled and 28-week extended treatment period) for the improvement of glycemic control in pediatric patients with type 2 diabetes mellitus (Trial 8) [see *Clinical Studies (14.5)*], 37/58 (64%) of BYDUREON-treated patients had high titer anti-exenatide antibodies (high titer was defined as ≥ 625) at any time during the study. In this trial, 60% of these patients had a peak high-titers at approximately Week 12. BYDUREON-treated pediatric patients that had high titer anti-exenatide antibodies appeared to have had a lower glycemic response compared to BYDUREON-treated pediatric patients who did not have anti-exenatide antibodies during this trial [see *Warnings and Precautions (5.6)*].

There is insufficient data to assess whether the observed anti-exenatide antibodies in pediatric patients had an effect on the incidence or severity of hypersensitivity reactions or injection site reactions.

The potential for development of antibodies cross-reactive with endogenous GLP-1 and glucagon has not been evaluated in BYDUREON-treated pediatric patients.

Anti-Drug Antibody Effects on Pharmacokinetics in Adult and Pediatric Patients

The percentage of exenatide-treated adult and pediatric patients (age ≥ 11 years) with high anti-exenatide antibody titers (≥ 625) that had steady state exenatide plasma concentrations below limit of quantitation was higher than in those with low anti-exenatide antibody titers (< 625).

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Thyroid C-cell tumors have been observed in rats and mice with GLP-1 receptor agonists.

A 2-year carcinogenicity study was conducted with exenatide extended-release, the active component of BYDUREON, in male and female rats at doses of 0.3, 1.0, and 3.0 mg/kg (2-, 9-, and 26-times human systemic exposure at the maximum recommended human dose (MRHD) of 2 mg/week BYDUREON based on plasma exenatide AUC, respectively) administered by subcutaneous injection every other week. In this study there was an increased incidence of C-cell adenomas and C-cell carcinomas at all doses. An increase in benign fibromas was seen in the skin subcutis at injection sites of males given 3 mg/kg. No treatment-related injection-site fibrosarcomas were observed at any dose. The human relevance of these findings is currently unknown.

Carcinogenicity of exenatide extended-release has not been evaluated in mice.

Exenatide, the active ingredient in BYDUREON, was not mutagenic or clastogenic, with or without metabolic activation, in the Ames bacterial mutagenicity assay or chromosomal aberration assay in Chinese hamster ovary cells. Exenatide was negative in the *in vivo* mouse micronucleus assay.

In mouse fertility studies with exenatide, the active ingredient in BYDUREON, at twice-daily subcutaneous doses of 6, 68, or 760 mcg/kg/day, males were treated for 4 weeks prior to and throughout mating, and females were treated 2 weeks prior to mating and throughout mating until gestation day 7. No adverse effect on fertility was observed at 760 mcg/kg/day, a systemic exposure 148 times the human exposure resulting from the recommended dose of 2 mg/week, based on AUC.

14 CLINICAL STUDIES

14.1 Overview of Clinical Trials

BYDUREON has been studied in adult patients with type 2 diabetes mellitus as monotherapy and in combination with metformin, a sulfonylurea, a thiazolidinedione, a combination of metformin and a sulfonylurea, a combination of metformin and a thiazolidinedione, in combination with a SGLT2 inhibitor on a background of metformin, and in combination with basal insulin [see *Clinical Studies (14.2, 14.3)*].

Cardiovascular (CV) outcomes were studied in the EXSCEL trial that enrolled adult patients with type 2 diabetes mellitus and multiple risk factors for CV disease or established CV disease [see *Clinical Studies (14.4)*].

BYDUREON has been studied in pediatric patients aged 10 years and older with type 2 diabetes mellitus as monotherapy and in combination with metformin, a sulfonylurea, and in combination with basal insulin with or without metformin [see *Clinical Studies (14.5)*].

14.2 Glycemic Control Trials in Adults with Type 2 Diabetes Mellitus - Monotherapy

BYDUREON Monotherapy versus Metformin, Sitagliptin, and Pioglitazone

A 26-week, randomized, comparator-controlled trial was conducted to compare the safety and efficacy of BYDUREON to metformin, sitagliptin, and pioglitazone in adult patients with type 2 diabetes whose glycemic control was inadequate with diet and exercise (Trial 1 - NCT00676338). Patients were randomly assigned to receive BYDUREON 2 mg given subcutaneously once every seven days (weekly), titrated metformin from 1000 to 2500 mg/day, sitagliptin 100 mg/day or titrated pioglitazone from 30 to 45 mg/day, all dosed according to approved labeling.

A total of 820 patients were studied: 552 (67%) were Caucasian, 102 (12%) were East Asian, 71 (9%) were West Asian, 65 (8%) were Hispanic, 25 (3.0%) were Black, 4 (0.5%) were Native American, and 1 was classified otherwise. The mean baseline HbA_{1c} was 8.5%.

The primary endpoint was change in HbA_{1c} from baseline to Week 26 (or the last value at time of early discontinuation).

Efficacy Results:

Treatment with BYDUREON 2 mg once weekly (QW) resulted in mean HbA_{1c} reduction that was statistically significantly greater compared to sitagliptin 100 mg/day. The mean reduction in HbA_{1c} was non-inferior compared with metformin 1000-2500 mg/day (mean dose 2077 mg/day at study endpoint). Non-inferiority of BYDUREON 2 mg QW to pioglitazone 30-45 mg/day (mean dose 40 mg/day at study endpoint) in reducing HbA_{1c} after 26 weeks of treatment was not demonstrated (the mean change from baseline in HbA_{1c} after 26 weeks was -1.6% with BYDUREON and -1.7% with pioglitazone). The non-inferiority margin was set at +0.3% in this study. The results for the primary endpoint at 26 weeks are summarized in Table 5.

The proportion of patients with a Week 26 value achieving HbA_{1c} of less than 7% at Week 26 were 56%, 52%, 40%, and 55% for BYDUREON, metformin, sitagliptin, and pioglitazone, respectively. Patients who did achieve HbA_{1c} goal <7% and discontinued before Week 26 were not included as responders. The mean changes from baseline to Week 26 for fasting serum glucose were -41 mg/dL, -36 mg/dL, -20 mg/dL, and -46 mg/dL, and for body weight were -2.0 kg, -2.0 kg, -0.8 kg, and +1.5 kg for BYDUREON, metformin, sitagliptin, and pioglitazone, respectively.

Table 5: Results of 26-Week Trial of BYDUREON Monotherapy versus Metformin, Sitagliptin, and Pioglitazone in Patients with Type 2 Diabetes Mellitus (Trial 1)

	BYDUREON 2 mg QW	Metformin 1000-2500 (mean dose 2077) mg/day	Sitagliptin 100 mg/day	Pioglitazone 30-45 (mean dose 40) mg/day
Intent-to-Treat Population (N)	248	246	163	163
HbA_{1c} (%)				
Mean Baseline	8.4	8.6	8.4	8.5
Mean Change at Week 26*	-1.6	-1.5	-1.2	-1.7
Difference from metformin* [Bonferroni-adjusted 98.3% CI]	-0.05 [-0.26, 0.17]			
Difference from sitagliptin* [Bonferroni-adjusted 98.3% CI]	-0.39 [†] [-0.63, -0.16]			
Difference from pioglitazone* [Bonferroni-adjusted 98.3% CI]	0.16 [-0.08, 0.41]			

N = number of patients in each treatment group; mean change is least squares mean change; QW = once weekly. The primary efficacy analysis was adjusted for multiple comparisons and a two-sided 98.3% confidence interval was utilized to assess difference between treatments.

HbA_{1c} change data at 26 weeks were available from 86%, 87%, 85%, and 82% of the randomized subjects in the BYDUREON, metformin, sitagliptin, and pioglitazone groups, respectively.

* Least squares means were obtained using a mixed model repeated measure analysis with treatment, pooled country, visit, baseline HbA_{1c} value, and treatment by visit interaction as fixed effects, and subject as a random effect.

† p<0.001, treatment vs comparator.

14.3 Glycemic Control Trials in Adults with Type 2 Diabetes Mellitus - Combination Therapy

BYDUREON versus Sitagliptin and Pioglitazone, All as Add-on to Metformin Therapy

A 26-week double-blind comparator-controlled trial was conducted to compare the safety and efficacy of BYDUREON to sitagliptin and pioglitazone in patients with type 2 diabetes whose glycemic control was inadequate with metformin therapy (Trial 2 - NCT00637273). Patients were randomly assigned to receive BYDUREON 2 mg given subcutaneously once every 7 days (weekly), sitagliptin 100 mg/day or pioglitazone 45 mg/day, in addition to their existing metformin therapy.

A total of 491 patients were studied 168 (34.2%) were Caucasian, 143 (29.1%) were Hispanic, 119 (24.2%) were Asian, 52 (10.6%) were Black, 3 (0.6%) were Native American, and 6 (1.2%) were classified otherwise. The mean baseline HbA_{1c} was 8.5%.

The primary endpoint was change in HbA_{1c} from baseline to Week 26 (or the last value at time of early discontinuation).

Efficacy Results:

In this study, treatment with BYDUREON 2 mg QW resulted in a statistically significant mean HbA_{1c} reduction compared to sitagliptin 100 mg/day. There was a numerically greater reduction in HbA_{1c} with BYDUREON compared to pioglitazone, but there was not sufficient evidence to conclude superiority of BYDUREON 2 mg QW to pioglitazone 45 mg/day in reducing HbA_{1c} after 26 weeks of treatment. Results for the primary endpoint at 26 weeks are summarized in Table 6.

The proportion of patients with a Week 26 value achieving HbA_{1c} of less than 7% at Week 26 were 46%, 30%, and 39% for BYDUREON, sitagliptin, and pioglitazone, respectively. Patients who did achieve an HbA_{1c} goal <7% and discontinued before Week 26 were not included as responders. The mean changes from baseline to Week 26 for fasting serum glucose were -32 mg/dL, -16 mg/dL and -27 mg/dL, and for body weight were -2.3 kg, -0.8 kg and +2.8 kg for BYDUREON, sitagliptin, and pioglitazone, respectively.

Table 6: Results of 26-Week Trial of BYDUREON versus Sitagliptin and Pioglitazone, All as Add-On to Metformin Therapy in Patients with Type 2 Diabetes Mellitus (Trial 2)

	BYDUREON 2 mg QW	Sitagliptin 100 mg/day	Pioglitazone 45 mg/day
Intent-to-Treat Population (N)	160	166	165
HbA_{1c} (%)			
Mean Baseline	8.6	8.5	8.5
Mean Change at Week 26*	-1.5	-0.9	-1.2
Difference from sitagliptin* [95% CI]	-0.63 [-0.89, -0.37]		

Table 6: Results of 26-Week Trial of BYDUREON versus Sitagliptin and Pioglitazone, All as Add-on to Metformin Therapy in Patients with Type 2 Diabetes Mellitus (Trial 2)

	BYDUREON 2 mg QW	Sitagliptin 100 mg/day	Pioglitazone 45 mg/day
Difference from pioglitazone* [95% CI]	-0.32 [-0.57, -0.06]		

N = number of patients in each treatment group; mean change is least squares mean change; QW = once weekly.

* Least squares means were obtained using an ANCOVA model with treatment, baseline HbA_{1c} stratum, and country as fixed effects. Missing Week 26 data (28%, 18%, and 24% for the BYDUREON, sitagliptin, and pioglitazone groups, respectively) were imputed by the LOCF technique.

BYDUREON versus Insulin Glargine, Both as Add-on to Metformin or Metformin + Sulfonylurea Therapy

A 26-week open-label comparator-controlled trial was conducted to compare the safety and efficacy of BYDUREON to titrated insulin glargine in patients with type 2 diabetes whose glycemic control was inadequate with metformin or metformin plus sulfonylurea therapy (Trial 3 - NCT00641056). Patients were randomly assigned to receive BYDUREON 2 mg given subcutaneously once every 7 days (weekly) or insulin glargine once daily in addition to their existing oral antidiabetic therapy. Insulin glargine was dosed to a target fasting glucose concentration of 72 to 100 mg/dL.

The mean dose of insulin glargine was 10 units/day at baseline and 31 units/day at endpoint. At Week 26, 21% of insulin glargine treated patients were at fasting glucose goal.

A total of 456 patients were studied: 379 (83.1%) were Caucasian, 47 (10.3%) were Hispanic, 25 (5.5%) were East Asian, 3 (0.7%) were Black, and 2 (0.4%) were West Asian. Background therapy was either metformin (70%) or metformin plus sulfonylurea (30%). The mean baseline HbA_{1c} was 8.3%.

The primary endpoint was change in HbA_{1c} from baseline to Week 26 (or the last value at time of early discontinuation).

Efficacy Results:

Treatment with BYDUREON once weekly resulted in a mean reduction in HbA_{1c} from baseline at 26 weeks of -1.5%. The mean reduction in HbA_{1c} seen in insulin glargine arm at 26 weeks was -1.3%. The difference in observed effect size between BYDUREON and glargine in this trial excluded the pre-specified non-inferiority margin of +0.3%.

The proportion of patients with a Week 26 value achieving HbA_{1c} of less than 7% at Week 26 were 57% and 48% for BYDUREON and insulin glargine, respectively. Patients who did achieve an HbA_{1c} goal <7% and discontinued before Week 26 were not included as responders. The mean changes from baseline to Week 26 for fasting serum glucose in this study were -38 mg/dL and -50 mg/dL, and for body weight were -2.6 kg and +1.4 kg for BYDUREON and insulin glargine, respectively.

BYDUREON versus BYETTA, Both as Monotherapy or as Add-on to Metformin, a Sulfonylurea, a Thiazolidinedione, or Combination of Oral Agents

A 24-week, randomized, open-label trial was conducted to compare the safety and efficacy of BYDUREON to BYETTA in patients with type 2 diabetes and inadequate glycemic control with diet and

exercise alone or with oral antidiabetic therapy, including metformin, a sulfonylurea, a thiazolidinedione, or combination of two of those therapies (Trial 4 - NCT00877890). Patients were randomly assigned to receive BYDUREON 2 mg subcutaneously once every 7 days (weekly) or BYETTA (10 mcg subcutaneously twice daily), in addition to existing oral antidiabetic agents. Patients assigned to BYETTA initiated treatment with 5 mcg twice daily then increased the dose to 10 mcg twice daily after 4 weeks.

A total of 252 patients were studied: 149 (59%) were Caucasian, 78 (31%) Hispanic, 15 (6%) Black, and 10 (4%) Asian. Patients were treated with diet and exercise alone (19%), a single oral antidiabetic agent (47%), or combination therapy of oral antidiabetic agents (35%). The mean baseline HbA_{1c} was 8.4%.

The primary endpoint was change in HbA_{1c} from baseline to Week 24 (or the last value at time of early discontinuation).

Efficacy Results:

Treatment with BYDUREON 2 mg QW resulted in a statistically significantly greater mean HbA_{1c} reduction compared to BYETTA 10 mcg twice daily. Change in body weight was a secondary endpoint. Twenty-four-week study results are summarized in Table 7.

Reductions from mean baseline (97/94 kg) in body weight were observed in both BYDUREON (-2.3 kg) and BYETTA (-1.4 kg) treatment groups.

Table 7: Results of 24-Week Trial of BYDUREON versus BYETTA, Both as Monotherapy or as Add-On to Metformin, a Sulfonylurea, a Thiazolidinedione, or Combination of Oral Agents in Patients with Type 2 Diabetes Mellitus (Trial 4)

	BYDUREON 2 mg QW	BYETTA 10 mcg twice daily*
Intent-to-Treat Population (N)	129	123
HbA_{1c} (%)		
Mean Baseline	8.5	8.4
Mean Change at Week 24 [†]	-1.6	-0.9
Difference from BYETTA [†] [95% CI]	-0.7 [-0.9, -0.4] [‡]	
Percentage Achieving HbA_{1c} <7% at Week 24 (%)	58 [‡]	30
Fasting Plasma Glucose (mg/dL)		
Mean Baseline	173	168
Mean Change at Week 24	-25	-5
Difference from BYETTA [†] [95% CI]	-20 [-31, -10] [‡]	

N = number of patients in each treatment group; mean change is least squares mean change; QW = once weekly.

* BYETTA 5 mcg twice daily before the morning and evening meals for 4 weeks followed by 10 mcg twice daily for 20 weeks.

[†] Least squares (LS) means are adjusted for baseline HbA_{1c} strata, background antihyperglycemic therapy, and baseline value of the dependent variable (if applicable).

[‡] p<0.001, treatment vs comparator.

BYDUREON versus Liraglutide, Both as Add-on to Metformin, a Sulfonylurea, Metformin + Sulfonylurea, or Metformin + Pioglitazone Therapy

A 26-week open-label comparator-controlled trial was conducted to compare the safety and efficacy of BYDUREON to liraglutide in patients with type 2 diabetes whose glycemic control was inadequate with metformin, a sulfonylurea, metformin plus sulfonylurea, or metformin plus pioglitazone therapy (Trial 5 - NCT01029886). Patients were randomly assigned to receive BYDUREON 2 mg given subcutaneously once every 7 days (weekly) or liraglutide uptitrated from 0.6 mg/day to 1.2 mg/day, then 1.8 mg/day in addition to their existing oral antidiabetic therapy. Each titration was to be completed after at least one week, but could be delayed if the patient had severe nausea or vomiting as established by the investigator. Patients not tolerating the 1.8 mg/day dose of liraglutide by Week 4 were discontinued from the study.

A total of 911 patients were studied: 753 (82.7%) were Caucasian, 111 (12.2%) were Asian, 32 (3.5%) were American Indian or Alaska Native, 8 (0.9%) were Black, 6 (0.7%) were multiple races, and 1 (0.1%) was Pacific Islander. Background therapy was either a single oral antidiabetic agent (35%) or a combination of oral antidiabetic agents (65%). The mean baseline HbA_{1c} was 8.4%.

The primary endpoint was change in HbA_{1c} from baseline to Week 26 (or the last value at time of early discontinuation).

Efficacy Results:

Treatment with BYDUREON once weekly resulted in a mean reduction in HbA_{1c} from baseline at 26 weeks of -1.3%. The mean reduction in HbA_{1c} seen in the liraglutide arm at 26 weeks was -1.5%. The HbA_{1c} reduction with BYDUREON did not meet predefined non-inferiority criteria compared to liraglutide 1.8 mg/day. The non-inferiority margin was set at +0.25% in this study. Results for the primary endpoint at 26 weeks are summarized in Table 8.

The proportion of patients with a Week 26 value achieving HbA_{1c} of less than 7% at Week 26 were 48% and 56% for BYDUREON and liraglutide, respectively. Patients who did achieve an HbA_{1c} goal <7% and discontinued before Week 26 were not included as responders. The mean changes from baseline to Week 26 for fasting serum glucose were -32 mg/dL and -38 mg/dL, and for body weight were -2.7 kg and -3.6 kg for BYDUREON and liraglutide, respectively.

Table 8: Results of 26-Week Trial of BYDUREON versus Liraglutide, Both as Add-On to Metformin, a Sulfonylurea, Metformin + Sulfonylurea, or Metformin + Pioglitazone Therapy in Patients with Type 2 Diabetes Mellitus (Trial 5)

	BYDUREON 2 mg QW	Liraglutide 1.8 mg/day
Intent-to-Treat Population (N)	461	450
HbA_{1c} (%)		
Mean Baseline	8.5	8.4
Mean Change at Week 26*	-1.3	-1.5
Difference from liraglutide* [95% CI]	0.2 [0.08, 0.33]	

N = number of patients in each treatment group; mean change is least squares mean change; QW = once weekly. HbA_{1c} change data at 26 weeks were available from 85% and 86% of the randomized subjects in the BYDUREON and liraglutide groups, respectively.

* Least squares means were obtained using a mixed model repeated measure analysis with treatment, country, OAD stratum, baseline HbA_{1c} stratum, visit, baseline HbA_{1c}, and treatment by visit interaction as fixed effects, and subject

as a random effect.

BYDUREON in Combination with Dapagliflozin versus BYDUREON Alone and Dapagliflozin Alone, All as Add-On to Metformin

A 28-week double-blind comparator-controlled trial was conducted to compare the efficacy of BYDUREON and dapagliflozin (an SGLT2 inhibitor) to BYDUREON alone and dapagliflozin alone in patients with type 2 diabetes with inadequate glycemic control with metformin therapy (Trial 6 - NCT02229396). All patients entered a 1-week placebo lead-in period. Patients with HbA_{1c} \geq 8.0% and \leq 12% and on metformin at a dose of at least 1,500 mg per day were randomly assigned to receive either BYDUREON 2 mg given subcutaneously once every 7 days (weekly) plus dapagliflozin 10 mg once daily, BYDUREON 2 mg given subcutaneously once weekly, or dapagliflozin 10 mg once daily.

A total of 694 patients were studied; 580 (83.6%) were Caucasian, 96 (13.8%) were Black, 5 (0.7%) were Asian, 2 (0.3%) were American Indian or Alaska Native and 11 (1.6%) were classified otherwise. The mean baseline HbA_{1c} was 9.3%.

The primary endpoint was change in HbA_{1c} from baseline to Week 28.

Efficacy Results:

At Week 28, BYDUREON in combination with dapagliflozin provided statistically significantly greater reductions in HbA_{1c} (-1.77%) compared to BYDUREON alone (-1.42%, p=0.012) and dapagliflozin alone (-1.32%, p=0.001). BYDUREON in combination with dapagliflozin provided statistically significantly greater reductions in FPG (-57.35 mg/dL) compared to BYDUREON alone (-40.53, p <0.001) and dapagliflozin alone (-44.72 mg/dL, p=0.006).

BYDUREON versus Placebo, Both as Add-On to Basal Insulin or Basal Insulin + Metformin Therapy

A 28-week, double-blind, placebo-controlled trial was conducted to compare the safety and efficacy of BYDUREON to placebo when added to basal insulin glargine, with or without metformin, in patients with type 2 diabetes with inadequate glycemic control (Trial 7 - NCT02229383). Patients on sulfonylurea therapy discontinued sulfonylurea. Patients on metformin continued on the same dose of metformin. All patients initially entered an 8-week insulin dose-titration phase. Insulin glargine was to be titrated every 3 days with an aim of achieving a target fasting plasma glucose concentration of 72 to 99 mg/dL. Following the titration period, patients with HbA_{1c} \geq 7.0% and \leq 10.5% were then randomly assigned to receive either BYDUREON 2 mg given subcutaneously once every 7 days (weekly) or placebo once every 7 days (weekly).

A total of 460 patients were studied: 400 (87.0%) were White, 47 (10.2%) were Black or African American, 6 (1.3%) were Asian, 1 (0.2%) was American Indian or Alaska Native, 1 (0.2%) was Pacific Islander and 5 (1.1%) were classified otherwise.

The primary endpoint was the change in HbA_{1c} from baseline to Week 28.

Efficacy Results:

Compared to placebo, treatment with BYDUREON resulted in a statistically significant reduction in mean HbA_{1c} from baseline to Week 28 (Table 9).

The mean change in fasting plasma glucose from baseline to Week 28 was -12.50 mg/dL for BYDUREON and -2.26 mg/dL for placebo. The mean change from baseline to Week 28 in body weight was -0.92 kg for BYDUREON and +0.38 kg for placebo.

Table 9: Results of 28-Week Trial of BYDUREON versus Placebo, Both as Add-On to Insulin Glargine or Insulin Glargine + Metformin (Trial 7)

	BYDUREON 2 mg QW	Placebo QW
Intent-to-Treat Population (N)	231	229
Mean HbA_{1c} (%)		
Mean Baseline	8.53	8.53
Mean Change at Week 28*	-0.88 (0.070)	-0.24 (0.069)
Difference from Placebo [95% CI]	-0.64 [†] [-0.83, -0.45]	
Percentage Achieving HbA_{1c} <7.0% at Week 28 (%)[‡]	32.5 [†]	7.0

N = number of patients in each treatment group, CI=confidence interval, QW=once weekly.

Note: mean change is least squares mean change.

*Adjusted LS means and treatment group difference(s) in the change from baseline values at Week 28 using a multiple imputation method that models a “wash-out” for patients having missing data who discontinued treatment. ANCOVA was used with treatment, region, baseline HbA_{1c} stratum (<9.0% or ≥9.0%), and baseline SU-use stratum (yes vs. no) as fixed factors, and baseline value as a covariate.

[†] p-value <0.001 (adjusted for multiplicity).

[‡] Categories are derived from continuous measurements. All patients with missing endpoint data are imputed as non-responders. Treatment comparison is based on Cochran-Mantel-Haenszel (CMH) test stratified by baseline HbA_{1c} (<9.0% or ≥9.0%), and baseline SU-use stratum (yes vs. no). P-values are from the general association statistics.

Analyses include measurements post rescue therapy and post premature discontinuation of study medication.

14.4 EXSCEL Cardiovascular Outcomes Trial in Adult Patients with Type 2 Diabetes Mellitus

EXSCEL was a multinational, placebo-controlled, double-blind, randomized, parallel group pragmatic study that evaluated cardiovascular (CV) outcomes during treatment with BYDUREON in adult patients with type 2 diabetes and any level of CV risk when added to the current usual care (NCT01144338). A total of 14,752 adult patients were randomized 1:1 to either BYDUREON 2 mg given subcutaneously once weekly or placebo and followed as in routine clinical practice for a median of 38.7 months with a median treatment duration of 27.8 months.

Ninety six percent of the patients in both treatment groups completed the study in accordance with the protocol, and the vital status was known at the end of the study for 98.9% and 98.8% of the patients in the BYDUREON and placebo group, respectively. The mean age at study entry was 62 years (21 to 92 years with 8.5% of the patients ≥75 years). Approximately 62.0% of the patients were male, 75.8% were Caucasian, 9.8% were Asian, 6.0% were Black, and 20.5% were Hispanic or Latino. The mean BMI was 32.7 kg/m² and the mean duration of diabetes was 13.1 years. Approximately 49.3% had mild renal impairment (estimated glomerular filtration rate [eGFR] ≥60 to ≤89 mL/min/1.73 m²) and 21.6% had moderate renal impairment (eGFR ≥30 to ≤59 mL/min/1.73 m²).

The mean HbA_{1c} was 8.1%. At baseline, 1.5% of patients were not treated with either oral antidiabetic medications or insulin, 42.3% were treated with one oral antidiabetic medication and 42.4% were treated with two or more oral antidiabetic medications. Usage of oral antidiabetic medications included metformin (76.6%), sulfonylurea (36.6%), DPP-4 inhibitors (14.9%), thiazolidinediones (3.9%), and SGLT2 inhibitors (0.9%). Overall insulin usage was 46.3% (13.8% with insulin alone and 32.6% with insulin and one or more oral antidiabetic medications).

Overall, at baseline, 26.9% of patients did not have established cardiovascular (CV) disease, while 73.1% had established CV disease. The concomitant use of CV medications (e.g., ACE inhibitors, angiotensin receptor blockers, diuretics, beta blockers, calcium channel blockers, antithrombotic and anticoagulants, and lipid-lowering agents) was similar in the BYDUREON and placebo groups. At baseline, the mean systolic blood pressure was 135.5 mmHg, the mean diastolic blood pressure was 78.1 mmHg, the mean LDL was 95.0 mg/dL, and the mean HDL was 44.0 mg/dL.

The primary endpoint in EXSCEL was the time to first confirmed Major Adverse Cardiac Event (MACE) from randomization. MACE was defined as occurrence of either a cardiovascular (CV)-related death, or a nonfatal myocardial infarction (MI) or a nonfatal stroke. All-cause mortality, CV-related death, and fatal or nonfatal MI or stroke, hospitalization for acute coronary syndrome, and hospitalization for heart failure were also assessed as secondary endpoints.

A Cox proportional hazards model was used to test for non-inferiority against the pre-specified risk margin of 1.3 for the hazard ratio of MACE and superiority on MACE if non-inferiority was demonstrated. Type-1 error was controlled across multiple tests using a hierarchical testing strategy.

MACE Results:

BYDUREON did not increase the risk of MACE in adult patients with type 2 diabetes mellitus (HR: 0.91; 95% CI: 0.832, 1.004; P<0.001 for non-inferiority; P=0.06 for superiority). See results in Table 10 and Figure 2. The incidence of MACE in patients with and without established CV disease was 13.4% in the BYDUREON group versus 14.6% in the placebo group and 6.0% (BYDUREON) versus 5.9% (placebo), respectively. Five hundred and seven (507) patients (6.9%) died in the BYDUREON group versus 584 (7.9%) in the placebo group.


Table 10: Analysis of Primary Composite Endpoint MACE and Its Components in Adult Patients with Type 2 Diabetes (EXSCEL)

	BYDUREON N=7356	Placebo N=7396	HR* (95% CI)
MACE Composite of CV death, nonfatal MI or nonfatal stroke (time to first confirmed event)	839 (11.4%)	905 (12.2%)	0.91 (0.832, 1.004)
Cardiovascular Death	340 (4.6%)	383 (5.2%)	0.88 (0.76, 1.02)
Nonfatal Myocardial Infarction	466 (6.3%)	480 (6.5%)	0.96 (0.85, 1.09)
Nonfatal Stroke	169 (2.3%)	193 (2.6%)	0.86 (0.70, 1.06)

N=number of patients in each treatment group, HR=hazard ratio, CI=confidence interval, CV=cardiovascular, MI=myocardial infarction.

* HR (active/placebo) and CI are based on Cox proportional hazards regression model, stratified by established CV disease, with treatment group only as explanatory variable.

Figure 2: Time to First Adjudicated MACE in Adult Patients with Type 2 Diabetes (EXSCEL)



HR=hazard ratio, CI=confidence interval.

14.5 Glycemic Control Trial in Pediatric Patients 10 Years of Age and Older with Type 2 Diabetes Mellitus

The efficacy and safety of BYDUREON 2 mg given subcutaneously once weekly or placebo was evaluated in a 24-week randomized, double-blind, placebo-controlled, parallel-group study (Trial 8 - NCT01554618) in 82 patients aged 10 to 17 years with type 2 diabetes treated with diet and exercise alone or in combination with a stable dose of oral antidiabetic agents and/or insulin. This study included a 28-week open-label long-term extension in which patients in both treatment groups received BYDUREON 2 mg given subcutaneously once weekly after the 24-week controlled period.

In this trial, 88% of patients completed the assessment period. The mean age at study entry was 15.1 years (13% were 11 to 12 years old, 59% were 13 to 16 years old and 28% were older than 16 years).

Approximately 42% of the patients were male, 43% were White, 4% were Asian, 31% were Black or African American, 6% were American Indian or Alaska Native, and 44% were Hispanic or Latino. The mean body weight was 100.6 kg (67% were in ≥ 97 weight percentile group, 26% in ≥ 85 to < 97 and 6% in ≥ 3 to < 85), mean BMI was 36.4 kg/m² and the mean duration of diabetes was 2.3 years.

At baseline, mean HbA_{1c} was 8.17%, 12.2% of patients were not treated with either oral antidiabetic medications or insulin, 40.2% were treated with one oral antidiabetic medication and 1.2% were treated with two oral antidiabetic medications. 79.3% of patients were receiving metformin and 1.2% were receiving sulfonylurea. Overall insulin usage was 46.3% (8.5% with insulin alone and 37.8% with insulin and one other oral antidiabetic medication).

The primary endpoint of the study was the change in HbA_{1c} from baseline to Week 24.

Efficacy Results:

BYDUREON was superior to placebo in reducing HbA_{1c} (Table 11).

The mean changes from baseline to Week 24 for body weight were -0.17 kg with (SE 0.7) and +0.88 kg (SE 1.0) for BYDUREON and placebo, respectively.

Table 11: Results of 24-Week Trial of BYDUREON versus Placebo, Both alone or in Combination with Oral Antidiabetic Agents and/or Insulin in Pediatric Patients 10 Years of Age and Older with Type 2 Diabetes Mellitus (Trial 8)

	BYDUREON 2 mg QW	Placebo QW
Intent-to-Treat Population (N)	58	24
HbA_{1c} (%)		
Mean Baseline	8.13	8.28
Mean Change at Week 24 (SE)*	-0.25 (0.21)	0.45 (0.29)
Difference from Placebo (95% CI)	-0.71† (-1.42, 0)	
HbA_{1c}<7%		
Percentage achieved goal at Week 24	31.0	8.3
Fasting Plasma Glucose (mg/dL)		
Mean Baseline	165.2	170.5
Mean Change at Week 24 (SE)‡	-1.3 (8.1)	16.2 (11.9)

* Adjusted LS means and treatment group difference(s) in the change from baseline values at Week 24 using a multiple imputation method that models a “wash-out” for patients having missing data who discontinued treatment. ANCOVA was used with treatment, region as fixed factors, and baseline HbA_{1c} value as a covariate. SE – standard error.

† p<0.05

‡ Adjusted LS means in the change from baseline values at Week 24 using a multiple imputation method that models a “wash-out” for patients having missing data who discontinued treatment. ANCOVA was used with treatment, region, as fixed factors, and baseline FPG value as a covariate.

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

BYDUREON (exenatide) for extended-release injectable suspension is supplied as:

- BYDUREON Pen supplied in cartons that contain four single-dose pens and one spare needle (NDC 0310-6530-04). Each single-dose pen contains:
 - One single-dose pen containing 2 mg of exenatide (as a white to off-white powder) and 0.65 mL of diluent.
 - One custom needle (23G, 9/32") specific to this delivery system.

Do **not** substitute needles or any other components provided with BYDUREON.

Storage and Handling

- Store BYDUREON in the refrigerator at 36°F to 46°F (2°C to 8°C). BYDUREON can be kept at room temperature not to exceed 77°F (25°C) [see USP Controlled Room Temperature] for no more than a total of 4 weeks, if needed.
- Do not freeze BYDUREON. Do not use BYDUREON if it has been frozen.
- Protect from light.
- Administer immediately after the supplied BYDUREON is reconstituted with the supplied diluent [see *Dosage and Administration (2.2)*].

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide and Instructions for Use).

Thyroid C-cell Tumors

Inform patients that exenatide extended-release causes benign and malignant thyroid C-cell tumors in rats and that the human relevance of this finding has not been determined. Counsel patients to report symptoms of thyroid tumors (e.g., a lump in the neck, hoarseness, dysphagia, or dyspnea) to their physician [see *Boxed Warning and Warnings and Precautions (5.1)*].

Acute Pancreatitis

Inform patients treated with BYDUREON of the potential risk for pancreatitis. Explain that persistent severe abdominal pain that may radiate to the back, and which may or may not be accompanied by vomiting, is the hallmark symptom of acute pancreatitis. Instruct patients to discontinue BYDUREON promptly and contact their healthcare provider if persistent severe abdominal pain occurs [see *Warnings and Precautions (5.2)*].

Hypoglycemia with Concomitant Use of Insulin Secretagogues or Insulin

Inform patients that the risk of hypoglycemia is increased when BYDUREON is used in combination with an agent that induces hypoglycemia, such as a sulfonylurea or insulin. Educate patients on the signs and symptoms of hypoglycemia [see *Warnings and Precautions (5.3)*].

Acute Kidney Injury

Inform patients treated with BYDUREON of the potential risk for worsening kidney function and explain the associated signs and symptoms of renal impairment, as well as the possibility of dialysis as a medical intervention if renal failure occurs [see *Warnings and Precautions (5.4)*].

Hypersensitivity Reactions

Inform patients that serious hypersensitivity reactions have been reported during postmarketing use of exenatide. Inform patients that if symptoms of hypersensitivity reactions occur, stop taking BYDUREON and seek medical advice promptly [see *Warnings and Precautions (5.7)*].

Drug-Induced Thrombocytopenia

Inform patients that drug-induced immune-mediated thrombocytopenia has been reported during use of exenatide. Inform patients that if symptoms of thrombocytopenia occur, e.g. bleeding, stop taking BYDUREON and seek medical advice promptly [see *Warnings and Precautions (5.8)*].

Serious Injection-Site Reactions

Inform patients that there have been postmarketing reports of serious injection-site reactions with or without subcutaneous nodules, with the use of BYDUREON. Isolated cases of injection-site reactions required surgical intervention. Advise patients to seek medical advice if symptomatic nodules occur, or for any signs or symptoms of abscess, cellulitis, or necrosis [see *Warnings and Precautions (5.9)*].

Acute Gallbladder Disease

Inform patients of the potential risk for cholelithiasis or cholecystitis. Instruct patients to contact their physician if cholelithiasis or cholecystitis is suspected for appropriate clinical follow-up [see *Warnings and Precautions (5.10)*].

Instructions

Train patients on how to use BYDUREON properly prior to administration. Instruct patients and caregivers on proper mixing and injection technique to ensure the product is adequately mixed and a full dose is delivered. Instruct caregivers to assist pediatric patients with mixing and administration. Refer patients and caregivers to the accompanying Instructions for Use for complete administration instructions with illustrations [see *Dosage and Administration (2)*].

If a patient is currently taking BYETTA, it should be discontinued upon starting BYDUREON. Inform patients formerly on BYETTA who start BYDUREON that they may experience transient elevations in blood glucose concentrations, which generally improve within the first 2 weeks after initiation of therapy [see *Dosage and Administration (2.1)* and *Clinical Studies (14.2)*].

Manufactured for:

AstraZeneca Pharmaceuticals LP
Wilmington, DE 19850

By:

Amylin Ohio LLC
West Chester, OH 45071

BYDUREON is a registered trademark of the AstraZeneca group of companies.

MEDICATION GUIDE
BYDUREON® (by-DUR-ee-on)
(exenatide) for extended-release injectable suspension
for subcutaneous use

What is the most important information I should know about BYDUREON?

BYDUREON may cause serious side effects, including:

- **Possible thyroid tumors, including cancer.** Tell your healthcare provider if you get a lump or swelling in your neck, hoarseness, trouble swallowing, or shortness of breath. These may be symptoms of thyroid cancer. In studies with rats or mice, BYDUREON and medicines that work like BYDUREON caused thyroid tumors, including thyroid cancer. It is not known if BYDUREON will cause thyroid tumors or a type of thyroid cancer called medullary thyroid carcinoma (MTC) in people.
- Do not use BYDUREON if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC), or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).

What is BYDUREON?

- BYDUREON is an injectable prescription medicine that may improve blood sugar (glucose) in adults and children who are 10 years of age and older with type 2 diabetes mellitus and should be used along with diet and exercise.
- BYDUREON is not recommended as the first choice of medicine for treating diabetes.
- BYDUREON is not for use in people with type 1 diabetes.
- BYDUREON and BYDUREON BCise are long-acting forms of the medicine in BYETTA (exenatide). BYDUREON should not be used at the same time as BYETTA or BYDUREON BCise.
- It is not known if BYDUREON can be used in people who have had pancreatitis.
- It is not known if BYDUREON is safe and effective for use in children younger than 10 years of age.

Who should not use BYDUREON?

Do not use BYDUREON if:

- you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC) or if you have an endocrine system condition called Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).
- you have had a severe allergic reaction to exenatide or any of the ingredients in BYDUREON. See the end of this Medication Guide for a complete list of ingredients in BYDUREON.
Symptoms of a severe allergic reaction with BYDUREON may include:
 - swelling of your face, lips, tongue, or throat
 - fainting or feeling dizzy
 - problems breathing or swallowing
 - very rapid heartbeat
 - severe rash or itching
- you have a history of low blood platelet count from using exenatide medicines (drug-induced thrombocytopenia).

What should I tell my healthcare provider before using BYDUREON?

Before using BYDUREON, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had problems with your pancreas or kidneys.
- have severe problems with your stomach, such as slowed emptying of your stomach (gastroparesis) or problems with digesting food.
- are pregnant or plan to become pregnant. BYDUREON may harm your unborn baby. Tell your healthcare provider if you become pregnant while using BYDUREON. Talk to your healthcare provider about the best way to control your blood sugar if you plan to become pregnant or while you are pregnant.
- are breastfeeding or plan to breastfeed. It is not known if BYDUREON passes into your breast milk. You should talk with your healthcare provider about the best way to feed your baby while using BYDUREON.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. BYDUREON may affect the way some medicines work and some medicines may affect the way BYDUREON works.

Before using BYDUREON, talk to your healthcare provider about low blood sugar and how to manage it.

Especially tell your healthcare provider if you take:

- other medicines to treat diabetes, including insulin or sulfonylureas.
- a water pill (diuretic).
- a blood pressure medicine.
- warfarin.
- a pain medicine.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I use BYDUREON?

- Read the **Instructions for Use** that comes with BYDUREON.

- Use BYDUREON exactly as your healthcare provider tells you to.
- BYDUREON should be injected right away after you prepare your dose.
- **Your healthcare provider should show you how to use BYDUREON before you use it for the first time.**
- Caregivers should help children with mixing and injecting BYDUREON.
- BYDUREON is injected under the skin (subcutaneously) of your stomach (abdomen), thigh, or upper arm. **Do not** inject BYDUREON into a muscle (intramuscularly) or vein (intravenously).
- **Use BYDUREON 1 time each week on the same day each week at any time of the day.**
- BYDUREON may be taken with or without food.
- If you miss a dose of BYDUREON, take the missed dose as soon as possible if there are at least 3 days (72 hours) until your next scheduled dose. If there are less than 3 days remaining, skip the missed dose and take your next dose on the regularly scheduled day. **Do not** take 2 doses of BYDUREON within 3 days of each other.
- You may change the day of the week as long as your last dose was given 3 or more days before.
- **Do not** mix insulin and BYDUREON together in the same injection.
- You may give an injection of BYDUREON and insulin in the same body area (such as, your stomach area), but not right next to each other.
- Change (rotate) your injection site with each weekly injection. **Do not** use the same site for each injection.

What are the possible side effects of BYDUREON?

BYDUREON may cause serious side effects, including:

- See "[What is the most important information I should know about BYDUREON?](#)"
- **inflammation of your pancreas (pancreatitis).** Stop using BYDUREON and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back.
- **low blood sugar (hypoglycemia).** Your risk for getting low blood sugar may be higher if you use BYDUREON with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin. **Signs and symptoms of low blood sugar may include:**
 - dizziness or light-headedness
 - sweating
 - confusion or drowsiness
 - headache
 - blurred vision
 - slurred speech
 - shakiness
 - fast heartbeat
 - anxiety, irritability, or mood changes
 - hunger
 - weakness
 - feeling jittery
- **kidney problems.** In people who have kidney problems, nausea, vomiting and diarrhea, may cause a loss of fluids (dehydration) which may cause kidney problems to get worse. These kidney problems include kidney failure. Dialysis or a kidney transplant may be needed.
 - **While taking BYDUREON:** Call your healthcare provider right away if you have nausea, vomiting, or diarrhea that does not go away.
- **stomach problems.** Other medicines like BYDUREON may cause severe stomach problems. It is not known if BYDUREON causes or worsens stomach problems.
- **serious allergic reactions.** Stop using BYDUREON and get medical help right away if you have any symptoms of a serious allergic reaction, including itching, rash, or difficulty breathing. See "[Who should not use BYDUREON?](#)"
- **low blood platelet count (drug-induced thrombocytopenia).** BYDUREON may cause the number of platelets in your blood to be reduced. When your platelet count is too low, your body cannot form blood clots. You could have serious bleeding that could lead to death. **Stop using BYDUREON and call your healthcare provider right away if you have unusual bleeding or bruising.** Your blood platelet count may continue to be low for about 10 weeks after stopping BYDUREON.
- **serious injection-site reactions.** Serious injection-site reactions, with or without bumps (nodules), have happened in some people who use BYDUREON. Some of these injection-site reactions have required surgery. Call your healthcare provider if you have any symptoms of an injection-site reaction, including severe pain, swelling, blisters, an open wound, a dark scab.
- **gallbladder problems.** Gallbladder problems have happened in some people who take BYDUREON. Tell your healthcare provider right away if you get symptoms of gallbladder problems, which may include:
 - pain in your upper stomach (abdomen)
 - fever
 - yellowing of skin or eyes (jaundice)
 - clay-colored stools

The most common side effects of BYDUREON may include nausea, diarrhea, headache, vomiting, constipation, itching at the injection site, a small bump (nodule) at the injection site, indigestion.

Nausea is most common when you first start using BYDUREON but decreases over time in most people as their body gets used to the medicine.

Talk to your healthcare provider about any side effect that bothers you or does not go away.

These are not all the possible side effects of BYDUREON. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Keep BYDUREON and all medicines out of the reach of children.

General information about the safe and effective use of BYDUREON.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use BYDUREON for a condition for which it was not prescribed. Do not give your BYDUREON to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about BYDUREON that is written for health professionals.

What are the ingredients in BYDUREON?

Contents of the powder:

Active ingredient: exenatide.

Inactive ingredients: polylactide-co-glycolide and sucrose.

Contents of liquid (diluent):

Inactive Ingredients: carboxymethylcellulose sodium, polysorbate 20, sodium phosphate monobasic monohydrate, sodium phosphate dibasic heptahydrate, sodium chloride, water for injection. Sodium hydroxide may be added during manufacture of BYDUREON Pen for pH adjustment.

BYDUREON is a registered trademark and BYETTA is a registered trademark of the AstraZeneca group of companies. All other marks are the marks of their respective owners.

Manufactured for:

AstraZeneca Pharmaceuticals LP
Wilmington, DE 19850

By:

Amylin Ohio LLC
West Chester, OH 45071

For more information about BYDUREON, go to www.BYDUREON.com or call 1-800-236-9933.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: December 2022

INSTRUCTIONS FOR USE
BYDUREON® (by-DUR-ee-on) Pen
(exenatide) for extended-release injectable suspension
for subcutaneous use
once-weekly




Figure A

Before using Bydureon Pen, your healthcare provider should show you how to use it the right way.

Caregivers should help children with mixing and injecting BYDUREON.

Read the Instructions for Use before you start using Bydureon Pen and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

Getting ready

Bydureon Pen is not for self-injection by people who are blind or cannot see well.

Supplies needed to give your Bydureon Pen injection (not all supplies are included):

- 1 Bydureon single-use "Pen" tray that contains:
 - 1 Bydureon Pen
 - 1 custom needle
- a clean flat surface
- alcohol swab
- sharps container for throwing away used needles and Pens. See "**Properly dispose of your Pen**" at the end of these instructions.



How should I store Bydureon?

- Store your Bydureon Pens in the refrigerator at 36°F to 46°F (2°C to 8°C).
- Protect pens from light until you are ready to prepare and use your dose.
- **Do not** use pens past the expiration date.
- **Do not** freeze Bydureon. **Do not** use Bydureon if it has been frozen.
- Keep Bydureon in its sealed tray until ready for use.
- If needed, you can keep your Bydureon Pen out of the refrigerator at 68°F to 77°F (20°C to 25°C) for up to 4 weeks.

Keep Bydureon Pen, and all medicines, out of the reach of children.

Step 1: Prepare your Bydureon Pen

Let your Pen come to room temperature.


- Remove 1 Pen from the refrigerator and let it stand at room temperature for at least 15 minutes.
- **Do not** use a pen past its expiration date.



Wash your hands.


Open the tray.

- Pull up on the corner tab.
- Remove the Pen and needle.
 - **Do not** use your Pen or needle if any parts are broken or missing.



Check the liquid in your Pen.

- **Check the liquid** inside the inspection window. It should be clear and free of particles. **Do not** use the Pen if the liquid is colored, has particles, or is not clear. Throw it away and get a new one.
 - You may see bubbles in the liquid, this is normal.



Peel off the paper tab from the needle cover.




Attach the needle to the Pen.

- Screw the needle onto the Pen by pushing and twisting clockwise until it is tight. **Do not** remove the needle cover yet.



Step 2: Mix your dose
Combine the medicine.

- While holding the pen straight up, **slowly** turn the knob. **Stop** when you hear the click and the green label disappears.



Firmly tap your Pen to mix.

- Hold your Pen by the end with the orange label and **tap the Pen firmly against the palm of your hand**.
 - **DO NOT** twist the white knob.
 - **ROTATE** your Pen every 10 taps.
 - **You may need to tap your Pen 80 times or more.**



Check the Bydureon mix.

- **Hold your Pen up to the light and look through both sides of the mixing window.** The solution should have **no clumps** and be uniformly cloudy (**see Figure B**).




Figure B

- **To get your full dose, Bydureon must be mixed well.**
- **If Bydureon is not mixed well, keep tapping your Pen longer and more firmly until it is mixed well.**
- **Do not give your Bydureon injection unless your Bydureon is mixed well.**

Stop. Do Not proceed unless your medicine is mixed well.

To get your full dose the medicine must be mixed well. If it is not mixed well, tap longer and more firmly.

Check the Bydureon mix again.

- **Compare both sides of the mixing window to the photos below** by holding your Pen against the page. Pay attention to the **bottom surface**. If you **do not see clumps** you are ready to inject (**see Figure C**).




Figure C


If you have any questions or are not sure if your Bydureon is mixed well, call 1-800-236-9933 for help.

Step 3: Inject your dose

Important: After the medicine is mixed well, you must inject your dose right away. You cannot save it for later use.


Choose your injection site.

- The recommended injection sites for Bydureon are your stomach (abdomen), thigh, or back of the arm.
 - Each week you can use the same area of your body but choose a different injection site in that area.
 - Gently wipe the site you choose with an alcohol swab (not included).




Twist knob to release injection button.

- Hold your **Pen with the needle pointing straight up** and turn the white knob until the orange label disappears and the injection button is released. **Do not** push the injection button yet.



Remove the needle cover.

- Pull the needle cover straight off. **Do not** twist the needle cover.
 - You may see a few drops of liquid on the needle or in the cover.



Inject your Bydureon.

- Insert the needle into your skin.
- Press the injection button with your thumb until you hear a “click”. Keep holding **the button down and slowly count to 10 to get your full dose.**



Properly dispose of your used Pen.

- Put your used needles and Pens in an FDA-cleared sharps disposal container right away after use. **Do not** throw away (dispose of) loose needles and Pens in your household trash. See **Common Questions and Answers** for additional disposal information.



Common Questions and Answers:**1. How do I know that the Bydureon is mixed well?**

The Bydureon is mixed well when the liquid looks cloudy from both sides of the window. You should not see any clumps in the liquid. It may help to hold your Pen up to the light to see in the window. If you see clumps of any size keep tapping your Pen firmly against the palm of your hand until mixed.

2. I am having trouble mixing my dose. What should I do?

Remember, before preparing your dose, leave your Pen out of the refrigerator for at least 15 minutes. This will let your Pen warm up to room temperature. It will be easier to mix Bydureon if your Pen is at room temperature.

Be sure you are holding your Pen at the end with the knob and the orange label. This will help you grip your Pen better and tap it more firmly against your palm.

It may also help to tap the mixing window on both sides against your palm. If you see any clumps, keep tapping.

3. After I mix Bydureon, how long can I wait before taking the injection?

You must inject your dose of Bydureon right after mixing it. If you do not inject your Bydureon right away, small clumps of medicine may form in your Pen and you may not get your full dose.

4. I am ready to inject my dose. What should I do if I see air bubbles in the Pen?

It is normal for air bubbles to be in your Pen. Bydureon is injected into your skin (subcutaneously). Air bubbles will not harm you or affect your dose with this type of injection.

5. What should I do if I cannot push the injection button all the way in when trying to inject my dose?

Check that you have fully screwed on the pen needle. Also be sure you twisted the knob until it stopped, the orange label disappeared, and the injection button appears.

If you still cannot push the button in, this may mean that the needle is clogged. Remove the needle from your skin and replace it with the spare needle from the carton. Review how to attach the needle. Then choose a different injection site and finish taking the injection.

If you still cannot push the button all the way in, remove the needle from your skin. Use a puncture-resistant container to throw away the pen with the needle still attached.

If you have problems giving your Bydureon Pen injection or have any questions call 1-800-236-9933 for more instructions.

6. How do I know if I injected my full dose?

To be sure you get your full dose, press the injection button with your thumb until you hear a "click". After the "click", continue to hold the needle in your skin for 10 seconds. This will allow enough time for you to get your full dose.

7. What if I do not have an FDA-cleared sharps disposal container?

Do not throw away (dispose of) loose needles and Pens in your household trash.

- If you do not have an FDA-cleared sharps disposal container, you may use a household container that is:
 - made of a heavy-duty plastic,
 - can be closed with a tight-fitting, puncture-resistant lid, without sharps being able to come out,
 - upright and stable during use,
 - leak-resistant, and

