



Biochar reduces yield-scaled emissions of reactive nitrogen gases from vegetable soils across China

Changhua Fan, Hao Chen, Bo Li, and Zhengqin Xiong

Jiangsu Key Laboratory of Low Carbon Agriculture and GHGs Mitigation, College of Resources and Environmental Sciences, Nanjing Agricultural University, Nanjing 210095, China

Correspondence to: Zhengqin Xiong (zqxiong@njau.edu.cn)

Received: 10 November 2016 – Discussion started: 15 December 2016

Revised: 14 April 2017 – Accepted: 1 May 2017 – Published: 13 June 2017

Abstract. Biochar amendment to soil has been proposed as a strategy for sequestering carbon, mitigating climate change and enhancing crop productivity. However, few studies have compared the general effect of different feedstock-derived biochars on the various gaseous reactive nitrogen emissions (GNrEs) of N_2O , NO and NH_3 simultaneously across the typical vegetable soils in China. A greenhouse pot experiment with five consecutive vegetable crops was conducted to investigate the effects of two contrasting biochars, namely wheat straw biochar (B_w) and swine manure biochar (B_m) on GNrEs, vegetable yield and gaseous reactive nitrogen intensity (GNrI) in four typical soils which are representative of the intensive vegetable cropping systems across mainland China: an Acrisol from Hunan Province, an Anthrosol from Shanxi Province, a Cambisol from Shandong Province and a Phaeozem from Heilongjiang Province. Results showed that remarkable GNrE mitigation induced by biochar occurred in Anthrosol and Phaeozem, whereas enhancement of yield occurred in Cambisol and Phaeozem. Additionally, both biochars decreased GNrI through reducing N_2O and NO emissions by 36.4–59.1 and 37.0–49.5 % for B_w (except for Cambisol), respectively, and by improving yield by 13.5–30.5 % for B_m (except for Acrisol and Anthrosol). Biochar amendments generally stimulated the NH_3 emissions with greater enhancement from B_m than B_w . We can infer that the biochar's effects on the GNrEs and vegetable yield strongly depend on the attributes of the soil and biochar. Therefore, in order to achieve the maximum benefits under intensive greenhouse vegetable agriculture, both soil type and biochar characteristics should be seriously considered before conducting large-scale biochar applications.

1 Introduction

Agriculture accounts for an estimated emission of 4.1 Tg N yr^{-1} for nitrous oxide (N_2O) and 3.7 Tg N yr^{-1} for nitric oxide (NO), contributing 60 and 10 %, respectively, to the total global anthropogenic emissions, largely due to increases of nitrogen (N) fertilizer application in cropland (Ciais, 2013). The concentration of atmospheric N_2O , a powerful, long-lived, greenhouse gas, has increased from 270 parts per billion by volume (ppbv) in the preindustrial era to ~ 324 ppbv (Ussiri and Lal, 2013); it has 265 times the global warming potential of carbon dioxide (CO_2) on a 100-year horizon (IPCC, 2013) and also causes depletion of the ozone layer in the atmosphere (Ravishankara et al., 2009). In contrast, NO_x , which is mainly emitted as NO, does not directly affect the earth's radiative balance but catalyzes the production of tropospheric ozone (O_3), which is a greenhouse gas associated with detrimental effects on human health (Anenberg et al., 2012) and crop production (Avnery et al., 2011). Finally, ammonia (NH_3) volatilization is one of the major N loss pathways (Harrison and Webb, 2001) as well, with up to 90 % coming from agricultural activities (Misselbrook et al., 2000; Boyer et al., 2002). As a natural component and a dominant atmospheric alkaline gas, NH_3 plays an important role in atmospheric chemistry and ambient aerosol formation (Langridge et al., 2012; Wang et al., 2015b). In addition to nutrient enrichment (eutrophication) of terrestrial and aquatic systems and global acidification of precipitation, NH_3 has also been shown to be a major factor in the formation of atmospheric particulate matter and secondary aerosols (Kim et al., 2006; Pinder et al., 2007), leading to potentially adverse effects on human and ecosystem health such as visibility degradation and

threats to biodiversity (Powelson et al., 2008; Behera et al., 2013).

In China, vegetable production devotes an area of approximately 24.7×10^6 ha, equivalent to 12.4 % of the total available cropping area, and the production represented 52 % of the world vegetable production in 2012 (FAO, 2015). Intensified vegetable cultivation in China is characterized by high N application rates, high cropping index and frequent farm practices. Annual N fertilizer inputs for intensively managed vegetable cultivation are 3–6 times higher than in cereal grain cultivation in China (Ju et al., 2006; Diao et al., 2013; Wang et al., 2015a). As a result, great concern exists about excess N fertilizer application and low N use efficiency in intensive vegetable fields in China (Deng et al., 2013; Diao et al., 2013; Li et al., 2017). Meanwhile, intensive vegetable agriculture is considered to be an important source of N_2O (Xiong et al., 2006; Jia et al., 2012; Li et al., 2015b; Zhang et al., 2015) and NO production (Mei et al., 2009). Moreover, NH_3 volatilization is another important N pathway in fertilized soil, resulting in large losses of soil–plant N (Pacholski et al., 2008; Zhang et al., 2011). Therefore, the reduction of reactive N loss is key to meet the joint challenges of high production and acceptable environmental consequences from intensive vegetable production (Zhang et al., 2013).

Biochar is the dark-colored, carbon (C)-rich residue of pyrolysis or gasification of plant biomass under oxygen (O_2)-limited conditions, specifically produced for use as a soil amendment (Sohi, 2012). The amendment of agricultural ecosystems with biochar has been proposed as an effective countermeasure for climate change (Smith, 2016). These additions have been suggested to increase soil carbon storage (Mukherjee and Zimmerman, 2013; Stavi and Lal, 2013), decrease greenhouse gas emissions (Li et al., 2017), and improve soil fertility and crop production (Major et al., 2010; Liu et al., 2013; Jeffery et al., 2017). However, some recent studies have reported no difference or even an increase in soil N_2O emissions induced by biochar application for various soils (Saarnio et al., 2013; Wang et al., 2015a). Furthermore, NH_3 volatilization was enhanced by biochar application in pasture soil (Clough et al., 2010), vegetable soil (Sun et al., 2014) and paddy soil in the wheat-growing season (Zhao et al., 2014). Additionally, crop productivity responses to biochar amendments differed among various biochars (Cayuela et al., 2014). These inconsistent results suggest that current biochar application to soil is not a “one-size-fits-all paradigm” because of the variation in the physical and chemical characteristics of the different biochars, soil types and crop species (Field et al., 2013; Cayuela et al., 2014). Moreover, limited types of biochar (Spokas and Reicosky, 2009) and soil (Sun et al., 2014) were involved in the experiments in previous studies. Thus, the evaluation of the different types of biochar under the typical soils is imperative to gain a comprehensive understanding of potential interactions before the large-scale application of biochars.

Therefore, a greenhouse pot experiment was conducted in an effort to investigate the effects of different types of biochar on gaseous reactive nitrogen emissions (GNrEs), namely N_2O , NO and NH_3 , simultaneously in four intensively cropped vegetable soils across main vegetable production areas of mainland China. We hypothesized that (1) biochar amendment could affect GNrEs, vegetable yield and yield-scaled gaseous reactive nitrogen emissions, namely gaseous reactive nitrogen intensity (GNrI) in vegetable soils across mainland China and that (2) those influences would vary among biochar and soil types. Overall, the objectives of this research were to gain a comprehensive insight into the effects of two contrasting biochars on the GNrEs, vegetable yield and GNrI in intensively managed vegetable production in China.

2 Materials and methods

2.1 Experimental soil and biochar

Four typical greenhouse vegetable cultivation sites with a long history (more than 10 years) of conventional cultivation were selected from Northeast, Northwest, Central and East China (Fig. S1): (1) a Phaeozem from Jiamusi ($46^{\circ}48'N$, $130^{\circ}12'E$) in Heilongjiang Province, (2) an Anthrosol from Yangling ($34^{\circ}18'N$, $108^{\circ}2'E$) in Shanxi Province, (3) an Acrisol from Changsha ($28^{\circ}32'N$, $113^{\circ}23'E$) in Hunan Province and (4) a Cambisol from Shouguang ($36^{\circ}56'N$, $118^{\circ}38'E$) in Shandong Province (FAO and ISRIC, 2012). Those four types of vegetable soil represented a range of differences in physicochemical properties and regions (Table S1 in the Supplement). Soil samples were manually collected from the cultivated layer (0–20 cm) after the local vegetable harvest in April 2015. The samples were air-dried and passed through a 5 mm stainless steel mesh sieve and homogenized thoroughly. Any visible roots and organic residues were removed manually before being packed with the necessary amount of soil to achieve the initial field bulk density. Each pot received 15 kg of $105^{\circ}C$ dry-weight-equivalent fresh soil. For each of the biochar amendment pot, 282.6 g pot^{-1} sieved biochar (2 mm) was mixed with the soil thoroughly before the experiment, which was equivalent to a 40 t ha^{-1} biochar dose (dry weight). No more biochar was added later in the experimental period.

The two types of biochar that were used in this experiment are derived from two common agricultural wastes in China: wheat straw and swine manure, hereafter referred to as B_w and B_m , respectively (Table S1). The B_w was produced at the Sanli New Energy Company in Henan, China, by pyrolysis and thermal decomposition at 400 – $500^{\circ}C$. The B_m was produced through thermal decomposition at $400^{\circ}C$ by the State Key Laboratory of Soil Science and Sustainable Agricultural, Institute of Soil Science, Chinese Academy of Sciences. In accordance with Lu (2000), soil organic car-

bon (SOC) was measured by wet digestion with H_2SO_4 – $\text{K}_2\text{Cr}_2\text{O}_7$, total nitrogen (TN) was determined by semi-micro Kjeldahl digestion, and soil texture was determined with the pipette method. The soil pH and biochar pH were measured in deionized water at a volume ratio of 1 : 2.5 (soil to water) with a PHS-3C mv pH^{-1} detector (Shanghai Kangyi Inc., China). Biochar content of hydrogen (H) was measured by elemental analysis after dry combustion (Euro EA, Hekatech GmbH, Wegberg, Germany). The oxygen content of biochar was measured with the same device after pyrolysis of the sample at 1000°C followed by reduction of the evolved O_2 to CO and quantified by GC-TCD. The soil nitrate (NO_3^- -N) and ammonium (NH_4^+ -N) were measured following the two-wavelength ultraviolet spectrometry and indophenol blue method, respectively, using an ultraviolet spectrophotometer (Hitachi, UV-2900, Tokyo, Japan). Electric conductivity (EC) was measured by using a Mettler Toledo instrument (FE30-K, Shanghai, China) at a 1 : 5 ($w : v$) soil to water ratio. Cation exchange capacity (CEC) was determined using the $\text{CH}_3\text{COONH}_4$ method. Dissolved organic carbon (DOC) was extracted from 5 g of the biochar/soil with an addition of 50 mL deionized water and measured by a TOC analyzer (TOC-2000/3000, Metash Instruments Co., Ltd., Shanghai, China). Ash content was measured by heating the biochars at 750°C for 4 h. The specific surface area of the biochar material was tested using the Brunauer–Emmett–Teller (BET) method, from which the N adsorption–desorption isotherms at 77 K were measured by an automated gas adsorption analyzer ASAP2000 (Micromeritics, Norcross, GA) with +5 % accuracy. Scanning electron microscopy (SEM) imaging analysis was conducted using a HITACHI S-3000N scanning electron microscope.

2.2 Experimental setup and management

The pot experiments were performed at the greenhouse experimental station of Nanjing Agricultural University, China. Five vegetable crops were grown successively in the four vegetable soils during the experimental period. For each type of soil, three treatments with three replicates were arranged in a random design: urea without biochar (N), and urea with wheat straw biochar ($N + B_w$), urea with swine manure biochar ($N + B_m$). In addition, phosphate and potassium fertilizers in the form of calcium magnesium phosphate and potassium chloride, together with urea, were broadcasted and mixed with soil thoroughly prior to sowing the vegetables. No topdressing events occurred because of the frequent cultivation and short growth period for the leafy vegetables. Based on the vegetable growth, all pots received equal amounts of water and no precipitation. Detailed information on the pot management practices is provided in Table S2.

Each pot consists of a 30 cm \times 30 cm (height \times diameter) cylinder made of polyvinyl chloride (PVC). The top of each pot was surrounded by a special water-filled trough collar, which allowed a chamber to sit on the pot and prevent gas

exchange during the gas-sampling period. Small holes (diameter of 1 cm) at the bottom of the pots were designed for drainage. To prevent soil loss, a fine nylon mesh (<0.5 mm) was attached to the base of the soil cores before packing.

2.3 Measurement of N_2O , NO and NH_3

The NO and N_2O fluxes were measured simultaneously from each vegetable cultivation using a static opaque chamber method (Zheng et al., 2008; Yao et al., 2009). A square PVC chamber of 35 cm \times 35 cm \times 40 cm (length \times width \times height) was temporarily mounted on the pot for gas flux measurement. The chamber was coated with sponge and aluminum foil outside to prevent solar radiation heating the chamber. Gas samples for flux measurements were collected between 08:00 and 10:00 on each measuring day to minimize the influence of diurnal temperature variation. Gas fluxes were usually measured once a week and every other day for 1 week following fertilizer application. To measure the N_2O flux, four samples were collected from the headspace chamber using 20 mL polypropylene syringes at 0, 10, 20 and 30 min after chamber closure. The gas concentrations in the samples were analyzed within 12 h after sampling using an Agilent 7890A gas chromatograph equipped with an electron capture detector (ECD) for N_2O detection. Argon–methane (5 %) was used as the carrier gas at a flow rate of 40 mL min^{-1} . The column and ECD temperatures were maintained at 40 and 300°C , respectively. The gas chromatography configurations described by Wang et al. (2013) were adopted for the gas concentration analysis. N_2O flux was calculated using the linear increases in gas concentration with time. Sample sets were rejected unless they yielded a linear regression value of $R^2 > 0.90$.

For each NO flux measurement, gas samples were collected from the same chamber that was used for the N_2O flux measurements (Yao et al., 2009). Before closing the chamber, an approximately 1.0 L gas sample from the headspace of each chamber was extracted into an evacuated sampling bag (Delin Gas Packing Co., Ltd., Dalian, China), and this measurement was regarded as time 0 min for NO analysis. After 30 min under chamber enclosure conditions (i.e., after the N_2O sample collections were completed), another headspace gas sample with the same volume was extracted from each chamber into another evacuated bag. Within 1 h after sampling, NO concentrations were analyzed by a model 42i chemiluminescence NO–NO– NO_x analyzer (Thermo Environmental Instruments Inc., Franklin, MA, USA). The NO fluxes were derived from the concentration differences between the two collected samples. The NO_x analyzer was calibrated by a model 146i dynamic dilution calibrator system at the end of each crop-growing season.

The mean flux of N_2O or NO during the experiment period is the average of all measured fluxes weighted by the interval between two neighboring measurements (Xiong et al., 2006).

The cumulative N_2O flux was calculated as the product of the mean flux and the entire duration.

The NH_3 volatilization was determined using the ventilation method (Zhao et al., 2010). The phosphoglycerol-soaked sponge was replaced every day after each fertilization event for approximately 1 week. The phosphoglycerol-soaked sponges used to collect the NH_3 samples were immediately extracted with 300 mL potassium chloride (KCl) solution (1 mol L^{-1}) for 1 h. The concentration of NH_4^+-N was measured using the indophenol blue method at 625 nm (Sororzano, 1969) by ultraviolet spectrophotometry (Hitachi, UV-2900, Tokyo, Japan, with 0.005 absorbance of photometric accuracy). The cumulative seasonal NH_3 volatilization was the sum of the daily emissions during the measurement period.

Cumulative fluxes of N_2O , NO and NH_3 were added to calculate total gaseous reactive nitrogen gas emissions (GNrEs):

$$\text{GNrEs} = \text{cumulative } N_2O + \text{cumulative NO} + \text{cumulative } NH_3 \text{ emissions (kg N ha}^{-1}\text{)}. \quad (1)$$

2.4 Auxiliary measurements

Simultaneously with the determination of trace gas fluxes, the air temperature and the soil temperature at a depth of 5 cm were measured using thermally sensitive probes at each sampling date. Soil water content was also measured using a portable water detector (model TZS-1K, Zhejiang Top Instrument Corporation Ltd., China) by the frequency domain reflectometer method at a depth of 5 cm. Measured soil water contents (v/v) were converted to water-filled pore space (WFPS) with the following equation:

$$\text{WFPS} = \frac{\text{volumetric water content (cm}^3 \text{ cm}^{-3}\text{)}}{\text{total soil porosity (cm}^3 \text{ cm}^{-3}\text{)}}. \quad (2)$$

Here, total soil porosity = $[1 - (\text{soil bulk density (g cm}^{-3}\text{)} / 2.65)]$ with an assumed soil particle density of $2.65 \text{ (g cm}^{-3}\text{)}$. The total soil bulk density was determined with the cutting ring method according to Lu (2000).

After each vegetable crop reached physiological maturity, the fresh vegetable yield was measured by weighing the whole aboveground and belowground biomass in each pot. Gaseous reactive nitrogen intensity (GNrI) was calculated using the following equation:

$$\text{GNrI} = \text{GNrEs} / \text{vegetable fresh yield (kg N t}^{-1} \text{ yield)}. \quad (3)$$

After the 1-year pot experiment, a soil sample from each pot was blended carefully. One subsample was stored at 4°C for determination of microbial biomass carbon (MBC), potential nitrification rate (PNR) and denitrification enzyme activity (DEA) within 3 days. Another subsample was air-dried for analysis of SOC, TN, pH and EC. MBC was determined by substrate-induced respiration using gas chromatog-

raphy (Anderson and Domsch 1978). PNR was measured using the chlorate inhibition soil-slurry method as previously described (Kurolo et al., 2005) with modifications (Hu et al., 2016). DEA was quantified as described by Smith and Tiedje (1979).

2.5 Data processing and statistics

Two-way ANOVA was used to analyze the effects of the biochar type, soil type and their interactions on soil properties; N_2O , NO and NH_3 emissions; vegetable yield; GNrEs; and GNrI throughout the experimental period. Multiple comparisons among the treatments were assessed using Tukey's HSD test. Significant differences were considered at $P < 0.05$. All statistical analyses were performed using JMP ver. 7.0 (SAS Institute, Cary, NC, USA, 2007). Pearson's correlation analysis was used to determine whether there were significant interrelationships between N_2O / NO and PNR or DEA in each soil, using SPSS for Windows, version 18.0 (SPSS Inc., Chicago, USA).

3 Results

3.1 Soil responses to biochar amendment

Appreciable differences in all observed soil properties existed among soil types (Table 1), reflecting the wide variations in soil characteristics across mainland China. Additionally, biochar amendments had significant influences on all the soil properties (Table 1, $p < 0.05$). Compared with N treatments, biochar amendments increased the SOC, TN and EC by 20.4–135.0, 0.5–21.2 and 2.4–38.1%, respectively, across all the soils. Compared with B_w , B_m amendment increased SOC and TN by 5.8–20.5 and 9.5–14.2% ($p < 0.05$), respectively, whereas EC values were higher by 3.3–21.5% induced by B_w than B_m amendment over all soils. Additionally, biochar amendments significantly increased soil pH by 0.27–0.64 and 0.08–0.10 units compared with N treatment in Acrisol and Anthrosol soils ($p < 0.05$), respectively, and B_m performed better than B_w on increasing soil pH in Acrisol. Furthermore, biochar amendments tended to increase MBC in Cambisol and Phaeozem, and B_m increased MBC relative to B_w in all soils.

As shown in Fig. 1, no consistent effects on PNR and DEA were observed with biochar amendments across all soils. Compared with N treatment, biochar amendments significantly increased PNR in Phaeozem but had no effect on Cambisol (Fig. 1a). Compared with B_w , B_m amendment significantly increased PNR in Acrisol and Anthrosol. Moreover, compared with N, biochar amendments reduced DEA in most soils, significantly in Anthrosol and Phaeozem by an average of 40.1 and 37.8% (Fig. 1b, $p < 0.05$), respectively. In comparison with B_w , enhancements in DEA were observed by 42.5 and 74.4% with B_m amendment in Acrisol and Anthrosol, respectively ($p < 0.05$).

Table 1. Soil organic carbon (SOC), soil total nitrogen (TN), soil pH, electric conductivity (EC) and microbial biomass carbon (MBC) as affected by different treatments across the four vegetable soils.

| Soil | Treatment | SOC (g kg ⁻¹) | TN (g kg ⁻¹) | pH | EC (ds m ⁻¹) | MBC (mg kg ⁻¹) |
|----------------|---------------------------------|---------------------------|--------------------------|--------------|--------------------------|----------------------------|
| Acrisol | <i>N</i> | 8.0 ± 0.8c | 1.37 ± 0.12b | 4.37 ± 0.04c | 1.76 ± 0.21b | 1353 ± 119a |
| | <i>N</i> + <i>B_w</i> | 15.6 ± 0.5b | 1.47 ± 0.07b | 4.64 ± 0.04b | 2.43 ± 0.31a | 1173 ± 49b |
| | <i>N</i> + <i>B_m</i> | 18.8 ± 0.6a | 1.64 ± 0.04a | 5.01 ± 0.03a | 2.00 ± 0.32ab | 1234 ± 50ab |
| Anthrosol | <i>N</i> | 9.7 ± 0.7c | 1.55 ± 0.04b | 7.53 ± 0.02b | 1.74 ± 0.27b | 490 ± 9a |
| | <i>N</i> + <i>B_w</i> | 15.6 ± 0.8b | 1.62 ± 0.06b | 7.61 ± 0.05a | 2.25 ± 0.22a | 495 ± 16a |
| | <i>N</i> + <i>B_m</i> | 17.5 ± 1.1a | 1.79 ± 0.03a | 7.63 ± 0.01a | 1.96 ± 0.06ab | 504 ± 18a |
| Cambisol | <i>N</i> | 7.9 ± 0.1b | 1.13 ± 0.04b | 7.70 ± 0.08a | 0.85 ± 0.03b | 535 ± 13b |
| | <i>N</i> + <i>B_w</i> | 14.2 ± 0.6a | 1.20 ± 0.04b | 7.66 ± 0.03a | 0.92 ± 0.04a | 554 ± 10ab |
| | <i>N</i> + <i>B_m</i> | 15.5 ± 1.4a | 1.37 ± 0.06a | 7.71 ± 0.03a | 0.87 ± 0.02ab | 573 ± 12a |
| Phaeozem | <i>N</i> | 29.9 ± 0.5b | 2.19 ± 0.04b | 6.91 ± 0.05a | 0.83 ± 0.03b | 921 ± 44b |
| | <i>N</i> + <i>B_w</i> | 36.0 ± 1.5a | 2.20 ± 0.03b | 6.92 ± 0.06a | 0.95 ± 0.03a | 988 ± 56b |
| | <i>N</i> + <i>B_m</i> | 38.1 ± 1.8a | 2.41 ± 0.01a | 6.94 ± 0.04a | 0.92 ± 0.06a | 1242 ± 196a |
| ANOVA results | | | | | | |
| Biochar | *** | *** | *** | *** | * | |
| Soil | *** | *** | *** | *** | *** | |
| Biochar × soil | * | n.s. | *** | n.s. | ** | |

Data shown are means ± SD of three replicates. See Fig. 1 for treatment codes. Different letters within the same column indicate significant differences among treatments within the same soil at $p < 0.05$ level. *** Significant at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; n.s.: not significant.

3.2 Seasonal variations in N₂O and NO emissions

The dynamics of N₂O fluxes from all N-applied treatments in the four vegetable soils were relatively consistent and followed a sporadic and pulse-like pattern that was accompanied with fertilization, tillage and irrigation (Fig. 2). In addition, peak N₂O fluxes varied greatly. Most of the N₂O emissions occurred during the amaranth and tung choy growing periods, and there were several small emission peaks during the spinach and coriander herb growing periods due to lower N application rate (Table S2), soil temperature and water content (Fig. S2). The highest peaks of N₂O emissions from Acrisol, Anthrosol, Cambisol and Phaeozem were 4133.7, 1784.0, 432.4 and 1777.2 μg N m⁻² h⁻¹, respectively. Although biochar (*B_w* and *B_m*) application did not significantly alter the seasonal pattern of the N₂O fluxes, they greatly lowered some peaks of N₂O emissions in the Anthrosol and Phaeozem by 8.7–74.4 and 23.6–73.6 %, respectively (Fig. 2b and d).

Clearly, the NO fluxes demonstrated similar seasonal dynamics to the N₂O fluxes (Fig. 3). Some relatively high peak NO fluxes were still observed in the spinach and coriander herb planting seasons even though relatively low temperatures occurred during these periods. The NO fluxes ranged from −44.6 to 377.6 μg N m⁻² h⁻¹ across all soil types. Furthermore, some NO peaks were significantly weakened with the *B_w* and *B_m* in the Acrisol (Fig. 3a).

3.3 Cumulative N₂O, NO and NH₃ emissions

Cumulative N₂O emissions varied greatly among soil types (Table 3a, $p < 0.05$), from 1.97 to 31.56 kg N ha⁻¹. Biochar




Figure 1. Potential nitrification rate (PNR) and denitrification enzyme activity (DEA) under different treatments in Acrisol, Anthrosol, Cambisol and Phaeozem. The three treatments with each soil were urea without biochar (*N*), urea with wheat straw biochar (*N* + *B_w*) and urea with swine manure biochar (*N* + *B_m*). Bars indicate standard deviation (mean + SD, $n = 3$). Different letters above the bars indicate significant differences among the different treatments within the same soil, at $p < 0.05$.

amendments had significant influences on the cumulative




Figure 2. Temporal dynamics of soil N_2O ($\mu\text{g N m}^{-2} \text{h}^{-1} \pm \text{SD}$, $n = 3$) fluxes under different treatments in Acrisol (a), Anthrosol (b), Cambisol (c) and Phaeozem (d) with five consecutive vegetable crops. The inserted panels describe the N_2O fluxes during the last two cropping seasons. The solid arrows indicate fertilization. See Fig. 1 for treatment codes.

N_2O emissions (Table 2, $p < 0.001$). In comparison with the N treatment, the effect of biochar amendment on N_2O emissions differed between soil types (Table 3a), indicating significant interactions between biochar and soil types (Table 2, $p < 0.001$). Additionally, B_w amendment decreased N_2O emissions by 11.8–38.4% across all the soils compared to B_m , indicating that B_w performed better mitigation effects than B_m across all the soils, significantly in Acrisol (Table 3a, $p < 0.05$). The values of cumulative NO emissions were much smaller than those of N_2O emissions, with a remarkable variation in 0.20–8.99 kg N ha^{-1} across all soils (Table 3b). Biochar amendments had pronounced effects on NO emissions (Table 2, $p < 0.001$), but their effects differed between vegetable soils (Table 3b), which suggested significant interactions between biochar and soil types (Table 2, $p < 0.001$). Compared with B_m , B_w amendment significantly reduced NO emissions in Anthrosol and Phaeozem (Table 3b, $p < 0.05$). Moreover, N_2O emissions had positive relationships with DEA both in Anthrosol and Phaeozem, and were affected positively with PNR in Acrisol (Table 4).

Additionally, NO emissions had positive correlations with both PNR and DEA in Anthrosol. However, neither N_2O nor NO emissions were influenced significantly by PNR and DEA in Cambisol.

As is shown in Table 3c, the cumulative NH_3 emissions fluctuated greatly from 4.72 to 7.57 kg N ha^{-1} across all the soils. Biochar amendments produced no significant influences on the NH_3 emissions relative to N treatment in most soils (Table 3c). A tendency was found for the cumulative NH_3 emissions in B_m to be higher than those in the B_w treatment, although this difference was not remarkable within each soil. Additionally, stimulation effects were consistently present after the first fertilization event in each type of soil (Fig. 4).

3.4 Vegetable yield and gaseous reactive N intensity during the five-vegetable crop rotation

The vegetable yields for the five consecutive vegetable crops are presented in Table 3e. Pronounced differences existed




Figure 3. Temporal dynamics of soil NO ($\mu\text{g N m}^{-2} \text{h}^{-1} \pm \text{SD}$, $n = 3$) fluxes under different treatments in Acrisol (a), Anthrosol (b), Cambisol (c) and Phaeozem (d) with five consecutive vegetable crops. The solid arrows indicate fertilization. See Fig. 1 for treatment codes.

among all soils (Table 3e, $p < 0.05$). Additionally, biochar amendments exerted no significant effects on vegetable yield (Table 2). Compared with the N treatment, biochar amendments were prone to increase vegetable yield in Cambisol and Phaeozem against Acrisol and Anthrosol (Table 3e), denoting pronounced interactions between soil and biochar (Table 2, $p < 0.05$). Compared with B_m , B_w amendment lowered total yield over all the soils (Table 3e), significantly in Acrisol and Cambisol ($p < 0.05$).

Table 3f presents the GNrI during the whole experiment period, with a pronounced variation among soils ($p < 0.05$). The GNrI was greatly affected by biochar amendment during the whole experiment period (Table 2, $p < 0.01$). Compared to N treatment, biochar amendments reduced the GNrI by 4.3–27.8 % across all soils, significantly in Anthrosol and Phaeozem (Table 3f, $p < 0.05$). Moreover, there were no remarkable differences between B_w and B_m throughout all soils.

4 Discussion

4.1 Biochar effects on GNrEs across different soil types

The effects of biochar amendment on the N_2O and NO emissions may be positive, negative or neutral, largely depending on the soil condition and the inherent characteristics of the biochar (Spokas and Reicosky, 2009; Nelissen et al., 2014). In our study, effects of two biochars on the N_2O and NO emissions did not show a consistent trend across the four typical vegetable soils (Table 3a, b). In agreement with Cayuela et al. (2014), who reported that the role of biochar in mitigating N_2O emission was maximal in soils close to pH neutral, remarkable mitigation effects were observed in Anthrosol and Phaeozem with the biochar amendments (Table 3a). These findings potentially resulted from the effects of the biochars on soil aeration, C/N ratio and pH, which affected the N dynamics and N cycling processes (Zhang et al., 2010; Ameloot et al., 2015). In line with Obia et al. (2015), biochar decreased NO emissions in low-pH Acrisol (Table 3b), probably by stimulating denitrification enzyme activity, which then resulted in less NO accumula-




Figure 4. Cumulative ammonia (NH₃) emissions from the Acrisol (a), Anthrosol (b), Cambisol (c) and Phaeozem (d) during the four nitrogen fertilization events *F*: every N fertilization event. The bars indicate the standard deviation of the mean (kg N ha⁻¹ ± SD, *n* = 3) of each treatment for the sum of the four *N* fertilization events. See Fig. 1 for treatment codes. Different letters above the bars indicate significant differences among the different treatments for each soil, at *p* < 0.05.

tion relative to N₂ production. Moreover, the liming effects of biochar may have prevented the chemical decomposition of NO₂⁻ to NO (Islam et al., 2008), leaving only enzymatically produced NO to accumulate.

Different from the other soils in our experiment, neither N₂O nor NO emissions from the Cambisol were significantly influenced by PNR or DEA. This finding suggests that processes other than nitrification and denitrification might play vital roles. Besides nitrification and denitrification, nitrifiers denitrification (Wrage et al., 2001) and heterotrophic nitrification (Zhu et al., 2011) can be important processes for producing N₂O/NO as well, especially in vegetable soils with low pH, low carbon content and high N content (Wrage et al., 2001). Ma et al. (2015) speculated that nitrifier denitrification was the main process producing N₂O in the North China Plain (Cambisol within this region). In addition, surplus N input in vegetable systems probably masked the beneficial effects of the biochar addition on the N transformation (Wang et al., 2015a). Therefore, future research needs to study the underlying mechanism of how biochar affects those processes.

Different biochars may not produce universal influences on N₂O emissions for the same soil due to the distinct properties of the biochar (Spokas and Reicosky, 2009). In the current study, overall, in comparison with B_m, the B_w amendment had more effective mitigation effects on N₂O and NO emissions (Table 3a, b), largely due to the following reasons. First, compared to B_w, the contents of TN and DOC were

80 and 40% higher in B_m (Table S1), respectively, which might supply extra N or C source for heterotrophic nitrification in the acidic Acrisol, leading B_m to being ineffective for reducing the N₂O emissions (Table 3a). This result was in accordance with Li et al. (2015a), who observed that biochar amendment had no significant influence on the cumulative N₂O emissions, and even higher N₂O emissions occurred with biochar addition. Additionally, as shown in Fig. 1, B_m was more prone to stimulate PNR and DEA, thus displaying lower mitigation ability than B_w. Second, compared with B_m, the C/N ratio was approximately 2-fold higher in B_w (Table S1), presumably leading to more inorganic nitrogen being immobilized in biochar with a higher C/N ratio (Ameloot et al., 2015), decreasing the available N for microorganisms. Last, as presented in Fig. S3 and Table S1, B_w had more pores and surface area, having a better advantage over B_m in absorbing NO accordingly. Others have found that the lower mitigation capacity of high-N biochars (e.g., manures or biosolids) is probably due to the increased N release in the soil from the biochar (Schouten et al., 2012). To our knowledge, very few studies have investigated biochar effects on NO emissions (Nelissen et al., 2014; Obia et al., 2015), and the mechanisms through which biochar influence NO emissions have not yet been elucidated. Therefore, more research is needed to clarify the underlying mechanisms of biochar on NO emission.

Intensively managed soils receiving fertilizer such as urea or anhydrous NH₃ and ruminant urine patches are potential

Table 2. Two-way ANOVA for the effects of biochar (B_c) and soil (S) types on cumulative N_2O , NO and NH_3 emissions, gaseous reactive nitrogen emissions (GNrEs), vegetable yield and gaseous reactive nitrogen intensity (GNrI) during the entire sampling period.

| Factors | DF | N ₂ O emission | | | NO emission | | | NH ₃ emission | | | GNrEs | | | Vegetable yield | | | GNrI | | |
|----------------|----|---------------------------|-------|-----|-------------|-------|-----|--------------------------|-----|------|--------|-------|-----|-----------------|-------|------|------|-------|------|
| | | SS | F | P | SS | F | P | SS | F | P | SS | F | P | SS | F | P | SS | F | P |
| B_c | 2 | 271.9 | 65.1 | *** | 46.4 | 174.7 | *** | 0.5 | 0.8 | n.s. | 380.5 | 86.4 | *** | 76.2 | 3.2 | n.s. | 0.1 | 7.9 | ** |
| S | 3 | 1429.9 | 228.1 | *** | 152.2 | 382.1 | *** | 4.1 | 3.8 | * | 2322.6 | 351.5 | *** | 4316.9 | 123.3 | *** | 2.3 | 110.3 | *** |
| $B_c \times S$ | 6 | 179.3 | 14.3 | *** | 33.4 | 41.9 | *** | 1.4 | 0.7 | n.s. | 234.5 | 17.7 | *** | 230.4 | 3.3 | * | 0.1 | 1.6 | n.s. |
| Model | 11 | 4009.7 | 174.5 | *** | 225.3 | 154.3 | *** | 29.1 | 7.5 | *** | 5290 | 218.3 | *** | 15962.0 | 124.4 | *** | 5.8 | 77.0 | *** |
| Error | 24 | 50.1 | | | 3.2 | | | 8.5 | | | 52.9 | | | 280.0 | | | 0.2 | | |

SS: the sum of squares, F value: the ratio of mean squares of two independent samples. P value: the index of differences between the control group and the experimental group. *, ** and *** indicate significance at $p < 0.05$, $p < 0.01$ and $p < 0.001$, respectively. n.s.: not significant.

hot spots for NH_3 formation, and the use of biochar is expected to retain NH_3-N under these conditions (Clough and Condron, 2010). Our results show that the effects of biochar amendments on NH_3 volatilization largely depend on soil characteristics and biochar types. Soil texture is an important factor impacting NH_3 transfer and release. High clay contents in the Anthrosol (Table S1) likely limited soil porosity; thus, the addition of porous biochar could enhance the soil aeration, promoting NH_3 volatilization (Sun et al., 2014). Additionally, it is worth noting that cumulative NH_3 emissions were slightly higher in soils with the B_m than those with the B_w amendment (Fig. 4 and Table 3c) and that difference could presumably be attributed to less surface area and the much higher pH of B_m (Fig. S3 and Table S1), resulting in weak adsorption and great liming effects.

4.2 Biochar effects on vegetable yield and GNrI across different soil types

The application of biochar is usually intended to increase crop yields, and evidence suggests that this may be successful (Schulz et al., 2013; Li et al., 2017; Jeffery et al., 2017). Due to its liming effect, biochar helps to improve the supply of essential macro- and micronutrients for plant growth (Chan and Xu, 2009; Major et al., 2010). Enhancement of vegetable yield with biochar amendment occurred in Cambisol and Phaeozem (Table 3e). Additionally, the effects of B_m and B_w on vegetable yield differed, which was probably due to large differences in physicochemical characteristics between the two biochars (Jeffery et al., 2017). First, compared to B_w , B_m has a higher DOC content (Table S1), through which more nutrients may be directly introduced to the soil (Rajkovich et al., 2012). Secondly, besides their large amount of plant-available nutrients (Hass et al., 2012), biochars produced with manure have been generally considered significant for improving soil fertility by promoting soil structure development (Joseph et al., 2010), with the result that B_m was found superior to B_w in vegetable production enhancement in our case (Table 3e). As biochar effects on vegetable yield were variable, both biochar properties and soil conditions and crop species ought to be taken into account comprehensively before applying biochar to a certain soil condition.

However, no promotion of yield was observed with biochar amendments in Acrisol and Anthrosol. We speculate that the lack of biochar effects on yield was caused by exacerbated soil salinity, which inhibited the uptake of nutrients and water (Ju et al., 2006; Zhou et al., 2010) and the growth of the soil microorganisms (Setia et al., 2011). Compared with other biochar (Jia et al., 2012), the higher amounts of ash in B_w and B_m may contain high salts, which would result in soil salinity (Hussain et al., 2017). After the addition of the two salt-rich biochars, the EC values of Acrisol and Anthrosol vegetable soils increased, which might reach the limits to tolerance for the leafy vegetables (Shannon and Grieve,

Table 3. Cumulative gaseous nitrogen (N₂O, NO and NH₃) emissions, gaseous reactive nitrogen emissions (GNrEs), vegetable yield and gaseous reactive nitrogen intensity (GNrI) under the different treatments across the four soils.

| Treatments | Acrisol | Anthrosol | Cambisol | Phaeozem |
|--|----------------|----------------|----------------|-----------------|
| (a) Cumulative N ₂ O emissions (kg N ha ⁻¹) | | | | |
| <i>N</i> | 30.59 ± 3.15aA | 7.83 ± 0.60aB | 2.52 ± 0.37aC | 7.10 ± 1.91aB |
| <i>N</i> + <i>B_w</i> | 19.45 ± 2.43bA | 3.20 ± 0.28bB | 1.97 ± 0.21aB | 3.45 ± 0.86bB |
| <i>N</i> + <i>B_m</i> | 31.56 ± 1.35aA | 3.63 ± 0.62bB | 2.26 ± 0.58aB | 4.01 ± 0.68bB |
| (b) Cumulative NO emissions (kg N ha ⁻¹) | | | | |
| <i>N</i> | 8.99 ± 1.01aA | 1.27 ± 0.15aB | 0.20 ± 0.08aC | 0.97 ± 0.11aBC |
| <i>N</i> + <i>B_w</i> | 4.54 ± 0.60bA | 0.80 ± 0.13bB | 0.33 ± 0.19aB | 0.52 ± 0.03bB |
| <i>N</i> + <i>B_m</i> | 3.87 ± 0.30bA | 1.16 ± 0.17aB | 0.21 ± 0.10aC | 0.94 ± 0.03aB |
| (c) Cumulative NH ₃ emissions (kg N ha ⁻¹) | | | | |
| <i>N</i> | 4.72 ± 0.27aB | 5.79 ± 0.54bA | 6.34 ± 0.51aA | 5.67 ± 0.42aA |
| <i>N</i> + <i>B_w</i> | 5.09 ± 0.38aB | 6.83 ± 0.74abA | 7.35 ± 0.75aA | 6.24 ± 0.49aAB |
| <i>N</i> + <i>B_m</i> | 5.32 ± 0.42aB | 7.57 ± 0.57aA | 7.37 ± 1.11aA | 6.48 ± 0.43aAB |
| (d) GNrEs (kg N ha ⁻¹) | | | | |
| <i>N</i> | 44.30 ± 3.13aA | 14.89 ± 1.33aB | 9.06 ± 0.80aC | 13.74 ± 1.67aB |
| <i>N</i> + <i>B_w</i> | 29.08 ± 2.21bA | 10.82 ± 1.14bB | 9.64 ± 0.88aB | 10.21 ± 0.92bB |
| <i>N</i> + <i>B_m</i> | 40.76 ± 1.66aA | 12.36 ± 0.74bB | 9.84 ± 0.49aC | 11.42 ± 0.27bBC |
| (e) Vegetable yield (t ha ⁻¹) | | | | |
| <i>N</i> | 35.20 ± 2.52aB | 25.29 ± 3.90aC | 39.09 ± 2.03bB | 75.65 ± 5.84bA |
| <i>N</i> + <i>B_w</i> | 29.05 ± 2.35bC | 23.57 ± 1.74aC | 44.53 ± 3.74bB | 76.95 ± 4.04abA |
| <i>N</i> + <i>B_m</i> | 34.93 ± 2.87aC | 26.30 ± 2.63aD | 51.00 ± 3.18aB | 85.89 ± 3.29aA |
| (f) GNrI (kg N t ⁻¹ yield) | | | | |
| <i>N</i> | 1.27 ± 0.18aA | 0.59 ± 0.08aB | 0.23 ± 0.02aC | 0.18 ± 0.04aC |
| <i>N</i> + <i>B_w</i> | 1.01 ± 0.12aA | 0.46 ± 0.05bB | 0.22 ± 0.04aC | 0.13 ± 0.02bC |
| <i>N</i> + <i>B_m</i> | 1.17 ± 0.15aA | 0.47 ± 0.04bB | 0.19 ± 0.01aC | 0.13 ± 0.01bC |

Data shown are means ± SD of the three replicates. See Fig. 1 for treatment codes. Different lowercase letters within the same column indicate significant differences among treatments within the same soil at $p < 0.05$ level. Different capital letters within the same row indicate significant differences among soil types within the same treatment at $p < 0.05$ level.

Table 4. The correlations between N₂O or NO emission and PNR or DEA in each soil.

| Item | Acrisol | | Anthrosol | | Cambisol | | Phaeozem | |
|------------------|---------|-------|-----------|-------|----------|------|----------|-------|
| | PNR | DEA | PNR | DEA | PNR | DEA | PNR | DEA |
| N ₂ O | 0.75* | 0.66 | 0.49 | 0.76* | -0.10 | 0.16 | -0.82** | 0.70* |
| NO | 0.62 | -0.29 | 0.79* | 0.69* | -0.54 | 0.01 | -0.63 | 0.22 |

Asterisks indicate 0.05-level significances (* $p < 0.05$) and 0.01-level significances (** $p < 0.01$), $n = 9$.

1998). Here, we assessed two feedstock-derived biochar effects on GNrI in typical cultivated vegetable soils across mainland China. Overall, biochar amendments reduced GNrI over all the soils, with the magnitude largely depending on soil type. Remarkable reduction in GNrI had been detected due to the efficient mitigation induced by biochar in Anthrosol and Phaeozem (Table 3f). Overall, *B_w* was superior to *B_m* in mitigating the GNrEs, while *B_m* performed better

in vegetable yield enhancement (Table 3d and e). Therefore, the mitigation efficacies on GNrI were not notably different between *B_w* and *B_m* amendments across the four soils.

5 Conclusions

The study demonstrated that biochar amendments mostly reduced N_2O and NO emissions and slightly increased the NH_3 emissions from four soils that are representative of vegetable cropping systems across mainland China. In contrast, biochar amendments did not result in consistent effects on yield, with treatment effects that were both biochar- and soil-specific. Additionally, biochar amendments did decrease GNrI in intensive vegetable soils across mainland China. Furthermore, B_w was superior to B_m in mitigating the GNrEs and the B_m performed better in crop yield throughout all soils. Consequently, both soil type and biochar characteristics need to be seriously considered before large-scale biochar application under certain regions of intensive vegetable production.

Data availability. No data sets were used in this article.

The Supplement related to this article is available online at <https://doi.org/10.5194/bg-14-2851-2017-supplement>.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We would like to express our heartfelt gratitude to the editor and anonymous reviewers for their constructive comments that greatly improved the manuscript. This work was jointly supported by the National Natural Science Foundation of China (41471192), the Special Fund for Agro-Scientific Research in the Public Interest (201503106) and the Ministry of Science and Technology (2013BAD11B01).

Edited by: K. J. van Groenigen

Reviewed by: three anonymous referees

References

- Ameloot, N., Sleutel, S., Das, K. C., Kanagaratnam, J., and Neve, S. D.: Biochar amendment to soils with contrasting organic matter level: effects on N mineralization and biological soil properties, *Glob. Change Biol.*, 7, 135–144, 2015.
- Anderson, J. and Domsch, K.: A physiological method for the quantitative measurement of microbial biomass in soils, *Soil Biol. Biochem.*, 10, 215–221, 1978.
- Anenberg, S. C., Schwartz, J., Shindell, D., Amann, M., Faluvegi, G., Klimont, Z., Janssensmaenhout, G., Pozzoli, L., Van, D. R., and Vignati, E.: Global Air Quality and Health Co-benefits of Mitigating Near-Term Climate Change through Methane and Black Carbon Emission Controls, *Environ. Health Persp.*, 120, 831–839, 2012.
- Avnery, S., Mauzerall, D. L., Liu, J., and Horowitz, L. W.: Global crop yield reductions due to surface ozone exposure: 1. Year 2000 crop production losses and economic damage, *Atmos. Environ.*, 45, 2284–2296, 2011.
- Behera, S. N., Sharma, M., Aneja, V. P., and Balasubramanian, R.: Ammonia in the atmosphere: a review on emission sources, atmospheric chemistry and deposition on terrestrial bodies, *Environ. Sci. Pollut. R.*, 20, 8092–8131, 2013.
- Boyer, E. W., Goodale, C. L., Jaworski, N. A., and Howarth, R. W.: Anthropogenic nitrogen sources and relationships to riverine nitrogen export in the northeastern USA, in: *The Nitrogen Cycle at Regional to Global Scales*, Springer, 2002.
- Cayuela, M., Van Zwieten, L., Singh, B., Jeffery, S., Roig, A., and Sánchez-Monedero, M.: Biochar's role in mitigating soil nitrous oxide emissions: A review and meta-analysis, *Agr. Ecosyst. Environ.*, 191, 5–16, 2014.
- Chan, K. Y. and Xu, Z.: Biochar: nutrient properties and their enhancement, *Biochar for environmental management*, *Sci. Technol.*, 1, 67–84, 2009.
- Ciais, P.: Carbon and other biogeochemical cycles: Final draft underlying scientific technical assessment, IPCC Secretariat, Geneva, 2013.
- Clough, T. J. and Condrón, L. M.: Biochar and the nitrogen cycle: introduction, *J. Environ. Qual.*, 39, 1218–1223, 2010.
- Clough, T. J., Bertram, J. E., Ray, J. L., Condrón, L. M., O'Callaghan, M., Sherlock, R. R., and Wells, N. S.: Unweathered Wood Biochar Impact on Nitrous Oxide Emissions from a Bovine-Urine-Amended Pasture Soil, *Soil Sci. Soc. Am. J.*, 74, 852–860, 2010.
- Deng, J., Zhou, Z., Zheng, X., and Li, C.: Modeling impacts of fertilization alternatives on nitrous oxide and nitric oxide emissions from conventional vegetable fields in southeastern China, *Atmos. Environ.*, 81, 642–650, 2013.
- Diao, T., Xie, L., Guo, L., Yan, H., Lin, M., Zhang, H., Lin, J., and Lin, E.: Measurements of N_2O emissions from different vegetable fields on the North China Plain, *Atmos. Environ.*, 72, 70–76, 2013.
- FAO, IIASA, ISRIC, and ISSCAS: Harmonized World Soil Database Version 1.2, 2012.
- Field, J. L., Keske, C. M. H., Birch, G. L., Defoort, M. W., and Cotrufo, M. F.: Distributed biochar and bioenergy coproduction: a regionally specific case study of environmental benefits and economic impacts, *Glob. Change Biol. Bioenergy*, 5, 177–191, 2013.
- Food and Agriculture Organization (FAO): FAOSTAT (Food and Agriculture Organization Statistical Data) Statistical Yearbook Vol. 4, available at: <http://faostat.fao.org>, last access: 12 August 2015
- Harrison, R. and Webb, J.: A review of the effect of N fertilizer type on gaseous emissions, *Adv. Agron.*, 73, 65–108, 2001.
- Hass, A., Gonzalez, J. M., Lima, I. M., Godwin, H. W., Halvorson, J. J., and Boyer, D. G.: Chicken manure biochar as liming and nutrient source for acid Appalachian soil, *J. Environ. Qual.*, 41, 1096–1106, 2012.
- Hu, H. W., Macdonald, C. A., Trivedi, P., Anderson, I. C., Zheng, Y., Holmes, B., Bodrossy, L., Wang, J. T., He, J. Z., and Singh, B. K.: Effects of climate warming and elevated CO_2 on autotrophic nitrification and nitrifiers in dryland ecosystems, *Soil Biol. Biochem.*, 92, 1–15, 2016.
- Hussain, M., Farooq, M., Nawaz, A., Al-Sadi, A. M., Solaiman, Z. M., Alghamdi, S. S., Ammara, U., Yong, S. O., and Siddique, K.

- H. M.: Biochar for crop production: potential benefits and risks, *J. Soil. Sediment.*, 17, 685–716, 2017.
- IPCC: Climate Change 2013: The Physical Science Basis: working group I contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Stockholm, 2013.
- Islam, A., Chen, D., White, R. E., and Weatherley, A. J.: Chemical decomposition and fixation of nitrite in acidic pasture soils and implications for measurement of nitrification, *Soil Biol. Biochem.*, 40, 262–265, 2008.
- Jeffery, S., Abalos, D., Prodana, M., Bastos, A., van Groenigen, J. W., Hungate, B., and Verheijen, F.: Biochar boosts tropical but not temperate crop yields, *Environ. Res. Lett.*, 12, 053001, 2017.
- Jia, J., Li, B., Chen, Z., Xie, Z., and Xiong, Z.: Effects of biochar application on vegetable production and emissions of N₂O and CH₄, *Soil Sci. Plant Nutr.*, 58, 503–509, 2012.
- Joseph, S. D., Campsarbertain, M., Lin, Y., Munroe, P., Chia, C. H., Hook, J., Van, Z. L., Kimber, S., Cowie, A., and Singh, B. P.: An investigation into the reactions of biochar in soil, *Aust. J. Soil Res.*, 48, 501–515, 2010.
- Ju, X. T., Kou, C. L., Zhang, F. S., and Christie, P.: Nitrogen balance and groundwater nitrate contamination: comparison among three intensive cropping systems on the North China Plain, *Environ. Pollut.*, 143, 117–125, 2006.
- Kim, J. Y., Song, C. H., Ghim, Y. S., Won, J. G., Yoon, S. C., Carmichael, G. R., and Woo, J. H.: An investigation on NH₃ emissions and particulate NH₄⁺–NO₃[–] formation in East Asia, *Atmos. Environ.*, 40, 2139–2150, 2006.
- Kurola, J., Salkinoja-Salonen, M., Aarnio, T., Hultman, J., and Romantschuk, M.: Activity, diversity and population size of ammonia-oxidising bacteria in oil-contaminated landfarming soil, *FEMS Microbiol. Lett.*, 250, 33–38, 2005.
- Langridge, J. M., Lack, D., Brock, C. A., Bahreini, R., Middlebrook, A. M., Neuman, J. A., Nowak, J. B., Perring, A. E., Schwarz, J. P., and Spackman, J. R.: Evolution of aerosol properties impacting visibility and direct climate forcing in an ammonia-rich urban environment, *J. Geophys. Res.-Atmos.*, 117, 2240–2260, 2012.
- Li, B., Fan, C. H., Xiong, Z. Q., Li, Q. L., and Zhang, M.: The combined effects of nitrification inhibitor and biochar incorporation on yield-scaled N₂O emissions from an intensively managed vegetable field in southeastern China, *Biogeosciences*, 12, 2003–2017, <https://doi.org/10.5194/bg-12-2003-2015>, 2015a.
- Li, B., Fan, C. H., Zhang, H., Chen, Z. Z., Sun, L. Y., and Xiong, Z. Q.: Combined effects of nitrogen fertilization and biochar on the net global warming potential, greenhouse gas intensity and net ecosystem economic budget in intensive vegetable agriculture in southeastern China, *Atmos. Environ.*, 100, 10–19, 2015b.
- Li, B., Bi, Z., and Xiong, Z.: Dynamic responses of nitrous oxide emission and nitrogen use efficiency to nitrogen and biochar amendment in an intensified vegetable field in southeastern China, *Glob. Change Biol. Bioenergy*, 9, 400–413, 2017.
- Liu, X., Zhang, A., Ji, C., Joseph, S., Bian, R., Li, L., Pan, G., and Paz-Ferreiro, J.: Biochar's effect on crop productivity and the dependence on experimental conditions – a meta-analysis of literature data, *Plant Soil*, 373, 583–594, 2013.
- Lu, R.: Methods of soil and agro-chemical analysis, China Agricultural Science and Technology Press, Beijing, 127–332, 2000 (in Chinese).
- Ma, L., Shan, J., and Yan, X.: Nitrite behavior accounts for the nitrous oxide peaks following fertilization in a fluvo-aquic soil, *Biol. Fert. Soils*, 51, 563–572, 2015.
- Major, J., Lehmann, J., Rondon, M., and Goodale, C.: Fate of soil-applied black carbon: downward migration, leaching and soil respiration, *Glob. Change Biol.*, 16, 1366–1379, 2010.
- Mei, B. L., Zheng, X. H., Xie, B. H., Dong, H. B., Zhou, Z. X., Rui, W., Jia, D., Feng, C., Tong, H. J., and Zhu, J. G.: Nitric oxide emissions from conventional vegetable fields in southeastern China, *Atmos. Environ.*, 43, 2762–2769, 2009.
- Misselbrook, T. H., Weerden, T. J. V. D., Pain, B. F., Jarvis, S. C., Chambers, B. J., Smith, K. A., Phillips, V. R., and Demmers, T. G. M.: Ammonia emission factors for UK agriculture, *Atmos. Environ.*, 34, 871–880, 2000.
- Mukherjee, A. and Zimmerman, A. R.: Organic carbon and nutrient release from a range of laboratory-produced biochars and biochar–soil mixtures, *Geoderma*, 193/194, 122–130, 2013.
- Nelissen, V.: Effect of different biochar and fertilizer types on N₂O and NO emissions, *Soil Biol. Biochem.*, 70, 244–255, 2014.
- Obia, A., Cornelissen, G., Mulder, J., and Dörsch, P.: Effect of Soil pH Increase by Biochar on NO, N₂O and N₂ Production during Denitrification in Acid Soils, *Plos One*, 10, 359–367, 2015.
- Pacholski, A., Cai, G. X., Fan, X. H., Ding, H., Chen, D., Nieder, R., and Roelcke, M.: Comparison of different methods for the measurement of ammonia volatilization after urea application in Henan Province, China, *J. Plant Nutr. Soil Sci.*, 171, 361–369, 2008.
- Pinder, R. W., Adams, P. J., and Pandis, S. N.: Ammonia emission controls as a cost-effective strategy for reducing atmospheric particulate matter in the Eastern United States, *Environ. Sci. Technol.*, 41, 380–386, 2007.
- Powelson, D. S., Addiscott, T. M., Benjamin, N., Cassman, K. G., de Kok, T. M., Van, G. H., L'Hirondel, J. L., Avery, A. A., and Van, K. C.: When does nitrate become a risk for humans?, *J. Environ. Qual.*, 37, 291–295, 2008.
- Rajkovich, S., Enders, A., Hanley, K., Hyland, C., Zimmerman, A. R., and Lehmann, J.: Corn growth and nitrogen nutrition after additions of biochars with varying properties to a temperate soil, *Biol. Fertil. Soil.*, 48, 271–284, 2012.
- Ravishankara, A. R., Daniel, J. S., and Portmann, R. W.: Nitrous oxide (N₂O): the dominant ozone-depleting substance emitted in the 21st century, *Science*, 326, 123–125, 2009.
- Saarnio, S., Heimonen, K., and Kettunen, R.: Biochar addition indirectly affects N₂O emissions via soil moisture and plant N uptake, *Soil Biol. Biochem.*, 58, 99–106, 2013.
- Schouten, S., Groenigen, J. W. V., Oenema, O., and Cayuela, M. L.: Bioenergy from cattle manure? Implications of anaerobic digestion and subsequent pyrolysis for carbon and nitrogen dynamics in soil, *Glob. Change Biol. Bioenergy*, 4, 751–760, 2012.
- Schulz, H., Dunst, G., and Glaser, B.: Positive effects of composted biochar on plant growth and soil fertility, *Agron. Sustain. Dev.*, 33, 817–827, 2013.
- Setia, R., Marschner, P., Baldock, J., Chittleborough, D., and Verma, V.: Relationships between carbon dioxide emission and soil properties in salt-affected landscapes, *Soil Biol. Biochem.*, 43, 667–674, 2011.
- Shannon, M. C. and Grieve, C. M.: Tolerance of vegetable crops to salinity, *Sci. Hortic.-Amsterdam*, 78, 5–38, 1998.

- Smith, M. S. and Tiedje, J. M.: Phases of denitrification following oxygen depletion in soil, *Soil Biol. Biochem.*, 11, 261–267, 1979.
- Smith, P.: Soil carbon sequestration and biochar as negative emission technologies, *Glob. Change Biol.*, 51, 574–575, 2016.
- Sohi, S. P.: Agriculture. Carbon storage with benefits, *Science*, 338, 1034–1035, 2012.
- Sororzano, L.: Determination of ammonia in natural waters by the phenolhypochlorite method, *Limnol. Oceanogr.*, 14, 799–801, 1969.
- Spokas, K. A. and Reicosky, D. C.: Impacts of sixteen different biochars on soil greenhouse gas production, *Ann. Environ. Sci.*, 3, 179–193, 2009.
- Stavi, I. and Lal, R.: Agroforestry and biochar to offset climate change: a review, *Agron. Sustain. Dev.*, 33, 81–96, 2013.
- Sun, L., Li, L., Chen, Z., Wang, J., and Xiong, Z.: Combined effects of nitrogen deposition and biochar application on emissions of N₂O, CO₂ and NH₃ from agricultural and forest soils, *Soil Sci. Plant Nutr.*, 60, 254–265, 2014.
- Ussiri, D. and Lal, R.: *The Role of Nitrous Oxide on Climate Change*, Springer Netherlands, 2013.
- Wang, J., Chen, Z., Ma, Y., Sun, L., Xiong, Z., Huang, Q., and Sheng, Q.: Methane and nitrous oxide emissions as affected by organic–inorganic mixed fertilizer from a rice paddy in southeast China, *J. Soil. Sediment.*, 13, 1408–1417, 2013.
- Wang, J., Chen, Z., Xiong, Z., Chen, C., Xu, X., Zhou, Q., and Kuzyakov, Y.: Effects of biochar amendment on greenhouse gas emissions, net ecosystem carbon budget and properties of an acidic soil under intensive vegetable production, *Soil Use Manage.*, 31, 375–383, 2015a.
- Wang, S., Nan, J., Shi, C., Fu, Q., Gao, S., Wang, D., Cui, H., Sainzlopez, A., and Zhou, B.: Atmospheric ammonia and its impacts on regional air quality over the megacity of Shanghai, China, *Scientific Reports*, 5, 15842, <https://doi.org/10.1038/srep15842>, 2015b.
- Wrage, N., Velthof, G., Van Beusichem, M., and Oenema, O.: Role of nitrifier denitrification in the production of nitrous oxide, *Soil Biol. Biochem.*, 33, 1723–1732, 2001.
- Xiong, Z., Xie, Y., Xing, G., Zhu, Z., and Butenhoff, C.: Measurements of nitrous oxide emissions from vegetable production in China, *Atmos. Environ.*, 40, 2225–2234, 2006.
- Yao, Z., Zheng, X., Xie, B., Mei, B., Wang, R., Klaus, B. B., Zhu, J., and Yin, R.: Tillage and crop residue management significantly affects N-trace gas emissions during the non-rice season of a subtropical rice-wheat rotation, *Soil Biol. Biochem.*, 41, 2131–2140, 2009.
- Zhang, A., Cui, L., Pan, G., Li, L., Hussain, Q., Zhang, X., Zheng, J., and Crowley, D.: Effect of biochar amendment on yield and methane and nitrous oxide emissions from a rice paddy from Tai Lake plain, China, *Agr. Ecosyst. Environ.*, 139, 469–475, 2010.
- Zhang, F., Chen, X., and Vitousek, P.: Chinese agriculture: An experiment for the world, *Nature*, 497, 33–35, 2013.
- Zhang, M., Fan, C. H., Li, Q. L., Li, B., Zhu, Y. Y., and Xiong, Z. Q.: A 2-yr field assessment of the effects of chemical and biological nitrification inhibitors on nitrous oxide emissions and nitrogen use efficiency in an intensively managed vegetable cropping system, *Agr. Ecosyst. Environ.*, 201, 43–50, 2015.
- Zhang, Y., Luan, S., Chen, L., and Shao, M.: Estimating the volatilization of ammonia from synthetic nitrogenous fertilizers used in China, *J. Environ. Manag.*, 92, 480–493, 2011.
- Zhao, L. M., Wu, L. H., Dong, C. J., and Li, Y. S.: Rice yield, nitrogen utilization and ammonia volatilization as influenced by modified rice cultivation at varying nitrogen rates, *Agr. Sci.*, 01, 10–16, 2010.
- Zhao, X., Wang, J., Wang, S., and Xing, G.: Successive straw biochar application as a strategy to sequester carbon and improve fertility: A pot experiment with two rice/wheat rotations in paddy soil, *Plant Soil*, 378, 279–294, 2014.
- Zheng, X., Mei, B., Wang, Y., Xie, B., Wang, Y., Dong, H., Xu, H., Chen, G., Cai, Z., and Yue, J.: Quantification of N₂O fluxes from soil–plant systems may be biased by the applied gas chromatograph methodology, *Plant Soil*, 311, 211–234, 2008.
- Zhou, J. B., Chen, Z. J., Liu, X. J., Zhai, B. N., and Powlson, D. S.: Nitrate accumulation in soil profiles under seasonally open “sunlight greenhouses” in northwest China and potential for leaching loss during summer fallow, *Soil Use Manage.*, 26, 332–339, 2010.
- Zhu, T., Zhang, J., and Cai, Z.: The contribution of nitrogen transformation processes to total N₂O emissions from soils used for intensive vegetable cultivation, *Plant Soil*, 343, 313–327, 2011.