

therapeutic levels (2.5–15 mcg/ml), it has been reported that Lamotrigine is neuroprotective and improves cognition. At the time of overdose, our patient had a Lamotrigine level of 21.5 mcg/ml. There is limited literature on cognitive effect of supra-therapeutic levels of Lamotrigine. As such, a causal relationship cannot be determined from a single case report. Also in differentials to consider are schizophrenia and seizures from lamotrigine withdrawal.

**FUNDING ACKNOWLEDGEMENTS:** No funding.

### 134 Improvements in Clinical Global Impression of Change With Deutetrabenazine Treatment in Tardive Dyskinesia From the ARM-TD and AIM-TD Studies

Hubert H. Fernandez, MD<sup>1</sup>; Mat D. Davis, PhD<sup>2</sup>; Stewart A. Factor, DO<sup>3</sup>; Robert A. Hauser, MD, MBA<sup>4</sup>; L. Fredrik Jarskog, MD<sup>5</sup>; Joohi Jimenez-Shahed, MD<sup>6</sup>; Rajeev Kumar, MD, FRCPC<sup>7</sup>; Stanislaw Ochudlo, MD, PhD<sup>8</sup>; William G. Ondo, MD<sup>9</sup>; and Karen E. Anderson, MD<sup>10</sup>

<sup>1</sup> Cleveland Clinic, Cleveland, Ohio, USA

<sup>2</sup> Teva Pharmaceutical Industries, Frazer, Pennsylvania, USA

<sup>3</sup> Emory University, Atlanta, Georgia, USA

<sup>4</sup> University of South Florida Parkinson's Disease and Movement Disorders Center, Tampa, Florida, USA

<sup>5</sup> University of North Carolina School of Medicine, Chapel Hill, North Carolina, USA

<sup>6</sup> Baylor College of Medicine, Houston, Texas, USA

<sup>7</sup> Rocky Mountain Movement Disorders Center, Englewood, Colorado, USA

<sup>8</sup> University Clinical Center of Silesian Medical University, Katowice, Poland

<sup>9</sup> Methodist Neurological Institute, Houston, Texas, USA

<sup>10</sup> Georgetown University, Washington, District of Columbia, USA

**ABSTRACT:** Introduction: Tardive dyskinesia (TD) is an involuntary movement disorder that is often irreversible, can affect any body region, and can be debilitating. In the ARM-TD and AIM-TD studies, deutetrabenazine treatment demonstrated statistically and clinically significant reductions in Abnormal Involuntary Movement Scale (AIMS) scores at Week 12 compared with placebo (primary endpoint).

**OBJECTIVE:** To evaluate the efficacy of deutetrabenazine, as measured by the Clinical Global Impression of Change (CGIC) scale, in patients with TD from the pooled ARM-TD and AIM-TD (24 and 36 mg/day doses) data sets, as compared with the pooled placebo cohort.

**METHODS:** ARM-TD and AIM-TD were 12-week, randomized, double-blind, placebo-controlled studies that evaluated the safety and efficacy of deutetrabenazine for the treatment of TD. The key secondary endpoint of each study was the proportion of patients "much improved" or "very much improved" (treatment success) at Week 12 on the CGIC.

**RESULTS:** At Week 12, the odds of treatment success among patients treated with deutetrabenazine (n = 152) was more than double that of patients given placebo (n = 107; odds ratio: 2.12; P = 0.005). In a categorical analysis of CGIC ratings, patients treated with deutetrabenazine showed greater improvement than patients given placebo (P = 0.003). Patients treated with deutetrabenazine also had a significantly better treatment response than those given placebo (least-squares mean CGIC score treatment difference: -0.4; P = 0.006).

**CONCLUSIONS:** Deutetrabenazine treatment led to statistically and clinically significant improvements in TD symptoms based on the CGIC result, suggesting that clinicians were able to recognize the benefit in patients treated with deutetrabenazine.

Presented at: The International Congress of Parkinson's Disease and Movement Disorders; June 4–8, 2017; Vancouver, British Columbia, Canada.

**FUNDING ACKNOWLEDGEMENTS:** These studies were funded by Teva Pharmaceutical Industries, Petach Tikva, Israel.

### 135 Use of Pimavanserin in Combination With Selective Serotonin Reuptake Inhibitors (SSRIs)

James Norton, PhD<sup>1</sup>; Doral Fredericks, PharmD, MBA<sup>2</sup>; Kathy Chi-Burris, MPH<sup>3</sup>; and Randy Owen, MD<sup>4</sup>

<sup>1</sup> Sr. Dir. Medical Affairs, ACADIA Pharmaceuticals Inc., San Diego, CA

<sup>2</sup> Vice President Medical Affairs, ACADIA Pharmaceuticals Inc., San Diego, CA

<sup>3</sup> Sr. Director Biostatistics, ACADIA Pharmaceuticals Inc., San Diego, CA

<sup>4</sup> Vice President Clinical Development, ACADIA Pharmaceuticals Inc., San Diego, CA

**ABSTRACT:** Study Objective: Psychosis is common in Parkinson's disease (PD) and increases in both frequency and severity with disease duration. It is associated with increased morbidity/mortality, complicates management of motor symptoms and often leads to long-term care placement. Pimavanserin is a selective 5-HT<sub>2A</sub> inverse agonist/antagonist approved in the U.S. for treatment of hallucinations and delusions associated