

PDBeFold (SSM: Secondary Structure Matching)

<http://pdbe.org/fold/>

This PDBe tutorial introduces the PDBeFold service, an interactive service for comparing protein structures in 3D. This service provides:

- pairwise and multiple comparison and 3D alignment of protein structures
- examination of a protein structure for similarity with the whole Protein Data Bank (PDB) archive or SCOP.
- best C α -alignment of compared structures
- download and visualisation of best-superposed structures using various graphical packages

PDBeFold structure alignment is based on identification of residues occupying “equivalent” geometrical positions. In other words, unlike sequence alignment residue type is neglected. The PDBeFold service is a very powerful structure alignment tool which can perform both **pairwise** and **multiple** three dimensional alignment. In addition to this there are various options by which the results of the structural alignment query can be sorted. The results of the **Secondary Structure Matching** can be sorted based on the Q score (C α -alignment), P score (taking into account RMSD, number of aligned residues, number of gaps, number of matched Secondary Structure Elements and the SSE match score), Z score (based on Gaussian Statistics), RMSD and %Sequence Identity.


It is hoped that at the end of this tutorial users will be able to use PDBeFOLD for the analysis of their own uploaded structures or entries already in the PDB archive.

How to use this tutorial


Please keep this tutorial open in a browser or PDF viewer and then start the tutorial by opening a new instance of your internet browser and follow the steps outlined in the tutorial using that browser instance

Tutorial

PDBeFOLD may be accessed from multiple locations on the PDBe website. From the PDBe home page (<http://pdbe.org/>), there are two access points for the program as shown below.



The link highlighted in blue may be used to automatically get all related structures to a given PDB code. In order to start the webservice, click on the link shown in red above. This will open up a introductory page concerning PDBeFOLD that contains additional information, tips and help regarding the webservice. On this page, now click on the “Start PDBeFOLD” button.



This should start up the actual webservice that will in the form of a submission form as shown below.


Submission Form for pairwise **3D alignment**
 multiple
[explanation of input](#)

Query	Target
Source: <input type="text" value="PDB entry"/>	Source: <input type="text" value="All PDB archive"/>
PDB code: <input type="text" value="1sar"/> view	
Select chains: <input type="text" value="*"/> Find chains	
Chains: <input type="text" value="*(all)"/>	
Lowest acceptable match (%) <input type="text" value="70"/>	Lowest acceptable match (%) <input type="text" value="70"/>
<input checked="" type="checkbox"/> match individual chains	<input checked="" type="checkbox"/> best matches only
<input checked="" type="checkbox"/> match connectivity	<input checked="" type="checkbox"/> unique matches only
<input checked="" type="checkbox"/> if no matches within acceptability limits found, show <i>some</i> of the close ones	
Precision: <input type="text" value="normal"/>	Sort by: <input type="text" value="Q-score"/> Viewer: <input type="text" value="Jmol"/>
<input type="button" value="Submit your query"/> <input type="button" value="Back to Home Page"/>	

The form contains parameters that may be adjusted or tweaked in order to customize the results as required. The “Query” contains information regarding the structure/structures that need to be compared. This could be an existing PDB

code, an uploaded coordinate file, SCOP entry or pair of existing PDB entries that need to be compared. In addition to this, if an entry contains more than one chain, the chain that has to be compared can be chosen. Similarly the target for the search could be another PDB entry, uploaded coordinate file or SCOP set or a set of files. The “lowest acceptable match” boxes tell the program what cut-off to use in the matching process. The default is 70% both for the query and target, meaning that in order to list a comparison as a match, at least 70% of the secondary structure of the query must match 70% of the target structure. These settings can be adjusted. Setting this to a lower value will result in a large number of hits and the opposite may result in only identical structures found.

We will use the PDB entry 2MJP (<http://pdbe.org/2mjp>) (STRUCTURE-BASED IDENTIFICATION OF THE BIOCHEMICAL FUNCTION OF A HYPOTHETICAL PROTEIN FROM METHANOCOCCUS JANNASCHII: MJ0226) as an example for demonstrating PDBefold.



The image shows the PDBefold submission form and a protein structure visualization. The form is titled "Submission Form for pairwise 3D alignment" and includes options for "multiple" and "explanation of input". It is divided into "Query" and "Target" sections. The Query section has "Source: PDB entry", "PDB code: 2mjp", "Chains: *(all)", and "Lowest acceptable match (%): 70". The Target section has "Source: All PDB archive" and "Lowest acceptable match (%): 70". There are checkboxes for "match individual chains", "match connectivity", "best matches only", and "unique matches only". The form also includes "Precision: normal", "Sort by: Q-score", and "Viewer: Jmol". A blue arrow points from the "PDB code" field to a 3D ribbon diagram of a protein structure, which is colored in yellow and pink.

Click on the “Submit your query” button. This will start a alignment job on the farm which will take upto 2-3 minutes depending on server load. Once the results are calculated, a new page showing the same will be displayed on the browser (Similar to that shown below).

##	Q	P	Z	Rmsd	N _{align}	N _g	% _{seq}	% _{sse}	Match	% _{sse}	N _{res}	x	Title
1	1.00	26.8	15.5	0.00	184	0	100	100	2mjp:A	100	184	<input type="checkbox"/>	STRUCTURE-BASED IDENTIFICATION OF THE BIOCHEMICAL FUNCTION OF A HYPOTHETICAL PROTEIN FROM METHANOCOCCUS JANNASCHII MJ0226
2	1.00	25.4	15.0	0.20	184	0	100	100	1b78:A	100	184	<input type="checkbox"/>	STRUCTURE-BASED IDENTIFICATION OF THE BIOCHEMICAL FUNCTION OF A HYPOTHETICAL PROTEIN FROM METHANOCOCCUS JANNASCHII MJ0226
3	0.95	22.3	14.1	0.49	182	1	99	93	1b78:B	93	184	<input type="checkbox"/>	STRUCTURE-BASED IDENTIFICATION OF THE BIOCHEMICAL FUNCTION OF A HYPOTHETICAL PROTEIN FROM METHANOCOCCUS JANNASCHII MJ0226
4	0.95	20.0	13.3	0.55	182	1	99	86	2mjp:B	100	184	<input type="checkbox"/>	STRUCTURE-BASED IDENTIFICATION OF THE BIOCHEMICAL FUNCTION OF A HYPOTHETICAL PROTEIN FROM METHANOCOCCUS JANNASCHII MJ0226
5	0.85	15.3	11.6	1.14	182	2	50	93	2dvn:B	100	186	<input type="checkbox"/>	STRUCTURE OF PH1917 PROTEIN WITH THE COMPLEX OF IMP FROM PYROCOCCUS HORIKOSHII
6	0.68	9.1	8.9	1.92	181	3	49	93	1v7r:A	100	186	<input type="checkbox"/>	STRUCTURE OF NUCLEOTIDE TRIPHOSPHATE PYROPHOSPHATASE FROM PYROCOCCUS HORIKOSHII OT3
7	0.67	8.5	8.6	1.96	181	3	49	93	2dvo:A	100	185	<input type="checkbox"/>	STRUCTURE OF PH1917 PROTEIN WITH THE COMPLEX OF ITP FROM PYROCOCCUS HORIKOSHII
8	0.67	9.0	8.9	1.95	181	3	49	93	2dvn:A	100	186	<input type="checkbox"/>	STRUCTURE OF PH1917 PROTEIN WITH THE COMPLEX OF IMP FROM PYROCOCCUS HORIKOSHII
9	0.66	6.7	7.7	1.93	177	5	50	79	2dvp:A	92	184	<input type="checkbox"/>	STRUCTURE OF NTPASE FROM PYROCOCCUS HORIKOSHII
10	0.65	7.9	8.3	2.08	181	3	49	93	2e5x:A	100	185	<input type="checkbox"/>	STRUCTURE OF NUCLEOTIDE TRIPHOSPHATE PYROPHOSPHATASE FROM PYROCOCCUS HORIKOSHII OT3
11	0.65	8.7	8.8	1.42	168	7	35	71	2car:B	83	194	<input type="checkbox"/>	CRYSTAL STRUCTURE OF HUMAN INOSINE TRIPHOSPHATASE
12	0.64	8.6	8.8	1.48	170	7	35	71	2car:A	83	196	<input type="checkbox"/>	CRYSTAL STRUCTURE OF HUMAN INOSINE TRIPHOSPHATASE
13	0.64	8.8	8.8	1.41	168	7	35	71	2i5d:A	83	195	<input type="checkbox"/>	CRYSTAL STRUCTURE OF HUMAN INOSINE TRIPHOSPHATE PYROPHOSPHATASE
14	0.63	5.4	7.0	1.81	173	6	35	71	1vp2:A	91	189	<input type="checkbox"/>	CRYSTAL STRUCTURE OF PUTATIVE XANTHOSINE TRIPHOSPHATE PYROPHOSPHATASE/HAM1 PROTEIN HOMOLOG (TM0159) FROM THERMOTOGA MARITIMA AT 1.78 Å RESOLUTION

The results here are sorted based on Q-score (Quality of alignment, with 1 being the highest score) for this entry. The pairwise alignment result between **2mjp** and **2e5x** is highlighted in the above figure. There is a **49% amino acid sequence identity** between the two proteins, whereas they have **93% secondary structure identity**. Clicking on the number link on the left hand side of the page will return residue-by-residue description about the structural alignment between the two proteins. *Note: as the PDB archive continuously adds new entries every week, it is possible that this result may not appear in the same position as shown.*

Match 11 of 29

[Back to match list](#)
[first match](#)
[<<](#)
[>>](#)
[last match](#)
[Back to query](#)

Query PDB 2mjp:A				Alignment				Target PDB 2e5x:A			
N _{res}	% _{res}	N _{SSE}	% _{SSE}	Q	P	RMSD	N _{align}	N _{res}	% _{res}	N _{SSE}	% _{SSE}
184	98	14	93	0.650	7.93	2.081	181	185	98	13	100
STRUCTURE-BASED IDENTIFICATION OF THE BIOCHEMICAL FUNCTION OF A HYPOTHETICAL PROTEIN FROM METHANOCOCCUS JANNASCHII MJ0226				% _{seq}	Z	N _{SSE}	N _{gaps}	STRUCTURE OF NUCLEOTIDE TRIPHOSPHATE PYROPHOSPHATASE FROM PYROCOCCUS HORIKOSHII OT3			
				48.6	8.35	13	3				

[view](#) [download](#)

[view superposed](#)

[view](#) [download](#)


[superpose whole entries](#)

Viewer: [Jmol](#)

Secondary Structure Alignment

2mjp:A **S**H**S**H**S**H**S**H**S**S**H**H**H**H
 2e5x:A **S**H**S**H**S**H**S**S**-S**H**H**H

Click on the “View Superposed” button to view the secondary structure alignment.



The two structures 2MJP and 2E5X appear share a high level of secondary structure identity despite only having <50% sequence identity. This indicates that the two proteins belong to the same structural family.

A little further down the page are the summary of the alignment showing the secondary structure elements that match between the two structures with residue ranges and the matrix required to move and align the target to the query structure.

Query PDB 2mjp:A

1	SD	4	A	ILE	11	ALA	14
2	H1	11	A	PRO	18	LEU	28
3	SD	4	A	ILE	36	ILE	39
4	H1	18	A	THR	49	LYS	66
5	SD	10	A	VAL	69	VAL	78
6	H1	9	A	TYR	88	ILE	96
7	H1	10	A	ILE	96	LEU	105
8	SD	12	A	ASN	112	ASP	123
9	SD	13	A	GLY	126	VAL	138
11	SD	3	A	PHE	155	PRO	157
12	H5	5	A	THR	163	MET	167
13	H1	6	A	THR	168	SER	173
14	H1	16	A	SER	176	ASP	191

Target PDB 2e5x:A

<->	1	SD	5	A	LYS	2	ILE	6
<->	2	H1	14	A	ASN	9	THR	22
<->	3	SD	5	A	GLU	26	LEU	30
<->	4	H1	14	A	LYS	41	LYS	54
<->	5	SD	10	A	PHE	61	ILE	70
<->	6	H1	9	A	TYR	80	ILE	88
<->	7	H1	10	A	ILE	88	MET	97
<->	8	SD	12	A	ARG	104	ILE	115
<->	9	SD	13	A	LYS	118	ILE	130
<->	10	SD	3	A	PHE	146	PRO	148
<->	11	H5	5	A	THR	154	MET	158
<->	12	H1	9	A	THR	159	SER	167
<->	13	H1	18	A	SER	167	LEU	184

SCOP: domain [33223](#) , family [c.51.4.1](#)

[PDBe Atlas](#) | [PDBe Motif](#) | [OCA](#)

[PDBe Atlas](#) | [PDBe Motif](#) | [OCA](#)

[GeneCensus](#) | [FSSP](#) | [3Dee](#) | [CATH](#) | [PDBsum](#)

[GeneCensus](#) | [FSSP](#) | [3Dee](#) | [CATH](#) | [PDBsum](#)

[view](#)

[download sequence](#)

[view superposed](#)

[view](#)

[download sequence](#)

Rotation-translation matrix

(to be applied to the target)

Download the page content

[>>](#) in plain text

[>>](#) in XML

-0.586	-0.334	-0.738	X	46.884
-0.714	0.643	0.276	Y	-30.053
0.383	0.689	-0.616	Z	-19.129

If you scroll down the page, there is a 3D structural alignment between the residues from the corresponding PDB entries. This alignment shows the extent of superposition between similar secondary structure folds. The **red** background indicates that matched residues have same residue name, **cyan** background is used for other matched residues. Unmatched residues are indicated by **black** background. The letters H and S represent helix and β -strands respectively.


3D Structural alignment

[notations](#)

PDB 2mjp:A	SI	Dist. (Å)	PDB 2e5x:A
			- A:MET 1
+ A:LYS 10	:::	2.20	S+ A:LYS 2
S- A:ILE 11	:::	2.05	S- A:ILE 3
S- A:TYR 12	:::	1.69	S- A:PHE 4
S- A:PHE 13	:::	1.35	S- A:PHE 5
S- A:ALA 14	:	1.19	S- A:ILE 6
· A:THR 15	:::	1.83	· A:THR 7
- A:GLY 16		2.95	· A:SER 8
+ A:ASN 17	:::	3.82	H+ A:ASN 9
H+ A:PRO 18	:::	4.87	H+ A:PRO 10
H+ A:ASN 19		5.09	H- A:GLY 11
H+ A:LYS 20	:::	3.70	H+ A:LYS 12
H- A:ILE 21	:::	3.77	H- A:VAL 13
H+ A:LYS 22	:::	4.19	H+ A:ARG 14
H+ A:GLU 23	:::	3.13	H+ A:GLU 15

The residue-by-residue mapping provides a very useful tool to analyse the structure function relationship between the two entries.

Lets now concentrate on the proteins themselves to get more information about them. You can go to the summary pages for these entries at the PDBe in new browser windows or tabs. Go to <http://www.pdbe.org/2mjp> in one window and <http://www.pdbe.org/2e5x> in another. On each of these pages, click on the “Ligands” link from the left-hand sidebar.





- Summary
- Details
- Experiment
- EDS
- Structure

PDB entry: 2mjp

STRUCTURE-BASED IDENTIFICATION OF THE BIOCHEMICAL FUNCTION OF A HYPOTHETICAL PROTEIN FROM

Bound ligands

Ligand		<p>ANP PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER 1 instances in this entry Formula: $C_{10}H_{17}N_6O_{12}P_3$ interactions - interaction statistics - HIC-Up</p>
--------	---	--






- Summary
- Details
- Experiment
- EDS
- Structure
 - Primary
 - Secondary
 - Tertiary
 - Quaternary
- Taxonomy
- Citation
- Ligands
- Cross references
- Visualisation
- Downloads

PDB entry: 2e5x

Structure of nucleotide triphosphate pyrophosphatase from *pyrococcus horikoshii* OT3

Bound ligands

Ligand		<p>EDO 1,2-ETHANEDIOL 2 instances in this entry Formula: $C_2H_6O_2$ interactions - interaction statistics - HIC-Up</p>
Ligand		<p>ITT INOSINE 5'-TRIPHOSPHATE 1 instances in this entry Formula: $C_{10}H_{15}N_4O_{14}P_3$ interactions - interaction statistics - HIC-Up</p>
Ligand		<p>NA SODIUM ION 3 instances in this entry Formula: Na interactions - interaction statistics</p>

As you can see from the above screenshots, both entries contain nucleotide-type ligands. The ligand bound to 2MJP is ANP (PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER). For 2E5X, the bound nucleotide is ITT (INOSINE 5'-TRIPHOSPHATE). Given that both entries share a high degree of structural similarity and also bind nucleotide-like ligands, do they also share the same or similar binding sites?

It should be possible answer that question by looking at the ligand interaction results for these two hetgroups from the corresponding entries. Click on the

Protein Data Bank in Europe

<http://pdbe.org>

PDBeFOLD Tutorial

“interactions” link for each of the two ligands as shown above. This takes you to the details of the residue interactions in both entries using another PDBe service (PDBeMotif: <http://pdbe.org/motif/>).

2MJP

bound molecules		
No. Motif/Active site	Ligand	Environment
PKC_PHOSPHO_SITE	A 176+178	
slinofil	A 176+180	
betaturn	A 149+152	
1	A 151+154	ANP 500A ASN 17A GLU 23A SER 89A ASP 152A HIS 177A ARG 178A PHE 115A HOH 586A ASN 19A LYS 20A SER 74A GLY 75A LYS 90A PHE 149A
niche	A 152+155	
alphabelamotif	A 151+155	
siturn	A 15+17	
niche	A 15+17	

2E5X

bound molecules		
No. Motif/Active site	Ligand	Environment
1	ITT 201A NA 301A NA 302A THR 7A LYS 12A SER 66A ARG 169A HOH 414A HOH 450A HOH 529A HOH 551A ASN 9A SER 82A PHE 140A GLY 141A TYR 142A SER 8A ASP 65A	

Now going back to the residue-by-residue 3D-mapping results provided by PDBefold, we can make some very interesting observations. For example ASN 17 and LYS 20 interacts with the ligand ANP in 2MJP, aligns with ASN 9 and LYS 12 from PDB entry 2E5X. Both ASN 9 and LYS 12 are also involved in interactions with ITT in a similar manner.

•	A:THR 15	::•	1.83		•	A:THR 7
-	A:GLY 16		2.95		•	A:SER 8
+	A:ASN 17	::•	3.82		H+	A:ASN 9
H+	A:PRO 18	::•	4.87		H+	A:PRO 10
H+	A:ASN 19		5.09		H-	A:GLY 11
H+	A:LYS 20	::•	3.70		H+	A:LYS 12
H-	A:ILE 21	::	3.77		H-	A:VAL 13
H+	A:LYS 22	::	4.19		H+	A:ARG 14
H+	A:GLU 23	::•	3.13		H+	A:GLU 15
H-	A:ALA 24	:	2.22		H-	A:VAL 16

In this way if we compare the rest of the residues present in the binding environment for ANP and ITT, it will be very clear that both the proteins try to adopt similar binding environment in order to interact with a nucleotide ligand. This observation also indicates that the hypothetical protein represented by the PDB entry 2MJP is potential pyrophosphatase similar to the protein under PDB

entry 2E5X. Therefore in this case, the function of the protein molecule dictates its overall structural fold rather than its sequence identity.

Go back to all the results of the structural comparison of 2MJP with the PDB.

9	0.66	8.2	8.5	2.04	181	3	49	93	2zti:A	100	184	<input checked="" type="checkbox"/>	STRUCTURES OF DIMERIC NONSTANDARD NUCLEOTIDE TRIPHOSPHATE PYROPHOSPHATASE FROM PYROCOCCUS HORIKOSHII OT3: FUNCTIONAL SIGNIFICANCE OF INTERPHOTOMER CONFORMATIONAL CHANGES
10	0.66	6.7	7.7	1.93	177	5	50	79	2dvp:A	92	184	<input type="checkbox"/>	STRUCTURE OF NTPASE FROM PYROCOCCUS HORIKOSHII
11	0.65	7.9	8.3	2.08	181	3	49	93	2e5x:A	100	185	<input type="checkbox"/>	STRUCTURE OF NUCLEOTIDE TRIPHOSPHATE PYROPHOSPHATASE FROM PYROCOCCUS HORIKOSHII OT3
12	0.65	8.7	8.8	1.42	168	7	35	71	2car:B	83	194	<input checked="" type="checkbox"/>	CRYSTAL STRUCTURE OF HUMAN INOSINE TRIPHOSPHATASE
13	0.64	8.6	8.8	1.48	170	7	35	71	2car:A	83	196	<input checked="" type="checkbox"/>	CRYSTAL STRUCTURE OF HUMAN INOSINE TRIPHOSPHATASE
14	0.64	8.8	8.8	1.41	168	7	35	71	215d:A	83	195	<input type="checkbox"/>	CRYSTAL STRUCTURE OF HUMAN INOSINE TRIPHOSPHATE PYROPHOSPHATASE
15	0.63	5.4	7.0	1.81	173	6	35	71	1vp2:A	91	189	<input type="checkbox"/>	CRYSTAL STRUCTURE OF PUTATIVE XANTHOSINE TRIPHOSPHATE PYROPHOSPHATASE/HAM1 PROTEIN HOMOLOG (TM0159) FROM THERMOTOGA MARITIMA AT 1.78 Å RESOLUTION
16	0.63	5.3	7.0	1.87	174	5	34	71	1vp2:B	91	189	<input type="checkbox"/>	CRYSTAL STRUCTURE OF PUTATIVE XANTHOSINE TRIPHOSPHATE PYROPHOSPHATASE/HAM1 PROTEIN HOMOLOG (TM0159) FROM THERMOTOGA MARITIMA AT 1.78 Å RESOLUTION
17	0.62	7.9	8.5	1.78	169	8	35	71	2j4e:G	77	186	<input checked="" type="checkbox"/>	THE ITP COMPLEX OF HUMAN INOSINE TRIPHOSPHATASE
18	0.61	8.2	8.6	1.76	169	7	34	71	2j4e:E	83	190	<input type="checkbox"/>	THE ITP COMPLEX OF HUMAN INOSINE TRIPHOSPHATASE
19	0.60	3.8	6.8	1.81	177	7	33	71	1k7k:A	91	207	<input type="checkbox"/>	CRYSTAL STRUCTURE OF RDGB- INOSINE TRIPHOSPHATE PYROPHOSPHATASE FROM E. COLI
20	0.60	3.4	6.7	1.79	176	6	32	71	2pyu:A	83	208	<input checked="" type="checkbox"/>	STRUCTURE OF THE E. COLI INOSINE TRIPHOSPHATE PYROPHOSPHATASE RDGB IN COMPLEX WITH IMP

The checkbox shown on the results page can be used to select entries to submit for multiple alignment from the pairwise alignment pages. Check a few of these results and choose “Submit for Multiple Alignment” from the bottom of the results page. This will do a multiple alignment between the chosen entries and the results will be shown similar to that seen for the pairwise alignment. You may choose the “View Superposed” button to see all your chosen entries aligned.

Multiple Alignment Results http://www.ebi.ac.uk/msd-srv/ssm/cgi-bin/ssmserver?qs_jm

Back to query Download XML Download text


##		Structure	N _{res}	N _{SSE}	Consensus scores	
					RMSD	Q-score
1	<input checked="" type="checkbox"/>	PDB 2mjp:A	184	14	0.9524	0.8393
2	<input checked="" type="checkbox"/>	PDB 2zti:A	184	13	1.5800	0.7233
3	<input checked="" type="checkbox"/>	PDB 2car:B	194	12	1.3319	0.7320
4	<input checked="" type="checkbox"/>	PDB 2j4e:G	186	13	1.1986	0.7882
5	<input checked="" type="checkbox"/>	PDB 2pyu:A	208	12	1.5801	0.6398

Number of aligned residues 170 Overall RMSD
Number of aligned SSEs 7 Overall Q-score

Select all Unselect all view superposed download FASTA
 superpose whole entries Viewer: J

Secondary Structure Alignment


[SSF details](#) PDB 2min:A -S_HsH_SH_SshshhH



As can be seen all our chosen entries appear to share the same overall structure.

Multiple Alignments using PDBeFOLD

The multiple alignment results page shows structural alignment among the 4 entries that were uploaded (1b78, 2dvn, 2e5x and 1v7r). You can also view the superposed entries in a graphics viewer by pressing the “Show Superposed” button.



The central domain region in all proteins appears to be highly similar. Analysis of the residue-by-residue mapping data (similar way it was done in the pairwise alignment before) also indicates high degree of similarity observed in the active binding site (nucleotide binding) of all these proteins.

3D Structural alignment

1b78	2dvn	2e5x:A	1v7r:A
A: MET 1	A: MET 1	MET 1	MET 1
A: LYS 10	A: LYS 2	LYS 2	LYS 2
A: ILE 11	A: ILE 3	ILE 3	ILE 3
A: TYR 12	A: PHE 4	PHE 4	PHE 4
A: PHE 13	A: PHE 5	PHE 5	PHE 5
A: ALA 14	A: ILE 6	ILE 6	ILE 6
A: THR 15	A: THR 7	THR 7	THR 7
A: GLY 16	A: SER 8	SER 8	SER 8
A: ASN 17	A: ASN 9	ASN 9	ASN 9
A: PRO 18	A: PRO 10	PRO 10	PRO 10
A: ASN 19	A: GLY 11	GLY 11	GLY 11
A: LYS 20	A: LYS 12	LYS 12	LYS 12
A: ILE 21	A: VAL 13	VAL 13	VAL 13
A: LYS 22	A: ARG 14	ARG 14	ARG 14
A: GLU 23	A: GLU 15	GLU 15	GLU 15
A: ALA 24	A: VAL 16	VAL 16	VAL 16
A: ASN 25	A: ALA 17	ALA 17	ALA 17
A: ILE 26	A: ASN 18	ASN 18	ASN 18
A: ILE 27	A: PHE 19	PHE 19	PHE 19
A: LEU 28	A: LEU 20	LEU 20	LEU 20
A: LYS 29	A: GLY 21	GLY 21	GLY 21


These results obtained from PDBfold service provide encouraging prospects of understanding possible roles of a hypothetical protein or structural genomics proteins whose function is yet to be determined.

OTHER EXAMPLES

Secondary structure alignments can often show relationships that are not immediately obvious from sequence identity alone. Here are a few examples which you may find interesting.

- a) Alpha-lactalbumin (PDB entry 1A4V <http://pdbe.org/1a4v>).

Start a PDBeFOLD comparison for all entries in the PDB archive against 1A4V. Once the results are shown, scroll to the bottom of the page and sort by %seq instead of Q-score.



Now scroll to the last page and choose one of the results from the last page.

660	0.76	6.5	7.5	1.45	122	5	34	80	21x2:A	100	129	<input type="checkbox"/>	THE THREE DIMENSIONAL STRUCTURE OF TURKEY EGG WHITE LYSOZYME AT 2.2 ANGSTROMS RESOLUTION
661	0.57	2.9	5.1	2.16	114	6	33	70	2h5x:A	100	122	<input type="checkbox"/>	CRYSTALLOGRAPHIC STRUCTURE OF DIGESTIVE LYSOZYME 1 FROM MUSCA DOMESTICA BOUND TO CHITOTETRAOSE AT 1.92 A RESOLUTION
662	0.55	2.9	5.1	2.23	115	5	31	70	3cb7:B	100	126	<input type="checkbox"/>	THE CRYSTALLOGRAPHIC STRUCTURE OF THE DIGESTIVE LYSOZYME 2 FROM MUSCA DOMESTICA AT 1.9 ANG.

Entry 3cb7 has 31% sequence identity but 70% structure similarity between the two proteins. Look at the details of the match. The structures are highly similar.

Match 662 of 662


Query PDB 1a4v:A				Alignment				Target PDB		
N _{res}	% _{res}	N _{SSE}	% _{SSE}	Q	P	RMSD	N _{align}	N _{res}	% _{res}	N _{...}
123	93	10	70	0.549	2.94	2.233	115	126	91	
ALPHA-LACTALBUMIN				% _{seq}	Z	N _{SSE}	N _{gaps}	THE CRYSTALLIC STRUCTURE OF THE LYSOZYME 2 FROM MUSCA DOMESTICA AT 1.9 ANGSTROMS RESOLUTION		
				31.3	5.08	7	5			

superpose whole entries

Secondary Structure Alignment

```


1a4v:A  HhHssHHHHH
3cb7:B  H-H--HHHHH
    
```



- b) Eosinophil Major Basic Protein (PDB entry 1H8U <http://pdbe.org/1h8u>). Do a search for structural similarity for PDB entry 1H8U against the whole PDB archive. Once the results are in, resort the results by %seq identity as in the previous example. Scroll to the last page.

191	0.39	5.7	7.4	1.65	85	7	14	88	1kcg:B	70	123	<input type="checkbox"/>	NRG2D IN COMPLEX WITH ULBP3
192	0.38	0.6	3.1	2.01	93	8	14	75	1kg0:C	75	136	<input type="checkbox"/>	STRUCTURE OF THE EPSTEIN-BARR VIRUS GP42 PROTEIN BOUND TO THE MHC CLASS II RECEPTOR HLA-DR1
193	0.43	5.9	7.5	1.89	94	8	14	88	1mpu:A	78	128	<input type="checkbox"/>	CRYSTAL STRUCTURE OF THE FREE HUMAN NKG2D IMMUNORECEPTOR

The last hit in the list is PDB entry 1MPU. This entry has 14% sequence identity with our query structure while sharing 88% structural identity. Look at the details of this hit and view the superposed entries as previously.



Scroll down the details page and look at the residue-by-residue listing. All the CYS residues between the two structures are conserved and at the same equivalent positions. These residues form the disulphide bonds in the two structures that keep the scaffold of the protein intact. As a matter of fact, both these proteins belong to a large family of sugar-binding proteins called c-type lectins. All c-type lectins share the same overall structure constrained by the disulphide bonds, and not all proteins in this family actually bind sugars.

- c) PDB entry 1TIM and 2O55. Do a pairwise alignment between 1TIM and 2O55 using the pairwise alignment form. The results will show that the two proteins share 4% sequence identity and 70% structural similarity.

Structure Alignment Results

[explanation of output](#)

Query: pdb entry **1tim:A**, 247 residues
STRUCTURE OF TRIOSE PHOSPHATE ISOMERASE FROM CHICKEN MUSCLE


Examined 1 entry (1 chain),
Match 1 of 1.

[Back to query](#)
[resort results](#)

##	Scoring			Rmsd	N _{align}	N _g	%seq	Query		Target (PDB entry)				Title
	Q	P	Z					%sse	Match	%sse	N _{res}	x		
1	0.19	-0.0	3.1	3.62	171	14	4	70	2o55:A	82	254	<input type="checkbox"/>	CRYSTAL STRUCTURE OF A PUTATIVE GLYCEROPHOSPHODIESTER PHOSPHODIESTERASE FROM GALDIERIA SULPHURARIA	

Examined 1 entry (1 chain),
Match 1 of 1.

[Back to query](#)



Both proteins have minimal sequence identity and yet belong to the same fold class (the TIM barrel).

This ends our tutorial on PDBeFOLD. We hope you found this useful and will be able to use this tool in your future research and analysis. It is hopefully clear from the examples given in this tutorial that fold space is more restricted than sequence space and most proteins tend to fall broadly into one of the many fold classes already found in the PDB. If you need to get in touch with the PDBe regarding any aspect of the programme, please email pdbehel@ebi.ac.uk and we will try to assist you in any way possible.