

LINEAR RECURSION RELATIONS — LESSON SEVEN ANALYZING LINEAR RECURSION SEQUENCES

Frequently one encounters problems such as the following: Find the next three terms in the following sequences:

$$1, 3, 5, 7, 9, 11, \dots$$

$$3, 4, 7, 11, 18, 29, \dots$$

$$3, 6, 12, 24, 48, 96, \dots$$

As has been pointed out many times, the solution to such problems is highly indeterminate. It is "obvious" that the general term of the first sequence is

$$T_n = 2n - 1$$

But

$$T_n = 2n - 1 + (n - 1)(n - 2)(n - 3)(n - 4)(n - 5)(n - 6)V_n$$

where V_n is the n^{th} term of any sequence of finite quantities would do just as well. Similarly for the other cases.

Or looking at the matter from the standpoint of linear recursion relations, the six numbers in each case might be the first six terms of a linear recursion relation of the sixth order. Hence any infinite number of possibilities arises.

How can the problem be made more specific? Possibly, one might say: Find the expression for the n^{th} term of a linear recursion relation of minimum order. Whether this is sufficient to handle all instances of this type is an open question, but it would seem to take care of the present cases.

The solutions in the three instances listed above are:

$$T_{n+1} = 2T_n - T_{n-1}$$

$$T_{n+1} = T_n + T_{n-1}$$

$$T_{n+1} = 2T_n$$

LINEAR RECURSION RELATIONS

If a sequence has terms which were derived from a polynomial expression in n , this expression can be found by the method of differences. As was pointed out in the first lesson, if the terms derive from a polynomial of degree k , the k^{th} differences are constant and the $(k + 1)^{\text{st}}$ difference is zero. A simple method of reconstituting the polynomial is to use Newton's Interpolation Formula:

$$(1) \quad f(n) = \frac{\Delta^k f(0)}{k!} n^{(k)} + \frac{\Delta^{k-1} f(0)}{(k-1)!} n^{(k-1)} + \dots + \frac{\Delta f(0)}{1!} n^{(1)} + f(0)$$

where $\Delta^k f(0)$ is the k^{th} difference taken at the zero value and $n^{(k)}$ is the factorial $n(n-1)(n-2)\dots(n-k+1)$ of k terms.

Example. Determine the polynomial of lowest degree which fits the following set of values.

n	$f(n)$	$\Delta f(n)$	$\Delta^2 f(n)$	$\Delta^3 f(n)$
0	6			
1	11	5	32	
2	48	37	50	18
3	135	87	68	18
4	290	155	86	18
5	531	241	104	18
6	876	345	122	18
7	1343	467	140	18
8	1950	607		

Using Newton's Interpolation Formula,

$$f(n) = \frac{18}{3!} n(n-1)(n-2) + \frac{32}{2!} n(n-1) + 5n + 6$$

$$f(n) = 3n^3 + 7n^2 - 5n + 6.$$

LINEAR RECURSION RELATIONS

Suppose that we have a sequence whose terms are the sum of the terms of two sequences: (1) A sequence whose values derive from a polynomial: (2) A sequence whose terms form a geometric progression. Is it possible to determine the components of this sequence?

Imagine that the terms of the sequence have been separated into their two component parts. Then on taking differences, the effect of the polynomial will eventually become nil. How does a geometric progression function under differencing? This can be seen from the table below.

a	$a(r - 1)$	$a(r - 1)^2$	$a(r - 1)^3$
ar	$ar(r - 1)$	$ar(r - 1)^2$	$ar(r - 1)^3$
ar^2	$ar^2(r - 1)$	$ar^2(r - 1)^2$	$ar^2(r - 1)^3$
ar^3	$ar^3(r - 1)$	$ar^3(r - 1)^2$	$ar^3(r - 1)^3$
ar^4	$ar^4(r - 1)$	$ar^4(r - 1)^2$	$ar^4(r - 1)^3$

Clearly, differencing a geometric progression produces a geometric progression with the same ratio. By examining the form of the leading term, one can readily deduce the value of a , the initial term of the geometric progression as well.

Example.

POLYNOMIAL AND GEOMETRIC PROGRESSION COMBINED

n	T_n				
1	4				
2	16	12			
3	70	54	42		
4	224	154	100	58	
5	616	392	238	138	80
6	1624	1008	616	378	240
7	4346	2722	1714	1098	720
8	12040	7644	4972	3258	2160
9	34444	22404	14710	9738	6480

In the last column, one has a geometric progression with ratio 3, but not in the previous column. Hence the polynomial that was combined with the geometric

LINEAR RECURSION RELATIONS

progression was of degree 3. For the geometric progression, $r = 3$ and

$$a \times 2^4 = 80, \text{ so that } a = 5 .$$

Eliminating the effect of the geometric progression from the leading terms gives:

$$58 - 2^3 \times 5 = 18$$

$$42 - 2^2 \times 5 = 22$$

$$12 - 2 \times 5 = 2$$

$$4 - 5 = -1$$

To apply Newton's Formula, we have to go back to the zero elements by extrapolation.

$$\begin{aligned} \Delta^3 f(0) &= 18, \quad \Delta^2 f(0) = 22 - 18 = 4, \quad \Delta f(0) = 2 - 4 = -2 \\ f(0) &= -1 - (-2) = 1 . \end{aligned}$$

Hence

$$f(n) = \frac{18}{3!} n(n-1)(n-2) + \frac{4}{2!} n(n-1) - 2n + 1$$

$$f(n) = 3n^3 - 7n^2 + 2n + 1 .$$

Hence the term of the sequence has the form:

$$T_n = 3n^3 - 7n^2 + 2n + 1 + 5 \times 3^{n-1} .$$

The recursion relation for this term can be readily found by the methods of the previous lesson.

POLYNOMIAL AND FIBONACCI SEQUENCE

If we know that the terms of a sequence are formed by combining the elements of a polynomial sequence and a Fibonacci sequence, we have a situation similar to the previous case. For whereas the polynomial element vanishes

LINEAR RECURSION RELATIONS

on taking a sufficient number of differences, the Fibonacci element persists. This can be seen from the following table.

n	T_n	ΔT_n	$\Delta^2 T_n$	$\Delta^3 T_n$
1	T_1	T_0		
2	T_2	T_1		
3	T_3	T_2	T_{-1}	T_{-2}
4	T_4	T_3	T_0	T_{-1}
5	T_5	T_4	T_2	T_0
6	T_6	T_5	T_3	T_1
7	T_7	T_6	T_4	T_2
8	T_8			

Example.

n	T_n				
1	2				
2	8	6			
3	35	27	21	23	
4	106	71	44	17	-6
5	238	132	61	22	5
6	453	215	83	21	-1
7	772	319	104	25	4
8	1220	448	129	28	3
9	1825	605	157		

The last column has a Fibonacci property, but the previous column does not. Hence the polynomial must have been of degree three. We identify the first terms of the Fibonacci sequence as being 3, the zero term as 4, the term with -1 subscript as -1, etc. The effect of these terms can be eliminated from the leading edge of the table to give: $23 - 5 = 18$; $21 - (-1) = 22$; $6 - 4 = 2$; $2 - 3 = -1$. Calculating the zero differences as before, the final form of the term to be found is:

$$T_n = 3n^3 - 7n^2 + 2n + 1 + V_n$$

LINEAR RECURSION RELATIONS

where $V_1 = 3$, $V_2 = 7$, and $V_{n+1} = V_n + V_{n-1}$.

PROBLEMS

1. Determine the polynomial for which $f(1) = -4$; $f(2) = 22$; $f(3) = 100$; $f(4) = 200$; $f(5) = 532$; $f(6) = 946$; $f(7) = 1532$; $f(8) = 2320$.

2. The following sequence of values correspond to terms T_1, T_2 , etc. of a sequence which is the sum of a polynomial and a Fibonacci sequence: 0, 4, 12, 29, 53, 87, 132, 192, 272, 391. Determine the polynomial and the Fibonacci sequence components.

3. The values: 13, 72, 227, 526, 1023, 1784, 2899, 4506, 6839 include a polynomial component and a geometric progression component. Determine the general form of the term of the sequence.

4. The sequence values: 4, 14, 12, 22, 20, 30, 28, 38, 36, \dots combine a polynomial and a geometric progression. Determine the general form of the term of the sequence.

5. The sequence values: 7, 19, 45, 109, 219, 395, 653, 1017, 1515 have a polynomial and a Fibonacci component. Determine the general form of the polynomial and find the Fibonacci sequence.

6. The following sequence has its terms the sum of corresponding terms of a geometric and arithmetic progression: 41, 73, 150, 407, 1384, 5241, 20618, 82075, 327852, 1310909, \dots . Identify the component sequences.

7. The following sequence has its terms the sum of corresponding terms of an arithmetic progression and a Fibonacci sequence: 8, 17, 24, 34, 45, 60, 80, 109, 152, 218, \dots . Identify the component sequences.

8. The following sequence has its terms the sum of two sequences, one geometric and the other with a general term that can be represented by a polynomial: 10, 26, 71, 201, 584, 1724, 5133, 15347, 45974, 137878, \dots . Determine the two component sequences.

9. The sequence 7, 8, 15, 33, 61, 103, 162, 245, 362, 530, \dots is the sum of a Fibonacci sequence and a sequence that can be represented by a polynomial. Find the two sequences.

10. The sequence 11, 23, 40, 75, 139, 262, 497, 951, 1832, 3551, \dots is the sum of a Fibonacci sequence and a geometric progression. Determine the two component sequences.



(See page 59 for answers to problems.)