

# SDEs in large dimension and numerical methods

## Part 1: Sampling the canonical distribution

T. Lelièvre

CERMICS - Ecole des Ponts ParisTech & Materials project-team - INRIA



RICAM Winterschool, December 2016

# Motivation

The aim of molecular dynamics simulations is to understand the relationships between the **macroscopic properties** of a molecular system and its **atomistic** features. In particular, one would like to evaluate numerically macroscopic quantities from models at the microscopic scale.

Many applications in various fields: biology, physics, chemistry, materials science.

Various models: discrete state space (kinetic Monte Carlo, Markov State Model) or **continuous state space** (Langevin).

The basic ingredient: a **potential**  $V$  which associates to a configuration  $(\mathbf{x}_1, \dots, \mathbf{x}_N) = \mathbf{x} \in \mathbb{R}^{3N_{atom}}$  an energy  $V(\mathbf{x}_1, \dots, \mathbf{x}_{N_{atom}})$ . The dimension  $d = 3N_{atom}$  is large (a few hundred thousand to millions).

## Empirical force field

Typically,  $V$  is a sum of potentials modelling interaction between two particles, three particles and four particles:

$$V = \sum_{i < j} V_1(\mathbf{x}_i, \mathbf{x}_j) + \sum_{i < j < k} V_2(\mathbf{x}_i, \mathbf{x}_j, \mathbf{x}_k) + \sum_{i < j < k < l} V_3(\mathbf{x}_i, \mathbf{x}_j, \mathbf{x}_k, \mathbf{x}_l).$$


For example,

$$V_1(\mathbf{x}_i, \mathbf{x}_j) = V_{LJ}(|\mathbf{x}_i - \mathbf{x}_j|)$$

where

$$V_{LJ}(r) = 4\epsilon \left( \left( \frac{\sigma}{r} \right)^{12} - \left( \frac{\sigma}{r} \right)^6 \right)$$

is the Lennard-Jones potential.



# Dynamics

Newton equations of motion:

$$\begin{cases} d\mathbf{X}_t = M^{-1}\mathbf{P}_t dt \\ d\mathbf{P}_t = -\nabla V(\mathbf{X}_t) dt \end{cases}$$

## Dynamics

Newton equations of motion + thermostat: Langevin dynamics:

$$\begin{cases} d\mathbf{X}_t = M^{-1}\mathbf{P}_t dt \\ d\mathbf{P}_t = -\nabla V(\mathbf{X}_t) dt - \gamma M^{-1}\mathbf{P}_t dt + \sqrt{2\gamma\beta^{-1}}d\mathbf{W}_t \end{cases}$$

where  $\gamma > 0$ . Langevin dynamics is ergodic wrt  $\mu(d\mathbf{x}) \otimes Z_p^{-1} \exp\left(-\beta \frac{\mathbf{p}^t M^{-1} \mathbf{p}}{2}\right) d\mathbf{p}$  with

$$d\mu = Z^{-1} \exp(-\beta V(\mathbf{x})) d\mathbf{x}$$

where  $Z = \int \exp(-\beta V(\mathbf{x})) d\mathbf{x}$  is the partition function and  $\beta = (k_B T)^{-1}$  is proportional to the inverse of the temperature.

## Dynamics

Newton equations of motion + thermostat: Langevin dynamics:

$$\begin{cases} d\mathbf{X}_t = M^{-1}\mathbf{P}_t dt \\ d\mathbf{P}_t = -\nabla V(\mathbf{X}_t) dt - \gamma M^{-1}\mathbf{P}_t dt + \sqrt{2\gamma\beta^{-1}}d\mathbf{W}_t \end{cases}$$

where  $\gamma > 0$ . Langevin dynamics is ergodic wrt  $\mu(d\mathbf{x}) \otimes Z_p^{-1} \exp\left(-\beta \frac{\mathbf{p}^t M^{-1} \mathbf{p}}{2}\right) d\mathbf{p}$  with

$$d\mu = Z^{-1} \exp(-\beta V(\mathbf{x})) d\mathbf{x}$$

where  $Z = \int \exp(-\beta V(\mathbf{x})) d\mathbf{x}$  is the partition function and  $\beta = (k_B T)^{-1}$  is proportional to the inverse of the temperature.

In the following, we focus on the *overdamped Langevin* (or gradient) dynamics

$$d\mathbf{X}_t = -\nabla V(\mathbf{X}_t) dt + \sqrt{2\beta^{-1}}d\mathbf{W}_t,$$

which is also ergodic wrt  $\mu$ .

## Macroscopic quantities of interest

These dynamics are used to compute macroscopic quantities:

- (i) **Thermodynamic quantities** (averages wrt  $\mu$  of some observables): stress, heat capacity, free energy,...

$$\mathbb{E}_{\mu}(\varphi(\mathbf{X})) = \int_{\mathbb{R}^d} \varphi(\mathbf{x}) \mu(d\mathbf{x}) \simeq \frac{1}{T} \int_0^T \varphi(\mathbf{X}_t) dt.$$





- (ii) **Dynamical quantities** (averages over trajectories): diffusion coefficients, viscosity, transition rates,...

$$\mathbb{E}(\mathcal{F}((\mathbf{X}_t)_{t \geq 0})) \simeq \frac{1}{M} \sum_{m=1}^M \mathcal{F}((\mathbf{X}_t^m)_{t \geq 0}).$$

**Difficulties:** (i) high-dimensional problem ( $N \gg 1$ ); (ii)  $\mathbf{X}_t$  is a **metastable process** and  $\mu$  is a multimodal measure.

# Metastability: energetic and entropic barriers

## A two-dimensional schematic picture




- 
- Slow convergence of trajectorial averages
  - Transitions between metastable states are **rare events**




## A toy model for solvation

Influence of the solvation on a dimer conformation [Dellago, Geissler].



Compact state.




Stretched state.


The particles interact through a pair potential: truncated LJ for all particles except the two monomers (black particles) which interact through a double-well potential. A slow variable is the distance between the two monomers.

# A toy example in material sciences


## The 7 atoms Lennard Jones cluster in 2D.




(a)  $C_0$ ,  $V = -12.53$



(b)  $C_1$ ,  $V = -11.50$



(c)  $C_2$ ,  $V = -11.48$



(d)  $C_3$ ,  $V = -11.40$


Figure: Low energy conformations of the Lennard-Jones cluster.

→ simulation

# Simulations of biological systems

## Unbinding of a ligand from a protein

(Diaminopyridine-HSP90, Courtesy of SANOFI)



Elementary time-step for the molecular dynamics =  $10^{-15}$  s  
Dissociation time = 0.5 s

**Challenge:** bridge the gap between timescales

# Outline


Outline of this part:

1. Definition of the **free energy** associated to a reaction coordinate.
2. **Thermodynamics integration**: A free energy computation method based on stochastic processes with constraints.
3. **Adaptive biasing techniques**: Free energy computation methods based on biased stochastic processes.

*Mathematical tools: delta measure and co-area formula, Entropy techniques and Logarithmic Sobolev Inequalities.*

Underlying question: how to properly define and quantify metastability ? Various answers: (i) rate of convergence to equilibrium; (ii) exit time from metastable states; (iii) decorrelation time; (iv) asymptotic variance of estimators.

# Reaction coordinate and free energy




## Reaction coordinate

We suppose in the following that **we know** a slow variable of **dimension 1**:  $\xi(\mathbf{X}_t)$ , where  $\xi : \mathbb{R}^d \rightarrow \mathbb{T}$  is a so-called reaction coordinate.

This reaction coordinate will be used to efficiently sample the canonical measure using two techniques: (i) constrained dynamics (thermodynamic integration) or (ii) biased dynamics (adaptive importance sampling technique).

**Free energy** will play a central role.

For example, in the 2D simple examples:  $\xi(x, y) = x$ .



## Free energy

Let us introduce two probability measures associated to  $\mu$  and  $\xi$ :

- The image of the measure  $\mu$  by  $\xi$ :

$$\xi_*\mu(dz) = \exp(-\beta A(z)) dz$$

where the free energy  $A$  is defined by:

$$A(z) = -\beta^{-1} \ln \left( \int_{\Sigma(z)} e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x}) \right),$$

with  $\Sigma(z) = \{\mathbf{x}, \xi(\mathbf{x}) = z\}$  is a (smooth) submanifold of  $\mathbb{R}^d$ , and  $\delta_{\xi(\mathbf{x})-z}(d\mathbf{x}) dz = d\mathbf{x}$ .

- The probability measure  $\mu$  conditioned to  $\xi(\mathbf{x}) = z$ :

$$\mu_{\Sigma(z)}(d\mathbf{x}) = \frac{\exp(-\beta V(\mathbf{x})) \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})}{\exp(-\beta A(z))}.$$

## Free energy (2d case)

In the simple case  $\xi(x, y) = x$ , we have:

- The image of the measure  $\mu$  by  $\xi$ :

$$\xi_*\mu(dx) = \exp(-\beta A(x)) dx$$

where the free energy  $A$  is defined by:

$$A(x) = -\beta^{-1} \ln \left( \int_{\Sigma(x)} e^{-\beta V(x,y)} dy \right)$$

and  $\Sigma(x) = \{(x, y), y \in \mathbb{R}\}$ .

- The probability measure  $\mu$  conditioned to  $\xi(x, y) = x$ :

$$\mu_{\Sigma(x)}(dy) = \frac{\exp(-\beta V(x, y)) dy}{\exp(-\beta A(x))}.$$



## The delta measure and the co-area formula

- The measure  $\delta_{\xi(\mathbf{x})-z}$  is defined by: for all test functions  $\varphi : \mathbb{T} \rightarrow \mathbb{R}$  and  $\psi : \mathbb{R}^d \rightarrow \mathbb{R}$ ,

$$\int_{\mathbb{R}^d} \varphi \circ \xi(\mathbf{x}) \psi(\mathbf{x}) d\mathbf{x} = \int_{\mathbb{T}} \varphi(z) \left( \int_{\Sigma(z)} \psi(\mathbf{x}) \delta_{\xi(\mathbf{x})-z}(d\mathbf{x}) \right) dz.$$

- The measure  $\delta_{\xi(\mathbf{x})-z}$  can be understood using a regularization procedure: for any test function  $\psi : \mathbb{R}^d \rightarrow \mathbb{R}$ ,

$$\int_{\Sigma(z)} \psi(\mathbf{x}) \delta_{\xi(\mathbf{x})-z}(d\mathbf{x}) = \lim_{\epsilon \rightarrow 0} \int_{\mathbb{R}^d} \psi(\mathbf{x}) \delta^\epsilon(\xi(\mathbf{x}) - z) d\mathbf{x}$$

where  $\lim_{\epsilon \rightarrow 0} \delta^\epsilon = \delta$  (Dirac mass at zero).

- The measure  $\delta_{\xi(\mathbf{x})-z}$  is related to the Lebesgue measure on  $\Sigma(z)$  through:

$$\delta_{\xi(\mathbf{x})-z} = |\nabla \xi|^{-1} d\sigma_{\Sigma(z)}.$$

This is the **co-area formula**. We thus have:

$$A(z) = -\beta^{-1} \ln \left( \int_{\Sigma(z)} e^{-\beta V} |\nabla \xi|^{-1} d\sigma_{\Sigma(z)} \right).$$

## Free energy: Remarks

- $A$  is the **free energy** associated with the **reaction coordinate** or collective variable  $\xi$  (angle, length, ...). The aim of many molecular dynamic simulations is to compute  $A$ .
- $A$  is defined up to an additive constant, so that it is enough to compute free energy differences, or the derivative of  $A$  (the **mean force**).
- $A(z) = -\beta^{-1} \ln Z_{\Sigma(z)}$  and  $Z_{\Sigma(z)}$  is the partition function associated with the conditioned probability measures:  

$$\mu_{\Sigma(z)} = Z_{\Sigma(z)}^{-1} e^{-\beta V} |\nabla \xi|^{-1} d\sigma_{\Sigma(z)}.$$


- If  $U = \int_{\Sigma(z)} V Z_{\Sigma(z)}^{-1} e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})$  and

$$S = -k_B \int_{\Sigma(z)} \ln \left( Z_{\Sigma(z)}^{-1} e^{-\beta V} \right) Z_{\Sigma(z)}^{-1} e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x}), \text{ then}$$

$$A = U - TS \text{ (since } \beta^{-1} = k_B T \text{).}$$

## Free energy on a simple example


What is free energy ? The simple example of the solvation of a dimer. (Profiles computed using thermodynamic integration.)




The density of the solvent molecules is lower on the left than on the right. At high (resp. low) density, the compact state is more (resp. less) likely. The “free energy barrier” is higher at high density than at low density. Related question: interpretation of the free energy barrier in terms of dynamics ?

# Free energy calculation techniques


There are many free energy calculation techniques:




(a) Thermodynamic integration.



(b) Histogram method.




(c) Non equilibrium dynamics.



(d) Adaptive dynamics.

# Thermodynamic integration



## Ingredient 1: the mean force

Thermodynamic integration is based on two ingredients:

Ingredient 1: The derivative  $A'(z)$  can be obtained by sampling the conditional probability measure  $\mu_{\Sigma(z)}$  (*Sprink, Ciccotti, Kapral, Vanden-Eijnden, E, den Otter, ...*)

$$\begin{aligned}
 A'(z) &= Z_{\Sigma(z)}^{-1} \int_{\Sigma(z)} \left( \frac{\nabla V \cdot \nabla \xi}{|\nabla \xi|^2} - \beta^{-1} \operatorname{div} \left( \frac{\nabla \xi}{|\nabla \xi|^2} \right) \right) e^{-\beta V} |\nabla \xi|^{-1} d\sigma_{\Sigma(z)} \\
 &= \int_{\Sigma(z)} f d\mu_{\Sigma(z)}
 \end{aligned}$$

where  $f = \frac{\nabla V \cdot \nabla \xi}{|\nabla \xi|^2} - \beta^{-1} \operatorname{div} \left( \frac{\nabla \xi}{|\nabla \xi|^2} \right)$ . Another equivalent expression:

$$A'(z) = Z_{\Sigma(z)}^{-1} \int_{\Sigma(z)} \frac{\nabla \xi}{|\nabla \xi|^2} \cdot \left( \nabla \tilde{V} + \beta^{-1} \mathbf{H} \right) \exp(-\beta \tilde{V}) d\sigma_{\Sigma(z)}$$

where  $\tilde{V} = V + \beta^{-1} \ln |\nabla \xi|$  and  $\mathbf{H} = -\nabla \cdot \left( \frac{\nabla \xi}{|\nabla \xi|} \right) \frac{\nabla \xi}{|\nabla \xi|}$  is the mean curvature vector.

## Ingredient 1: the mean force

In the simple case  $\xi(x, y) = x$ , remember that

$$A(x) = -\beta^{-1} \ln \left( \int_{\Sigma(x)} e^{-\beta V(x,y)} dy \right),$$

so that

$$\begin{aligned} A'(x) &= \frac{\int_{\Sigma(x)} \partial_x V e^{-\beta V(x,y)} dy}{\int_{\Sigma(x)} e^{-\beta V(x,y)} dy} \\ &= \int_{\Sigma(x)} \partial_x V d\mu_{\Sigma(x)}. \end{aligned}$$

## Ingredient 1: the mean force

*Proof in the general case* :  $A'(z) = -\beta^{-1} \frac{d}{dz} \frac{\int_{\Sigma(z)} \exp(-\beta V) \delta_{\xi(\mathbf{x})-z}(\mathbf{d}\mathbf{x})}{\int_{\Sigma(z)} \exp(-\beta V) \delta_{\xi(\mathbf{x})-z}(\mathbf{d}\mathbf{x})}$

and

$$\begin{aligned}
 & \int_{\mathbb{T}} \left( \int_{\Sigma(z)} \exp(-\beta V) \delta_{\xi(\mathbf{x})-z}(\mathbf{d}\mathbf{x}) \right)' \phi(z) dz \\
 &= - \int_{\mathbb{T}} \int_{\Sigma(z)} \exp(-\beta V) \delta_{\xi(\mathbf{x})-z}(\mathbf{d}\mathbf{x}) \phi'(z) dz \\
 &= - \int_{\mathbb{T}} \int_{\Sigma(z)} \exp(-\beta V) \phi' \circ \xi \delta_{\xi(\mathbf{x})-z}(\mathbf{d}\mathbf{x}) dz \\
 &= - \int_{\mathbb{R}^d} \exp(-\beta V) \phi' \circ \xi d\mathbf{x} = - \int_{\mathbb{R}^d} \exp(-\beta V) \nabla(\phi \circ \xi) \cdot \frac{\nabla \xi}{|\nabla \xi|^2} d\mathbf{x} \\
 &= \int_{\mathbb{R}^d} \nabla \cdot \left( \exp(-\beta V) \frac{\nabla \xi}{|\nabla \xi|^2} \right) \phi \circ \xi d\mathbf{x} \\
 &= \int_{\mathbb{T}} \int_{\Sigma(z)} \left( -\beta \frac{\nabla V \cdot \nabla \xi}{|\nabla \xi|^2} + \nabla \cdot \left( \frac{\nabla \xi}{|\nabla \xi|^2} \right) \right) \exp(-\beta V) \delta_{\xi(\mathbf{x})-z}(\mathbf{d}\mathbf{x}) \phi(z) dz
 \end{aligned}$$



## Ingredient 2: constrained dynamics

Ingredient 2: It is possible to sample the conditioned probability measure  $\mu_{\Sigma(z)} = Z_{\Sigma(z)}^{-1} \exp(-\beta \tilde{V}) d\sigma_{\Sigma(z)}$  by considering the following **rigidly constrained dynamics**:

$$\text{(RCD)} \quad \begin{cases} d\mathbf{X}_t = -\nabla \tilde{V}(\mathbf{X}_t) dt + \sqrt{2\beta^{-1}} d\mathbf{W}_t + \nabla \xi(\mathbf{X}_t) d\Lambda_t \\ d\Lambda_t \text{ such that } \xi(\mathbf{X}_t) = z \end{cases}$$

The Lagrange multiplier writes  $d\Lambda_t = d\Lambda_t^m + d\Lambda_t^f$ , with

$$d\Lambda_t^m = -\sqrt{2\beta^{-1}} \frac{\nabla \xi}{|\nabla \xi|^2}(\mathbf{X}_t) \cdot d\mathbf{W}_t \text{ and}$$

$$d\Lambda_t^f = \frac{\nabla \xi}{|\nabla \xi|^2} \cdot \left( \nabla \tilde{V} + \beta^{-1} \mathbf{H} \right) (\mathbf{X}_t) dt = f(\mathbf{X}_t) dt$$

## Ingredient 2: constrained dynamics

Equivalently, the rigidly constrained dynamics writes:

$$(RCD) \quad d\mathbf{X}_t = P(\mathbf{X}_t) \left( -\nabla \tilde{V}(\mathbf{X}_t) dt + \sqrt{2\beta^{-1}} d\mathbf{W}_t \right) + \beta^{-1} \mathbf{H}(\mathbf{X}_t) dt$$

where  $P(\mathbf{x})$  is the orthogonal projection operator on  $T_x(\Sigma(\xi(\mathbf{x})))$ :

$$P(\mathbf{x}) = \text{Id} - \mathbf{n}(\mathbf{x}) \otimes \mathbf{n}(\mathbf{x}),$$

with  $\mathbf{n}$  the unit normal vector:  $\mathbf{n}(\mathbf{x}) = \frac{\nabla \xi}{|\nabla \xi|}(\mathbf{x})$ .

(RCD) can also be written using the [Stratonovitch product](#):

$$d\mathbf{X}_t = -P(\mathbf{X}_t) \nabla \tilde{V}(\mathbf{X}_t) dt + \sqrt{2\beta^{-1}} P(\mathbf{X}_t) \circ d\mathbf{W}_t.$$

One can check that  $\xi(\mathbf{X}_t)$  is constant if  $\mathbf{X}_t$  satisfies (RCD).

## Ingredient 2: constrained dynamics

[G. Ciccotti, TL, E. Vanden-Einjen, 2008] Assume wlg that  $z = 0$ . The probability  $\mu_{\Sigma(0)}$  is the **unique invariant measure** with support in  $\Sigma(0)$  for (RCD).

**Proposition:** Let  $\mathbf{X}_t$  be the solution to (RCD) such that the law of  $\mathbf{X}_0$  is  $\mu_{\Sigma(0)}$ . Then, for all smooth function  $\phi$  and for all time  $t > 0$ ,

$$\mathbb{E}(\phi(\mathbf{X}_t)) = \int \phi d\mu_{\Sigma(0)}.$$

*Proof:* Introduce the infinitesimal generator and apply **the divergence theorem on submanifolds** :  $\forall \phi \in \mathcal{C}^1(\mathbb{R}^d, \mathbb{R}^d)$ ,

$$\int \operatorname{div}_{\Sigma(0)}(\phi) d\sigma_{\Sigma(0)} = - \int \mathbf{H} \cdot \phi d\sigma_{\Sigma(0)},$$

where  $\operatorname{div}_{\Sigma(0)}(\phi) = \operatorname{tr}(P\nabla\phi)$ .

## Thermodynamic integration

Using the two ingredients above,  $A'(z) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(\mathbf{X}_t) dt$ , where  $\mathbf{X}_t$  satisfies (RCD) and  $\xi(\mathbf{X}_0) = z$ . The free energy profile is then obtained by **thermodynamic integration**:

$$A(z) - A(0) = \int_0^z A'(z) dz \simeq \sum_{i=0}^K \omega_i A'(z_i).$$

Notice that there is actually no need to compute  $f$  in practice since **the mean force can be obtained by averaging the Lagrange multipliers**:

$$A'(z) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T d\Lambda_t = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T d\Lambda_t^f$$

since  $d\Lambda_t = d\Lambda_t^m + d\Lambda_t^f$ , with  $d\Lambda_t^m = -\sqrt{2\beta^{-1}} \frac{\nabla \xi}{|\nabla \xi|^2}(\mathbf{X}_t) \cdot d\mathbf{W}_t$  and  $d\Lambda_t^f = f(\mathbf{X}_t) dt$ .

## Discretization of (RCD)

The two following schemes are consistent with (RCD):

$$(S1) \left\{ \begin{array}{l} \mathbf{X}_{n+1} = \mathbf{X}_n - \nabla \tilde{V}(\mathbf{X}_n) \Delta t + \sqrt{2\beta^{-1}} \Delta \mathbf{W}_n + \lambda_n \nabla \xi(\mathbf{X}_{n+1}), \\ \text{with } \lambda_n \in \mathbb{R} \text{ such that } \xi(\mathbf{X}_{n+1}) = 0, \end{array} \right.$$

$$(S2) \left\{ \begin{array}{l} \mathbf{X}_{n+1} = \mathbf{X}_n - \nabla \tilde{V}(\mathbf{X}_n) \Delta t + \sqrt{2\beta^{-1}} \Delta \mathbf{W}_n + \lambda_n \nabla \xi(\mathbf{X}_n), \\ \text{with } \lambda_n \in \mathbb{R} \text{ such that } \xi(\mathbf{X}_{n+1}) = 0, \end{array} \right.$$

where  $\Delta \mathbf{W}_n = \mathbf{W}_{(n+1)\Delta t} - \mathbf{W}_{n\Delta t}$ . The constraint is exactly satisfied (important for longtime computations). An approximation of  $A'(0) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T d\Lambda_t$  is:

$$\lim_{T \rightarrow \infty} \lim_{\Delta t \rightarrow 0} \frac{1}{T} \sum_{n=1}^{T/\Delta t} \lambda_n = A'(0).$$

## Error analysis

[Faou, TL, Mathematics of Computation, 2010] Using classical techniques (Talay-Tubaro like proof), one can check that the ergodic measure  $\mu_{\Sigma(0)}^{\Delta t}$  sampled by the Markov chain  $(\mathbf{X}_n)_{n \geq 0}$  is an approximation of order one of  $\mu_{\Sigma(0)}$ : for all smooth functions  $g : \Sigma(0) \rightarrow \mathbb{R}$ ,

$$\left| \int_{\Sigma(0)} g d\mu_{\Sigma(0)}^{\Delta t} - \int_{\Sigma(0)} g d\mu_{\Sigma(0)} \right| \leq C \Delta t.$$

## Rigidly and softly constrained dynamics

Another way to constrain the overdamped Langevin dynamics to  $\Sigma(0)$  is to add a constraining potential (soft constraint):

$$d\mathbf{X}_t^\eta = -\nabla V(\mathbf{X}_t^\eta) dt - \frac{1}{2\eta} \nabla(\xi^2)(\mathbf{X}_t^\eta) dt + \sqrt{2\beta^{-1}} d\mathbf{W}_t$$

One can show that  $\lim_{\eta \rightarrow 0} \mathbf{X}_t^\eta = \mathbf{X}_t$  (in  $L_{t \in [0, \tau]}^\infty(L_\omega^2)$ -norm) where  $\mathbf{X}_t$  satisfies (RCD). Notice that we used  $V$  and not  $\tilde{V}$  in the softly constrained dynamics.

The statistics associated with the dynamics where the constraints are rigidly imposed and the dynamics where the constraints are softly imposed are different: “a stiff spring  $\neq$  a rigid rod” (*van Kampen, Hinch,...*).

## Back to the sampling of $\mu$

TI yields a way to compute  $\int_{\mathbb{R}^d} \phi d\mu$ :

$$\begin{aligned}
 \int_{\mathbb{R}^d} \phi d\mu &= Z^{-1} \int_{\mathbb{R}^d} \phi e^{-\beta V} d\mathbf{x} \\
 &= Z^{-1} \int_{\mathbb{T}} \int_{\Sigma(z)} \phi e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x}) \\
 &= Z^{-1} \int_{\mathbb{T}} \frac{\int_{\Sigma(z)} \phi e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})}{\int_{\Sigma(z)} e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})} \int_{\Sigma(z)} e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x}) dz \\
 &= \left( \int_{\mathbb{T}} e^{-\beta A(z)} dz \right)^{-1} \int_{\mathbb{T}} \left( \int_{\Sigma(z)} \phi d\mu_{\Sigma(z)} \right) e^{-\beta A(z)} dz
 \end{aligned}$$

where, we recall,  $\Sigma(z) = \{\mathbf{x}, \xi(\mathbf{x}) = z\}$ ,

$A(z) = -\beta^{-1} \ln \left( \int_{\Sigma(z)} e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x}) \right)$  and

$\mu_{\Sigma(z)} = e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x}) / \int_{\Sigma(z)} e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})$ .



## Generalization to Langevin dynamics

Interests: (i) Newton's equations of motion are more "natural"; (ii) leads to numerical schemes which sample the constrained measure without time discretization error; (iii) seems to be more robust wrt the timestep choice.

$$\begin{cases} dq_t = M^{-1} p_t dt \\ dp_t = -\nabla V(q_t) dt - \gamma M^{-1} p_t dt + \sqrt{2\gamma\beta^{-1}} dW_t + \nabla \xi(q_t) d\lambda_t \\ \xi(q_t) = z. \end{cases}$$

The probability measure sampled by this dynamics is

$$\mu_{T^*\Sigma(z)}(dqdp) = Z^{-1} \exp(-\beta H(q, p)) \sigma_{T^*\Sigma(z)}(dqdp)$$

where  $H(q, p) = V(q) + \frac{1}{2} p^T M^{-1} p$ .

## Generalization to Langevin dynamics

The marginal of  $\mu_{T^*\Sigma(z)}(dqdp)$  in  $q$  writes:

$$\nu_{\Sigma(z)}^M = \frac{1}{Z} \exp(-\beta V(q)) \sigma_{\Sigma(z)}^M(dq) \neq \frac{1}{Z} \exp(-\beta V(q)) \delta_{\xi(q)-z}(dq).$$

Thus, the “free energy” which is naturally computed by this dynamics is

$$A^M(z) = -\beta^{-1} \ln \left( \int_{\Sigma(z)} \exp(-\beta V(q)) \sigma_{\Sigma(z)}^M(dq) \right).$$

The original free energy may be recovered from the relation: for  $G_M = \nabla \xi^T M^{-1} \nabla \xi$ ,

$$A(z) - A^M(z) = -\beta^{-1} \ln \left( \int_{\Sigma(z)} \det(G_M)^{-1/2} d\nu_{\Sigma(z)}^M \right).$$

## Generalization to Langevin dynamics

Moreover, one can check that:

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T d\lambda_t = (A^M)'(z).$$

**Discretization:** A natural numerical scheme is obtained by a splitting technique:

- 1/2 midpoint Euler on the fluctuation-dissipation part,
- 1 Verlet step on the Hamiltonian part (RATTLE scheme) and
- 1/2 midpoint Euler on the fluctuation-dissipation part.

## Generalization to Langevin dynamics

$$\left\{ \begin{array}{l} p^{n+1/4} = p^n - \frac{\Delta t}{4} \gamma M^{-1}(p^n + p^{n+1/4}) + \sqrt{\frac{\Delta t}{2}} \sigma G^n + \nabla \xi(q^n) \lambda^{n+1/4}, \\ \nabla \xi(q^n)^T M^{-1} p^{n+1/4} = 0, \end{array} \right.$$

$$\left\{ \begin{array}{l} p^{n+1/2} = p^{n+1/4} - \frac{\Delta t}{2} \nabla V(q^n) + \nabla \xi(q^n) \lambda^{n+1/2}, \\ q^{n+1} = q^n + \Delta t M^{-1} p^{n+1/2}, \end{array} \right.$$

$$\left\{ \begin{array}{l} \xi(q^{n+1}) = z, \end{array} \right.$$

$$\left\{ \begin{array}{l} p^{n+3/4} = p^{n+1/2} - \frac{\Delta t}{2} \nabla V(q^{n+1}) + \nabla \xi(q^{n+1}) \lambda^{n+3/4}, \\ \nabla \xi(q^{n+1})^T M^{-1} p^{n+3/4} = 0, \end{array} \right.$$

$$\left\{ \begin{array}{l} p^{n+1} = p^{n+3/4} - \frac{\Delta t}{4} \gamma M^{-1}(p^{n+3/4} + p^{n+1}) + \sqrt{\frac{\Delta t}{2}} \sigma G^{n+1/2} \\ \quad + \nabla \xi(q^{n+1}) \lambda^{n+1}, \\ \nabla \xi(q^{n+1})^T M^{-1} p^{n+1} = 0, \end{array} \right.$$

and  $\lim_{T \rightarrow \infty} \lim_{\Delta t \rightarrow 0} \frac{1}{T} \sum_{n=1}^{T/\Delta t} (\lambda^{n+1/2} + \lambda^{n+3/4}) = (A^M)'(z)$ .

## Generalization to Langevin dynamics

Using the symmetry of the Verlet step, it is easy to **add a Metropolization step** to the previous numerical scheme, thus removing the time discretization error. Indeed, the proposal  $(q^n, p^n) \mapsto (q^{n+1}, -p^{n+1})$  is symmetric, so that the Metropolis Hastings acceptance ratio is simply  $\exp(-\beta(H(q^{n+1}, p^{n+1}) - H(q^n, p^n)))$ .

For this modified scheme, one can prove that

$$\lim_{\Delta t \rightarrow 0} \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{n=1}^{T/\Delta t} \left( \lambda^{n+1/2} + \lambda^{n+3/4} \right) = (A^M)'(z).$$

## Generalization to Langevin dynamics

By choosing  $M = \Delta t \gamma / 4 = \text{Id}$ , this leads to an original sampling scheme in the configuration space ([generalized Hybrid Monte Carlo scheme](#)).

Notice that it is not clear how to use such a Metropolization step for the dynamics (RCD) since the proposal kernel is not symmetric, and does not admit any simple analytical expression.

**Algorithm:** Let us introduce  $R_{\Delta t}$  which is such that, if  $(q^n, p^n) \in T^*\Sigma(z)$ , and  $|p^n|^2 \leq R_{\Delta t}$ , one step of the RATTLE scheme is well defined (*i.e.* there exists a unique solution to the constrained problem).

Then the GHMC scheme writes ( $M = \Delta t \gamma / 4 = \text{Id}$ ):

## Generalization to Langevin dynamics

Consider an initial configuration  $q^0 \in \Sigma(z)$ . Iterate on  $n \geq 0$ ,

1. Sample a random vector in the tangent space  $T_{q^n}\Sigma(z)$  ( $\nabla\xi(q^n)^T p^n = 0$ ):

$$p^n = \beta^{-1/2} P(q^n) G^n,$$

where  $(G^n)_{n \geq 0}$  are i.i.d. standard random Gaussian variables, and compute the energy  $E^n = \frac{1}{2}|p^n|^2 + V(q^n)$  of the configuration  $(q^n, p^n)$ ;

2. If  $|p^n|^2 > R_{\Delta t}$ , set  $E^{n+1} = +\infty$  and go to (3); otherwise perform one integration step of the RATTLE scheme:

$$\begin{cases} p^{n+1/2} = p^n - \frac{\Delta t}{2} \nabla V(q^n) + \nabla\xi(q^n) \lambda^{n+1/2}, \\ \tilde{q}^{n+1} = q^n + \Delta t p^{n+1/2}, \\ \xi(\tilde{q}^{n+1}) = z, \\ \tilde{p}^{n+1} = p^{n+1/2} - \frac{\Delta t}{2} \nabla V(\tilde{q}^{n+1}) + \nabla\xi(\tilde{q}^{n+1}) \lambda^{n+1}, \\ \nabla\xi(\tilde{q}^{n+1})^T \tilde{p}^{n+1} = 0; \end{cases}$$

## Generalization to Langevin dynamics

3. If  $|\tilde{p}^{n+1}|^2 > R_{\Delta t}$ , set  $E^{n+1} = +\infty$ ; otherwise compute the energy  $E^{n+1} = \frac{1}{2}|\tilde{p}^{n+1}|^2 + V(\tilde{q}^{n+1})$  of the new phase-space configuration. Accept the proposal and set  $q^{n+1} = \tilde{q}^{n+1}$  with probability

$$\min\left(\exp(-\beta(E^{n+1} - E^n)), 1\right);$$

otherwise, reject and set  $q^{n+1} = q^n$ .


It can be checked that the probability measure

$$\nu_{\Sigma(z)}^M = \frac{1}{Z} \exp(-\beta V(q)) \sigma_{\Sigma(z)}^M(dq)$$

is invariant for the Markov Chain  $(q^n)_{n \geq 0}$ .



# Adaptive biasing techniques





# Adaptive biasing techniques

We suppose again that we know a slow variable of dimension 1:  $\xi(\mathbf{X}_t)$ , where  $\xi : \mathbb{R}^d \rightarrow \mathbb{T}$  is a so-called reaction coordinate.

This reaction coordinate will be used to bias the dynamics (adaptive importance sampling technique), using the free energy  $A$  associated with the reaction coordinate  $\xi$ .

For example, in the 2D simple examples:  $\xi(x, y) = x$ .



## Adaptive biasing techniques

The bottom line of adaptive methods is the following: for “well chosen”  $\xi$  the potential  $V - A \circ \xi$  is less rugged than  $V$ . Indeed, by construction  $\xi_* \exp(-\beta(V - A \circ \xi)) = 1_{\mathbb{T}}$ .

Problem:  $A$  is unknown ! Idea: use a time dependent potential of the form

$$\mathcal{V}_t(\mathbf{x}) = V(\mathbf{x}) - A_t(\xi(\mathbf{x}))$$


where  $A_t$  is an approximation at time  $t$  of  $A$ , given the configurations visited so far.

Hopes:

- build a dynamics which goes quickly to equilibrium,
- compute free energy profiles.





Wang-Landau, ABF, metadynamics: *Darve, Pohorille, Hénin, Chipot, Laio, Parrinello, Wang, Landau,...*

# Free energy biased dynamics (1/2)



A 2D example of a free energy biased trajectory: **energetic barrier**.

# Free energy biased dynamics (2/2)



A 2D example of a free energy biased trajectory: **entropic barrier**.

## Updating strategies

How to update  $A_t$  ? Two methods depending on whether  $A'_t$  (Adaptive Biasing Force) or  $A_t$  (Adaptive Biasing Potential) is approximated.

To avoid geometry problem, an extended configurational space  $(\mathbf{x}, z) \in \mathbb{R}^{n+1}$  may be considered, together with the **meta-potential**:

$$V^k(\mathbf{x}, z) = V(\mathbf{x}) + k(z - \xi(\mathbf{x}))^2.$$

Choosing  $(\mathbf{x}, z) \mapsto z$  as a reaction coordinate, the associated free energy  $A^k$  is close to  $A$  (in the limit  $k \rightarrow \infty$ , up to an additive constant).

Adaptive algorithms used in molecular dynamics fall into one of these four possible combinations [TL, M. Rousset, G. Stoltz, J Chem Phys, 2007]:

	$A'_t$	$A_t$
$V$	ABF	Wang-Landau
$V^k$	...	metadynamics

# The ABF method

For the **Adaptive Biasing Force** (ABF) method, the idea is to use the formula

$$A'(z) = \frac{\int_{\Sigma(z)} \left( \frac{\nabla V \cdot \nabla \xi}{|\nabla \xi|^2} - \beta^{-1} \operatorname{div} \left( \frac{\nabla \xi}{|\nabla \xi|^2} \right) \right) e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})}{\int_{\Sigma(z)} e^{-\beta V} \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})}$$
$$= \int_{\Sigma(z)} f d\mu_{\Sigma(z)} = \mathbb{E}_{\mu}(f(\mathbf{X}) | \xi(\mathbf{X}) = z).$$

The **mean force**  $A'(z)$  is the average of  $f$  with respect to  $\mu_{\Sigma(z)}$ .

## The ABF method

In the simple case  $\xi(x, y) = x$ , remember that

$$A(x) = -\beta^{-1} \ln \left( \int e^{-\beta V(x, y)} dy \right),$$

so that

$$\begin{aligned} A'(x) &= \frac{\int_{\Sigma(x)} \partial_x V e^{-\beta V(x, y)} dy}{\int_{\Sigma(x)} e^{-\beta V(x, y)} dy} \\ &= \int \partial_x V d\mu_{\Sigma(x)}. \end{aligned}$$

Notice that actually, whatever  $A_t$  is,

$$A'(z) = \frac{\int_{\Sigma(z)} f e^{-\beta(V - A_t \circ \xi)} \delta_{\xi(x) - z}(d\mathbf{x})}{\int_{\Sigma(z)} e^{-\beta(V - A_t \circ \xi)} \delta_{\xi(x) - z}(d\mathbf{x})}.$$



## The ABF method

Thus, we would like to simulate:

$$\begin{cases} d\mathbf{X}_t = -\nabla(V - A \circ \xi)(\mathbf{X}_t) dt + \sqrt{2\beta^{-1}} d\mathbf{W}_t, \\ A'(z) = \mathbb{E}_\mu(f(\mathbf{X}) | \xi(\mathbf{X}) = z) \end{cases}$$

but  $A$  is unknown...

## The ABF method

The ABF dynamics is then:

$$\begin{cases} d\mathbf{X}_t = -\nabla(V - A_t \circ \xi)(\mathbf{X}_t) dt + \sqrt{2\beta^{-1}} d\mathbf{W}_t, \\ A'_t(z) = \mathbb{E}(f(\mathbf{X}_t) | \xi(\mathbf{X}_t) = z). \end{cases}$$

## The ABF method

The ABF dynamics is then:

$$\begin{cases} d\mathbf{X}_t = -\nabla(V - A_t \circ \xi)(\mathbf{X}_t) dt + \sqrt{2\beta^{-1}} d\mathbf{W}_t, \\ A'_t(z) = \mathbb{E}(f(\mathbf{X}_t) | \xi(\mathbf{X}_t) = z). \end{cases}$$

The associated (nonlinear) Fokker-Planck equation writes:

$$\begin{cases} \partial_t \psi = \operatorname{div} (\nabla(V - A_t \circ \xi)\psi + \beta^{-1} \nabla \psi), \\ A'_t(z) = \frac{\int f \psi \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})}{\int \psi \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})}, \end{cases}$$

where  $\mathbf{X}_t \sim \psi(t, \mathbf{x}) d\mathbf{x}$ .

## The ABF method

The ABF dynamics is then:

$$\begin{cases} d\mathbf{X}_t = -\nabla(V - A_t \circ \xi)(\mathbf{X}_t) dt + \sqrt{2\beta^{-1}} d\mathbf{W}_t, \\ A'_t(z) = \mathbb{E}(f(\mathbf{X}_t) | \xi(\mathbf{X}_t) = z). \end{cases}$$



The associated (nonlinear) Fokker-Planck equation writes:

$$\begin{cases} \partial_t \psi = \operatorname{div} (\nabla(V - A_t \circ \xi)\psi + \beta^{-1} \nabla \psi), \\ A'_t(z) = \frac{\int f \psi \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})}{\int \psi \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})}, \end{cases}$$



where  $\mathbf{X}_t \sim \psi(t, \mathbf{x}) d\mathbf{x}$ .

Questions: Does  $A'_t$  converge to  $A'$ ? What did we gain compared to the original gradient dynamics?

# Back to the 2D example




Left: the 2D potential – energetic barrier; Right: average exit time from the left well




The ABF trajectory (right: zoom on the first 2500 iterations)

## Back to the toy example for solvation



Compact state.



Stretched state.

The reaction coordinate  $\xi$  is the distance between the two monomers.  $\rightarrow$  **simulation**

## Longtime convergence and entropy (1/3)

Recall the original gradient dynamics:

$$d\mathbf{Q}_t = -\nabla V(\mathbf{Q}_t) dt + \sqrt{2\beta^{-1}} d\mathbf{W}_t.$$

The associated (linear) Fokker-Planck equation writes:

$$\partial_t \phi = \text{div} (\nabla V \phi + \beta^{-1} \nabla \phi).$$

where  $\mathbf{Q}_t \sim \phi(t, \mathbf{q}) d\mathbf{q}$ .

The metastable behaviour of  $\mathbf{Q}_t$  is related to the multimodality of  $\mu$ , which can be quantified through the **rate of convergence of  $\phi$  to  $\phi_\infty = Z^{-1} \exp(-\beta V)$** .

A classical approach for partial differential equations (PDEs):  
entropy techniques.

## Longtime convergence and entropy (2/3)

Notice that the Fokker-Planck equation rewrites

$$\partial_t \phi = \beta^{-1} \operatorname{div} \left( \phi_\infty \nabla \left( \frac{\phi}{\phi_\infty} \right) \right).$$

Let us introduce **the entropy**:

$$E(t) = H(\phi(t, \cdot) | \phi_\infty) = \int \ln \left( \frac{\phi}{\phi_\infty} \right) \phi.$$

We have (Csiszár-Kullback inequality):

$$\|\phi(t, \cdot) - \phi_\infty\|_{L^1} \leq \sqrt{2E(t)}.$$



## Longtime convergence and entropy (3/3)

$$\begin{aligned}
 \frac{dE}{dt} &= \int \ln \left( \frac{\phi}{\phi_\infty} \right) \partial_t \phi \\
 &= \beta^{-1} \int \ln \left( \frac{\phi}{\phi_\infty} \right) \operatorname{div} \left( \phi_\infty \nabla \left( \frac{\phi}{\phi_\infty} \right) \right) \\
 &= -\beta^{-1} \int \left| \nabla \ln \left( \frac{\phi}{\phi_\infty} \right) \right|^2 \phi =: -\beta^{-1} I(\phi(t, \cdot) | \phi_\infty).
 \end{aligned}$$

If  $V$  is such that the following **Logarithmic Sobolev inequality** (LSI( $R$ )) holds:  $\forall \phi$  pdf,

$$H(\phi | \phi_\infty) \leq \frac{1}{2R} I(\phi | \phi_\infty)$$

then  $E(t) \leq E(0) \exp(-2\beta^{-1}Rt)$  and thus  $\phi$  converges to  $\phi_\infty$  exponentially fast with rate  $\beta^{-1}R$ .

**Metastability**  $\iff$  small  $R$

## Efficiency of thermodynamic integration

With thermodynamic integration, the conditional measures  $\mu_{\Sigma(z)}$  are sampled rather than the original Gibbs measure  $\mu$ . The long-time behaviour of the constrained dynamics (RCD) will be essentially limited by the LSI constant  $\rho(z)$  of the conditional measures  $\mu_{\Sigma(z)}$  (to be compared with the LSI constant  $R$  of the original measure  $\mu$ ). For well-chosen  $\xi$ ,  $\rho(z) \gg R$ , which explains the efficiency of the whole procedure.

## Convergence of ABF (1/4)

A convergence result [TL, M. Rousset, G. Stoltz, *Nonlinearity* 2008]: Recall the ABF Fokker-Planck equation:

$$\begin{cases} \partial_t \psi = \operatorname{div} (\nabla(V - A_t \circ \xi)\psi + \beta^{-1} \nabla \psi), \\ A'_t(z) = \frac{\int f \psi \delta_{\xi(x)-z}(dx)}{\int \psi \delta_{\xi(x)-z}(dx)}. \end{cases}$$

Suppose:

(H1) “Ergodicity” of the microscopic variables: the conditional probability measures  $\mu_{\Sigma(z)}$  satisfy a LSI( $\rho$ ),

(H2) Bounded coupling:  $\|\nabla_{\Sigma(z)} f\|_{L^\infty} < \infty$ ,

then

$$\|A'_t - A'\|_{L^2} \leq C \exp(-\beta^{-1} \min(\rho, r)t).$$

The rate of convergence is limited by:

- the rate  $r$  of convergence of  $\bar{\psi} = \int \psi \delta_{\xi(x)-z}(dx)$  to  $\overline{\psi_\infty}$ ,
- the LSI constant  $\rho$  (the real limitation).

## Convergence of ABF (2/4)

In summary:

- Original gradient dynamics:  $\exp(-\beta^{-1}Rt)$  where  $R$  is the LSI constant for  $\mu$ ;
- ABF dynamics:  $\exp(-\beta^{-1}\rho t)$  where  $\rho$  is the LSI constant for the conditioned probability measures  $\mu_{\Sigma(z)}$ .

If  $\xi$  is well chosen,  $\rho \gg R$ : the free energy can be computed very efficiently.

Two ingredients of the proof:

(1) The marginal  $\bar{\psi}(t, z) = \int \psi(t, \mathbf{x}) \delta_{\xi(\mathbf{x})-z}(d\mathbf{x})$  satisfies a closed PDE:

$$\partial_t \bar{\psi} = \beta^{-1} \partial_{z,z} \bar{\psi} \text{ on } \mathbb{T},$$

and thus,  $\bar{\psi}$  converges towards  $\overline{\psi_\infty} \equiv \mathbf{1}$ , with exponential speed  $C \exp(-4\pi^2 \beta^{-1} t)$ . (Here,  $r = 4\pi^2$ ).

## Convergence of ABF (3/4)

(2) The total entropy can be decomposed as [N. Grunewald, F. Otto, C. Villani, M. Westdickenberg, Ann. IHP, 2009]:

$$E = E_M + E_m$$

where

The total entropy is  $E = H(\psi|\psi_\infty)$ ,

The macroscopic entropy is  $E_M = H(\bar{\psi}|\bar{\psi}_\infty)$ ,

The microscopic entropy is

$$E_m = \int H\left(\psi(\cdot|\xi(\mathbf{x}) = z) \middle| \psi_\infty(\cdot|\xi(\mathbf{x}) = z)\right) \bar{\psi}(z) dz.$$



We already know that  $E_M$  goes to zero: it remains only to consider  $E_m$ ...

## Convergence of ABF (4/4)

Other results based on this set of assumptions:

- [TL, JFA 2008] LSI for the cond. meas.  $\mu_{\Sigma(z)}$   
 + LSI for the marginal  $\bar{\mu}(dz) = \xi_* \mu(dz)$   
 + bdd coupling ( $\|\nabla_{\Sigma(z)} f\|_{L^\infty} < \infty$ )  $\implies$  LSI for  $\mu$ .
- [F. Legoll, TL, Nonlinearity, 2010] Effective dynamics for  $\xi(\mathbf{Q}_t)$ . Uniform control in time:

$$H(\mathcal{L}(\xi(\mathbf{Q}_t))|\mathcal{L}(z_t)) \leq C \left( \frac{\|\nabla_{\Sigma(z)} f\|_{L^\infty}}{\rho} \right)^2 H(\mathcal{L}(\mathbf{Q}_0)|\mu).$$



## Discretization of ABF

Discretization of adaptive methods can be done using two (complementary) approaches:

- Use empirical means over many replicas (interacting particle system):

$$\mathbb{E}(f(\mathbf{X}_t) | \xi(\mathbf{X}_t) = z) \simeq \frac{\sum_{m=1}^N f(\mathbf{X}_t^{m,N}) \delta^\alpha(\xi(\mathbf{X}_t^{m,N}) - z)}{\sum_{m=1}^N \delta^\alpha(\xi(\mathbf{X}_t^{m,N}) - z)}.$$

This approach is easy to parallelize, flexible (selection mechanisms) and efficient in cases with multiple reactive paths. [TL, M. Rousset, G. Stoltz, 2007; C. Chipot, TL, K. Minoukadeh, 2010 ; TL, K. Minoukadeh, 2010]

- Use trajectorial averages along a single path:

$$\mathbb{E}(f(\mathbf{X}_t) | \xi(\mathbf{X}_t) = z) \simeq \frac{\int_0^t f(\mathbf{X}_s) \delta^\alpha(\xi(\mathbf{X}_s) - z) ds}{\int_0^t \delta^\alpha(\xi(\mathbf{X}_s) - z) ds}.$$

The longtime behavior is much more difficult to analyze.

# ABF: Current developments and open problems

- Avoid the computation of  $\xi$ : extended-ABF
- Projection on a gradient of the mean force (Helmholtz decomposition)
- Reaction coordinates in larger dimension: exchange bias, separated representations
- Extension of the analysis to the Langevin dynamics
- Extension of the analysis to approximations of the mean force or the free energy based on time averages



# Other techniques to compute thermodynamic quantities

Other algorithms which are used in MD to sample efficiently  $\mu$ :

- **Umbrella sampling and statistical reconstruction:** Histogram methods
- **Out of equilibrium methods:** fluctuation relations *à la* Jarzynski-Crooks
- **Modify the dynamics:** Metropolis Hastings algorithms with well-chosen proposals, non-reversible perturbations,...
- **Interacting replicas techniques:** Parallel tempering, Replica exchange dynamics, ...

# References

Monographs on numerical methods in MD:

- Allen, M., Tildesley, D.: Computer Simulation of Liquids. Oxford University Press (1989)
- Frenkel, D., Smit, B.: Understanding Molecular Simulation: From Algorithms to Applications. Oxford University Press (2001)
- Hairer, E., Lubich, C., Wanner, G.: Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations, Springer-Verlag (2006).
- Leimkuhler, B., Reich, S.: Simulating Hamiltonian Dynamics. Cambridge University Press (2005).
- Rapaport, D.: The Art of Molecular Dynamics Simulation. Cambridge University Press (2004)
- Tuckerman, M.: Statistical Mechanics: Theory and Molecular Simulation. Oxford University Press (2010).

