

cumsp — Graph cumulative spectral distribution

[Description](#)
[Options](#)
[Also see](#)

[Quick start](#)
[Remarks and examples](#)

[Menu](#)
[Methods and formulas](#)

[Syntax](#)
[Reference](#)

Description

`cumsp` plots the cumulative sample spectral-distribution function evaluated at the natural frequencies for a (dense) time series.

Quick start

Plot cumulative sample spectral-distribution function for `y` using `tsset` data

```
cumsp y
```

Same as above, and create `newv` containing the cumulative distribution estimates

```
cumsp y, generate(newv)
```

Menu

Statistics > Time series > Graphs > Cumulative spectral distribution

Syntax

```
cumsp varname [if] [in] [, options]
```

<i>options</i>	Description
Main	
<code><u>generate</u>(<i>newvar</i>)</code>	create <i>newvar</i> holding distribution values
Plot	
<code><i>cline_options</i></code>	affect rendition of the plotted points connected by lines
<code><i>marker_options</i></code>	change look of markers (color, size, etc.)
<code><i>marker_label_options</i></code>	add marker labels; change look or position
Add plots	
<code><u>addplot</u>(<i>plot</i>)</code>	add other plots to the generated graph
Y axis, X axis, Titles, Legend, Overall	
<code><i>twoway_options</i></code>	any options other than <code>by()</code> documented in [G-3] twoway_options

You must `tsset` your data before using `cumsp`; see [TS] [tsset](#). Also, the time series must be dense (nonmissing with no gaps in the time variable) in the sample specified.

varname may contain time-series operators; see [U] [11.4.4 Time-series varlists](#).

Options

Main

`generate(newvar)` specifies a new variable to contain the estimated cumulative spectral-distribution values.

Plot

`cline_options` affect the rendition of the plotted points connected by lines; see [G-3] [cline_options](#).

`marker_options` specify the look of markers. This look includes the marker symbol, the marker size, and its color and outline; see [G-3] [marker_options](#).

`marker_label_options` specify if and how the markers are to be labeled; see [G-3] [marker_label_options](#).

Add plots

`addplot(plot)` provides a way to add other plots to the generated graph; see [G-3] [addplot_option](#).

Y axis, X axis, Titles, Legend, Overall


`twoway_options` are any of the options documented in [G-3] [twoway_options](#), excluding `by()`. These include options for titling the graph (see [G-3] [title_options](#)) and for saving the graph to disk (see [G-3] [saving_option](#)).

Remarks and examples

► Example 1

Here we use the international airline passengers dataset (Box et al. 2016, Series G). This dataset has 144 observations on the monthly number of international airline passengers from 1949 through 1960. In the cumulative sample spectral distribution function for these data, we also request a vertical line at frequency $1/12$. Because the data are monthly, there will be a pronounced jump in the cumulative sample spectral-distribution plot at the $1/12$ value if there is an annual cycle in the data.

```
. use https://www.stata-press.com/data/r18/air2
(TIMESLAB: Airline passengers)
. cumsp air, xline(.083333333)
```



The cumulative sample spectral-distribution function clearly illustrates the annual cycle.

◀

Methods and formulas

A time series of interest is decomposed into a unique set of sinusoids of various frequencies and amplitudes.

A plot of the sinusoidal amplitudes versus the frequencies for the sinusoidal decomposition of a time series gives us the spectral density of the time series. If we calculate the sinusoidal amplitudes for a discrete set of “natural” frequencies ($1/n, 2/n, \dots, q/n$), we obtain the periodogram.

Let $x(1), \dots, x(n)$ be a time series, and let $\omega_k = (k-1)/n$ denote the natural frequencies for $k = 1, \dots, \lfloor n/2 \rfloor + 1$ where $\lfloor \cdot \rfloor$ indicates the greatest integer function. Define

$$C_k^2 = \frac{1}{n^2} \left| \sum_{t=1}^n x(t) e^{2\pi i(t-1)\omega_k} \right|^2$$

A plot of nC_k^2 versus ω_k is then called the periodogram.

The sample spectral density may then be defined as $\hat{f}(\omega_k) = nC_k^2$.

If we let $\hat{f}(\omega_1), \dots, \hat{f}(\omega_Q)$ be the sample spectral density function of the time series evaluated at the frequencies $\omega_j = (j - 1)/Q$ for $j = 1, \dots, Q$ and we let $q = \lfloor Q/2 \rfloor + 1$, then

$$\hat{F}(\omega_k) = \frac{\sum_{i=1}^k \hat{f}(\omega_j)}{\sum_{i=1}^q \hat{f}(\omega_j)}$$

is the sample spectral-distribution function of the time series.

Reference

Box, G. E. P., G. M. Jenkins, G. C. Reinsel, and G. M. Ljung. 2016. *Time Series Analysis: Forecasting and Control*. 5th ed. Hoboken, NJ: Wiley.

Also see

[TS] **corrgram** — Tabulate and graph autocorrelations

[TS] **pergram** — Periodogram

[TS] **tsset** — Declare data to be time-series data

Stata, Stata Press, and Mata are registered trademarks of StataCorp LLC. Stata and Stata Press are registered trademarks with the World Intellectual Property Organization of the United Nations. StataNow and NetCourseNow are trademarks of StataCorp LLC. Other brand and product names are registered trademarks or trademarks of their respective companies. Copyright © 1985–2023 StataCorp LLC, College Station, TX, USA. All rights reserved.



For suggested citations, see the FAQ on [citing Stata documentation](#).